K-mer based method for finding sequence type-specific PCR primers for *Listeria monocytogenes*

Reidar Andreson, Age Brauer, Lauris Kaplinski, Merle Külaots, Signe Saumaa, Ants Kurg, Maido Remm.

Institute of Molecular and Cell Biology, University of Tartu, Riia 23, Tartu 51010, Estonia

Abstract

Listeria monocytogenes is a deadly foodborne pathogen for which rapid identification of outbreak strains is critical. Yet current sequence typing methods such as multi locus sequence typing or whole-genome sequencing are time consuming and resource intensive. In this study, we developed a k-mer-based computational approach to pinpoint genomic regions unique to specific *L. monocytogenes* sequence types and designed polymerase chain reaction primers targeting each type's signature sequences. We screened thousands of genomes to identify robust type-specific markers, then validated the resulting primer sets on 51 isolates representing 17 sequence types in both singleplex and multiplex PCR assays. The primers successfully identified their target sequence types with 100% sensitivity and specificity across the tested isolates, even when assays were combined in multiplex format. This k-mer guided primer design strategy enables rapid and low-cost sequence typing of *L. monocytogenes* for routine screening and outbreak response. It bridges the gap between lengthy genomic sequencing workflows and the need for immediate field ready strain identification in food safety or other public health contexts.

Keywords: *Listeria monocytogenes*, k-mers, MLST, multiplex PCR, food safety

1 Introduction

Listeria monocytogenes is a facultatively anaerobic Gram-positive foodborne bacterium that is causing fatal disease listeriosis with high mortality rates up to 30% (Mead et al., 1999). Infections affect mostly elderly population or immunocompromised patients, but also pregnant women and newborns (Farber and Peterkin, 1991). *L. monocytogenes* can easily adapt to harsh environments with high salt concentrations (Burall et al., 2015), cold temperatures (Muchaamba et al., 2021), acidity (Cheng et al., 2015) and chemical treatments (Duze et al., 2021). The bacterium uses many genes (roughly seven percent of its genome) for adapting to specific environmental conditions (Glaser et al., 2001). It is also shown that *L. monocytogenes* is naturally resistant to multiple types of antibiotics and therefore rapid finding of the source of outbreaks and identification of pathogenic strains quickly is critical (Olaimat et al., 2018).

Although various alternative methods ranging from serotyping to banding patterns (e.g., RFLPs, AFLPs) have been implemented over the years, currently the most popular method

for genotyping major clonal groups and subtypes of *L. monocytogenes* is through the multi
locus sequence typing (MLST) scheme (Ragon et al., 2008; Nyarko and Donnelly, 2015). The
MLST proof-of-concept method is based on identification of DNA sequences of a small
number of hypervariable conserved housekeeping genes and comparing their mutation
profiles (Maiden et al., 1998; Salcedo et al., 2003). Ongoing development of next generation
sequencing technologies has made it possible to identify full genomes of large number
bacterial isolates during foodborne disease outbreaks. Furthermore, gene-by-gene allelic
profiling of core genome genes or the core genome MLST (cgMLST) has been developed to
become the main diagnostic tool for strain typing (Maiden et al., 2013; Ruppitsch et al.,
2015). It is even shown that the discriminatory power of cgMLST clearly exceeds pulsed-
field gel electrophoresis (PFGE) typing, which is often considered as the gold standard for
subtyping of *L. monocytogenes* (Lüth et al., 2018).
56
However, there are some major challenges of cgMLST typing due to lack of standardized
nomenclature to exchange data between agencies within and between different countries
(Moura et al., 2016). During foodborne disease outbreaks it is crucial to timely track the
source of bacterial contamination (Jackson et al., 2016). Sending samples into public health
agencies to run whole genome sequencing may take weeks to get results. Also, the cost of
sequencing and storing data is another criterion that needs to be considered when analyzing
large quantities of samples on a regular basis.
64
In contrast, k-mer based classifiers such as Kraken2 (Wood et al., 2019), StrainScan (Liao et
al., 2023) or StrainSeeker (Roosaare et al., 2017) are optimized for taxonomic/strain
identification from read k-mers and not for detecting MLST-specific regions from genome.
Kraken2 uses exact k-mer matches and lowest-common-ancestor assignment to label reads
against a reference database that is fast and sensitive for detection, but it does not output
standardized allelic profiles. StrainSeeker similarly maps sample-specific k-mer sets onto a
user supplied guide tree to report the best matching strain/clade rather than alleles. StrainScan
extends this idea with a tree-based k-mer index to resolve known strains in metagenomic or
isolate data. However, it again relies on database encoded strain signatures rather than calling
alleles at defined housekeeping loci. Moreover, the performance of k-mer classifiers is tightly
coupled to the completeness and composition of their reference databases and can degrade
when the sample's strain diverges from available references. These conditions can complicate
their use for generating portable MLST genotype designations.
78
Recent tools illustrate adjacent ideas, but do not directly solve MLST type-specific primer
design. PUPpy (Ghezzi et al., 2024) automates discovery of phylogenetically unique targets
within a defined community and then designs microbe- or group-specific primers for strain
level detection and absolute quantification. It operates on coding sequences provided by the
user and reports specific assays rather than alleles or specific markers tied to MLST loci.
Another tool called varVAMP (Fuchs et al., 2025) focuses on highly variable viral genomes,
using an MSA-derived consensus and a k-mer-based enumeration around Primer3 (Kõressaar
et al., 2018) to create degenerate and tiled primer schemes for full-genome sequencing and
qPCR. However, it is aimed at pan-specific coverage across variants, not discrete type-
specific diagnostics. SURE-Pipe[1] focuses on shared/unique region extraction by orchestrating
BLAST/BEDTools (Quinlan and Hall, 2010) within a Nextflow workflow to report regions
unique to a target versus non-targets. It's general purpose is for genome comparison and does
not natively couple uniqueness calls to type-specific nomenclature or downstream multiplex
primer selection.
93

94 In this work, we introduce a k-mer based approach to identify MLST type-specific regions
95 from bacterial whole genomes and create unique PCR primers for each type. We have used
96 51 *L. monocytogenes* samples covering 17 different sequence types to experimentally test the
97 success rate of given primers in singleplex and multiplex form.
98

2 Materials and methods

2.1 Bacterial reads and isolates

For identifying type-specific regions of *L. monocytogenes* we needed genomic reads from the
ENA database (O'Cathail et al., 2025). In total, 28545 isolate samples with Illumina paired-
end reads were downloaded. These samples included raw sequencing data from various
projects around the world. For experimental validation, we had DNA samples from 51
bacterial isolates of 17 different MLST types (designated ST) (ST5, ST7, ST8, ST9, ST29,
ST37, ST87, ST101, ST121, ST155, ST173, ST177, ST425, ST451, ST551, ST580, ST1247).
For each type, there was DNA of three isolates, a total of 51 samples (Table 1). DNA samples
of isolates were obtained from The National Centre for Laboratory Research and Risk
Assessment (LABRIS, Estonia), where the MLST types of these samples had been previously
determined by whole genome sequencing data analysis. All these *L. monocytogenes* samples
were collected from Estonia. Among them were ST1247 isolates, which received widespread
resonance in local and foreign press (Mäesaar et al., 2021).
113

2.2 Genome assembly and determination of MLST types

To determine the MLST types of downloaded public strains, we used the computational
workflow described below. First, the reads were cleaned of low quality nucleotides, and
adapter sequences were removed. For this, the *fastp* program (Chen et al., 2018) was used
with the following parameters: minimum read length 50 nucleotides (*--length_required 50*),
average quality 30 (*-cut_mean_quality 30*), *--cut_tail, --cut_front*, and *--*
detect_adapter_for_pe. In the next step, the cleaned reads were assembled with *SPAdes*
(Prjibelski et al., 2020) into genomic contig sequences using the *--isolate* parameter. Samples
with the longest contig of <50,000 nucleotides and an average sequencing depth of less than
20 were filtered out from the results. The final step was the determination of the MLST type
by the MLST[2] program, for which the default parameters and the PubMLST (Jolley et al.,
2018) dataset containing >2300 different MLST types of *L. monocytogenes* was used.
126

2.3 Phylogenetic tree of MLST types

A prerequisite for the construction of a tree is the creation of distance matrix based on
sequence comparisons. For this, we used the *dashing* program (Baker and Langmead, 2019),
that created a distance matrix based on sequencing reads. When running the program, the
following parameters were changed: the length of the k-mers 32 nucleotides (*-k 32*), *--sketch-*
size 20, *--use-range-minhash*, and *--full-mash-dist*. From the created distance matrix, a
phylogenetic tree in newick format was compiled using the *quicktree* (Howe et al., 2002)
program (with default parameters). The results were visualized with the MEGA11 program
(Tamura et al., 2021).
136

2.4 Search for type-specific k-mers

As a first step, it was necessary to create a list of k-mers for each sample of ENA reads. To do
this, we used the GenomeTester4 package (Kaplinski et al., 2015) program *glistmaker*, for

which we set the length of the k-mer to 32 nucleotides (--*wordlength 32*). Previous work by
our group has shown that 32 could be optimal for finding more specific k-mers (Raime and
Remm, 2018).

To create a common list of non-targets, the GenomeTester4 package program *glistcompare*
was used with parameters --*union* and a frequency limit value of 5 (--*cutoff 5*) for k-mers. The
latter is necessary to exclude very rare (<5 times) k-mers, which are most likely sequencing
errors.

Similarly to the above, using *glistcompare* a list of k-mers was created with the parameters --
intersection and the frequency limit value of k-mers 5 (--*cutoff 5*). Those k-mers that were not
represented in all samples were removed. This was necessary to eliminate k-mers, which are
present only in a particular sample and are therefore not characteristic of all members of a
given type.

Next, we compared the common list of each type with the combined list of non-targets to
exclude k-mers that are also present in types other than the target. To do this, we used the
glistcompare with the parameter --*difference*.

2.5 Design of type-specific PCR primers

Type-specific k-mers were mapped to their genomic coordinates using NCBI BLAST
(Altschul, 1997). The package program *blastn* was used with an alignment percentage of 100
(-*perc_identity 100*) and a query sequence coverage percentage of 100 (-*qcov_hsp_perc*
100). The resulting HSP coordinates for each k-mer were recorded for downstream
processing.

K-mers whose mapped coordinates overlapped or were directly adjacent on the same
contig/strand were merged into contiguous genomic intervals as type-specific regions.
Candidate intervals were restricted to sizes compatible with PCR using a product length
window of 60–1000 nt. Intervals shorter than 60 nt (e.g. single isolated k-mer hits) and >1000
nt were excluded from primer design.

For every retained region, PCR primers were designed with Primer3 using following settings:
primer length: min 18, opt 20, max 22 nt; product size range: 60–1000 nt; maximum allowed
difference between primer pair melting temperatures (Tm): 3 $^\circ$C. All other Primer3
parameters were left at their defaults.

To assess potential cross-reactivity, all candidate primers were aligned to the non-target
genome set using *blastn*. In this step, we permitted sequence mismatches, but required full-
length primer coverage: -*qcov_hsp_perc 100* (no -*perc_identity* constraint). All HSPs and
their genomic coordinates were retained for pairwise evaluation. A pair was retained if: (a) at
least one primer had no non-target hit, or (b) both primers hit but not in opposing orientations
1–1000 nt apart (no plausible amplicon), or (c) both primers hit at 1–1000 nt spacing but at
least one had a mismatch within the 3′ terminal 6 nt. Only pairs meeting more than one of
these conditions was kept.

2.6 DNA extraction

The genomic DNA of *L. monocytogenes* isolates were purified with GeneJET Genomic DNA
purification Kit (Thermo Scientific) by using "Gram-positive Bacteria genomic DNA

Purification Protocol" according to manufacturer's protocol, except for the first step. As
isolates of *L. monocytogenes* were obtained growing on blood agar, the bacterial cells were
first collected from agar by using an inoculation loop and after that were suspended in Gram-
positive bacteria lysis buffer. All procedures were done in strict rules and conditions of
Biosafety Lab 2 in Institute of Molecular and Cell Biology, University of Tartu.
194

2.7 PCR experiments

To check the specific PCR primers designed, we conducted laboratory experiments. We had
the possibility to use DNA samples from 17 different types of isolates and each type in three
replicates. In total, it had formed 51 singleplex reactions in which only one pair of primers is
used at a time. Additionally, the success rate of universal primers was controlled under the
same reaction conditions. Ideally, only a single product (band on the gel) should arise with
primers of the corresponding type and other types should not show any yield in given
reaction. Universal primer pair should ideally work with all samples.
203

204 **Singleplex solution and program:** 10 μl reaction volume:10x buffer (Solis Biodyne) 1μl; 25
205 mM MgCl$_2$ (Solis Biodyne) 0,6 μl; 2.5 mM dNTP mix (Solis Biodyne) 1.2 μl; Hot FirePol
206 (Solis Biodyne) (5U/μl) 0.1 μl; MQ water 5,1 μl; Primers (10nM (5 F+5 R))1 μl; DNA
207 (5ng/μl) 1 μl. PCR program: 95°C 15 min; 95°C 30 s, 60°C 30 s - 30x,72°C 40 s; 72°C 10
208 min.
209

In multiplex PCR experiments, the desired number of primer pairs are mixed in one reaction,
which allows saving the costs and time of the reagents. As a result, the gel image should
develop several bands of the correct length instead of one correct band. Since in some cases
bands of several products are expected to be performed on the track of a single gel because of
a multiplex test, it is extremely important that the lengths of the products are different enough
to achieve visual resolution. Therefore, we have checked with the MultiPLX (Kaplinski et al.,
2005) program (parameters -cutoff1 -3.0 -cutoff2 -10 -cutoff3 -13.0) that primer pairs do not
create strong secondary structures among themselves, which can negatively affect the success
of PCR, and are divided into groups with separable product sizes.
219

220 **Multiplex solution and program:** 10 μl reaction volume: 10x buffer (Solis Biodyne) 1μl; 25
221 mM MgCl$_2$ (Solis Biodyne) 0,6 μl; 2.5 mM dNTP mix (Solis Biodyne) 1.2 μl; Hot FirePol
222 (Solis Biodyne) (5U/μl) 0,1 μl; MQ water 4.5 μl; Primers (pairs mixed together) (10nM)
223 1.5μl; Primer Univ_6 (10nM) 0,1 μl; DNA (5ng/μl) 1 μl. PCR program: 95°C 15 min; 95°C
224 30 s, 58°C 30 s - 31x, 72°C 40 s; 72°C 10 min.
225

3 Results

3.1 Identifying type-specific k-mers from genomes

In order To find unique type-specific k-mers for *L. monocytogenes* samples, we downloaded
28,545 public ENA Illumina paired-end read sets. These were quality filtered and adapter
trimmed, resulting high quality reads suitable for the assembly. Genome assemblies were
generated with SPAdes and screened. Datasets with insufficient contiguity or depth were
excluded from downstream analysis. For the retained samples, sequence types were
successfully assigned using the pubMLST scheme for *L. monocytogenes*. The MLST type
remained undetermined for 980 out of 28,545 and 1252 different types were determined in
total. Samples for which the species were determined to be something else than *L.*

monocytogenes were removed from the dataset. The final number of samples remaining was
27,206 (Supplementary Table S1).
238
Next, we decided to construct a phylogenetic tree of isolates based on reads to see if the
isolates with the same MLST type cluster together in a clade or are more broadly distributed.
In the latter case, it may be more difficult to find type-specific regions as genomes share
common regions with many different types. Since with all 27,206 samples this data analysis
would be too time consuming and difficult to visualize later, we decided to take ten random
samples for each type (out of 1252), which occurred at least 10 times in the dataset. In
addition, we checked that all 17 types of our experimental dataset were included. In total,
1168 samples of various types were used to build a phylogenetic tree.
247
The phylogenetic tree shows the distribution of sequence types along the branches, with the
17 sequence types of interest highlighted in different colors (Supplementary Figure S1).
Fifteen of these 17 types cluster relatively compactly. The exceptions are ST37 and ST173,
which are distributed across several major branches. ST9 and ST580 form a distinct group,
and a similar pattern is observed for ST8, ST551 and ST1247. In this case, such groupings
are consistent with their presence in the corresponding clonal complexes CC9 and CC8. A
clonal complex is defined as a set of sequence types grouped according to the inferred
evolutionary descent of isolates under the assumption of a shared common ancestor. The
name of the complex typically reflects the putative ancestral type (e.g. ST8 for CC8). Overall,
this grouping pattern suggests that it should be feasible to identify type-specific genomic
regions and to design corresponding PCR primers.
259
260   From 27,206 ENA read sets, we enumerated 32-nt k-mers for each sample and used these to
261   assemble two complementary resources: a comprehensive non-target background and type-
262   specific core repertoires. Figure 1 shows the structure of the workflow, by which it was
263   possible to create lists of k-mers of samples from targets and non-targets. One of the types
264   was chosen as the target and the rest remained among the non-targets.
265
Aggregating all non-target samples produced a unified background of k-mers while filtering
out very rare instances (occurring <5 times), thereby minimizing the influence of sequencing
errors. Secondly, for each of the 17 target types, we derived a stringent "core" set by retaining
only those k-mers observed (≥5 instances) in every sample of that type, which removed k-
mers confined to individual specimens and not representative of the given sequence type.
Finally, subtracting the non-target background from each type's core yielded target-specific k-
mer panels. These k-mers are absent from all non-target samples yet ubiquitous across the
corresponding target type. Given workflow produced high confidence type defining marker
sets suitable for downstream detection and classification.
275
276   3.2 Merging k-mers to type-specific regions and PCR primer design

277   To find type-specific genome regions for which PCR primers could be designed, specific k-
278   mers were mapped on genome sequence and the corresponding coordinates of hits were
279   recorded. The PCR primer design workflow is graphically described in Figure 2.
280
281   The k-mers with genomic positions in the overlay were merged into longer regions based on
282   coordinates. Considering the recommended minimum (roughly two primer lengths) and
283   maximum (<1000 nucleotide) product lengths for the PCR methodology, we set the range to

284   60-1000. Shorter regions or single k-mers were removed from the selection. As a result, the
285   median value of specific regions across all types was 551.
286
For each specific region, we tried to design PCR primers with Primer3. Allowing for near
identical matches, we realigned all type-specific primers against non-target genomes using
full-length (100%) coverage. This secondary screen flagged a subset of primers with potential
cross reactivity due to single nucleotide differences, while the remainder showed no full-
length hits in non-targets. Primers with such near perfect alignments were marked for
exclusion or further scrutiny, and the non-flagged set was carried forward as the high
specificity candidate pool.
294
The result files were analysed and the final primer pairs had to meet the following conditions.
The pair was named suitable, (a) if at least one primer mate did not give alignments to any
non-target genome of the type; (b) if both primers in the pair did provide alignment from a
non-target type's genome, but not a distance between 1-1000 nucleotides; (c) if both primers
in the pair did provide alignment from a non-target type's genome and a distance between 1
and 1000 nucleotides, but the difference (mutation) lies within the last six nucleotides at the
3' end of at least one primer.
302
When we used the full list of samples from ENA database (27,206), problems occurred with
some of the target types as very few specific k-mers remained. This may be due to
contaminated data, where most of the reads in one sample are indeed from cells of one MLST
type, but to a lesser extent from another type. Since our methodology detects all k-mers
present in the reads, samples with contaminated reads can end up as false positives among
non-targets or targets. In addition, there is always the possibility that sequence type
contamination will occur in the process of sequencing, when several samples of *L.*
monocytogenes are analysed in the same laboratory.
311
Therefore, we decided to reduce the amount of data and selected up to 10 random samples
(with a total number of 155) from each target type from all ENA data and reran the entire
previous workflow with them. This approach made sense, because we had to use only the
DNA of these 17 types for experimental control of type-specificity. For types ST173, ST580
and ST1247, we had to use a smaller number of samples than ten, since there were simply no
more data in the database. As a result of the analysis, we managed to find type-specific k-
mers with a median value of 47000 for 17 ST-s and two CC-s (Table 2). The list of primer
sequences and their properties are shown in Supplementary Table S2.
320

3.3 Singleplex PCR experiments

The results of the singleplex PCR tests can be seen in Figure 3. A total of 83 different primer
pairs were tested on 51 different DNA samples. Three pairs were universal primers (Univ_3,
Univ_6, Univ_7), 14 pairs were clonal-complex-specific primer pairs, and the remaining 66
were type-specific. The green boxes represent the correct result (a decent band on the gel)
that was supposed to come out (true positive). Yellow colours indicate a false positive result,
and white indicates the absence of the expected reaction. For types, there must be 3 green
boxes for each pair of primers, for universal ones all must be green, and for clonal complex-
specific primers, the types belonging to the corresponding complex must be green (9 DNA
for CC8, 6 for CC9). For each type and clonal complex, a smaller number of primers were
first tested and if pairs with 100% specificity and sensitivity were not detected, new primers
were picked from the list of candidate primers for testing. A total of 4233 PCR experiments

were carried out, of which 2813 (~67%) were successful, i.e. gave the expected result (true
positive).
335
Closer look at the specificity and sensitivity of primers (Supplementary Table S3) shows that
corresponding median values for all types are 89% and 100%. Specificity indicates the ability
of the primer pair to distinguish a particular type from the rest, that is the primer pair must
not give a PCR product on any other type. Sensitivity indicates how well the primer pair
identified the correct type. Only five pairs of primers did not develop the correct positive
PCR product with the DNA of all three corresponding types. In 55% of types, specificity was
lower than 100%, for which false positive products with DNA of other types were formed.
Based on 10 random ENA samples, it was not possible to find a primer pair with 100%
specificity for the ST37 type. For CC8, the best primer pair had 98% specificity and 100%
sensitivity (CC8_7). In all other cases we found at least one perfectly working PCR primer
pair. From the tested universal primers, the one called Univ_6 had 100% experimental
success rate for both specificity and sensitivity.
348
We also wanted to analyse genomic regions for all designed PCR primers to see, if there are
some shared properties. For that we used *blastx* with *L. monocytogenes* reference strain EGD-
e genome and protein sequences. Only 11% of the blast hits corresponded to coding regions
with known or annotated functions, whereas about 89% were in proteins annotated as
hypothetical, putative, or with no characterized function. The hits clustering in surface
proteins and regulatory elements may point to repetitive DNA elements in those proteins
rather than in unique, highly conserved housekeeping genes. Additionally, a search of the
primer alignment coordinates and identifiers revealed no hits within or near MLST (*abcZ*,
bglA, *cat*, *dapE*, *dat*, *ldh, lhkA*) loci.
358

3.4 5-plex PCR experiments

As the first option for multiplexing, we tried to match a maximum of 5-plex pairs (4-5 pairs
of primers in one reaction). To do this, based on the results of previous singleplex
experiments, we selected primer pairs whose product lengths would be sufficiently
distinguishable from each other (Figure 4A). As a result, we formed three different groups of
primers, with five pairs of primers in two and four in one. For the reaction, mixtures of 4 or 5
pairs of primers were mixed and experiments were carried out on the DNA of all 51 isolates.
366
The results of the 5-plex multiplex PCR experiment can be seen in Figure 4B. The first group
(Group I) gives correct results with six types (CC9 gives with two different types), the second
group (Group II) gives with seven types (CC8 with three different ones), and the third group
(Group III) with four types. The sensitivity of these groups is 100%, all prescribed types are
correctly recognized. In the case of Group II, there was a problem with single isolate (yellow
box), which is why the specificity with this group is 98%. Such a result was expected, since it
was seen from the singleplex experiments that CC8_7 primer pair gives a nonspecific
reaction with the 7th isolate. Summing up, it can be said that the 5-plex experiments were as
successful as the singleplex experiments.
376

3.5 15-plex PCR experiment

Based on the success of the previous 5-plex experiments, we decided to take it a step further
and try the multiplex PCR reaction with all 14 pairs of primers. Accordingly, we had to
slightly change the list of suitable primers, where we replaced the CC8_7 primer pair with
CC8_6 variant, since the length of a given product (131 nucleotides) was more easily

distinguishable from ST1247 (belongs to the CC8 clonal complex). We mixed 14 primer pairs
with the best working universal primer pair (Univ_6) and checked whether PCR products are
formed for all 51 isolates and whether they are also of the correct length (Supplementary
Table S4).
386
The result of the 15-plex multiplex PCR experiment can be seen in the gel image in Figure 5.
In the central part of the gel is located a line of straight bands with a universal primer. In the
case of CC8, there are three bands on tracks 49-51, where one is a product of a universal
primer pair with a length of 430 nt, above it a product of the ST1247 type with a length of
409 nt and underneath a product of the CC8 clonal complex with a length of 131. On all other
tracks a band of both a universal product and a type-specific product can be seen.
393

4 Discussion

This study presents and experimentally validates a k-mer driven workflow for discovering
MLST type-specific genomic regions in *L. monocytogenes* and designing PCR assays that
function in both singleplex and multiplex formats. Using 27,206 public WGS datasets to
define type-specific k-mers and 51 locally sourced isolates across 17 MLST sequence types
for laboratory validation, we achieved robust type discrimination with median sensitivity of
100% and median specificity of 89% for candidate primer pairs, with several perfect
performers per type or clonal complex. These results demonstrate that allele independent k-
mer set operations can yield portable, interpretable, and laboratory ready screening markers
even when closely related types share substantial genome content.
404
Our algorithm is distinctive in three ways. First, it uses strict set logic on k-mer inventories
like intersection within type and union of non-targets to isolate regions that are both
ubiquitous in the type and absent from others and these properties translate directly into
reliable PCR targets. Second, it places candidate k-mers onto assemblies with exact match
constraints to form contiguous primer sized intervals avoiding fragile point mutation
signatures. Third, it demonstrates multiplex readiness by incorporating primer
thermodynamics and cross interaction checks from the outset. The result is a pipeline that
transforms MLST labels into deployable type-specific assays without requiring whole-
genome alignment or special SNP modelling.
414
For routine screening in food production environments and public health triage, turnaround
time and per-sample cost are decisive (Tran et al., 2023). Centralized WGS workflows
typically require ~5–10 days from sample to report when library prep, batching, sequencing,
and analysis are considered (Beukers et al., 2021). This can be often longer in public health
institutions as their pipelines are optimized for cost over speed. In contrast, PCR-based assays
routinely deliver results within 12–24 hours, can be automated at scale, and are far cheaper
per sample. These enable high-throughput and rapid decision making in line clearance or
targeted sanitation, for example. Thus, the development and usage of multiplex PCR panels
offers a pragmatic balance between speed, cost, and epidemiologic resolution.
424
Our approach depends on the representativeness and quality of public data used to identify
type-specific regions. While our curation mitigated some contamination and imbalance,
continued updates to the non-target set and routine rescreening of candidate regions will be
necessary as new diversity is deposited. PCR experiment performance can also be affected by
DNA source (food vs clinical), DNA quality and thermocycler conditions. The additional *in*
silico checks with an updated screenings against expanding *L. monocytogenes* genomes and

optimizations of wet-lab protocol can further reduce some sporadic false positive results.
Although we have validated 5-plex and 15-plex formats with high success, higher-plex panels
may require some pipeline or protocol redesign to preserve clear gel separation or transition
to qPCR readouts.
435

5 Conclusion

We have linked *L. monocytogenes* MLST labels to k-mer defined type-specific regions and
thus created a practical route from population nomenclature to rapid type identification. The
singleplex and multiplex results across 17 MLST types indicate that this strategy is viable for
front line industrial screening and outbreak triage. Although periodic off-target rescreening is
advisable as databases grow, given approach yields portable, interpretable, and multiplex
PCR assays that performed well across diverse list of MLST types. Given approach can be
easily extended to additional *Listeria* lineages and other bacterial taxa.
445

Data availability statement

448 The ENA identifiers used in this study can be found in Supplementary Table S1. All custom
449 scripts used in this work are available at https://github.com/bioinfo-ut/MLST-
450 specific_genomic_regions.
451

Author contributions

RA: Investigation, Software, Methodology, Validation, Visualization, Writing – original draft.
AB: Conceptualization, Methodology, Writing – review & editing. LK: Conceptualization,
Investigation, Writing – review & editing. MK: Methodology, Visualization, Writing – review
& editing. SS: Methodology, Validation, Writing – review & editing. AK: Conceptualization,
Investigation, Writing – review & editing. MR: Conceptualization, Investigation, Funding
acquisition, Project administration, Supervision, Writing – review & editing.
460

Funding

462
463 This work was funded by Estonian Research Council through the project EAG129
464 "Diagnostic test for fast identification of Listeria monocytogenes sequence types" and co-
465 funded by European Union and Ministry of Education and Research via project TemTA-35
466 "Surveillance of food safety, origin and composition using DNA sequencing".
467

Acknowledgements

469
470 We thank Toomas Kramarenko from LABRIS for providing the isolates of *L.monocytogenes*.
471 Some parts of the data analysis was performed using the facilities of the High-Performance
472 Computing Center at the University of Tartu (https://doi.org/10.23673/PH6N-0144).
473

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Footnotes

[1]https://github.com/BPaul-bioinfoLAB/SURE-Pipe
[2]https://github.com/tseemann/mlst

References

Altschul, S. (1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res*. 25, 3389–3402. doi: 10.1093/nar/25.17.3389

Baker, D. N., and Langmead, B. (2019). Dashing: fast and accurate genomic distances with HyperLogLog. *Genome Biol*. 20, 265. doi: 10.1186/s13059-019-1875-0

Beukers, A. G., Jenkins, F., and Van Hal, S. J. (2021). Centralised or Localised Pathogen Whole Genome Sequencing: Lessons Learnt From Implementation in a Clinical Diagnostic Laboratory. *Front. Cell. Infect. Microbiol*. 11, 636290. doi: 10.3389/fcimb.2021.636290

Burall, L. S., Simpson, A. C., Chou, L., Laksanalamai, P., and Datta, A. R. (2015). A novel gene, lstC, of Listeria monocytogenes is implicated in high salt tolerance. *Food Microbiol*. 48, 72–82. doi: 10.1016/j.fm.2014.12.008

Chen, S., Zhou, Y., Chen, Y., and Gu, J. (2018). fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* 34, i884–i890. doi: 10.1093/bioinformatics/bty560

Cheng, C., Yang, Y., Dong, Z., Wang, X., Fang, C., Yang, M., et al. (2015). Listeria monocytogenes varies among strains to maintain intracellular pH homeostasis under stresses by different acids as analyzed by a high-throughput microplate-based fluorometry. *Front. Microbiol*. 6. doi: 10.3389/fmicb.2015.00015

Duze, S. T., Marimani, M., and Patel, M. (2021). Tolerance of Listeria monocytogenes to biocides used in food processing environments. *Food Microbiol*. 97, 103758. doi: 10.1016/j.fm.2021.103758

Farber, J. M., and Peterkin, P. I. (1991). Listeria monocytogenes, a food-borne pathogen. *Microbiol. Rev*. 55, 476–511. doi: 10.1128/mr.55.3.476-511.1991

Fuchs, J., Kleine, J., Schemmerer, M., Kreibich, J., Maier, W., Battur, N., et al. (2025). varVAMP: degenerate primer design for tiled full genome sequencing and qPCR. *Nat. Commun*. 16, 5067. doi: 10.1038/s41467-025-60175-9

Ghezzi, H., Fan, Y. M., Ng, K. M., Burckhardt, J. C., Pepin, D. M., Lin, X., et al. (2024).
PUPpy: a primer design pipeline for substrain-level microbial detection and absolute
quantification. *mSphere* 9, e00360-24. doi: 10.1128/msphere.00360-24

Glaser, P., Frangeul, L., Buchrieser, C., Rusniok, C., Amend, A., Baquero, F., et al. (2001).
Comparative Genomics of *Listeria* Species. *Science* 294, 849–852. doi:
10.1126/science.1063447

Howe, K., Bateman, A., and Durbin, R. (2002). QuickTree: building huge Neighbour-Joining
trees ofprotein sequences. *Bioinformatics* 18, 1546–1547. doi:
10.1093/bioinformatics/18.11.1546

Jackson, B. R., Tarr, C., Strain, E., Jackson, K. A., Conrad, A., Carleton, H., et al. (2016).
Implementation of Nationwide Real-time Whole-genome Sequencing to Enhance
Listeriosis Outbreak Detection and Investigation. *Clin. Infect. Dis.* 63, 380–386. doi:
10.1093/cid/ciw242

Jolley, K. A., Bray, J. E., and Maiden, M. C. J. (2018). Open-access bacterial population
genomics: BIGSdb software, the PubMLST.org website and their applications.
Wellcome Open Res. 3, 124. doi: 10.12688/wellcomeopenres.14826.1

Kaplinski, L., Andreson, R., Puurand, T., and Remm, M. (2005). MultiPLX: automatic
grouping and evaluation of PCR primers. *Bioinformatics* 21, 1701–1702. doi:
10.1093/bioinformatics/bti219

Kaplinski, L., Lepamets, M., and Remm, M. (2015). GenomeTester4: a toolkit for performing
basic set operations - union, intersection and complement on k-mer lists. *GigaScience*
4, 58. doi: 10.1186/s13742-015-0097-y

Kõressaar, T., Lepamets, M., Kaplinski, L., Raime, K., Andreson, R., and Remm, M. (2018).
Primer3_masker: integrating masking of template sequence with primer design
software. *Bioinformatics* 34, 1937–1938. doi: 10.1093/bioinformatics/bty036

Liao, H., Ji, Y., and Sun, Y. (2023). High-resolution strain-level microbiome composition
analysis from short reads. *Microbiome* 11, 183. doi: 10.1186/s40168-023-01615-w

Lüth, S., Kleta, S., and Al Dahouk, S. (2018). Whole genome sequencing as a typing tool for
foodborne pathogens like Listeria monocytogenes – The way towards global
harmonisation and data exchange. *Trends Food Sci. Technol.* 73, 67–75. doi:
10.1016/j.tifs.2018.01.008

Mäesaar, M., Mamede, R., Elias, T., and Roasto, M. (2021). Retrospective Use of Whole-
Genome Sequencing Expands the Multicountry Outbreak Cluster of Listeria
monocytogenes ST1247. *Int. J. Genomics* 2021, 1–5. doi: 10.1155/2021/6636138

Maiden, M. C. J., Bygraves, J. A., Feil, E., Morelli, G., Russell, J. E., Urwin, R., et al. (1998).
Multilocus sequence typing: A portable approach to the identification of clones within
populations of pathogenic microorganisms. *Proc. Natl. Acad. Sci.* 95, 3140–3145. doi:
10.1073/pnas.95.6.3140

Maiden, M. C. J., Van Rensburg, M. J. J., Bray, J. E., Earle, S. G., Ford, S. A., Jolley, K. A., et
al. (2013). MLST revisited: the gene-by-gene approach to bacterial genomics. *Nat.*
Rev. Microbiol. 11, 728–736. doi: 10.1038/nrmicro3093

Mead, P. S., Slutsker, L., Dietz, V., McCaig, L. F., Bresee, J. S., Shapiro, C., et al. (1999).
Food-Related Illness and Death in the United States. *Emerg. Infect. Dis.* 5, 607–625.
doi: 10.3201/eid0505.990502

Moura, A., Criscuolo, A., Pouseele, H., Maury, M. M., Leclercq, A., Tarr, C., et al. (2016).
Whole genome-based population biology and epidemiological surveillance of Listeria
monocytogenes. *Nat. Microbiol.* 2, 16185. doi: 10.1038/nmicrobiol.2016.185

Muchaamba, F., Stephan, R., and Tasara, T. (2021). Listeria monocytogenes Cold Shock
Proteins: Small Proteins with A Huge Impact. *Microorganisms* 9, 1061. doi:
10.3390/microorganisms9051061

Nyarko, E. B., and Donnelly, C. W. (2015). *Listeria monocytogenes*□: Strain Heterogeneity,
Methods, and Challenges of Subtyping. *J. Food Sci.* 80. doi: 10.1111/1750-
3841.13133

O'Cathail, C., Ahamed, A., Burgin, J., Cummins, C., Devaraj, R., Gueye, K., et al. (2025).
The European Nucleotide Archive in 2024. *Nucleic Acids Res.* 53, D49–D55. doi:
10.1093/nar/gkae975

Olaimat, A. N., Al□Holy, M. A., Shahbaz, H. M., Al□Nabulsi, A. A., Abu Ghoush, M. H.,
Osaili, T. M., et al. (2018). Emergence of Antibiotic Resistance in *Listeria*
monocytogenes Isolated from Food Products: A Comprehensive Review. *Compr. Rev.*
Food Sci. Food Saf. 17, 1277–1292. doi: 10.1111/1541-4337.12387

Prjibelski, A., Antipov, D., Meleshko, D., Lapidus, A., and Korobeynikov, A. (2020). Using
SPAdes De Novo Assembler. *Curr. Protoc. Bioinforma.* 70, e102. doi:
10.1002/cpbi.102

Quinlan, A. R., and Hall, I. M. (2010). BEDTools: a flexible suite of utilities for comparing
genomic features. *Bioinformatics* 26, 841–842. doi: 10.1093/bioinformatics/btq033

Ragon, M., Wirth, T., Hollandt, F., Lavenir, R., Lecuit, M., Le Monnier, A., et al. (2008). A
New Perspective on Listeria monocytogenes Evolution. *PLoS Pathog.* 4, e1000146.
doi: 10.1371/journal.ppat.1000146

Raime, K., and Remm, M. (2018). Method for the Identification of Taxon-Specific k-mers
from Chloroplast Genome: A Case Study on Tomato Plant (Solanum lycopersicum).
Front. Plant Sci. 9, 6. doi: 10.3389/fpls.2018.00006

Roosaare, M., Vaher, M., Kaplinski, L., Möls, M., Andreson, R., Lepamets, M., et al. (2017).
StrainSeeker: fast identification of bacterial strains from raw sequencing reads using
user-provided guide trees. *PeerJ* 5, e3353. doi: 10.7717/peerj.3353

Ruppitsch, W., Pietzka, A., Prior, K., Bletz, S., Fernandez, H. L., Allerberger, F., et al. (2015).
Defining and Evaluating a Core Genome Multilocus Sequence Typing Scheme for
Whole-Genome Sequence-Based Typing of Listeria monocytogenes. *J. Clin.*
Microbiol. 53, 2869–2876. doi: 10.1128/JCM.01193-15

Salcedo, C., Arreaza, L., Alcalá, B., De La Fuente, L., and Vázquez, J. A. (2003).
Development of a Multilocus Sequence Typing Method for Analysis of *Listeria*
monocytogenes Clones. *J. Clin. Microbiol.* 41, 757–762. doi: 10.1128/JCM.41.2.757-
762.2003

Tamura, K., Stecher, G., and Kumar, S. (2021). MEGA11: Molecular Evolutionary Genetics
Analysis Version 11. *Mol. Biol. Evol.* 38, 3022–3027. doi: 10.1093/molbev/msab120

Tran, M., Smurthwaite, K. S., Nghiem, S., Cribb, D. M., Zahedi, A., Ferdinand, A. D., et al.
(2023). Economic evaluations of whole-genome sequencing for pathogen
identification in public health surveillance and health-care-associated infections: a
systematic review. *Lancet Microbe* 4, e953–e962. doi: 10.1016/S2666-
5247(23)00180-5

Wood, D. E., Lu, J., and Langmead, B. (2019). Improved metagenomic analysis with Kraken
2. *Genome Biol.* 20, 257. doi: 10.1186/s13059-019-1891-0

Figure legends

Figure 1. The flowchart of identifying type-specific k-mers.

Figure 2. The flowchart of the design of *L. monocytogenes* type-specific PCR primers.

Figure 3. Singleplex PCR experiments. Two columns in the Y axis describe the isolate
number and given MLST type accordingly. In X axis, "ST" describes type-specific primer
pair and "CC" clonal complex specific primer pair. "Univ" states for universal primer pair
that must give a positive result for all samples. TP – true positive, TN – true negative, FP –
false positive, FN – false negative.

Figure 4. 5-plex PCR experiments. In section A, three different multiplex groups were
formed that contain primer pairs with product sizes ranging from 71 to 680 nucleotides. In

section B, two columns in the Y axis describe the isolate number and given MLST type accordingly. In X axis, "Group I-III" describes three different PCR primer pair groups. "Univ" states for universal primer pair that must give a positive result for all samples. TP – true positive, TN – true negative, FP – false positive, FN – false negative.

Figure 5. 15-plex PCR experimental results. 51 sample DNA-s were screened with 15-plex primer pairs. Single continuous band in the represents the product from the universal primer pair (Univ_6) as a positive control of *L. monocytogenes'* DNA presence. 50bp states as a DNA step ladder.

Tables

Table 1. **DNA isolates used for the validation and indications of their types.**

Isolate ID	MLST type	Clonal complex	Isolate ID	MLST type	Clonal complex
1	ST87	CC87	25	ST551	CC8
2	ST87	CC87	26	ST551	CC8
3	ST87	CC87	27	ST551	CC8
4	ST8	CC8	28	ST580	CC9
5	ST8	CC8	29	ST580	CC9
6	ST8	CC8	30	ST580	CC9
7	ST9	CC9	31	ST177	CC177
8	ST9	CC9	32	ST177	CC177
9	ST9	CC9	33	ST177	CC177
10	ST37	CC37	34	ST451	CC11
11	ST37	CC37	35	ST451	CC11
12	ST37	CC37	36	ST451	CC11
13	ST155	CC155	37	ST173	CC19
14	ST155	CC155	38	ST173	CC19
15	ST155	CC155	39	ST173	CC19
16	ST101	CC101	40	ST425	CC90
17	ST101	CC101	41	ST425	CC90
18	ST101	CC101	42	ST425	CC90
19	ST121	CC121	43	ST7	CC7
20	ST121	CC121	44	ST7	CC7
21	ST121	CC121	45	ST7	CC7
22	ST5	CC5	46	ST29	CC29
23	ST5	CC5	47	ST29	CC29
24	ST5	CC5	48	ST29	CC29
			49	ST1247	CC8
			50	ST1247	CC8
			51	ST1247	CC8

Table 2: *L. monocytogenes* **type-specific sequences.**

MLST type	CC type	# of ENA samples	# of randomly selected samples	# of type-specific k-mers	# of selected specific regions (>60 nt)	# of PCR primer pairs (product size range 60-1000 nt)	# of PCR primer pairs after filtering
ST5	CC5	3119	10	161122	1928	984	17
ST7	CC7	1431	10	64594	687	307	13
ST8	CC8	569	10	153	1	0	0
ST9	CC9	1445	10	47	0	0	0
ST29	CC29	252	10	39010	440	206	12
ST37	CC37	269	10	46426	444	202	12
ST87	CC87	345	10	103565	1302	704	12
ST101	CC101	187	10	32327	429	195	11
ST121	CC121	623	10	100178	828	492	12
ST155	CC155	1367	10	85948	627	296	14
ST173	CC19	5	5	107450	689	319	14
ST177	CC177	13	10	57056	574	262	17
ST425	CC90	24	10	41904	444	201	12
ST451	CC11	163	10	52729	694	315	11
ST551	CC8	35	10	789	18	3	0
ST580	CC9	2	2	1694	24	6	0
ST1247	CC8	8	8	618	11	5	5
CC8*	CC8	612	28	65936	581	253	10
CC9*	CC9	1447	12	47007	533	214	13

632 *MLST types marked with coloured background, for which no suitable PCR primer pairs were found, were
633 assembled into one group of clonal complexes named CC8 and CC9. For ST1247, separate primers, specific to
634 this type, were also tested.

SRR00001.fastq
SRR00002.fastq
SRR00003.fastq
...

Cleaned reads from non-target samples

SRR00010.fastq
SRR00011.fastq
...

Cleaned reads from target samples

GenomeTester4 – identify k-mers (*glistmaker*)

GenomeTester4 – identify k-mers (*glistmaker*)

GenomeTester4 – create a union of all k-mers (*glistcompare*)

GenomeTester4 – create an intersect of k-mers (*glistcompare*)

List of k-mers from non-target samples

List of k-mers from target samples

GenomeTester4 – remove k-mers from target list that are shared by non-target list (*glistcompare*)

List of target-specific k-mers

A

PCR product sizes of primer pairs in 5-plex groups

Chart plotting PCR product size (nt) on the y-axis (0 to 900) against Primer pair ID on the x-axis.

Primer pair ID	PCR product size (nt)	Group
ST7_4	124	Group I
ST87_9	289	Group I
ST173_5	482	Group I
ST29_10	569	Group I
CC9_13	670	Group I
ST121_9	138	Group II
ST155_6	245	Group II
CC8_7	430	Group II
ST451_5	641	Group II
ST5_17	786	Group II
ST101_2	71	Group III
ST177_2	154	Group III
ST1247_1	409	Group III
ST425_6	680	Group III

15-plex

50bp 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 50bp

50bp 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 neg 50bp

Univ_6