Machine learning model predicts new-onset lower extremity deep vein thrombosis after pelvic fracture surgery and targeted diagnosis

Haoyuan Fu1,2*,Qi Dong,1,2*,Guoqiang LI1,2*, Kuo Zhao 1,2# ，Zhiyong Hou 1,2#

1 Department of Orthopedic Surgery，Third Hospital of Hebei Medical University，Shijiazhuang，Hebei，China..

2Orthopaedic Research Institute of Hebei Province，Third Hospital of Hebei Medical University，

Shijiazhuang，Hebei，China.

#Corresponding author: Dr. Zhiyong Hou，Address: Department of Orthopaedical Surgery，the Third Hospital

of Hebei Medical University，Shijiazhuang 050051，China. Mail address: drzyhou@hebmu.edu.cn

Dr. Kuo Zhao, Address : Department of Orthopaedical Surgery，the Third Hospital of Hebei Medical

University，Shijiazhuang 050051，China. Mail address: Surzhao@163.com

Abstract

Background: Postoperative new-onset deep vein thrombosis (PNO-DVT) of the lower extremities represents a prevalent and serious clinical complication following pelvic fractures, which substantially impedes patient rehabilitation and diminishes quality of life. Conventional risk assessment methodologies exhibit inherent limitations, rendering them inadequate for precise prediction and individualized management of DVT. In recent years, machine learning techniques have demonstrated significant advantages in data analysis, emerging as promising tools for predicting postoperative DVT risk. This study sought to investigate the predictive efficacy of machine learning models for the development of new-onset lower extremity deep vein thrombosis following pelvic fracture surgery.

Methods: Data from 745 patients who underwent pelvic fracture surgery at our hospital between January 2016 and December 2019 were collected. The analysis encompassed demographic information, general patient data, preoperative laboratory test results, surgical details, and scoring systems. Initially, the data were analyzed using univariate logistic regression, least absolute shrinkage and selection operator (LASSO) regression, and multivariate logistic regression to identify 12 independent risk factors, including age, HDL-C, and ApoB. Subsequently, the dataset was partitioned into a training set and a test set at a 7:3 ratio. Six models were employed for analysis, including logistic regression, support vector machine (SVM), random forest, XGBoost, LightGBM, and AdaBoost.

Conclusion: Comparative analysis of the six machine learning models revealed XGBoost exhibited the highest performance (AUC: 0.8633), followed by LightGBM (0.8349), random forest (0.8055), logistic regression (0.7503), SVM (0.7505), and AdaBoost (0.8179). Model sensitivity ranged from 0.3684 to 0.8421, and accuracy ranged from 0.6502 to 0.9238.

Keywords

DVT risk prediction; Machine learning; Model prediction; Postoperative complications

Background

Pelvic fractures represent a severe form of trauma associated with a broad spectrum of complex and diverse postoperative complications. Notably, the development of deep vein thrombosis (DVT) in the lower extremities is a critical factor contributing to elevated mortality and disability rates among these patients. Over the years, numerous studies have demonstrated that individuals with pelvic fractures are predisposed to a hypercoagulable state, attributed to factors such as prolonged immobilization, tissue damage, and postoperative inflammatory responses, thereby elevating the risk of DVT [1-2]. The incidence of lower limb DVT in patients with pelvic fractures has been reported to reach 20% or higher, with most cases occurring postoperatively, thereby significantly impacting patient rehabilitation outcomes and quality of life [3-4].

Traditional risk assessment for DVT predominantly relies on clinical experience and isolated clinical indicators, such as D-dimer levels, patient age, and body mass index, in conjunction with straightforward assessment scales, including the Wells score, Autar score, and Padua score. However, these methodologies exhibit notable limitations in practical application, characterized by inadequate predictive accuracy and specificity, impairing the ability to achieve early and precise risk stratification in clinical settings. For instance, while the Autar and Padua scores demonstrate high sensitivity in intensive care unit (ICU) populations, their low specificity contributes to a significant rate of misdiagnosis, thereby complicating the implementation of tailored preventive strategies [5-6]. Furthermore, there is a dearth of specialized risk assessment instruments for patients with pelvic fractures, as traditional

models are predominantly grounded in single disease contexts, failing to account for the

intricate pathological conditions associated with pelvic fractures[2].

With the rapid advancement of artificial intelligence and machine learning, risk prediction

models leveraging big data have demonstrated significant potential in the medical domain.

Machine learning is adept at processing extensive, multidimensional clinical datasets,

enabling automatic learning and identification of complex, nonlinear relationships, thereby

enhancing the accuracy and stability of risk assessments. Recent studies have developed

machine learning models to predict postoperative DVT risks in patients undergoing brain

trauma, tumor surgeries, and multiple injuries, achieving commendable predictive

performance. For instance, models utilizing algorithms such as Gradient Boosting Machine

(GBM) and LightGBM have reported area under the curve (AUC) values of 0.85 or higher in

predicting DVT following brain trauma and brain tumor surgeries, markedly outperforming

traditional statistical models[7-9]. Furthermore, integrating feature selection via Support

Vector Machine (SVM) with reinforcement learning algorithms has achieved an accuracy of

95.9% for DVT risk prediction, underscoring the distinct advantages of machine learning in

extracting critical risk factors and optimizing prediction models [10].

Previous studies on pelvic fracture patients have sought to develop risk prediction models

based on clinical variables. These include multifactorial regression models that integrate

factors such as emergency abdominal surgery, the Injury Severity Score (ISS), and indicators

of renal and liver function. The resulting risk assessment nomogram has demonstrated high

predictive accuracy, with an AUC of approximately 0.88, in both training and validation

cohorts, thereby serving as an effective tool for preoperative risk stratification of DVT[2].

Nonetheless, the comprehensive application of machine learning techniques to predict the

risk of new-onset lower limb DVT following pelvic fracture surgery remains nascent.

Accordingly, there is a pressing need to systematically incorporate large-scale, multicenter

datasets and integrate multimodal clinical information to develop a more precise, clinically

valuable prediction model with substantial clinical application potential.

Accordingly, a combination of traditional logistic regression and machine learning algorithms

was adopted in the present study to develop a predictive model for new-onset lower limb

DVT after pelvic fracture surgery, thereby enhancing clinical decision-making.

Methods

Data collection and processing

The study collected data from 857 patients diagnosed with pelvic fractures at the Third

Hospital of Hebei Medical University from January 2016 to December 2019. After excluding

112 patients with preoperative thrombosis, 745 patients who developed postoperative DVT

were included in the final analysis. The incidence of postoperative non-occlusive deep vein

thrombosis (PNO-DVT) was 8.5% (n=64/681), with a predominance of male patients

(74.8%, n=558/857).

Identification of risk factors for PNO-DVT

Independent risk factors were identified using a two-step analytical approach. Initially, a

single-factor logistic regression analysis was employed to select factors with a P-value ≤ 0.05.

Subsequently, the least absolute shrinkage and selection operator (LASSO) regression was

utilized to further refine the selection of influencing factors for postoperative thrombosis,

addressing potential collinearity issues. In the LASSO regression model, an increasing λ

value intensified the penalty term, driving the coefficients of the independent variables

towards zero. This regularization process effectively mitigated model overfitting. Based on

the nine variables identified through univariate analysis and LASSO regression, a

multivariate logistic regression was conducted to identify independent predictors of

postoperative thrombosis.

Model establishment

The dataset for model establishment was derived from selecting 12 significant features

identified through single-factor analysis. This dataset was then randomly partitioned into a

training set (70%) and a test set (30%). Given the approximately 1:10 ratio of positive to

negative samples, a notable class imbalance was present. To mitigate potential model bias

towards the negative samples, which could lead to a diminished recall rate, sampling

techniques were applied exclusively to the training set, leaving the test set unaltered. Model

evaluation was subsequently conducted to more accurately assess the predictive capabilities

of the models. Six machine learning algorithms, including logistic regression, SVM, random

forest, XGBoost, LightGBM, and AdaBoost, were employed to develop six distinct models.

All statistical analyses and data visualizations were executed using R version 4.5.1. The study

utilized the "xgboost", "e1071", "randomForest", "rpart", and "caret" packages for machine

learning modeling, and the "tidyverse", "pROC", "ggplot2", "gridExtra", "tibble", and

"shapviz" packages for data organization and visualization.

Results

1. Results of univariate logistic regression analysis

According to the univariate analysis of baseline characteristics in Table 1, age, hypertension,

kidney disease, rheumatoid disease, peripheral vascular disease, and body mass index (BMI)

were identified as potential risk factors for the development of postoperative thrombosis.

The univariate logistic regression analysis of surgical and blood transfusion data (Table 2)

identified several potential risk factors for postoperative thrombosis. These factors included

surgical approach (fracture reduction), type of bone graft, duration of the operation, intraoperative blood loss, intraoperative blood transfusion, method of blood transfusion, and the administration of suspended red blood cells, plasma, and cryoprecipitate coagulation factors.

The results of the univariate logistic regression analysis, presented in Table 3 of the preoperative laboratory data, indicated that Gamma-Glutamyl Transferase (GGT), High-Density Lipoprotein Cholesterol (HDL-C), Apolipoprotein B (ApoB), Total Carbon Dioxide (TCO2), Glucose (GLU), Glasgow Admission Prognostic Score (GAP), Eosinophil Count (EOS), and Mean Platelet Volume (MPV) represented potential risk factors for postoperative thrombosis.

Further analysis evaluated systemic inflammatory markers, including the Systemic Immune-Inflammation Index (SII), Neutrophil-to-Lymphocyte Ratio (NLR), and Platelet-to-Lymphocyte Ratio (PLR). However, none of these indicators demonstrated significant differences, as presented in Table 4.

Table 4: Univariate Analysis of Validation Index Scores

variables	data type	OR(95%CI)	P
SII (plt*neut/lymph)	CV	1(1-1)	0.475
NLR (neut/lymph)	CV	0.991(0.943-1.04)	0.704
PLR (plt/lymph)	CV	0.998(0.996-1)	0.128

2. LASSO regression for feature selection

LASSO regression was employed to further identify the factors influencing postoperative thrombosis. To mitigate collinearity in the data analysis, LASSO regression was used to select variables with P-values < 0.05 from the univariate analysis. As λ increased, the penalty

also increased, leading to the gradual shrinkage of the regression coefficients of the

independent variables towards zero. This regularization term effectively reduced model

overfitting. Figures 1 and 2 illustrate the variable selection process using LASSO regression

and the impact of cross-validation on overfitting. In Figure 1, each curve represents the

trajectory of a variable's coefficient, with the ordinate indicating the coefficient value and the

abscissa showing the parameter λ, which controls the regularization intensity. As λ varied, the

coefficients of the variables progressively approached zero, ultimately retaining only those

variables with non-zero coefficients. Figure 2 emonstrates the determination of the optimal λ

value via 10-fold cross-validation, indicated by a vertical dotted line. The left dotted line

illustrated that when Lambda was set to Lambda.min, 18 variables were selected, whereas the

right dotted line demonstrated that when Lambda was set to Lambda.1se, 13 variables were

selected. In this study, Lambda.1se (i.e., Lambda = 0.01641399) was selected as the optimal

penalty coefficient for the model. Following the LASSO regression analysis, variables with

non-zero regression coefficients were identified, namely age, nephropathy, peripheral

vascular disease, BMI, fracture reduction, operation time, intraoperative blood loss, de-

erythrocytized RBC, HDL-C group, ApoB group, GLU group, GAP group, and MPV group,

as presented in Table 5.

Figure 1: Feature selection process using Least Absolute Shrinkage and Selection Operator (LASSO) regression.

Figure 2: Graph of the Optimal λ Selection for Cross-Validation

172   Table 5: Lasso Result Coefficient Table

variables	coefficient
Age	0.1448
hypertension	0.0000
nephropathy	0.1127
rheumatoid disease	0.0000

peripheral vascular disease	0.1021
BMI	0.1217
fracture reduction	0.1159
Bone graft type	0.0000
time of operation	0.0492
Intraoperative blood loss	0.2299
Intraoperative blood transfusion	0.0000
Blood transfusion method	0.0000
red blood cells suspension	0.1791
Input plasma volume	0.0000
Cold precipitate coagulation factors	0.0000
GGT	0.0000
HDL-C	-0.1370
APOB	0.0610
TCO2	0.0000
GLU	0.0612
GAP	0.1014
E0S	0.0000
MPV	-0.0684

3.Multivariate Logistic Regression Analysis

Building on the nine variables identified through univariate analysis and LASSO regression, a multivariate logistic regression analysis was conducted to identify independent predictors of postoperative thrombosis. The assignment outcomes for the independent variables are

178 presented in the table below. Our analysis identified age, nephropathy, BMI, fracture

179 reduction,intraoperative blood loss, administration of red blood cell suspension, HDL-C,

180 ApoB, GAP, and MPV as independent risk factors for blood transfusion. The detailed results

181 of the regression analysis are presented in Table 6.

182 Table 6: Independent Predictors of Postoperative Thrombosis in Multivariate Logistic Regression Analysis

variables	β	standard	OR(95% CI)	P
Age	0.022	0.011	1.022(1.000-1.044)	0.046
nephropathy	2.519	0.923	12.415(2.033-75.826)	0.006
peripheral vascular disease	2.702	1.358	14.917(1.041-213.739)	0.047
BMI	0.092	0.043	1.096(1.007-1.193)	0.034
fracture reduction	1.859	0.851	6.420(1.210-34.065)	0.029
time of operation	0.001	0.002	1.001(0.998-1.004)	0.507
Intraoperative blood loss	0.000	0.0002	1.0005(1.0001-1.0010)	0.024
red blood cells suspension	0.000	0.0001	1.0003(1.0001-1.0005)	0.007
HDL-C < 0.9(Refer to 0.9-2)	-0.892	0.322	0.410(0.218-0.770)	0.006
APOB < 0.8((Refer to 0.8-1.2)	0.566	0.348	1.761(0.891-3.482)	0.104
APOB > 1.2((Refer to 0.8-1.2)	0.959	0.489	2.609(1.001-6.799)	0.050
GLU > 6.1(Refer to 3.9-6.1)	0.608	0.305	1.836(1.010-3.336)	0.046
GAP < 8((Refer to 8-16)	0.549	0.332	1.731(0.904-3.316)	0.098
GAP > 16(Refer to 8-16)	2.270	0.682	9.676(2.540-36.865)	0.001
MPV < 7.4(Refer to 7.4-11)	-0.977	0.478	0.376(0.147-0.961)	0.041
MPV > 11(Refer to 7.4-11)	-0.468	1.100	0.627(0.073-5.409)	0.671

Establishment and Analysis of Postoperative PNO-DVT Prediction Model Based on

Machine Learning Algorithms

1. Establishment of the prediction model and performance evaluation

The twelve significant features identified from the preceding univariate analysis were

selected as the dataset for the final model construction. The dataset was randomly split into a

training set (70%) and a test set (30%). Given that the ratio of positive to negative samples

was approximately 1:10, there was a notable imbalance. To mitigate the risk of the model

being biased towards the negative samples during prediction, which could result in an

unacceptably low recall rate, sampling was conducted on the training set; however, no

processing was applied to the test set. This approach ensured that the model evaluation more

accurately reflected the model's predictive capabilities. Six machine learning algorithms,

including logistic regression, SVM, random forest, XGBoost, LightGBM, and AdaBoost,

were employed to construct six distinct models. Among these, the XGBoost algorithm

achieved the highest AUC value of 0.8633, indicating superior predictive performance for

postoperative thrombosis within this dataset (Table 7). Consequently, XGBoost was

determined to be the optimal model(Figure 3). Figure 4 illustrates the feature importance

rankings for the random forest, XGBoost, LightGBM, and AdaBoost models.

Table 7: Performance Evaluation of Each Model's Prediction Model

Model Name	Accuracy	Sensitivity	Specificity	F1	AUC
logistic regression	0.6771	0.7368	0.6716	0.2800	0.7503
random forest	0.9238	0.3158	0.9804	0.4138	0.8055
XGBoost	0.8117	0.8421	0.8088	0.4324	0.8633
LightGBM	0.8879	0.3684	0.9363	0.3590	0.8349

SVM	0.6502	0.6842	0.6471	0.2500	0.7505
AdaBoost	0.9283	0.4211	0.9755	0.5000	0.8179

Figure 3: Comparison of ROC curves for each model

Figure 4: Model Feature Importance Chart

2. Explanatory analysis based on SHAP

Figures 4 and 5 provide visual representations of the feature importance rankings derived from SHAP values in the XGBoost model. Specifically, Figure 4 presents a swarm plot, while Figure 5 displays a feature importance plot. Both figures effectively illustrate the relative significance of each feature within the model. Age was identified as the most important predictor, followed by intraoperative blood loss, red blood cell suspension, BMI, GLU group, GAP group, reduction method group, MPV group, HDL-C group, ApoB group, kidney disease, and peripheral vascular disease(Figure 6).

Figure 5: Summary plot of SHAP (SHapley Additive exPlanations) values, illustrating the impact of the top

features on the model's output

Figure 6: Feature Importance Ranking Chart Based on Shap Values

3.Analysis based on SHAP waterfall chart and force-directed layout

Figure 7 demonstrates the utility of SHAP force plots for deconstructing individual

predictions. These visualizations elucidate the contribution of each feature to a single

sample's outcome, where the color spectrum represents the direction of influence: features in

yellow elevate the prediction score, while those in purple diminish it. The vertical axis

enumerates the specific variables and their respective values for the given case. To illustrate

this interpretability, force plots for representative positive and negative cases were generated.

Analysis of a high-risk (positive) patient revealed that the model's prediction was

predominantly driven by several key factors. The patient's age of 43 years was the most

substantial contributor, increasing the prediction score by +0.269. This was closely followed

by the administration of 2400 mL of packed red blood cells (+0.257) and an intraoperative

blood loss of 700 mL (+0.16). Membership in GLU Group 1 provided a further, though

smaller, positive influence (+0.103). Conversely, certain features moderated the overall risk.

Membership in the GAP Group 0 exerted a slight protective effect (negative impact),

partially offsetting the risk-enhancing factors. Other variables, such as HDL-C Group 0,

MPV Group 0, reduction method 2, and ApoB Group 2, demonstrated minimal influence on

the final prediction, despite their positive directional effect. The collective impact of the

remaining features was negligible.

The values associated with the features of negative sample patients are detailed as follows:

Age 35 exerted the most significant negative impact on the prediction value, decreasing it by

0.766. The administration of 400 units of red blood cell suspension reduced the prediction

value by 0.21, indicating a notable inhibitory effect. A BMI of 25.0 contributed a negative

impact of 0.159, further decreasing the prediction value. The MPV group value of 1 reduced

the prediction value by 0.0616, acting as a factor with a negative inhibitory effect.

Conversely, the GLU group value of 1, highlighted in yellow, positively influenced the

prediction value, albeit with a relatively small amplitude. Features such as the GAP group

value of 0 and intraoperative blood loss of 600, marked in purple, negatively affected the

prediction value, though their influence was relatively minor, as indicated by the figure.

Besides, features such as the HDL-C group value of 1 and the reduction method value of 2

yielded a discernible impact on the prediction value, with the degree of influence being

relatively small, as illustrated in the figure. The remaining 3 had a negligible impact on the

prediction value.

Figure 7: The Case's Attempt

Discussion

In recent years, the application of machine learning technology in the medical domain has notably expanded, particularly in thrombosis risk prediction, where it has demonstrated considerable benefits. Machine learning models can predict the likelihood of DVT after fractures or surgery, creating novel opportunities for tailored prevention. Empirical studies have shown that machine learning algorithms are proficient at handling substantial volumes of high-dimensional data, thereby facilitating the identification of novel risk factors and enhancing patient risk stratification[11].

Machine learning models have demonstrated significant efficacy in predicting DVT

following fractures. For example, a study developed several machine learning models to

forecast the incidence of DVT after tibial fracture surgery. Notably, the random forest model,

when integrated with recursive feature elimination (RFE), achieved an AUC of 0.99,

indicating exceptionally high predictive accuracy [12]. Furthermore, the support vector

machine model, when combined with Boruta or LASSO feature selection methods,

demonstrated superior calibration, underscoring the reliability of these models for risk

estimation [12].

In various surgical contexts, machine learning models have shown significant promise in

predicting DVT. For example, in the context of laparoscopic surgery, a study assessed 20

machine learning algorithms and developed a stacked model comprising the five most

effective algorithms. This model outperformed the conventional Caprini score in external

validation [13]. Besides, in the context of gastric cancer surgery, a machine learning model

utilizing imaging data has emerged as an innovative non-invasive predictive tool,

demonstrating enhanced efficacy in the early detection and management of DVT [14].

The use of machine learning to predict DVT extends beyond the postoperative context. In

patients experiencing spontaneous intracerebral hemorrhage, machine learning algorithms

have been employed to forecast the risk of lower extremity DVT during hospitalization. The

Light Gradient Boosting Machine (LGBM) model was found to exhibit a notable advantage

in DVT prediction[15]. Furthermore, in oncology patients undergoing chemotherapy, machine

learning models are utilized to evaluate the risk of catheter-related thrombosis. Despite the

clinical application being constrained by complexity, the Bayesian learning model offers a

simplified and robust alternative[16].

As the utilization of machine learning algorithms in the medical field continues to expand,

several researchers have endeavored to develop predictive models for DVT following

fracture surgery using these advanced techniques. In the context of predicting DVT after

various fractures or surgical procedures, numerous studies have demonstrated that machine

learning models exhibit good performance in identifying significant risk factors and

facilitating risk stratification. For instance, researchers have employed a range of algorithms

for predicting DVT after knee and hip replacement surgeries, including XGBoost, RF, and

SVM, integrating patient electronic health records (EHRs) to construct a highly accurate

predictive model, achieving an AUC exceeding 0.92, which markedly surpasses the

traditional Caprini score [17]. Similarly, in the prediction of DVT following spinal surgery, the

combination of Boruta and LASSO algorithms for feature selection enabled the random

forest model to perform exceptionally well, accurately identifying patients at high risk [18].

Furthermore, a study investigating DVT following surgical intervention for lower limb

fractures compared various machine learning models, including XGBoost, Random Forest,

and Logistic Regression. The results revealed that XGBoost exhibited superior performance

with an AUC of 0.979, indicating robust predictive capability[19].

Machine learning models offer substantial advantages over logistic regression in the context

of large sample datasets, a perspective corroborated by numerous studies. Primarily, machine

learning techniques excel at handling high-dimensional data and handling missing values.

Contemporary machine learning approaches, including penalized regression, tree-based

models, and deep learning, exhibit good performance in handling missing data within high-

dimensional contexts, mitigating non-response bias, and demonstrating superiority in both

simulation studies and practical applications[20]. Furthermore, in the analysis of large-scale

neuroimaging datasets, machine learning effectively mitigates overfitting issues by

integrating the predictive strengths of multiple classifiers, thereby enhancing the model's

generalization capabilities[21].

In the medical domain, machine learning models have demonstrated superior performance

compared to traditional logistic regression models in predicting patient readmission and

mortality rates following colon cancer surgery. Empirical research indicates that models

developed using machine learning algorithms, such as Random Forest and XGBoost, exhibit

significantly higher AUC values than logistic regression models, thereby reflecting enhanced

discriminatory capabilities[22]. Similarly, in the context of risk stratification for COVID-19

patients, machine learning models achieve substantially higher AUC values compared to

logistic regression models, suggesting greater accuracy in identifying patients at risk of

severe outcomes [23].

Moreover, machine learning models demonstrate superior capabilities for handling the effects

of large-scale datasets and the intricate interactions among variables. Empirical studies have

shown that machine learning models exhibit greater robustness to variations in dataset size,

whereas deep learning models can maintain strong performance even in the absence of

interaction terms [24]. Another study demonstrated that machine learning models could

enhance the prediction of post-surgical complications by reliably identifying high-risk

patients, yielding more precise prognostic data[25].

In this study, six predictive models were developed, including a logistic regression model, to

predict PNO-DVT. The logistic regression model achieved an AUC of 0.7503, ranking fourth

in performance. Notably, the XGBoost model demonstrated the best performance

(AUC=0.8633).

In patients with pelvic fractures, the severity of vascular injury is significantly correlated with

prognosis. Research indicates that, even in cases of relatively isolated pelvic injuries, the

severity of vascular injury is more strongly associated with clinical outcomes than the

anatomical complexity of the fracture itself [26]. Such vascular injuries not only elevate the

risk of hemorrhage but may also initiate a series of inflammatory responses, thereby

exacerbating the hypercoagulable state of the blood. In clinical practice, the management of

trauma stress and inflammatory responses is crucial for improving the prognosis of patients

with pelvic fractures. Research has shown that chronic psychological and social stress can

affect the immune response through the β-adrenergic receptor signaling pathway, thereby

influencing fracture healing[26]. Therefore, interventions targeting traumatic stress and

inflammatory responses may alleviate hypercoagulability and improve overall patient

prognosis. Accordingly, systemic inflammatory indicators SII, NLR, and PLR were

incorporated into our analysis. Ultimately, none of these indicators were identified as

significant predictors in this patient population.

Subsequently, SHAP values were employed to elucidate the probability of PNO-DVT as

predicted by the XGBoost model. The analysis of SHAP values revealed that age was the

most significant factor, followed by intraoperative blood loss, leukocyte removal, BMI, GLU

group, GAP group, reduction method group, MPV group, HDL-C group, ApoB group,

kidney disease, and peripheral vascular disease. Age and BMI, recognized as common

cardiovascular risk factors, have been consistently associated with thrombosis in numerous

studies. Current evidence indicates that older patients are at an increased risk for thrombotic

events, potentially due to decreased vascular elasticity and the presence of atherosclerosis [28].

Furthermore, an elevated BMI has been linked to metabolic syndrome and inflammatory

responses, both of which may contribute to the development of thrombosis [29].

Intraoperative blood loss and the volume of suspended red blood cells transfused

postoperatively are critical determinants influencing postoperative thrombosis. Significant

intraoperative blood loss can result in hemodynamic instability, whereas excessive

postoperative transfusion of suspended red blood cells may elevate blood viscosity, thereby

augmenting the risk of thrombosis [30]. These factors necessitate careful consideration in

postoperative management to mitigate the risk of thrombosis.

Among laboratory test indicators, MPV is recognized as a marker of platelet activation and is

closely associated with thrombosis. Empirical evidence suggests that patients with elevated

MPV are more susceptible to venous thromboembolism (VTE), potentially due to the

increased platelet volume, which enhances platelet reactivity and thus promotes thrombosis

[31-32]. Furthermore, integrating MPV with other hematological indicators can enhance the

accuracy of thrombosis risk prediction [33].

Elevated glucose levels are significantly correlated with the incidence of DVT. Research

indicates that hyperglycemia elevates the risk of thrombosis, potentially due to increased

blood viscosity and vascular endothelial dysfunction associated with high glucose levels [34].

Furthermore, individuals with diabetes are predisposed to thrombosis as a result of prolonged

hyperglycemia, substantiating that glucose represents a risk factor for DVT [35].

Secondly, triglycerides and HDL-C are integral to lipid metabolism. Elevated triglyceride

levels and reduced HDL-C levels have been linked to the incidence of DVT. Research

indicates that dyslipidemia can damage the vascular endothelium and trigger inflammatory

responses, thereby elevating the risk of thrombosis [36-37]. Furthermore, low-density

lipoprotein cholesterol (LDL-C) and the lymphocyte-to-lymphocyte ratio (LLR) have been

documented as predictors of DVT, underscoring the significant role of lipid metabolism

disorders in the pathogenesis of DVT [37].

Apolipoprotein B (APOB), the primary component of low-density lipoprotein, is significantly

associated with atherosclerosis and thrombosis. Research indicates that elevated levels of

APOB may elevate the risk of deep vein thrombosis (DVT) by promoting lipid deposition

and vascular endothelial damage [38].

In conclusion, our machine learning model identified age, intraoperative blood loss,

intraoperative and postoperative blood transfusion, body mass index (BMI), glucose (GLU),

and mean platelet volume (MPV) as the most critical predictors of postoperative non-

occlusive deep vein thrombosis (PNO-DVT) in patients with pelvic fractures across the entire

dataset. The collection of these biomarkers facilitates the prediction of postoperative PNO-

DVT risk, thereby informing clinical practice. Furthermore, we employed SHapley Additive

exPlanations (SHAP) to enhance the model's interpretability. SHAP values assist clinicians in

providing personalized interventions for patients at high risk of DVT, allowing for tailored

intervention strategies based on individual patient risk factors rather than a standardized

treatment approach for all patients. This personalized approach promotes the efficient

allocation of medical resources.

Conclusion

In this study, we employed six models, including logistic regression, SVM, random forest,

XGBoost, LightGBM, and AdaBoost, to establish a predictive model for new-onset DVT

following pelvic fracture surgery. Among these machine learning models, XGBoost

demonstrated superior performance, achieving the highest AUC of 0.8633. The integration of

machine learning models into clinical practice has the potential to enhance the prognosis of

patients with pelvic fractures by facilitating efficient risk stratification for post-operative

new-onset DVT.

Abbreviations

403 PNO-DVT   postoperative new-onset deep venous thrombosis

404 DVT   deep venous thrombosis

405 CV   continuous variable

Declarations

Ethics approval and consent to participate

According to the Helsinki Declaration, this study has been approved by the Ethics Committee

of the Third Hospital of Hebei Medical University. Before analysis, all data were anonymized,

and since it was a retrospective study design, obtaining patient consent was not necessary.

Consent for publication

All the authors have agreed to publish.

Availability of data and materials

Data cannot be published without patients' consent. Researchers who are interested for

academic need could contact the corresponding authors.

Competing interests

This paper has been uploaded to medRxiv as a preprint:

https://www.medrxiv.org/content/10.64898/2025.12.01.25341405v1.full

The authors declare that they have no competing interests.

Funding

This study was supported by Key R&D Program of the China Ministry of Science and

Technology (2024YFC2510600) and National Natural Science Foundation of China /

8220090241

Acknowledgements

Not applicable

Clinical trial number

Not applicable

Authors' information

Haoyuan Fu:fuhaoyuan2001@163.com;QiDong: 15755819518@163.com;Guoqiang Li:

drgqli@outlook.com;Kuo Zhao: Surzhao@163.com; Zhiyong Hou: drzyhou@hebmu.edu.cn

1 Department of Orthopedic Surgery，Third Hospital of Hebei Medical University，Shijiazhuang，Hebei，

China..

2Orthopaedic Research Institute of Hebei Province，Third Hospital of Hebei Medical University，

Shijiazhuang，Hebei，China.

#Corresponding author: Dr. Zhiyong Hou，Address: Department of Orthopaedical Surgery，the Third Hospital

of Hebei Medical University，Shijiazhuang 050051，China. Mail address: drzyhou@hebmu.edu.cn

Dr. Kuo Zhao, Address : Department of Orthopaedical Surgery，the Third Hospital of Hebei Medical

University，Shijiazhuang 050051，China. Mail address: Surzhao@163.com

References

[1]. Chen H, Sun L, Kong X. Risk assessment scales to predict risk of lower extremity deep vein thrombosis among multiple trauma patients: a prospective cohort study. BMC Emerg Med. 2023 Dec 5;23(1):144. doi: 10.1186/s12873-023-00914-7. PMID: 38053029; PMCID: PMC10696745.

[2]. Chen Y, He J, Pan X. Prediction of risk factors for preoperative deep vein thrombosis in patients with pelvic fracture. Front Surg. 2025 Apr 28;12:1585460. doi: 10.3389/fsurg.2025.1585460. PMID: 40356946; PMCID: PMC12066781.

[3]. Wang H, Wu G, Chen CY, Qiu YY, Xie Y. Percutaneous screw fixation assisted by hollow pedicle finder for superior pubic ramus fractures. BMC Surg. 2022 Jun 3;22(1):216. doi: 10.1186/s12893-022-01659-z. PMID: 35658934; PMCID: PMC9166495.

[4]. Yang CS, Tan Z. Construction and validation of a predictive model for preoperative lower extremity deep vein thrombosis risk in elderly hip fracture patients: An observational study. Medicine (Baltimore). 2024 Sep 20;103(38):e39825. doi: 10.1097/MD.0000000000039825. PMID: 39312315; PMCID: PMC11419451.

[5]. Orak F, Saadat M, Saki Malehi A, Behdarvandan A, Esfandiarpour F. Comparison of the Pauda and the Autar DVT Risk Assessment Scales in Prediction of Venous Thromboembolism in ICU Patients. Med J Islam Repub Iran. 2024 Apr 30;38:48. doi: 10.47176/mjiri.38.48. PMID: 39399622; PMCID: PMC11469705.

[6].   Shekarchian S, Notten P, Barbati ME, Van Laanen J, Piao L, Nieman F, Razavi MK,

Lao M, Mees B, Jalaie H. Development of a prediction model for deep vein thrombosis

in a retrospective cohort of patients with suspected deep vein thrombosis in primary care.

J Vasc Surg Venous Lymphat Disord. 2022 Sep;10(5):1028-1036.e3. doi:

10.1016/j.jvsv.2022.04.009. Epub 2022 May 27. PMID: 35644336.

[7].   Tang Z, Li N, Tian Y. A nomogram for predicting risk factors for lower limb deep

venous thrombosis in elderly postoperative patients with severe traumatic brain injury in

the intensive care unit. Phlebology. 2025 Jul;40(6):446-455. doi:

10.1177/02683555251332988. Epub 2025 Apr 10. PMID: 40205921.

[8].   Wu L, Zhao Y, Yao G, Li X, Zhao X. Prediction and Analysis of Risk Factors for Lower

Extremity Deep Vein Thrombosis After Craniotomy in Patients with Primary Brain

Tumors: A Machine Learning Approach. Turk Neurosurg. 2025;35(4):636-643. doi:

10.5137/1019-5149.JTN.47938-24.3. PMID: 40577511.

[9].   Qiu W, Cui P, Li S, Tang Z, Chen J, Wang J, Li Y. Machine learning models predict risk

of lower extremity deep vein thrombosis in hospitalized patients with spontaneous

intracerebral hemorrhage. Sci Rep. 2025 Jul 10;15(1):24932. doi: 10.1038/s41598-025-

10905-2. PMID: 40640445; PMCID: PMC12246041.

[10].   Li R, Chen S, Xia J, Zhou H, Shen Q, Li Q, Dong Q. Predictive modeling of deep vein

thrombosis risk in hospitalized patients: A Q-learning enhanced feature selection model.

Comput Biol Med. 2024 Jun;175:108447. doi: 10.1016/j.compbiomed.2024.108447.

Epub 2024 Apr 12. PMID: 38691912.

[11].   Chrysafi P, Lam B, Carton S, Patell R. From Code to Clots: Applying Machine Learning

to Clinical Aspects of Venous Thromboembolism Prevention, Diagnosis, and

Management. Hamostaseologie. 2024 Dec;44(6):429-445. doi: 10.1055/a-2415-8408.

Epub 2024 Dec 10. PMID: 39657652.

[12]. Baki H, Özçelik İB. Machine Learning-Based Prediction of Postoperative Deep Vein

Thrombosis Following Tibial Fracture Surgery. Diagnostics (Basel). 2025 Jul

16;15(14):1787. doi: 10.3390/diagnostics15141787. PMID: 40722536; PMCID:

PMC12293441.

[13]. Yang SZ, Peng MH, Lin Q, Guan SW, Zhang KL, Yu HB. A machine learning-based

predictive model for the occurrence of lower extremity deep vein thrombosis after

laparoscopic surgery in abdominal surgery. Front Surg. 2025 May 30;12:1502944. doi:

10.3389/fsurg.2025.1502944. PMID: 40520687; PMCID: PMC12162540.

[14]. Zeng Y, Chen Y, Zhu D, Xu J, Zhang X, Ying H, Song X, Zhou R, Wang Y, Yu F.

Machine learning assisted radiomics in predicting postoperative occurrence of deep

venous thrombosis in patients with gastric cancer. BMC Cancer. 2025 Feb 7;25(1):220.

doi: 10.1186/s12885-025-13630-1. PMID: 39920636; PMCID: PMC11806839.

[15]. Qiu W, Cui P, Li S, Tang Z, Chen J, Wang J, Li Y. Machine learning models predict risk

of lower extremity deep vein thrombosis in hospitalized patients with spontaneous

intracerebral hemorrhage. Sci Rep. 2025 Jul 10;15(1):24932. doi: 10.1038/s41598-025-

10905-2. PMID: 40640445; PMCID: PMC12246041.

[16]. An T, Han H, Xie J, Wang Y, Zhao Y, Jia H, Wang Y. Enhancing prediction and

stratifying risk: machine learning and bayesian-learning models for catheter-related

thrombosis in chemotherapy patients. BMC Cancer. 2025 Mar 27;25(1):552. doi:

10.1186/s12885-025-13946-y. PMID: 40148861; PMCID: PMC11948715.

[17]. Wang X, Xi H, Geng X, Li Y, Zhao M, Li F, Li Z, Ji H, Tian H. Artificial Intelligence-

Based Prediction of Lower Extremity Deep Vein Thrombosis Risk After Knee/Hip

Arthroplasty. Clin Appl Thromb Hemost. 2023 Jan-Dec;29:10760296221139263. doi:

10.1177/10760296221139263. PMID: 36596268; PMCID: PMC9830569.

[18]. Wu X, Wang Z, Zheng L, Yang Y, Shi W, Wang J, Liu D, Zhang Y. Construction and

verification of a machine learning-based prediction model of deep vein thrombosis

formation after spinal surgery. Int J Med Inform. 2024 Dec;192:105609. doi:

10.1016/j.ijmedinf.2024.105609. Epub 2024 Aug 30. PMID: 39260049.

[19]. Wei C, Wang J, Yu P, Li A, Xiong Z, Yuan Z, Yu L, Luo J. Comparison of different

machine learning classification models for predicting deep vein thrombosis in lower

extremity fractures. Sci Rep. 2024 Mar 22;14(1):6901. doi: 10.1038/s41598-024-57711-

w. PMID: 38519523; PMCID: PMC10960026.

[20]. Chen S, Xu C. Handling high-dimensional data with missing values by modern machine

learning techniques. J Appl Stat. 2022 May 1;50(3):786-804. doi:

10.1080/02664763.2022.2068514. PMID: 36819079; PMCID: PMC9930810.

[21]. Lanka P, Rangaprakash D, Dretsch MN, Katz JS, Denney TS Jr, Deshpande G.

Supervised machine learning for diagnostic classification from large-scale neuroimaging

datasets. Brain Imaging Behav. 2020 Dec;14(6):2378-2416. doi: 10.1007/s11682-019-

00191-8. PMID: 31691160; PMCID: PMC7198352.

[22]. Leonard G, South C, Balentine C, Porembka M, Mansour J, Wang S, Yopp A, Polanco

P, Zeh H, Augustine M. Machine Learning Improves Prediction Over Logistic

Regression on Resected Colon Cancer Patients. J Surg Res. 2022 Jul;275:181-193. doi:

10.1016/j.jss.2022.01.012. Epub 2022 Mar 11. PMID: 35287027.

[23]. Ye J, Hua M, Zhu F. Machine Learning Algorithms are Superior to Conventional

Regression Models in Predicting Risk Stratification of COVID-19 Patients. Risk Manag

Healthc Policy. 2021 Jul 29;14:3159-3166. doi: 10.2147/RMHP.S318265. PMID:

34349576; PMCID: PMC8328384.

[24]. Bailly A, Blanc C, Francis É, Guillotin T, Jamal F, Wakim B, Roy P. Effects of dataset

size and interactions on the prediction performance of logistic regression and deep

learning models. Comput Methods Programs Biomed. 2022 Jan;213:106504. doi:

10.1016/j.cmpb.2021.106504. Epub 2021 Oct 28. PMID: 34798408.

[25]. Michelsen C, Jørgensen CC, Heltberg M, Jensen MH, Lucchetti A, Petersen PB,

Petersen T, Kehlet H; Center for Fast-track Hip Knee Replacement Collaborative group;

Madsen F, Hansen TB, Gromov K, Jakobsen T, Varnum C, Overgaard S, Rathsach M,

Hansen L. Machine-learning vs. logistic regression for preoperative prediction of

medical morbidity after fast-track hip and knee arthroplasty-a comparative study. BMC

Anesthesiol. 2023 Nov 29;23(1):391. doi: 10.1186/s12871-023-02354-z. PMID:

38030979; PMCID: PMC10685559.

[26]. Wu YT, Cheng CT, Tee YS, Fu CY, Liao CH, Hsieh CH. Pelvic injury prognosis is

more closely related to vascular injury severity than anatomical fracture complexity: the

WSES classification for pelvic trauma makes sense. World J Emerg Surg. 2020 Aug

17;15(1):48. doi: 10.1186/s13017-020-00328-x. PMID: 32807185; PMCID:

PMC7433075.

[27]. Haffner-Luntzer M, Foertsch S, Fischer V, Prystaz K, Tschaffon M, Mödinger Y,

Bahney CS, Marcucio RS, Miclau T, Ignatius A, Reber SO. Chronic psychosocial stress

compromises the immune response and endochondral ossification during bone fracture

healing via β-AR signaling. Proc Natl Acad Sci U S A. 2019 Apr 23;116(17):8615-8622.

doi: 10.1073/pnas.1819218116. Epub 2019 Apr 4. PMID: 30948630; PMCID:

PMC6486758.

[28]. Rong X, Jiang L, Qu M, Yang S, Wang K, Jiang L. Risk factors and characteristics of

ischemic stroke in patients with immune thrombocytopenia: A retrospective cohort

study. J Stroke Cerebrovasc Dis. 2022 Oct;31(10):106693. doi:

10.1016/j.jstrokecerebrovasdis.2022.106693. Epub 2022 Aug 31. PMID: 36054971.

[29]. Hansen ES, Edvardsen MS, Aukrust P, Ueland T, Hansen JB, Brækkan SK, Morelli VM.

Combined effect of high factor VIII levels and high mean platelet volume on the risk of

future incident venous thromboembolism. J Thromb Haemost. 2023 Oct;21(10):2844-

2853. doi: 10.1016/j.jtha.2023.06.022. Epub 2023 Jun 29. PMID: 37393000.

[30]. Zhang YM, Chen W, Wei HL, Tang XH, Xie FH, Wang RX. Analysis of predictive

factors of thrombosis in autogenous arteriovenous fistula. J Vasc Access. 2024

Jul;25(4):1134-1139. doi: 10.1177/11297298221151135. Epub 2023 Jan 27. PMID:

36707987.

[31]. Edvardsen MS, Hansen ES, Hindberg K, Morelli VM, Ueland T, Aukrust P, Brækkan

SK, Evensen LH, Hansen JB. Combined effects of plasma von Willebrand factor and

platelet measures on the risk of incident venous thromboembolism. Blood. 2021 Dec

2;138(22):2269-2277. doi: 10.1182/blood.2021011494. PMID: 34161566.

[32]. Wang Z, Chen X, Wu J, Zhou Q, Liu H, Wu Y, Liu S, Liu Y. Low Mean Platelet

Volume is Associated with Deep Vein Thrombosis in Older Patients with Hip Fracture.

Clin Appl Thromb Hemost. 2022 Jan-Dec;28:10760296221078837. doi:

10.1177/10760296221078837. PMID: 35157546; PMCID: PMC8848069.

[33]. Jakobsen L, Frischmuth T, Brækkan SK, Hansen JB, Morelli VM. Joint Effect of

Multiple Prothrombotic Genotypes and Mean Platelet Volume on the Risk of Incident

Venous Thromboembolism. Thromb Haemost. 2022 Nov;122(11):1911-1920. doi:

10.1055/a-1863-2052. Epub 2022 May 26. PMID: 35617954.

[34]. Liu X, Li T, Xu H, Wang C, Ma X, Huang H, Hu Y, Chu H. Hyperglycemia may

increase deep vein thrombosis in trauma patients with lower limb fracture. Front

Cardiovasc Med. 2022 Sep 8;9:944506. doi: 10.3389/fcvm.2022.944506. PMID:

36158801; PMCID: PMC9498976.

[35]. Hang L, Haibier A, Kayierhan A, Abudurexiti T. Risk factors for deep vein thrombosis

of the lower extremity after total hip arthroplasty. BMC Surg. 2024 Sep 11;24(1):256.

doi: 10.1186/s12893-024-02561-6. PMID: 39261801; PMCID: PMC11389418.

[36]. Abdelmalik BHA, Leslom MMA, Gameraddin M, Alshammari QT, Hussien R, Alyami

MH, Salih M, Yousef M, Yousif E. Assessment of Lower Limb Deep Vein Thrombosis:

Characterization and Associated Risk Factors Using Triplex Doppler Imaging. Vasc

Health Risk Manag. 2023 May 4;19:279-287. doi: 10.2147/VHRM.S409253. PMID:

37168880; PMCID: PMC10166097.

[37]. Guo H, Li C, Wu H, Ma M, Zhu R, Wang M, Yang B, Pan N, Zhu Y, Wang J. Low-

density lipoprotein cholesterol-to-lymphocyte count ratio (LLR) is a promising novel

predictor of postoperative new-onset deep vein thrombosis following open wedge high

tibial osteotomy: a propensity score-matched analysis. Thromb J. 2024 Jul 16;22(1):64.

doi: 10.1186/s12959-024-00635-2. PMID: 39014396; PMCID: PMC11250942.

[38]. Gu H, Yang F, Xie H, Li M, Wang Z, Sheng L. Serum VEGF, P-Selectin, HDL-C,

Platelet Index, and Coagulation Function Index in Deep Vein Thrombosis after

Traumatic Fracture. Clin Lab. 2024 Feb 1;70(2). doi: 10.7754/Clin.Lab.2023.230425.

PMID: 38345981.

606   Table 1: Univariate Analysis of General Information

variables	targeted value	OR(95%CI)	P
age	CV	1.03(1.012-1.048)	0.001
hypertension	NO	Ref.	
	YES	2.029(1.055-3.903)	0.034
diabetes	NO	Ref.	
	YES	1.09(0.564-2.105)	0.798
cerebrovascular disease	NO	Ref.	
	YES	1.395(0.572-3.402)	0.464
cardiopathy	NO	Ref.	
	YES	1.128(0.389-3.268)	0.824
arrhythmia	NO	Ref.	
	YES	0.475(0.063-3.586)	0.471
respiratory failure	NO	Ref.	
	YES	0(0-Inf)	0.986
COPD	NO	Ref.	
	YES	0(0-Inf)	0.989
history of lung disease	NO	Ref.	
	YES	0(0-Inf)	0.987
pneumonia	NO	Ref.	
	YES	0(0-Inf)	0.986
spinal cord injury	NO	Ref.	
	YES	0(0-Inf)	0.986
tumour	NO	Ref.	
	YES	0(0-Inf)	0.984

cerebral trauma	NO	Ref.	
	YES	1.16(0.4-3.366)	0.784
hepatopathy	NO	Ref.	
	YES	0.936(0.279-3.136)	0.915
nephropathy	NO	Ref.	
	YES	6.649(1.552-28.493)	0.011
hyperthyroidism	NO	Ref.	
	YES	0(0-Inf)	0.989
rheumatoid disease	NO	Ref.	
	YES	10.952(1.516-79.096)	0.018
peripheral vascular disease	NO	Ref.	
	YES	21.935(1.961-245.341)	0.012
anemia	NO	Ref.	
	YES	0(0-Inf)	0.986
urinary tract infection	NO	Ref.	
	YES	0(0-Inf)	0.982
Surgical history	NO	Ref.	
	YES	0.688(0.359-1.32)	0.261
Paraplegia or quadriplegia	NO	Ref.	
	YES	4.361(0.829-22.945)	0.082
allergy	NO	Ref.	
	YES	1.265(0.638-2.507)	0.501
blood type	A	Ref.	
	B	1.529(0.744-3.144)	0.248

	AB	1.433(0.664-3.092)	0.359
	0	1.955(0.787-4.854)	0.149
Rh-negative Blood type	NO	Ref.	
	YES	0(0-Inf)	0.988
place of residence	County	Ref.	
	City	0.772(0.41-1.455)	0.424
	spring	Ref.	
	summer	1.066(0.554-2.052)	0.849
Hospitalization season	fall	0.714(0.344-1.482)	0.366
	winter	0.563(0.262-1.209)	0.141
	spring	Ref.	
	summer	1.006(0.515-1.964)	0.987
Surgical season	fall	0.852(0.419-1.733)	0.659
	winter	0.61(0.282-1.321)	0.210
Interval between admission and surgery	CV	0.987(0.942-1.036)	0.601
Total number of fractures	CV	1.053(0.93-1.193)	0.413
	1	Ref.	
	2	2.132(0.538-8.445)	0.281
Cause of injury	3	1.38(0.651-2.923)	0.401
	4	1.953(0.898-4.247)	0.091
	6	0(0-Inf)	0.981
	closed injury	Ref.	
Type of injury	open injury	1.353(0.663-2.76)	0.406
smoking history	NO	Ref.	

	YES	0.872(0.462-1.646)	0.673
	NO	Ref.	
Alcohol consumption history			
	YES	0.536(0.21-1.37)	0.193
height	CV	0.985(0.951-1.02)	0.387
weight	CV	1.014(0.994-1.036)	0.176
BMI	CV	1.104(1.027-1.186)	0.007

607   Cause of injury (1 = Simple fall, 2 = Injury from electric bike or bicycle, 3 = Car accident injury, 4 = Injury

608   from falling from a height, 5 = Injury from sharp objects, 6 = Other injuries)

627   Table 2: Univariate Analysis of Surgical and Blood Transfusion Information

variables	targeted value	OR(95%CI)	P
	0	Ref.	
	1	1513860.09(0-Inf)	0.995
Incision grade	2	1043423.998(0-Inf)	0.995
	3	1(0-Inf)	1.000
	1	Ref.	
anesthesia method	2	0.705(0.166-2.992)	0.636
	3	0.859(0.354-2.085)	0.737
Anesthesia group	1	Ref.	
	2	0.979(0.468-2.046)	0.954
fracture reduction	1	Ref.	
	2	5.715(1.377-23.717)	0.016
	1	Ref.	
fixed form	2	0.302(0.041-2.24)	0.241
	3	0(0-Inf)	0.989
internal fixator	0	Ref.	

	1	2.841(0.376-21.487)	0.312
	2	4.138(0.436-39.26)	0.216
	3	1.905(0.204-17.786)	0.572
	4	4.118(0.52-32.607)	0.180
	5	0(0-Inf)	0.991
	6	0(0-Inf)	0.992
	7	0(0-Inf)	0.992
	8	2.386(0.282-20.2)	0.425
bone grafting	NO	Ref.	
	YES	1.905(0.711-5.106)	0.200
Bone graft type	0	Ref.	
	1	8.288(1.812-37.914)	0.006
	2	0.884(0.204-3.825)	0.869
time of operation	CV	1.004(1.002-1.007)	<0.001
Intraoperative blood loss	CV	1.001(1-1.001)	<0.001
ASA	1	Ref.	

	2	0.747(0.345-1.616)	0.459
	3	0.871(0.378-2.01)	0.747
	4	1.083(0.121-9.684)	0.943
Intraoperative blood transfusion	CV	1(1-1.001)	0.008
	0	Ref.	
Blood transfusion method	1	1.35(0.382-4.772)	0.641
	2	1.895(1.059-3.391)	0.031
	3	2.181(0.95-5.005)	0.066
Surgery timing	1	Ref.	
	2	0.816(0.105-6.338)	0.846
Input suspended red blood cells	CV	1(1-1.001)	<0.001
The total volume of blood transfused	CV	0.97(0.046-20.657)	0.984
The amount of plasma transfused	CV	1.001(1-1.001)	0.009
The amount of platelets transfused	0	Ref.	
	1	1.417(0.317-6.34)	0.648
	2	0(0-Inf)	0.981

Albumin supplements	0	Ref.	
	1	1.243(0.686-2.254)	0.474

Anesthesia method 1 = Local anesthesia (including brachial plexus, cervical plexus nerve block) 2 = Spinal

anesthesia (including intradural, extradural anesthesia, and combined spinal-epidural anesthesia) 3 = General

anesthesia; Anesthesia grouping: 1 local anesthesia; 2 general anesthesia; Reduction method 1 = Closed

reduction 2 = Open reduction (or approach); Fixation method 1 = Internal fixation 2 = External fixation 3 =

Conservative; Internal fixation device 0 = None 1 = Plate 2 = Intramedullary nail 3 = Kirschner wire 4 = Simple

screw 5 = Total hip prosthesis 6 = Semi-hip prosthesis 7 = Rivet 8 = Others; Bone graft type 0 = No bone

grafting 1 = Autologous bone 2 = Allogeneic bone; Blood transfusion method 0 = No blood transfusion 1 =

Autologous blood transfusion 2 = Allogeneic blood transfusion 3 = Autologous + allogeneic blood transfusion;

Surgery timing 1 = Daytime surgery 2 = Nighttime surgery

654 Table 3: Univariate Analysis of Laboratory Information

variables	targeted	OR(95%CI)	P
TP(g/L)	≥65	Ref.	
	< 65	1.178(0.697-1.991)	0.540
ALB(references 30–50 g/L)	30-50	Ref.	
	< 30 or > 50	1.305(0.769-2.212)	0.324
GLOB(references 20–40 g/L)	20-40	Ref.	
	< 20 or > 40	0.832(0.423-1.635)	0.593
A/G(references 1.2–2.4)	1.2-2.4	Ref.	
	< 1.2	0.588(0.293-1.182)	0.136
	> 2.4		0.980
ALT(references 9–50 U/L)	9-50	Ref.	
	< 9 or > 50	1.438(0.857-2.415)	0.169
AST(references 15–40 U/L)	15-40	Ref.	
	< 15 or > 40	1.632(0.974-2.733)	0.063
TBIL（references 1.7-17.1μmol/L）	1.7-17.1	Ref.	
	> 17.1	0.872(0.477-1.593)	0.655
DBIL (references 0-6.8 μmol/L)	0-6.8	Ref.	
	> 6.8	1.115(0.666-1.865)	0.679
IBIL (references1.7-10.2μmoL/L)	1.7-10.2	Ref.	
	> 10.2	0.859(0.47-1.569)	0.620
ALP(references 45–125 U/L)	45-125	Ref.	
	< 45	1.143(0.553-2.363)	0.719

	> 125	1.667(0.887-3.133)	0.113
GGT(references 10–60 U/L)	10-60	Ref.	
	< 10 or > 60	1.693(1.001-2.864)	0.050
CHE(references 5–12 U/L)	5-12	Ref.	
	< 5 or > 12	0.728(0.416-1.272)	0.265
TBA(references 1–10 μmol/L)	1-10	Ref.	
	< 1 or > 10	1.426(0.62-3.279)	0.404
HCRP(references 0-3mg/L)	≤3	Ref.	
	> 3	1.842(0.776-4.375)	0.166
LDH(references 0-250 U/L)	< 250	Ref.	
	≥250	1.421(0.825-2.448)	0.205
TC(references 0-5.2mmol/L)	≤5.2	Ref.	
	> 5.2	0(0-Inf)	0.981
TG(references 0-1.7mmol/L)	≤1.7	Ref.	
	> 1.7	0.823(0.395-1.712)	0.601
HDL-C(references 0.9-2mmol/L)	0.9-2	Ref.	
	< 0.9 or > 2	0.582(0.347-0.975)	0.040
LDL-C(references0-3.1mmol/L)	≤3.1	Ref.	
	> 3.1	0.777(0.301-2.002)	0.601
VLDL(references2-30mg/dL)	2-30	Ref.	
	< 2 or > 30	0.888(0.426-1.85)	0.751
APOB（references 0.8-1.2g/L）	0.8-1.2	Ref.	
	< 0.8	1.3(0.722-2.341)	0.381
	> 1.2	2.823(1.229-6.484)	0.014

	137-147	Ref.	
Na+(references 137–147 mmol/L	< 137	0.936(0.559-1.569)	0.803
	> 147	2.597(0.281-23.96)	0.400
	3.5-5.3	Ref.	
K+(references 3.5–5.3 mmol/L)	< 3.5	2.517(0.781-8.109)	0.122
	> 5.3	0.947(0.55-1.631)	0.844
	99-110	Ref.	
CL-(references 99–110 mmol/L)	< 99	1.13(0.604-2.116)	0.701
	> 110	2.153(0.709-6.54)	0.176
	20-30	Ref.	
TCO2(references 20–30 mmol/L)	< 20	3.289(1.49-7.258)	0.003
	> 30	0(0-Inf)	0.984
GLU(references3.9-6.1mmol/L)	< 6.1	Ref.	
	> 6.1	1.797(1.072-3.014)	0.026
	57-97	Ref.	
CREA(references 57–97 mmol/L)	< 57	1.062(0.484-2.333)	0.880
	> 97	1.411(0.703-2.833)	0.333
	8-16	Ref.	
GAP(references 8-16 mmol/L)	< 8	1.216(0.696-2.123)	0.492
	> 16	6.754(2.155-21.165)	0.001
	208-428	Ref.	
UA(references 208–428 mmol/L)	< 208	0.752(0.424-1.331)	0.328
	> 428	1.003(0.406-2.473)	0.995
CA(references 2.11–2.52	2.11-2.52	Ref.	

mmol/L)	< 2.11	1.1(0.658-1.836)	0.717
	> 2.52	0(0-Inf)	0.985
	0.85-1.51	Ref.	
P(references 0.85–1.51 mmol/L)	< 0 . 85	0.207(0.028-1.531)	0.123
	> 1.51	1.114(0.51-2.432)	0.787
	0.75-1.02	Ref.	
Mg(references 0.75–1.02 mmol/L)	< 0.75	1.581(0.81-3.087)	0.180
	> 1.02	1.091(0.249-4.79)	0.908
WBC(references 3.5–9.5 × 109/L)	3.5-9.5	Ref.	
	< 3.5 or > 9.5	1.113(0.664-1.865)	0.684
NEUT(references 2.8–6.3 × 109/L)	2.8-6.3	Ref.	
	< 2.8 or > 6.3	0.907(0.534-1.541)	0.718
LYM(references 1.1–3.2 × 109/L)	1.1-3.2	Ref.	
	< 1.1 or > 3.2	1.177(0.544-2.544)	0.679
MON(references 0.1–0.6 × 109/L)	0.1-0.6	Ref.	
	< 0.1	0(0-Inf)	0.989
	> 0.6	0.989(0.532-1.839)	0.973
EOS(references 0.02–0.05 × 109/L)	0.02-0.05	Ref.	
	< 0.02	1.869(1.057-3.307)	0.032
	> 0.05	0(0-Inf)	0.987
BAS	> 0.06	Ref.	
	≤0.06	1.065(0.134-8.456)	0.952
NEUT(references 45–75%)	45-75	Ref.	
	< 45 or > 75	1.224(0.7-2.141)	0.479

	20-40	Ref.	
LYM%references 20–40%	< 20	0.948(0.563-1.597)	0.841
	> 40	2.625(0.286-24.117)	0.394
MON%(references 3–10%)	3-10	Ref.	
	< 3 or > 10	1.303(0.703-2.413)	0.401
E0S(references 0.02–0.05 × 109/L)	0.02-0.05	Ref.	
	< 0.02	1.869(1.057-3.307)	0.032
	> 0.05	0(0-Inf)	0.987
RBC	normal	Ref.	
	abnormal	1.569(0.697-3.53)	0.277
HGB(references > 110/120g/L)	normal	Ref.	
	abnormal	2.013(0.898-4.513)	0.089
HCT(references 40-50%)	40-50	Ref.	
	< 40	1.276(0.565-2.884)	0.558
	> 50	0(0-Inf)	0.985
MCV(references 82–100 fL)	82-100	Ref.	
	< 82	1.571(0.348-7.087)	0.556
	> 100	1.375(0.522-3.622)	0.519
MCH(references 27–34 pg)	27-34	Ref.	
	< 27	1.819(0.397-8.324)	0.441
	> 34	1.213(0.417-3.526)	0.723
MCHC(references 316–354 g/L)	316-354	Ref.	
	< 316	3.083(0.836-11.375)	0.091
	> 354	1.528(0.577-4.042)	0.394

	11.6–16.5	Ref.	
RDW(references 11.6–16.5%)	< 11.6	0.556(0.073-4.23)	0.571
	> 16.5	1.078(0.413-2.813)	0.877
	125-350	Ref.	
PLT(references 125–350 × 10^9/L)	< 125	0.577(0.174-1.915)	0.369
	> 350	0.853(0.432-1.685)	0.647
	7.4-11	Ref.	
MPV(references 7.4–11.0 fL)	< 7.4	0.404(0.181-0.905)	0.028
	> 11	0.567(0.074-4.356)	0.585
	0.16-0.43	Ref.	
PCT(references 0.16–0.43%)	< 0.16	0.598(0.298-1.202)	0.149
	> 0.43		0.985
	12-18.1	Ref.	
PDW(references 12.0–18.1%)	< 12	0.668(0.28-1.596)	0.364
	> 18.1	2.82(0.904-8.797)	0.074
PT	CV	0.911(0.754-1.101)	0.336
PTA	CV	1.004(0.99-1.019)	0.553
INR	CV	0.298(0.034-2.606)	0.274
APTT(references 28–42 S)	CV	0.949(0.882-1.021)	0.160
APTT-R(references 28–42 S)	CV	0.093(0.008-1.064)	0.056
TT(references 28–42 S)	CV	1.109(0.987-1.245)	0.082
FIB(references 28–42 S)	CV	0.855(0.704-1.038)	0.113
AT□(references 28–42 S)	CV	1.01(0.995-1.024)	0.208
D-Dimer(> 2.26 mg/L)	CV	1(0.966-1.035)	0.986

FDP	CV	1.006(0.999-1.013)	0.105
TT-R	CV	5.408(0.865-33.79)	0.071

Comparison of preoperative laboratory indicators between the two groups of patients. TP - total protein, ALB -

albumin, GLOB - globulin, A/G value (albumin/globulin), ALT - alanine aminotransferase, AST - aspartate

aminotransferase, TBIL - total bilirubin, DBIL - direct bilirubin, IBIL - indirect bilirubin, ALP - alkaline

phosphatase, GGT - γ-glutamyl transpeptidase, CHE - phenol red enzyme, TBA - total bile acid, HCRP - high-

sensitivity C-reactive protein, LDH - lactate dehydrogenase, CREA - creatinine, UA - uric acid, CA - calcium, P

- phosphorus, Mg - magnesium, BNP - brain natriuretic peptide, WBC - white blood cells, NEUT - neutrophils,

LYM - lymphocytes, MON - monocytes, EOS - eosinophils, BAS - basophils, RBC - red blood cells. Reference

range: for females, $3.5–5.0 \times 10^{12}$/L; for males, $4.0–5.5 \times 10^{12}$/L. HGB - hemoglobin, reference range: for

females, 110–150 g/L; for males, 120–160 g/L. HCT - hematocrit, 40–50%, MCV - mean cell volume, MCH -

mean corpuscular hemoglobin content, MCHC - mean corpuscular hemoglobin concentration, RDW - red cell

distribution width, PLT - platelets, $100–300 \times 10^{9}$/L, MPV - mean platelet volume. Platelet volume, PCT

(procalcitonin), PDW (platelet distribution width), PT (prothrombin time), PTA (prothrombin activity), INR

(international normalized ratio), APTT (activated partial thromboplastin time), APTT-R (ratio of activated

partial thromboplastin time), TT (thrombin time), TT-R (thrombin ratio), FIB (fibrinogen), ATIII (antithrombin

669   3).