Prospective genomic and epidemiologic surveillance

of *Klebsiella pneumoniae* in a tertiary NICU

Carolin Böhne[1#], Claas Baier[2,3], Jelena Erdmann[4], Ella Ebadi[2], Carina Zirkler[1], Marc

Lindenberg[2], Dirk Schlüter[2,5], Sabine Pirr[1,5], Corinna Peter[1], Bettina Bohnhorst[1], Leonard

Knegendorf[2,4#]

[1] Department of Pediatric Pulmonology, Allergology and Neonatology, Hannover Medical School

(MHH), Carl-Neuberg Str. 1, 30625 Hannover, Germany

[2] Institute for Medical Microbiology and Hospital Epidemiology, Hannover Medical School

(MHH), Carl-Neuberg Str. 1, 30625 Hannover, Germany

[3] Infection Prevention and Control Department, AGAPLESION Diakonieklinikum Rotenburg,

Elise-Averdieck-Str. 17, 27356 Rotenburg, Germany

[4] Institute for Molecular Bacteriology, TWINCORE Centre for Experimental and Clinical Infection

Research, Feodor-Lynen-Str. 7, 30625 Hannover, Germany

[5] Cluster of Excellence RESIST (EXC 2155), Hannover Medical School (MHH), Carl-Neuberg

Str. 1, 30625 Hannover, Germany

[#]Corresponding authors: Carolin Böhne, E-Mail: boehne.carolin@mh-hannover.de; Leonard

Knegendorf, E-Mail: knegendorf.leonard@mh-hannover.de;

Running title: Genomic Surveillance of Klebsiella in Neonatal ICU

Manuscript word count: 2,849
Abstract word count: 242

Abstract

Klebsiella pneumoniae complex (Kp) is a relevant neonatal pathogen colonizing preterm infants. While outbreak investigations often focus on multidrug-resistant strains, the epidemiology and genomic dynamics of wild-type Kp in nonoutbreak neonatal intensive care unit (NICU) settings remain elusive.

We conducted a 30-month (October 2021 to March 2024) cohort study with weekly active, unselective colonization surveillance of all NICU patients to identify risk factors for nosocomial Kp acquisition and drivers of transmission in a tertiary 21-bed NICU/intermediate care unit (IMC) in Germany.

Among 936 patients, 8.7% carried Kp, of which 70.4% were nosocomial. Very low birth weight (VLBW; <1500 g) was the only independent risk factor for nosocomial acquisition (adjusted odds ratio [aOR], 3.42; 95% CI, 1.29-9.32). Kp infections occurred in three Kp carriers (3.7%). Genomic analyses of at least the first isolate per patient (83 in total) revealed an oligoclonal population structure, with distinct sequence types driving temporally overlapping clusters. Ten genomic clusters (median size, four patients) were identified, with markedly higher odds in VLBW infants (aOR, 8.76; 95% CI, 2.45-34.16). Nosocomial cluster-assigned cases had higher rates and longer durations of noninvasive ventilation and peripheral venous catheter use. Cluster prevalence showed climate-associated variation, with a six-feature extreme gradient boosting (XGBoost) model identifying temperature and humidity among the strongest predictors ($R^2 = 0.80$).

Patient- and climate-associated parameters are main drivers of nosocomial wild-type Kp acquisition and cluster occurrence. Comprehensive surveillance and risk-adapted infection prevention and control support sustainable Kp control in VLBW infants.

Importance

Klebsiella pneumoniae complex is a relevant cause of colonization and infection in preterm and very low birth weight infants. While genomic analyses in neonatal care often focus on multidrug-resistant strains or defined outbreaks, little is known about the transmission dynamics of wild-type *K. pneumoniae* in routine clinical settings. This study integrates regular colonization screening with prospective whole-genome sequencing to characterize acquisition patterns in a high-risk neonatal intensive care environment. By combining clinical, microbiological, genomic, and environmental information, we show how specific patient factors and macroclimatic conditions influence the occurrence of genomic clusters. The findings illustrate how integrating genomic information into routine diagnostics can support earlier recognition of potential transmission events and inform targeted infection prevention and control measures in neonatal care.

List of abbreviations (alphabetical order)

aOR   adjusted odds ratio

BV-BRC   Bacterial and Viral Bioinformatics Resource Center

cgMLST   core genome multi-locus sequence typing

CI   confidence interval

ESBL   extended-spectrum β-lactamase

GA   gestational age

hvKp   hypervirulent *Klebsiella pneumoniae* complex

ICU   intensive care unit

IMC   intermediate care unit

IPC   infection prevention and control

Kp   *Klebsiella pneumoniae* complex

KRINKO   Commission for Infection Prevention and Hygiene in Healthcare and Nursing

at the Robert Koch Institute, Germany (*Kommission für Infektionsprävention in*

medizinischen Einrichtungen und in Einrichtungen und Unternehmen der

Pflege und Eingliederungshilfe)

MHH   Hannover Medical School (Medizinische Hochschule Hannover)

MRSA   Methicillin-resistant *Staphylococcus aureus*

NICU   neonatal intensive care unit

PHS   patient hospital stay

SHAP   Shapley Additive Explanations

SKA   split k-mer analysis

SNP   single nucleotide polymorphism

VLBW   very low birth weight

VRE   Vancomycin-resistant enterococci

WGS   whole genome sequencing

XGBoost   extreme gradient boosting

Introduction

The Gram-negative *Klebsiella pneumoniae complex* (Kp) is known for colonizing preterm infants in neonatal intensive care units (NICUs)[1] and is one of the pathogens frequently leading to neonatal infections[2,3]. Besides patient-related factors and the use of invasive devices and procedures[4-6], infection susceptibility is further influenced by pathogen-related attributes such as hypervirulent Kp isolates (hvKp)[7]. A gene-based grading system (Kleborate)[8] and a hypermucoviscous Kp phenotype (positive string test[9] or tellurite resistance[10]), may indicate increased virulence. Hypervirulence may also coincide with antibiotic resistance, including Carbapenem resistance[11].

Kp can persist in the environment (e.g. surfaces, medical products), thus being easily transmittable in NICUs[12,13] and resulting in several outbreak reports from NICUs worldwide[13-16], including Carbapenem-resistant Kp[17,18]. Since neonatal infections still represent a major cause of neonatal morbidity and mortality even in western, industrialized countries[19], close epidemiologic and molecular Kp monitoring in NICUs seems crucial from a clinical and infection prevention and control (IPC) perspective. In Germany, weekly microbiological colonization screening is mandatory in NICUs, following national recommendations of the Commission for Infection Prevention and Hygiene in Healthcare and Nursing (KRINKO)[20,21]. To clarify transmission pathways, whole-genome sequencing (WGS) of Kp[22], including typing methods such as core genome multi-locus sequence typing (cgMLST)[23,24], has been increasingly used. However, data on Kp transmission and infection apart from outbreak scenarios are still scarce. Therefore, the objective of our study was to identify patient-related risk factors for nosocomial Kp acquisition in a non-outbreak, tertiary NICU/intermediate care (IMC) setting. The integration of epidemiological Kp surveillance and prospective IPC measures with a routine phenotypic and genomic analysis of Kp isolates, irrespective of resistance profile or outbreak suspicion, enabled us to assess Kp cluster prevalence. Furthermore, we contextualized the genomic features of Kp isolates with public genomes and explored the associations of Kp cluster prevalence with intensive care level and climate conditions using interpretable machine learning.

116 This comprehensive approach aimed to improve our understanding of the pathogen-patient-

117 environment interactions as a basis for strengthening IPC strategies in the future.

Methods

Study type

We conducted a retrospective cohort study of our prospective surveillance program including all

patients admitted to the tertiary 21-bed NICU/IMC of Hannover Medical School (MHH) from

October 1, 2021, to March 31, 2024. The primary outcome was nosocomial Kp acquisition,

defined as detection of Kp on or after day 3 of NICU/IMC stay. The secondary outcome was the

prevalence of Kp clusters defined by genomic sequencing. Exploratory analyses compared

nosocomial vs community-acquired cases and cluster-assigned vs non-cluster-assigned

nosocomial cases. In addition, we characterized phenotypic and genomic features of all Kp

isolates. Ethical approval was obtained (numbers 9365_BO_K_2020 and 10372_BO_K_2022),

and informed consent was waived by the ethics committee. The study adhered to the

Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) reporting

guideline[25]. The ward setting, definitions, and data acquisition are detailed in the eMethods.

In brief, a patient hospital stay (PHS) was defined as continuous inpatient episode on the

NICU/IMC. The time at risk was defined as the number of days from admission to first detection

of Kp in any microbiological sample.

IPC concept and surveillance

Detailed information on our IPC concept, Kp surveillance program, and microbiological and

molecular diagnostics are available in the eMethods. In short, following microbiological

admission screening, all NICU patients and IMC patients with an actual weight <1500g received

weekly colonization screening of respiratory secretions and rectal swabs according to

KRINKO[20,21]. Microbiological diagnostics were conducted at the ISO 15189-accredited

microbiology laboratory of MHH, and each newly emerged Kp isolate was sequenced using

Illumina short-read technology.

Statistical analysis and machine learning

To reduce immortal-time bias, the *at-risk* population for nosocomial Kp acquisition was defined

as PHSs lasting ≥48 hours with negative Kp status until that time. We performed multivariable

logistic regression to estimate adjusted odds ratios (aOR) with 95% profile-likelihood confidence

intervals (CIs) for nosocomial acquisition, including gestational age (GA), very low birth weight

(<1500g, VLBW, categorical), and sex as covariates (reference categories: absence of VLBW,

female sex). Screening effort (number of surveillance samples per patient divided by the

individual time at risk) was also included as a covariate to account for potential detection bias.

An α level of .05 was used. Epidemiological and clinical variables were summarized as median

(interquartile range [IQR]) or number (percentage). Group differences were tested using the chi-

square test for categorical and Wilcoxon rank-sum test for continuous variables, with

significance at *P*<.05. GraphPad Prism (version 10.2.0; GraphPad Software, Massachusetts,

USA) was used for statistical analyses. Reporting of machine learning models followed the

TRIPOD+AI statement[26]. Detailed information on machine learning analyses is available in the

eMethods. Machine learning analyses modeled cluster prevalence as the outcome in relation to

patient-care metrics and climate factors. Hyperparameters were tuned to optimize model fit and

interpretability, and models were used in a hypothesis-generating rather than predictive

capacity. Model hyperparameters are detailed in eTable 1.

Results

Epidemiology and primary outcome

During the study period, 936 individual patients (median: 51 patients/month) were admitted to

the NICU/IMC, generating 1009 PHSs and 14 684 patient days (median PHS length, 6 days)

(**Table 1**). Microbiological diagnostics identified 81 Kp patients (prevalence, 8.7/100 patients;

incidence density, 5.6 PHS/1000 patient days). Numbers differ slightly between patients, PHSs,

and isolates due to multiple acquisitions in single patients (details in Table footnotes).

The mean (IQR) birth weight of all 936 NICU/IMC patients was 2503 (1610 – 3286) g, the mean

(IQR) GA was 36.3 (32.3 –39.0) weeks. VLBW infants accounted for 21.6% of all patients

(202/936) and were threefold more frequent in the Kp subgroup. Among Kp patients, 70.4%

(57/81) had nosocomial acquisition, of whom 66.7% (38/57) were VLBW infants. In multivariable

logistic regression, VLBW was the only strong risk factor for nosocomial Kp acquisition (aOR,

3.42; 95% CI, 1.29–9.32), whereas higher gestational age was weakly protective (aOR, .98;

95% CI, .97-.9986). After adjustment for screening effort, which was itself strongly associated

with Kp acquisition, the association of VLBW remained robust (**Table 1**). Exploratory analyses

showed that nosocomial cases, compared with community-acquired cases, had a significantly

higher proportion of VLBW infants, lower GA, and lower birth weight (eTable 2). During the time

at risk, nosocomial cases also showed a high proportion of peripheral venous catheters

(96.6%), noninvasive ventilation (75.9%), and intravenous antibiotic therapy (43.1%; eTable 2).

Three patients (3.7%) developed a Kp infection (**Table 2**). One patient developed infection

postoperatively following colonization acquired at a transferring hospital, whereas in the two

other patients the infection occurred during prolonged hospitalizations; both were profoundly

immunocompromised, including one solid organ transplant recipient.

188 Nosocomially acquired Kp and molecular cluster assignment

Eighty-three isolates from 79 patients were available for molecular analysis. 40 patients

(40/81 [49.4%]) were assigned to 10 genomic clusters, each comprising two to ten patients.

Cluster assignment was more frequent among nosocomial Kp patients (37/57 [64.9%]) than

among community-acquired Kp patients (3/24 [12.5%]). In multivariable analysis, VLBW was

associated with substantially higher odds of cluster assignment (aOR, 8.76; 95% CI, 2.45-

34.16; **Table 1**).

Exploratory subgroup analyses showed that nosocomial cluster-assigned cases had more

often a VLBW and had higher rates of noninvasive ventilation compared with non-cluster

assigned cases (**Table 3**)**.** Moreover, the duration of noninvasive ventilation and peripheral

venous catheterization was significantly longer in cluster-assigned cases (eTable 3).

200   Integrated phenotypic and genomic analysis

Phenotypic hypervirulence markers yielded one positive string test and two indeterminate

tellurite results; none belonged to a genomic cluster or caused infection. Kleborate analysis did

not identify hypervirulent strains. Three isolates within cgMLST cluster 8 carried yersiniabactin

and colibactin, but none caused infection (eTable 4). Most isolates (63/83) exhibited a wild-type

antimicrobial resistance profile; 10 had an extended-spectrum β-lactamase (ESBL) phenotype,

of which eight belonged to clusters (ST29 and ST353, four isolates each, all blaCTX-M-15-

positive; **Figure 1**). No carbapenem resistance was detected. All cgMLST clusters displayed

consistent phenotypic profiles, with a single discrepant isolate in cluster 1 (Split k-mer analysis

[SKA] distance >20 single nucleotide polymorphisms [SNPs]).

SNP comparison against public genomes (Bacterial and Viral Bioinformatics Resource Center

[BV-BRC] database)[27] revealed the closest match (48 SNPs) to a marine isolate from Norway[28];

this strain (ST866) was identified upon admission in a patient residing at the German North Sea

coast (eTable 5).

Throughout the study, molecular typing results were presented during regular interdisciplinary

IPC audits. The median time from sampling to completed WGS was 3.1 weeks (IQR, 2.6–4.6).

217   Association of climate and patient-care factors with

218   molecular cluster prevalence

Kp cluster prevalence showed seasonal variations (eFigure 1). To examine external drivers,

machine learning models were trained with climate variables and nurse-to-patient ratios,

including 4-week lagged values based on exploratory analyses suggesting temporal

associations.

Linear models (elastic net, lasso, and ridge) showed limited predictive ability (eTable 1). An

extreme gradient boosting (XGBoost) model using a reduced feature set provided the best fit

to the observed variance (R^2 = 0.80). A random forest model achieved a comparable fit (R^2 =

0.78) but required inclusion of 24 features (**Figure 2A;** full feature set in eMethods). The

largest discrepancy between observed and predicted prevalence occurred in August 2023,

mainly attributable to cluster 5 (ST20).

Feature importance was evaluated using gain-based metrics and Shapley Additive

Explanations (SHAP). A higher minimum outdoor temperature four weeks earlier (lagged)

consistently emerged as the top feature across seasons – weeks with little or no cooling

phases (warmer nights) were followed by higher cluster prevalence. Humidity showed a

complex, bidirectional influence on predicted outcomes across its observed range and

exerted stronger relative effects in winter and autumn, while the VLBW nurse-to-patient ratio,

correlated with higher predicted prevalence, contributed more modestly but stably over time

(**Figure 2B-C**, eFigure 2). Learning curves indicated moderate overfitting and potential gains

from additional data (eFigure 3).

239   Discussion

In this study, we comprehensively analyzed epidemiological and genomic Kp surveillance at a

tertiary NICU/IMC in a non-outbreak setting, including an assessment of risk factors for

nosocomial Kp acquisition and an evaluation of cluster prevalence using machine learning.

Kp was detected in ~8-9% of hospitalized patients, consistent with 2016-2018 data from our

unit[29], and about ten times higher than local MRSA prevalence[30]. In contrast, intestinal Kp

carriage was 40-60% in Indian VLBW preterm infants[31], while adult intensive care unit (ICU)

colonization rates ranged between 6-20%[32,33]. More than two thirds of Kp cases in our NICU

were nosocomial, consistent with studies showing that the initial microbiome in preterm infants

is shaped by the unit's environment, including other infants, surfaces, staff and parents[34-36].

Although VLBW patients accounted for only ~21% of all PHSs, VLBW was the only independent

factor for nosocomial acquisition, with 67% of nosocomial cases and 81% of cluster-associated

PHSs occurring in VLBW infants.

Kp is a leading cause of neonatal sepsis following intestinal colonization[37-39]. Known infection

risk factors include patient-related factors, invasive procedures, and prior antibiotic use[40],

although pathomechanisms remain elusive[32]. We observed increased rates and durations of

noninvasive ventilation and peripheral venous catheter use in our nosocomial cluster-associated

patients. Peripheral catheters are a recognized risk factor for nosocomial bloodstream infections

in VLBW patients[41] and have been addressed in German IPC recommendations[20]. However,

given the longer time at risk in nosocomial patients, these findings should be interpreted

cautiously, as they may reflect opportunity bias.

Most NICU Kp reports focus on multidrug-resistant or hypervirulent strains during outbreaks[12-

15,17]. In contrast, our colonization screening explicitly addresses also the carriage of susceptible

(wild-type) Kp isolates. No hypervirulent or carbapenem-resistant isolates were identified,

suggesting that classical Kp remains the predominant colonizer in NICUs[15,42,43]. Although

phenotypic markers such as string test and tellurite resistance may yield false positives, their

use enabled real-time triaging and IPC adjustments before sequencing results were available.

ESBL production was rare and largely confined to two clusters (ST29 and ST353), both

harboring blaCTX-M-15, consistent with sporadic import rather than endemic persistence.

Genomic consistency within clusters – observed with both cgMLST and SKA – supports

cgMLST-based IPC interventions without isolate-specific genotyping. However, the identification

of one isolate with divergent resistance within a cgMLST-defined cluster (SNP distance >20)

underscores the value of SNP-level analysis to refine cluster boundaries[44]. Interestingly, the

closest genomic match for one isolate (ST866) was a marine environmental strain from Norway,

illustrating potential environmental influx and the value of contextualizing local isolates against

public datasets.

While the spread of hospital-associated, carbapenem-resistant lineages of Kp is well-

documented[45], we found no clusters involving globally circulating lineages. Consistent with the

absence of hvKp/carbapenem resistance and rapid containment, the infection attack rate of

3.7% of all Kp patients (0.3% of admissions) was comparable to prior reports[6,39].

Kp was repeatedly introduced and also transmitted, but cluster size remained small (median,

four patients), suggesting that transmission was usually limited in size and duration, consistent

with timely IPC interventions. These included training/feedback, cohorting/isolation, encouraging

hand hygiene, intensified patient and environmental screening, and enhanced disinfection[12-15,17].

Audit-style feedback on epidemiological and genomic findings was progressively strengthened

during the study, reaching near-weekly intervals, with rising interdisciplinary participation.

Similar formats have proven effective in teaching IPC[46]. Clusters detected by molecular typing

were addressed systematically during audits and in scheduled or ad hoc interdisciplinary

training sessions, prompting targeted interventions to interrupt transmission chains and prevent

establishment of dominant 'ward clones'.

Six pairs of twins were included among cluster patients, highlighting elevated risks for patient-

to-patient or parent-to-patient transmission[35,36]. Though skin-to-skin (kangaroo) care in VLBW

infants facilitates bacterial exchange, it remains a standard practice that favors

neurodevelopment and microbiome stability[47,48]. From an IPC perspective, awareness of this

potential transmission route in preterm multiples is essential.

Seasonal variation in Kp cluster prevalence, despite stable NICU occupancy, prompted further

investigation using machine learning models. Tree-based models, particularly a reduced-feature

XGBoost model, explained up to 80% of variance in weekly prevalence, while linear models

failed to capture nonlinear relationships. These findings suggest that Kp transmission in NICUs

is shaped not only by patient-specific factors, but also by structural and environmental variables

with delayed effects.

A higher minimum daily outdoor temperature four weeks earlier was the most influential

predictor, consistent with multicenter studies linking temperature with bloodstream infection

incidence and environmental Kp abundance[49-52]. In particular, Schwab *et al.* reported that

nosocomial bloodstream infection rates in German ICUs were approximately 16% higher in

months with mean temperatures ≥20 °C compared with <5 °C, with the strongest association

observed for temperatures in the preceding month[49]. Additional drivers in our data included

relative humidity and increased intensive-care density/capacity, with the VLBW nurse-to-patient

ratio serving as a surrogate for patient-related risk. We hypothesize that climate conditions may

influence staff and visitor colonization on one hand and accumulation of Kp in the ward

environment (surfaces, devices) on the other hand. The importance of lagged features appears

biologically and operationally plausible given the interval from increased bacterial load to

colonization and detectability in surveillance cultures.

These models were exploratory and used to quantify associations, not to predict future risk.

Nonetheless, they offer a framework for hypothesis generation and provide actionable insights

for IPC planning, such as anticipating periods of increased transmission risk. Nonlinear

relationships, such as the bidirectional effect of humidity, underline the complexity of these

dynamics. Expansion of an ST20 cluster despite predictively unfavorable external influences

may suggest clone-specific transmissibility, although we could not find literature specifically

supporting this.

Because molecular results were delayed, findings mainly informed medium- to long-term IPC

strategies in our setting. Streamlined workflows (e.g. usage of Oxford Nanopore Sequencing

technologies[53,54] or more rapid Illumina sequencing) may allow more actionable short-term

interventions in the future.

324   Limitations

325   This single-center retrospective cohort study includes weekly colonization screening

326   mandated by German IPC regulations, which may increase awareness of bacterial carriage

327   and antibiotic use in preterm infants. Therefore, it may not be directly generalizable to other

328   NICU settings. Routine Kp sequencing was subject to a time lag, limiting its utility for

329   immediate IPC interventions. While we adjusted the primary analysis to reduce immortal time

330 bias, differences in observation periods between nosocomial and community-acquired cases

331 may have introduced opportunity bias. The machine learning models were not externally

332 validated, and prone to overfitting, requiring larger multicenter datasets for predictive use.

333 The study period was relatively short for drawing climate-related conclusions. In addition, the

334 effect of daily outdoor temperature, rather than indoor temperature, on cluster prevalence

335 was analyzed. Practical observations on the ward suggest that, despite the presence of

336 ventilation systems, outdoor temperature affected indoor thermal conditions. Additional

337 potential transmission vectors – such as staff and visitor carriage or environmental reservoirs

338 – were not systematically assessed.

340 Conclusions

341 In a tertiary NICU/IMC, prospective genomic surveillance within a structured IPC program

342 showed that Kp transmission was mainly driven by wild-type isolates. Infection rates were

343 uncommon and genomic clusters remained small, consistent with the effect of routine

344 screening and sequencing-informed audits with rapid feedback into practice. VLBW infants

345 were the main risk group for nosocomial acquisition and were disproportionately affected by

346 cluster-associated Kp. Cluster prevalence reflected the combined influence of VLBW patient

347 vulnerability, structural, and environmental influences, with higher rates observed following

348 periods of sustained warmer nights. These findings outline a scalable surveillance-to-action

349 approach that may help further limit transmission in highly vulnerable neonatal populations.

350 The identified time lag suggests a planning horizon for risk-triggered IPC: timely staff

351 refreshers, increased environmental disinfection, focused screening, and the preemptive

352 reverse isolation of infants at highest risk during anticipated higher-risk periods could be

353 measures worth evaluating their efficacy in future studies.

References

1. Moussa B, Oumokhtar B, Arhoune B, et al. Gut acquisition of Extended-spectrum β-lactamases-producing Klebsiella pneumoniae in preterm neonates: Critical role of enteral feeding, and endotracheal tubes in the neonatal intensive care unit (NICU). *PLoS One.* 2023;18(11):e0293949.

2. Magill SS, Edwards JR, Bamberg W, et al. Multistate point-prevalence survey of health care-associated infections. *N Engl J Med.* 2014;370(13):1198-1208.

3. Vogiantzi G, Metallinou D, Tigka M, Deltsidou A, Nanou CI. Bloodstream Infections in the Neonatal Intensive Care Unit: A Systematic Review of the Literature. *Cureus.* 2024;16(8):e68057.

4. Sampah MES, Hackam DJ. Dysregulated Mucosal Immunity and Associated Pathogeneses in Preterm Neonates. *Front Immunol.* 2020;11:899.

5. Humberg A, Fortmann I, Siller B, et al. Preterm birth and sustained inflammation: consequences for the neonate. *Semin Immunopathol.* 2020;42(4):451-468.

6. Wei X, Liang J, Zhang H, et al. Clinical features and risk factors of Klebsiella pneumoniae infection in premature infants: a retrospective cohort study. *BMC Infect Dis.* 2024;24(1):1311.

7. de Lajartre OB, Maamar A, Dejoies L, Delamaire F. Community-acquired hypervirulent Klebsiella pneumoniae invasive infection in critically ill patients who dramatically improved. *Eur J Clin Microbiol Infect Dis.* 2022;41(10):1275-1277.

8. Lam MMC, Wick RR, Watts SC, Cerdeira LT, Wyres KL, Holt KE. A genomic surveillance framework and genotyping tool for Klebsiella pneumoniae and its related species complex. *Nat Commun.* 2021;12(1):4188.

9. Watanabe N, Masuda A, Watari T, Otsuka Y, Yamagata K, Fujioka M. Evaluation of an optimal agar medium for detecting hypervirulent Klebsiella pneumoniae using string test. *Access Microbiol.* 2024;6(9).

10. Wu X, Zhan F, Zhang J, Chen S, Yang B. Identification of hypervirulent Klebsiella pneumoniae carrying terW gene by MacConkey-potassium tellurite medium in the general population. *Front Public Health.* 2022;10:946370.

11. Lan P, Jiang Y, Zhou J, Yu Y. A global perspective on the convergence of hypervirulence and carbapenem resistance in Klebsiella pneumoniae. *J Glob Antimicrob Resist.* 2021;25:26-34.

12. Shi Q, Zhao J, Wei L, et al. Transmission of ST45 and ST2407 extended-spectrum β-lactamase-producing Klebsiella pneumoniae in neonatal intensive care units, associated with contaminated environments. *J Glob Antimicrob Resist.* 2022;31:309-315.

13. Baek EH, Kim SE, Kim S, et al. Successful control of an extended-spectrum beta-lactamase-producing Klebsiella pneumoniae ST307 outbreak in a neonatal intensive care unit. *BMC Infect Dis.* 2020;20(1):166.

14. Priante E, Minotti C, Contessa C, et al. Successful Control of an Outbreak by Phenotypically Identified Extended-Spectrum Beta-Lactamase-Producing Klebsiella pneumoniae in a Neonatal Intensive Care Unit. *Antibiotics (Basel).* 2022;11(11).

15. Mavroidi A, Liakopoulos A, Gounaris A, et al. Successful control of a neonatal outbreak caused mainly by ST20 multidrug-resistant SHV-5-producing Klebsiella pneumoniae, Greece. *BMC Pediatr.* 2014;14:105.

16. Wisgrill L, Lepuschitz S, Blaschitz M, et al. Outbreak of Yersiniabactin-producing Klebsiella pneumoniae in a Neonatal Intensive Care Unit. *Pediatr Infect Dis J.* 2019;38(6):638-642.

17. Pathak A, Tejan N, Dubey A, et al. Outbreak of colistin resistant, carbapenemase (bla (NDM), bla (OXA-232)) producing Klebsiella pneumoniae causing blood stream infection among neonates at a tertiary care hospital in India. *Front Cell Infect Microbiol.* 2023;13:1051020.

18. Magobo RE, Ismail H, Lowe M, et al. Outbreak of NDM-1- and OXA-181-Producing
Klebsiella pneumoniae Bloodstream Infections in a Neonatal Unit, South Africa.
Emerg Infect Dis. 2023;29(8):1531-1539.
19. Li J, Shen L, Qian K. Global, regional, and national incidence and mortality of
neonatal sepsis and other neonatal infections, 1990-2019. *Front Public Health.*
2023;11:1139832.
20. KRINKO. Praktische Umsetzung sowie krankenhaushygienische und
infektionspra□ventive Konsequenzen des mikrobiellen Kolonisationsscreenings bei
intensivmedizinisch behandelten Fru□h- und Neugeborenen
Erga□nzende Empfehlung der KRINKO beim Robert Koch-Institut, Berlin, zur
Implementierung der Empfehlungen zur Pra□vention nosokomialer Infektionen bei
neonatologischen Intensivpflegepatienten mit einem Geburtsgewicht unter 1.500 g
aus dem Jahr 2007 und 2012. *Epidem Bull.* 2013;42:421-433.
21. Litz JE, Goedicke-Fritz S, Härtel C, Wagenpfeil G, Zemlin M, Simon A. Umsetzung
des mikrobiologischen Kolonisationsscreenings: Umfrage an 80 neonatologischen
Intensivstationen. *Epid Bull.* 2019;37:387 – 392.
22. Carlos CC, Masim MAL, Lagrada ML, et al. Genome Sequencing Identifies Previously
Unrecognized Klebsiella pneumoniae Outbreaks in Neonatal Intensive Care Units in
the Philippines. *Clin Infect Dis.* 2021;73(Suppl_4):S316-s324.
23. Gona F, Comandatore F, Battaglia S, et al. Comparison of core-genome MLST,
coreSNP and PFGE methods for Klebsiella pneumoniae cluster analysis. *Microb*
Genom. 2020;6(4).
24. Hallbäck ET, Johnning A, Myhrman S, et al. Outbreak of OXA-48-producing
Enterobacteriaceae in a neonatal intensive care unit in Western Sweden. *Eur J Clin*
Microbiol Infect Dis. 2023;42(5):597-605.
25. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The
Strengthening the Reporting of Observational Studies in Epidemiology (STROBE)
statement: guidelines for reporting observational studies. *Ann Intern Med.*
2007;147(8):573-577.
26. Collins GS, Moons KGM, Dhiman P, et al. TRIPOD+AI statement: updated guidance
for reporting clinical prediction models that use regression or machine learning
methods. *BMJ.* 2024;385:e078378.
27. Olson RD, Assaf R, Brettin T, et al. Introducing the Bacterial and Viral Bioinformatics
Resource Center (BV-BRC): a resource combining PATRIC, IRD and ViPR. *Nucleic*
Acids Res. 2023;51(D1):D678-d689.
28. Håkonsholm F, Hetland MAK, Svanevik CS, Lunestad BT, Löhr IH, Marathe NP.
Insights into the genetic diversity, antibiotic resistance and pathogenic potential of
Klebsiella pneumoniae from the Norwegian marine environment using whole-genome
analysis. *Int J Hyg Environ Health.* 2022;242:113967.
29. Baier C, Pirr S, Ziesing S, et al. Prospective surveillance of bacterial colonization and
primary sepsis: findings of a tertiary neonatal intensive and intermediate care unit. *J*
Hosp Infect. 2019;102(3):325-331.
30. Böhne C, Knegendorf L, Schwab F, et al. Epidemiology and infection control of
Methicillin-resistant Staphylococcus aureus in a German tertiary neonatal intensive
and intermediate care unit: A retrospective study (2013-2020). *PLoS One.*
2022;17(9):e0275087.
31. Sharma N, Chaudhry R, Panigrahi P. Quantitative and qualitative study of intestinal
flora in neonates. *J Glob Infect Dis.* 2012;4(4):188-192.
32. Martin RM, Cao J, Brisse S, et al. Molecular Epidemiology of Colonizing and Infecting
Isolates of Klebsiella pneumoniae. *mSphere.* 2016;1(5).
33. Gorrie CL, Mirceta M, Wick RR, et al. Gastrointestinal Carriage Is a Major Reservoir
of Klebsiella pneumoniae Infection in Intensive Care Patients. *Clin Infect Dis.*
2017;65(2):208-215.
34. Brooks B, Olm MR, Firek BA, et al. The developing premature infant gut microbiome
is a major factor shaping the microbiome of neonatal intensive care unit rooms.
Microbiome. 2018;6(1):112.

35.   Brooks B, Firek BA, Miller CS, et al. Microbes in the neonatal intensive care unit
resemble those found in the gut of premature infants. *Microbiome.* 2014;2(1):1.
36.   Conceição T, Aires de Sousa M, Miragaia M, et al. Staphylococcus aureus reservoirs
and transmission routes in a Portuguese Neonatal Intensive Care Unit: a 30-month
surveillance study. *Microb Drug Resist.* 2012;18(2):116-124.
37.   Schwartz DJ, Shalon N, Wardenburg K, et al. Gut pathogen colonization precedes
bloodstream infection in the neonatal intensive care unit. *Sci Transl Med.*
2023;15(694):eadg5562.
38.   Moftian N, Rezaei-Hachesu P, Arab-Zozani M, et al. Prevalence of gram-negative
bacteria and their antibiotic resistance in neonatal sepsis in Iran: a systematic review
and meta-analysis. *BMC Infect Dis.* 2023;23(1):534.
39.   Reichert F, Piening B, Geffers C, Gastmeier P, Bührer C, Schwab F. Pathogen-
Specific Clustering of Nosocomial Blood Stream Infections in Very Preterm Infants.
Pediatrics. 2016;137(4).
40.   Olsen AL, Reinholdt J, Jensen AM, Andersen LP, Jensen ET. Nosocomial infection in
a Danish Neonatal Intensive Care Unit: a prospective study. *Acta Paediatr.*
2009;98(8):1294-1299.
41.   Geffers C, Gastmeier A, Schwab F, Groneberg K, Rüden H, Gastmeier P. Use of
central venous catheter and peripheral venous catheter as risk factors for nosocomial
bloodstream infection in very-low-birth-weight infants. *Infect Control Hosp Epidemiol.*
2010;31(4):395-401.
42.   Mukherjee S, Mitra S, Dutta S, Basu S. Neonatal Sepsis: The Impact of Carbapenem-
Resistant and Hypervirulent Klebsiella pneumoniae. *Front Med (Lausanne).*
2021;8:634349.
43.   Hu Y, Yang Y, Feng Y, et al. Prevalence and clonal diversity of carbapenem-resistant
Klebsiella pneumoniae causing neonatal infections: A systematic review of 128
articles across 30 countries. *PLoS Med.* 2023;20(6):e1004233.
44.   Wang J, Feng Y, Zong Z. Worldwide transmission of ST11-KL64 carbapenem-
resistant Klebsiella pneumoniae: an analysis of publicly available genomes. *mSphere.*
2023;8(4):e0017323.
45.   David S, Reuter S, Harris SR, et al. Epidemic of carbapenem-resistant Klebsiella
pneumoniae in Europe is driven by nosocomial spread. *Nat Microbiol.*
2019;4(11):1919-1929.
46.   Mohammedi SB, Gillois P, Landelle C. Nursing students'knowledge and effectiveness
of teaching in infection prevention and control. *BMC Nurs.* 2025;24(1):850.
47.   Altit G, Hamilton D, O'Brien K. Skin-to-skin care (SSC) for term and preterm infants.
Paediatr Child Health. 2024;29(4):238-254.
48.   Hendricks-Muñoz KD, Xu J, Parikh HI, et al. Skin-to-Skin Care and the Development
of the Preterm Infant Oral Microbiome. *Am J Perinatol.* 2015;32(13):1205-1216.
49.   Schwab F, Gastmeier P, Hoffmann P, Meyer E. Summer, sun and sepsis-The
influence of outside temperature on nosocomial bloodstream infections: A cohort
study and review of the literature. *PLoS One.* 2020;15(6):e0234656.
50.   Anderson DJ, Richet H, Chen LF, et al. Seasonal variation in Klebsiella pneumoniae
bloodstream infection on 4 continents. *J Infect Dis.* 2008;197(5):752-756.
51.   Eber MR, Shardell M, Schweizer ML, Laxminarayan R, Perencevich EN. Seasonal
and temperature-associated increases in gram-negative bacterial bloodstream
infections among hospitalized patients. *PLoS One.* 2011;6(9):e25298.
52.   Aloufi AS. MaxEnt modeling of Klebsiella pneumoniae: predicting future distribution
and evaluating the risk for public health. *Geomatics, Natural Hazards and Risk.*
2024;15(1).
53.   Foster-Nyarko E, Cottingham H, Wick RR, et al. Nanopore-only assemblies for
genomic surveillance of the global priority drug-resistant pathogen, Klebsiella
pneumoniae. *Microb Genom.* 2023;9(2).
54.   Wagner GE, Dabernig-Heinz J, Lipp M, et al. Real-Time Nanopore Q20+ Sequencing
Enables Extremely Fast and Accurate Core Genome MLST Typing and Democratizes

Access to High-Resolution Bacterial Pathogen Surveillance. *J Clin Microbiol.* 2023;61(4):e0163122.

Acknowledgements

The authors thank all technical staff of the medical microbiology and hospital hygiene

laboratories involved in bacterial isolation, phenotypic virulence testing, strain collection, and

library preparation. In addition, the authors would like to acknowledge Dr. Ludwig Sedlacek

and PD Dr. Marius Vital, who were supervising the routine sequencing workflow in the

medical microbiology laboratory. Furthermore, the authors are very grateful to all clinical staff

and participating families who contributed to this study.

Conflict of Interest Statement

The authors have no conflicts of interest to declare.

Funding Sources

This study did not receive any external funding.

Author Contributions

CBO, CB, LK: conceptualization, methodology, data acquisition, formal analysis, validation,

writing original draft and review and editing

JE, ML: formal analysis, validation, draft review and editing

CZ: data acquisition, validation, draft review and editing

EE, DS: validation, draft review and editing

SP, CP, BB: validation, draft review and editing

Data Availability Statement

Patient data used in this study are confidential in accordance with the German Data Privacy

Act, the ethics committee, and the data protection commissioner of Hannover Medical

School. Patient-related information such as ward of admission, age, sex, underlying disease,

or length of stay constitutes indirect identifiers that could potentially allow re-identification. To

ensure confidentiality and protect privacy, only anonymized and aggregated data are

available for research use, in compliance with the Data Privacy Act. Interested researchers

may contact the corresponding authors to request access to anonymized data, subject to

approval by the data protection commissioner of Hannover Medical School. All isolate-level

metadata used in this study are provided in the Online Supplement (eTable 4-5). Whole-

genome sequencing data have been deposited in the European Nucleotide Archive (ENA)

under study accession PRJEB98673; individual sample accessions are provided in eTable 4.

Tables

Table 1. Epidemiological and clinical characteristics of all PHSs stratified by Kp status and molecular cluster assignment.

Parameter	All PHSs	PHSs without Kp (A)	PHSs with any Kp (B)	PHSs with nosocomial Kp (C)	PHSs with nosocomial, cluster-assigned Kp (D)	Adjusted OR [95% CI][c] (C vs. A)	Adjusted OR [95% CI][c] (D vs. A)
Total PHSs, n ([a])	1009 (936)	927 (859[b])	82 (81)	58 (57)	37 (37)	-	-
Length of PHSs, median [IQR], days	6.0 [2.4 – 15.2]	5.3 [2.2 – 13.6]	28.6 [9.1 – 50.7]	32.8 [17.1 – 62.4]	35.0 [26.2 – 57.4]	-	-
Sex, Male, n (%)	523/936 (55.9)	479/859 (55.8)	44/81 (54.3)	32/57 (56.1)	23/37 (62.2)	.84 [.47 – 1.50]	.60 [.29 – 1.23]
VLBW, n (%)	202/936 (21.6)	158/859 (18.4)	45/81 (55.6)	38/57 (66.7)	30/37 (81.1)	**3.42 [1.29 – 9.32]**	**8.76 [2.45 – 34.16]**
GA, median [IQR], days (weeks)	254 (36.3) [226 (32.3) – 273 (39.0)]	256 (36.6) [230 (32.9) – 274 (39.1)]	219 (31.3) [190 (27.1) – 249 (35.6)]	208 (29.7) [188 (26.9) – 237 (33.9)]	202 (28.9) [187 (26.7) – 221 (31.6)]	**.98 [.97 – .9986]**	.99 [.97 – 1.01]
Birth weight, median [IQR], grams	2503 [1610 – 3286]	2600 [1718 – 3340]	1470 [960 – 2225]	1300 [955 – 1878]	1220 [930 – 1478]	-	-
Screening effort [d]	-	-	-	-	-	**4.83 [2.01 – 11.10]**	**3.62 [1.05 – 10.78]**

Abbreviations and definitions: PHS, patient hospital stay; defined as a continuous inpatient episode on the NICU/IMC. Discharge or transfer ended

the PHS, readmission started a new PHS. Kp, *Klebsiella pneumoniae* complex; A Kp case was any PHS with ≥1 Kp-positive specimen.

Nosocomial Kp carriage was defined as acquisition on or after day three or later. The *at-risk set* analyzed in the logistic regression models

comprised patients hospitalized for ≥48 h and Kp negative until that time point (landmark definition). VLBW, very low birth weight; GA, gestational

age. [a] Numbers in parentheses indicate corresponding patients. [b] Four patients acquired Kp during a later PHS. [c] Adjusted odds ratios with 95%

profile-likelihood confidence intervals are from multivariable logistic regression at the patient level, comparing nosocomial Kp acquisition (C) or

nosocomial Kp with molecular cluster assignment (D) against no Kp detection (A). Because of collinearity, VLBW (<1500 g birth weight) and not

birth weight was retained in the multivariable model. Reference categories: male sex and absence of VLBW. [d] Screening effort was defined as the

number of surveillance samples per patient divided by the individual time at risk, i.e., length of stay in the NICU/IMC until Kp acquisition for positive

cases or until discharge for negative cases. Because it is not a baseline characteristic, no descriptive statistics are provided. Inclusion of screening

effort did not materially change the association of VLBW with Kp acquisition.

Table 2: Epidemiological and clinical characteristics of Kp infections.

Epidemiological Parameters	Patient A	Patient B	Patient C
GA, category	Early preterm	Early preterm	Term
Birth weight, grams	[1,500-2,499]	[1,500-2,499]	[2,500-4,200]
Time at risk[a], days	0[b]	48	56
Time until infection, days	5	48	56
Length of stay, days	11	86	85
Clinical parameters during time until infection			
Central venous catheter, days	5	33	56
Peripheral venous catheter, days	5	22	5
Invasive ventilation, days	1	14	56
Non-invasive ventilation, days	1	11	0
Transurethral catheter, days	0	9	10
Surgery	yes	no	yes

Abbreviations and definitions: GA, gestational age; Kp, *Klebsiella pneumoniae* complex; Kp

infection was defined as the presence of Kp in a diagnostic microbiological specimen (e.g.

blood culture), combined with ≥5 days of antibiotic treatment and clinical documentation.

[a] Time at risk refers to days from admission until acquisition of the Kp isolate. [b] Postoperative

infection following Kp colonization at a transferring hospital.

Table 3. Epidemiological and clinical parameters of nosocomial Kp PHSs stratified by

molecular cluster assignment.

Parameter	All nosocomial Kp PHSs (A)	Nosocomial Kp PHSs with molecular cluster assignment (B)	Nosocomial Kp PHSs without molecular cluster assignment (C)	*P*-value[c] (B vs. C)
Epidemiological parameters	**n=58**	**n=37**	**n=21**	
Total PHSs, n ([a]) [% of all PHSs (A)]	58 (57)[b]	37 (37) [63.8%]	21 (21) [36.2%]	-
Length of PHSs, median [IQR], days	32.8 [17.1 – 62.4]	35.0 [26.2 – 57.4]	24.7 [12.1 – 67.9]	.27
time at risk of PHSs, median [IQR], days	12.5 [6.8 – 25.5]	16.0 [9.0 – 30.0]	7.0 [4.0 – 21.5]	.06
Patient associated parameters	**n=57**	**n=37**	**n=21**	
VLBW, n (%)	38 (66.7)	30 (81.1)	8 (38.1)	**.0009**
GA, median [IQR], days (weeks)	208 (29.7) [188 (26.9) - 237 (33.9)]	202 (28.9) [187 (26.7) – 221 (31.6)]	238 (34.0) [189.5 (27.1) – 260.5 (37.2)]	**.02**
Birth weight, median [IQR], grams	1300 [955 – 1878]	1220 [930 – 1478]	1670 [985 – 2952]	**.01**
Sex, Male, n (%)	32 (56.1)	23 (62.2)	10 (47.6)	.28
Death, n (%)	6 (10.5)	6 (16.2)	0 (0)	.05
Clinical parameters per PHS	**n=58**	**n=37**	**n=21**	
Central venous catheter, n (%)	25 (43.1)	15 (40.5)	10 (47.6)	.60
Peripheral venous catheter, n (%)	56 (96.5)	37 (100)	19 (90.5)	.06
Invasive ventilation, n (%)	23 (39.7)	15 (40.5)	8 (38.1)	.85
Non-invasive ventilation, n (%)	44 (75.9)	32 (86.5)	12 (57.1)	**.01**
Transurethral catheter, n (%)	10 (17.2)	4 (10.8)	6 (28.6)	.08
Surgery, n (%)	8 (13.8)	5 (13.5)	3 (14.3)	.93
Systemic (intravenous) antibiotic therapy, n (%)	25 (43.1)	16 (43.2)	9 (42.9)	.97
Cefotaxime, n (%)	11 (18.9)	8 (21.6)	3 (14.3)	.49
Vancomycin, n (%)	8 (13.8)	3 (8.1)	5 (23.8)	.09
Meropenem, n (%)	9 (15.5)	6 (16.2)	3 (14.3)	.84
Tobramycin, n (%)	22 (37.9)	17 (45.9)	5 (23.8)	.09

Abbreviations: PHS, patient hospital stay; Kp, *Klebsiella pneumoniae* complex; VLBW, very

low birth weight; GA, gestational age.

[a] Corresponding number of single patients. [b] One patient initially carried a non-cluster-

assigned Kp strain and, after discharge and subsequent readmission to the neonatal

intensive care unit (NICU) during the same overall hospital stay, acquired a cluster-assigned

Kp strain. This resulted in two separate PHSs for that patient, one counted in column (B) and

one in column (C). Consequently, the number of single *patients* in column (A) totals 57,

whereas the sum of patients in columns (B) and (C) equals 58, consistent with the total

number of PHSs (A). [c] Chi-square test for categorical parameters and Wilcoxon rank sum

test for continuous parameters, respectively. Level of significance **P<.05**.

Figure Legends

Figure 1. Phylogeny, antimicrobial susceptibility, and genomic/phenotypic features of

83 Kp isolates.

Rows represent individual isolates ordered by a midpoint-rooted neighbor-joining

dendrogram based on core-genome multilocus sequence typing (cgMLST) allele distances

calculated in Ridom SeqSphere+ (left). The central heatmap displays EUCAST antimicrobial

susceptibility interpretations (susceptible [S], increased exposure [I], resistant [R]) for nine

agents: ampicillin/sulbactam (SAM), piperacillin/tazobactam (TZP), cefuroxime (CXM),

cefotaxime (CTX), ceftazidime (CAZ), meropenem (MEM), gentamicin (GEN), ciprofloxacin

(CIP), and trimethoprim/sulfamethoxazole (SXT). Left annotations: patient ID (duplicates

highlighted), acquisition mode (darkest = nosocomial; lightest = community), and cluster

assignment by MLST, cgMLST, and SKA2 (shared color key); unclustered ("singleton")

entries are shown in light gray. Right annotations: presence of extended-spectrum β-

lactamase (ESBL) loci and major virulence determinants (Yersiniabactin, Colibactin,

Aerobactin, Salmochelin, RmpADC; colored by allele). Phenotypic results for tellurite

resistance and string test shown as positive, negative, or unclear; light gray indicates tests

not performed.

Figure 2. Machine learning-based prediction of molecular cluster prevalence and

model interpretation.

(A) Weekly observed cluster prevalence (black line) vs predicted prevalence from Random

Forest (blue dotted line) and reduced-feature extreme gradient boosting (XGBoost) model

(orange line). Shaded area indicates bootstrapped 95% confidence interval of the

predictions.

(B) Global feature importance in the reduced-feature XGBoost model. Shapley Additive

Explanations (SHAP) beeswarm plot shows mean absolute SHAP values (ranked by

importance) and individual SHAP values per observation (colored by feature value). Grey

bars indicate gain-based feature importance.

(C) Seasonal variation in feature importance. Heatmap of mean absolute SHAP values

stratified by meteorological season.

Abbreviations: [Tmin (lagged)] minimum temperature, 4-week lag; [RH] current humidity;

[VLBW :2 (lagged)] very low birth weight (<1500g) patients treated with a 1:2 nurse-to-patient

ratio (1 nurse per 2 patients), 4-week lag; [Tmax (lagged)] maximum temperature, 4-week

lag; [HBW (lagged)] peak number of patients with birth weight >1500g ("higher birth weight")

treated, 4-week lag; [VLBW] current peak number of very low birth weight patients treated (all

nursing ratios). For definitions, see eMethods.