Genetic risk factors for postoperative complications after major surgery and shared genetic aetiology with non-postoperative phenotypes

Richard A. Armstrong,[1*] Paul Yousefi,[1,2] Ben Gibbison,[3] Golam M Khandaker,[1,2,4,5] Tom R Gaunt[1,2]

1. MRC Integrative Epidemiology Unit, Bristol Medical School, University of Bristol, Bristol, UK.

2. NIHR Bristol Biomedical Research Centre, University Hospitals Bristol and Weston NHS Foundation Trust and University of Bristol, Bristol, UK.

3. Bristol Medical School, University of Bristol, Bristol, UK.

4. Avon and Wiltshire Mental Health Partnership NHS Trust, Bristol, UK.

5. Centre for Academic Mental Health, Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK.

*Correspondence to: Richard A Armstrong

Email: ra17848@bristol.ac.uk

X: @drrichstrong

Short title: Genetic risk factors for postoperative complications

Abstract (242 words)

Background

Up to 15% of patients worldwide experience serious postoperative complications after major surgery. We explored the genetic basis of five common postoperative complications and whether they share genetic aetiology with their non-postoperative equivalents.

Methods

We performed case-control genome-wide association studies (GWAS) using UK Biobank data from >140,000 participants. The primary outcomes were postoperative acute kidney injury (AKI), atrial fibrillation (AF), myocardial infarction (MI), stroke and surgical site infection (SSI). Additionally, we assessed genetic correlation between UK Biobank postoperative outcomes, as well as between these and their non-postoperative (undifferentiated) equivalents from the FinnGen cohort. Finally, we assessed the association between observed postoperative outcomes and polygenic risk scores (PRS) for their non-postoperative equivalents.

Results

Among 8,472 postoperative complications in 140,563 eligible participants, three genome-wide significant risk loci for AF were identified on chromosomes 1 (*KCNN3*), 4 (*PITX2*) and 16 (*TRAF7:CASKIN1*). No genome-wide significant loci were found for AKI, MI, stroke or SSI. Strong genetic correlation between postoperative and non-postoperative AF was found (r_g = 0.69, p = 0.001). Increasing PRS quintiles for non-postoperative AF, MI, stroke and chronic kidney disease were associated with increased odds of developing the corresponding postoperative complication.

Conclusions

In addition to identifying a novel genomic risk loci associated with postoperative AF, we show that postoperative complications and their non-postoperative equivalents share genetic architecture. These findings suggest that postoperative complications may

represent the acute manifestation of underlying genetic vulnerability, which has implications for preoperative risk stratification and postoperative clinical management.

Introduction

Up to 15% of patients worldwide experience serious postoperative complications after major surgery with significant implications for morbidity, mortality and resource utilisation [1]. These complications represent the interplay between intervention-specific outcomes, the insult of surgery more generally, and underlying vulnerabilities in individuals. They are poorly understood from an aetiological perspective and have few clinically practical predictors to guide preventive treatment.

There is increasing recognition that genetic influences can impact postoperative outcomes and that perioperative care needs to adapt to the era of 'big data' in which 'traditional' risk factors can be combined with individualised data including genomics and other high-dimensional omics [2–4]. Coupled with this is evidence that a seemingly disparate range of postoperative complications may be due, in part, to an aberrant host response to the inflammatory insult of surgery [5–13]. This suggests there could be a common immune-mediated component which is generic rather than organ- or operation-specific.

Genome-wide association studies (GWAS) are a powerful approach to study such complex outcome phenotypes. They can provide insight into the underlying mechanisms and biological substrate whilst identifying genetic variants to be used for risk prediction or therapeutic targeting [14]. GWAS for postoperative complications have examined atrial fibrillation (AF) [15], venous thromboembolism (VTE) [16], acute kidney injury (AKI) [17], and a composite of myocardial infarction (MI), AF, stroke, AKI and delirium [18]. Most have found no associated variants at a genome-wide significance level but sample size has been a common limitation, as has the use of methods which can be unstable in the context of case-control imbalance [19].

To investigate genetic risk factors for postoperative complications and address the methodological limitations of previous studies we conducted a GWAS of postoperative complications after major surgery in the UK Biobank cohort, controlling for case-control imbalance. We focused on AKI, AF, acute MI, stroke and surgical site infection (SSI) as

common, possible immune-mediated complications. We then extended our analyses to explore whether these complications share genetic risk factors with one another, and whether the postoperative phenotypes are genetically related to their non-postoperative equivalents.

Methods

Study populations

The primary analysis was conducted in UK Biobank, a large-scale biomedical database and research resource containing genetic, lifestyle and health information from half a million UK participants recruited between 2006 and 2010 (http://www.ukbiobank.ac.uk). Linked health data is available for all participants with follow-up to 2022 [20]. The UK Biobank study was approved by the North-West Multi-centre Research Ethics Committee and all participants provided written informed consent. This research has been conducted under UK Biobank project number 128619.

FinnGen (freeze 9) was used as an external cohort for genetic correlation analyses. FinnGen is a large-scale genomics initiative that has analysed over 500,000 Finnish biobank samples and correlated genetic variation with health data to understand disease mechanisms and predispositions. The project is a collaboration between research organisations and biobanks within Finland and international industry partners [21].

Genotyping

We used UK Biobank genotype data that was previously generated using Affymetrix using two purpose-designed arrays. The genotype data had been quality controlled, phased and ~96 million genotypes imputed using the Haplotype Reference Consortium and UK10K haplotype resources. FinnGen genotyping used a ThermoFisher Axiom custom chip array with imputation to a reference dataset derived from Finnish whole-genome sequences. Further information on the methods and QC pipeline are available [21,22].

Inclusion and exclusion criteria

We defined eligible participants as those undergoing major, inpatient surgery after their date of enrolment in UK Biobank. Major surgery was defined by OPCS4 codes using the Bupa Schedule of Procedures and Abbott classification [23,24]. Exclusion criteria for all phenotypes included a previous diagnosis of the outcome of interest and planned day case surgery. For acute MI, cardiothoracic and cardiology procedures were excluded to reduce the risk of the diagnosis preceding the procedure (see Supplementary Methods for more detail on phenotype definition and exclusions).

Outcome definition

The primary outcome for each GWAS was a first diagnosis of the complication within a 30 day period following surgery. Controls were defined as those undergoing qualifying surgery but not having the outcome within 30 days. Outcome diagnoses were identified using the following International Classification of Diseases, Tenth Revision (ICD-10) codes: atrial fibrillation, I48.9; acute kidney injury, N17; myocardial infarction, I21; stroke, I61, I63 and I64; and surgical site infection, T81.4.

Genome-wide Association Study (GWAS)

Case-control GWAS adjusted for sex, genetic chip and the first ten genetic principal components was performed for each outcome using REGENIE (v3.2.2) [19]. Previously established MRC-IEU workflows and methods were used for quality control and ancestry definition [25]. Genome-wide significance was defined as $p < 5 \times 10^{-8}$ with an additional exploratory threshold of $p < 1 \times 10^{-6}$. Post-hoc analyses were performed using Functional Mapping and Annotation of Genome-Wide Association Studies (FUMA GWAS [26]). Independent significant SNPs were defined as those meeting the genome-wide significance threshold and independent of each other at a linkage disequilibrium (LD) threshold of $r^2 < 0.6$. Lead SNPs were identified from independent significant SNPs at $r^2 < 0.1$. Gene-based analysis was performed using Multi-marker Analysis of GenoMic Annotation (MAGMA [27]) with a genome-wide significance threshold of $p < 0.05/19,840$ (2.52×10^{-6}). For postoperative AF, age was included as an additional covariate [28] and

subgroup analyses were performed after excluding participants who underwent cardiothoracic surgery or cardiology procedures (as defined by treatment specialty).

Genetic correlation analyses between postoperative complications

The GWAS results were used to analyse genetic correlation across phenotypes. Global genetic correlation was assessed through cross-trait LD Score Regression [29,30] and Local Analysis of [co]Variant Association (LAVA) was used to analyse bivariate local genetic correlation in 2,495 predefined regions [31].

Genetic correlation analysis with non-postoperative phenotypes

GWAS summary statistics were obtained from the FinnGen study cohort (Freeze 9, https://r9.finngen.fi/) for non-postoperative equivalents of the outcome phenotypes, i.e. undifferentiated AF, AKI, MI and stroke, as well as chronic kidney disease (CKD). No summary statistics were available for SSI. The results of the primary GWAS analyses were used with this external dataset to analyse global and local genetic correlation between the postoperative and non-postoperative paired phenotypes as above.

Polygenic risk score analysis

Polygenic risk scores (PRS) were identified in the Polygenic Score Catalog (https://www.pgscatalog.org) for non-postoperative AF, AKI/CKD, MI and stroke. No PRS were available for SSI. The *pgsc_calc* workflow [32] was used to download, variant match and calculate individual polygenic scores for the UK Biobank cohort. Scores were included if at least 75% of variants present in the score were also present in the UK Biobank genomic data. Individual scores which passed matching were then centred and scaled before being summed and averaged. The average score was then converted into risk quintiles (1=lowest risk, 5=highest) to improve interpretability [33]. Odds ratios for the association of increasing quintile of PRS for the non-postoperative phenotype with the postoperative equivalent were estimated by logistic regression after adjustment for age and sex using the *glm* function in R (version 4.4.0).

Results

Participant characteristics

A total of 8,472 complications (AF, AKI, MI, SSI, or stroke) occurred in 140,563 eligible individuals. Case-control numbers for each outcome phenotype after excluding previous diagnoses (and cardiothoracic/cardiology procedures for AMI) are shown in Table 1. Most individuals (7,711) had a single complication with 701 and 60 individuals having two and three complications respectively. The most common combinations of complications were AF and AKI (228 individuals) and AKI and SSI (205 individuals).

Individuals who developed postoperative complications tended to be older, were more likely to be male, current or previous smokers and had higher rates of comorbidity (all p < 0.001) (Table 2). Compared to controls, cases were more likely to be emergency admissions (2,899 (34%) vs 27,690 (21%), p < 0.001) and to have undergone cardiothoracic surgical procedures (1,158 (14%) vs 4,211 (3.2%), p < 0.001). Additional details and tables of participant characteristics for each outcome phenotype can be found in supplementary tables 1-6.

Genome-wide association study

A total of 171 genome-wide significant variants were identified for postoperative AF, with nine independent significant SNPs across three genomic risk loci on chromosomes 1, 4 and 16 identified by FUMA GWAS (Figures 1-2; Table 3). The lead SNPs were rs36088503, 4:111697815_TG_T (GRCh37.p13) and rs145672192 respectively. The nearest genes were *KCNN3,* which reached genome-wide significance in a MAGMA gene-level analysis (p = 2.4 x 10^{-7}, Figure 3), *PITX2, RP11-777N19.1* and *TRAF7:CASKIN1*. A further 38 genes were mapped to these regions (supplementary table 7). No genome-wide significant SNPs were found for AKI, AMI, stroke or surgical site infection (supplementary figures 1–8). At an exploratory threshold of p < 1 x 10^{-6}, three independent variants were identified for both AKI and AMI, and eight for SSI (supplementary table 8).

Subgroup analyses were performed for postoperative AF in cardiothoracic and non-cardiothoracic surgery subgroups. No genome-wide significant SNPs or genes were identified for the cardiac surgery cohort (962 cases, 13,504 controls). For the non-cardiac cohort (1777 cases, 122,374 controls), six independent significant SNPs were identified in risk loci on chromosomes 2 and 4 from a total of 162 genome-wide significant variants (supplementary table 9). The lead SNPs were rs12052983 (Chr 2) and rs6843082 (Chr 4). Eighteen mapped genes were identified but none reached genome-wide significance on MAGMA testing.

Genetic correlation between postoperative complications in UK Biobank

No significant global genetic correlation was found between the different postoperative phenotypes in UK Biobank on cross-trait LD Score Regression analysis (supplementary table 10). Local genetic correlation between complications was therefore not assessed given the low SNP-based heritability z-scores and effective sample sizes [29].

Genetic correlation with between postoperative (UK Biobank) and non-postoperative (FinnGen) phenotypes

A strong global genetic correlation was observed between postoperative AF in UK Biobank and non-postoperative AF in the FinnGen cohort (r_g = 0.69, p = 0.001). Significant local correlation was confirmed in two regions near to the *PITX2* (Chr 1) and *KCNN3* (Chr 4) genes identified in the primary GWAS (Table 4). There was some suggestion of shared genetic liability between postoperative AKI and chronic renal failure, but the evidence for global genetic correlation was limited (r_g = 0.42, p = 0.05) and there was little evidence of local genetic correlation. The results for the other phenotypes did not support the presence of global genetic correlation between postoperative and non-postoperative settings (supplementary table 11) and were not analysed further.

Association of polygenic risk scores for related phenotypes (PGS Catalog) with postoperative complications (UK Biobank)

Polygenic risk scores were identified and calculated for non-postoperative AF (20 PRS), AMI (6 PRS) and stroke (25 PRS). No PRS were available for AKI but 20 scores were identified for chronic kidney disease (supplementary table 12). After adjusting for age and sex, increasing PRS quintile was associated with increased odds of postoperative complications for all phenotypes with the most pronounced effects seen for AF and AMI (Figure 4; supplementary table 13). In a subgroup analysis for AF stratified by cardiothoracic/non-cardiothoracic surgery, stronger effects were seen in the non-cardiothoracic cohort: PRS quintile 5, OR (95% CI) for non-cardiothoracic surgery, 6.5 (5.4–7.8), and for cardiothoracic surgery 2.6 (2.1–3.3) (supplementary table 14).

Discussion

In this study we used a UK-based cohort with linked healthcare data to explore the genetic basis of common postoperative complications. We identified three genomic loci associated with postoperative atrial fibrillation (AF) and evidence of shared genetic risk with non-postoperative AF. Similarly, we found common genetic risk factors between postoperative outcomes and their respective non-postoperative counterparts for both myocardial infarction and stroke.

Our study is better powered than previous GWAS of postoperative AF (number of cases/controls: 257/620 and 242/640) which failed to find any genome-wide significant results [34,35]. A comparable previous GWAS of postoperative AF using the UK Biobank cohort reported two variants in proximity to the *PITX2* gene on chromosome 4 [15]. However, our analysis builds on these findings by restricting eligibility to major surgical procedures, defining the outcome phenotype more strictly, and employing a method that is robust in the context of case-control imbalance [19]. We identified independent significant SNPs in a high degree of linkage disequilibrium (LD) with those variants, adding to the evidence implicating the 4q25 locus and *PITX2* gene, established genetic predictors of AF more broadly, in postoperative AF [36–38]. We also found additional risk loci on chromosomes 1 and 16. The 1q21 region has previously been associated with non-postoperative AF in GWAS [38] and post-cardiac surgery AF in a targeted genotyping analysis [37]. We found this region, including the *KCNN3* gene, to be a risk locus for postoperative AF with additional evidence of genetic correlation with non-postoperative AF. The third risk locus identified, 16p13 (lead SNP rs145672192), appears to be a novel finding. It is distinct from an established risk locus for AF more broadly, 16q22 (*ZFHX3*), and is not in LD with this region. The nearest genes are *TRAF7:CASKIN1* which are closely located and coregulated. They may contribute to the development of postoperative AF through roles in inflammatory modulation and signalling (*TRAF7*) or cardiac conduction and excitability (*CASKIN1*).

We found no genome-wide significant variants associated with postoperative AKI, MI, stroke or SSI. Whilst we were better powered than a previous GWAS including AKI, MI

and stroke [35] we were likely only powered to detect large effects. However, our sample sizes for AKI and SSI were similar to that of AF and so it may be that these outcomes are driven more by environmental or direct perioperative factors than underlying genetic susceptibility. This is supported by SNP-based heritability estimates of 22.1% for AF [39] versus <0.1% and 4.5% for AKI after cardiac and non-cardiac surgery respectively [40,41].

The relationship between postoperative and non-postoperative paired phenotypes was explored by genetic correlation and polygenic risk score analyses. Postoperative AF showed the strongest evidence of commonality with its non-postoperative counterpart supported by global, local and PRS-based analyses. The association between increasing PRS and the postoperative outcome was particularly marked in the non-cardiothoracic surgery cohort. Whilst the cardiothoracic cohort was smaller and so relatively underpowered, AF after cardiothoracic surgery has a distinct set of pathophysiological contributory mechanisms which make a reduced genetic influence a plausible finding. No PRS were available for postoperative AKI and so chronic kidney disease was used as the paired phenotype. We found suggestive evidence of global genetic correlation between AKI and CKD as well as increased risk of postoperative AKI with increasing PRS for CKD. Higher PRS quintiles for MI and stroke were also associated with increased risk of the relevant postoperative complication. Taken together, these findings may indicate a 'two-hit' paradigm in which the inflammatory insult of surgery precipitates the outcome in individuals with underlying genetic vulnerability for related chronic pathology.

There may therefore be a role for existing PRS in risk-stratification and preventive perioperative care. In the case of postoperative AF, a review of clinical practice across UK cardiac surgery centres found most did not risk-stratify patients or have a care package in place to prevent postoperative AF [42]. The use of PRS for non-postoperative AF could therefore be a useful adjunct in the pre-operative phase, highlighting those at highest risk who may benefit from a preventive package of care. There are also implications for the longer-term follow-up of acute postoperative complications. Given the presence of shared underlying genetic risk between these postoperative

phenotypes and their non-postoperative equivalents, there will be a proportion of individuals in whom the acute event is the initial presentation of a chronic condition.

This study does have limitations. Some factors which would be of clinical relevance are not collected in UK Biobank or the linked NHS healthcare data, for example ASA physical status grade or type of anaesthesia. Additionally, the UK Biobank cohort is not representative of the wider population, particularly due to differences in socioeconomic status and comorbid status [43]. The postoperative complications studied were defined by clinical coding data and were not prospectively screened for. This may have resulted in some cases being missed; however, this would dilute any observed effect and so suggests our findings are robust to this misclassification. Whilst efforts were taken to minimise the risk of the outcomes preceding surgery it is possible that some cases remain. The limited sample size, particularly for some outcomes, restricted the power of our primary analysis and the application of some genetic correlation methods. Finally, the external cohort used does not define postoperative phenotypes and so direct replication and validation was not possible.

In conclusion, we found multiple genomic risk loci associated with postoperative atrial fibrillation in a major surgery cohort, including a novel locus on chromosome 16 which would benefit from validation and further investigation. We demonstrate evidence of shared genetic aetiology between postoperative complications and their non-postoperative equivalents with implications for both pre-operative risk stratification, through the use of existing polygenic risk scores, and postoperative follow-up, as an acute complication may be indicator or first presentation of an underlying chronic condition.

Tables and figures

Table 1 Numbers of cases and controls for each outcome phenotype in UK Biobank

Outcome	Cases	Controls
Acute kidney injury	3,002	134,941
Atrial fibrillation	2,739	135,878
Surgical site infection	2,701	136,120
Stroke	569	137,842
Acute myocardial infarction	282	123,339

Table 2 Summary of participant characteristics for cases and controls. For individuals experiencing multiple complications, details of the first complication were used. For controls, details of the first eligible surgical procedure were used.

	Cases n = 8,472	Controls n = 132,091	p-value
Age (yr), mean (SD)	68 (6)	64 (8)	<0.001
Female sex, number (%)	3,524 (42)	74,283 (56)	<0.001
Ethnicity, number (%)			<0.001
Asian or Asian British	146 (1.7)	2,166 (1.6)	
Black or Black British	94 (1.1)	1,565 (1.2)	
Chinese	12 (0.1)	228 (0.2)	
Mixed	36 (0.4)	644 (0.5)	
Other ethnic group	54 (0.6)	935 (0.7)	
White	8,090 (96)	125,951 (96)	
Missing	40 (0.5)	602 (0.5)	
Smoking status, number (%)			<0.001
Current	1,138 (13)	14,151 (11)	
Never	3,697 (44)	66,825 (51)	
Previous	3,570 (42)	50,389 (38)	
Prefer not to answer or missing	67 (0.8)	726 (0.5)	
Charlson Comorbidity Index, median IQR)	1.00 (0.00-2.00)	0.00 (0.00-1.00)	<0.001
Admission method, number (%)			<0.001
Elective	5,242 (62)	101,732 (77)	
Emergency	2,899 (34)	27,690 (21)	
Other	331 (3.9)	2,669 (2)	
Surgical specialty, number (%)			<0.001
Cardiothoracic surgery	1,158 (14)	4,211 (3.2)	
General surgery	2,229 (26)	24,604 (19)	
Neurosurgery	371 (4.4)	4,902 (3.7)	
Other	1,865 (22)	34,937 (26)	
Trauma and orthopaedics	1,986 (23)	51,927 (39)	
Urology	624 (7.4)	10,080 (7.6)	

Vascular	239 (2.8)	1,430 (1.1)	

Numbers are number (proportion), mean (SD) or median (IQR). Additional details can be found in Supplementary table 1.

Table 3 Independent significant SNPs identified in case-control GWAS of postoperative atrial fibrillation

rsID*	Chr:Pos**	Nearest gene	Function	Effect allele	OR (95% CI)	p
-	4:111697815_TG_T	*RP11-777N19.1*	intergenic	TG	0.73 (0.68-0.78)	6.5×10^{-20}
rs200657990	4:111679136	*RP11-777N19.1*	intergenic	A	0.71 (0.65-0.77)	1.2×10^{-13}
rs10032150	4:111646618	*RP11-777N19.1*	intergenic	A	0.81 (0.76-0.86)	8.5×10^{-11}
-	4:111607460_ATT_A	*PITX2*	intergenic	ATT	0.78 (0.73-0.84)	3.3×10^{-10}
rs72667982	4:111673642	*RP11-777N19.1*	intergenic	A	0.81 (0.76-0.87)	2.4×10^{-9}
rs145672192	16:2227985	*TRAF7:CASKIN1*	UTR3	G	2.32 (1.67-3.24)	2.3×10^{-8}
rs4571403	4:111613368	*PITX2*	intergenic	G	1.19 (1.12-1.26)	3.1×10^{-8}
rs36088503	1:154812943	*KCNN3*	intronic	G	0.83 (0.77-0.88)	4.1×10^{-8}
rs723363	4:111724501	*RP11-777N19.1*	intergenic	T	1.18 (1.11-1.25)	4.4×10^{-8}

*rsID: Reference SNP cluster ID. **GRCh37.p13

Table 4 Regions of significant local genetic correlation between postoperative atrial fibrillation (in UK Biobank) and non-postoperative atrial fibrillation (in FinnGen)

Chr	Start	Stop	Rho	R^2	p
1	154685546	156813845	0.54 (0.36-0.74)	0.29 (0.13-0.55)	7.3×10^{-8}
4	109978983	111733579	0.58 (0.44-0.73)	0.34 (0.20-0.45)	2.1×10^{-13}

Rho is the estimate of the local genetic correlation between traits (moderate to strong positive correlation: genetic variants in this region may affect both traits in a similar way). R2 is the proportion of variance explained by the local genetic correlation, or how much of the trait variability can be account for by this region. Proportion of covariance between the traits that is explained by variation in this range.

Figure legends

Figure 1 Manhattan plot of genome-wide association study results for postoperative atrial fibrillation. The red line represents a genome-wide significant p value of 5×10^{-8}.

Figure 2 Q-Q plot for genome-wide association study of postoperative atrial fibrillation. Genomic inflation factor (λ) 1.04.

Figure 3 Manhattan plot of gene-based testing for postoperative atrial fibrillation

Figure 4 Association between postoperative complications and increasing polygenic risk score quintile for non-postoperative equivalent phenotype. Red line is odds ratio of 1. Error bars are 95% confidence interval.

Acknowledgements

This research has been conducted using the UK Biobank Resource under Application Number 128619. This includes linked data provided by patients and collected by the NHS as part of their care and support and data assets made available by National Safe Haven as part of the Data and Connectivity National Core Study, led by Health Data Research UK in partnership with the Office for National Statistics and funded by UK Research and Innovation (grant ref MC_PC_20058).

Quality Control filtering of the UK Biobank data used the process described by R.Mitchell, G.Hemani, T.Dudding, L.Corbin, S.Harrison, L.Paternoster[25].

This work was carried out using the computational and data storage facilities of the Advanced Computing Research Centre, University of Bristol (http://www.bristol.ac.uk/acrc/).

Funding

RAA is funded by a Wellcome Trust GW4-CAT PhD Programme for Health Professionals PhD Fellowship [316275/Z/24/Z]. RAA, PY, GMK and TRG are supported by the Medical Research Council Integrative Epidemiology Unit at the University of Bristol (RA, TG: MC_UU_00032/3; PY: MC_UU_00032/4; GMK: MC_UU_0032/6). GMK acknowledges additional funding from the Wellcome Trust (grant numbers: 201486/Z/16/Z and 201486/B/16/Z), the Medical Research Council (grant numbers: MR/W014416/1; MR/S037675/1; MR/Z50354X/1; and MR/Z503745/1. PY, TRG, and GMK are also supported by the UK National Institute for Health and Care Research (NIHR) Bristol Biomedical Research Centre (grant number: NIHR 203315). The views expressed are those of the authors and not necessarily those of the UK NIHR or the Department of Health and Social Care.

References

1. Dobson GP. Trauma of major surgery: A global problem that is not going away. *Int. J. Surg. Lond. Engl.* 2020; **81**: 47–54. https://doi.org/10.1016/j.ijsu.2020.07.017.

2. Shaw A. Exploring the perioptome: the role of genomics in thoracic surgery and anaesthesia. *Curr. Opin. Anesthesiol.* 2007; **20**: 32. https://doi.org/10.1097/ACO.0b013e32801271e4.

3. Levin MA, Joseph TT, Jeff JM et al. iGAS: A framework for using electronic intraoperative medical records for genomic discovery. *J. Biomed. Inform.* 2017; **67**: 80–9. https://doi.org/10.1016/j.jbi.2017.02.005.

4. Ajitsaria P, Eissa SZ, Kerridge RK. Risk Assessment. *Curr. Anesthesiol. Rep.* 2018; **8**: 1–8. https://doi.org/10.1007/s40140-018-0246-9.

5. Vlisides P, Mashour GA. Perioperative Stroke. *Can. J. Anaesth. J. Can. Anesth.* 2016; **63**: 193–204. https://doi.org/10.1007/s12630-015-0494-9.

6. Shu L, Aziz YN, de Havenon A et al. Perioperative Stroke: Mechanisms, Risk Stratification, and Management. *Stroke* 2025; **56**: 2798–809. https://doi.org/10.1161/STROKEAHA.125.051673.

7. Gaudino M, Andreotti F, Zamparelli R et al. The −174G/C Interleukin-6 Polymorphism Influences Postoperative Interleukin-6 Levels and Postoperative Atrial Fibrillation. Is Atrial Fibrillation an Inflammatory Complication? *Circulation* 2003; **108**: II–195. https://doi.org/10.1161/01.cir.0000087441.48566.0d.

8. Boos CJ, Anderson RA, Lip GYH. Is atrial fibrillation an inflammatory disorder? *Eur. Heart J.* 2006; **27**: 136–49. https://doi.org/10.1093/eurheartj/ehi645.

9. Korantzopoulos P, Letsas KP, Tse G, Fragakis N, Goudis CA, Liu T. Inflammation and atrial fibrillation: A comprehensive review. *J. Arrhythmia* 2018; **34**: 394–401. https://doi.org/10.1002/joa3.12077.

10. Kinsey GR, Li L, Okusa MD. Inflammation in Acute Kidney Injury. *Nephron Exp. Nephrol.* 2008; **109**: e102–7. https://doi.org/10.1159/000142934.

11. Rabb H, Griffin MD, McKay DB et al. Inflammation in AKI: Current Understanding, Key Questions, and Knowledge Gaps. *J. Am. Soc. Nephrol. JASN* 2016; **27**: 371–9. https://doi.org/10.1681/ASN.2015030261.

12. Frangogiannis NG, Smith CW, Entman ML. The inflammatory response in myocardial infarction. *Cardiovasc. Res.* 2002; **53**: 31–47. https://doi.org/10.1016/S0008-6363(01)00434-5.

13. Moyes LH, Leitch EF, McKee RF, Anderson JH, Horgan PG, McMillan DC. Preoperative systemic inflammation predicts postoperative infectious complications in patients

undergoing curative resection for colorectal cancer. *Br. J. Cancer* 2009; **100**: 1236–9. https://doi.org/10.1038/sj.bjc.6604997.

14. Bain CR, Shaw AD. Genetics and epigenetics in perioperative medicine. *Curr. Opin. Crit. Care* 2012; **18**: 548. https://doi.org/10.1097/MCC.0b013e328357af6d.

15. Christensen MA, Bonde A, Sillesen M. Genetic risk factors for postoperative atrial fibrillation—a nationwide genome-wide association study (GWAS). *Front. Cardiovasc. Med.* 2023; **10**. https://doi.org/10.3389/fcvm.2023.1040757.

16. Christensen M, Bonde A, Sillesen M. An assessment of the effect of the genotype on postoperative venous thromboembolism risk in 140,831 surgical patients. *Ann. Med. Surg.* 2021; **71**: 102938. https://doi.org/10.1016/j.amsu.2021.102938.

17. Db L, A L, L B et al. Limited clinical utility for GWAS or polygenic risk score for postoperative acute kidney injury in non-cardiac surgery in European-ancestry patients. *BMC Nephrol.* 2022; **23**. https://doi.org/10.1186/s12882-022-02964-8.

18. S W, C S, M G et al. Genome-wide association study of myocardial infarction, atrial fibrillation, acute stroke, acute kidney injury and delirium after cardiac surgery - a sub-analysis of the RIPHeart-Study. *BMC Cardiovasc. Disord.* 2019; **19**. https://doi.org/10.1186/s12872-019-1002-x.

19. Mbatchou J, Barnard L, Backman J et al. Computationally efficient whole-genome regression for quantitative and binary traits. *Nat. Genet.* 2021; **53**: 1097–103. https://doi.org/10.1038/s41588-021-00870-7.

20. : External Info : Data_providers_and_dates. https://biobank.ndph.ox.ac.uk/showcase/exinfo.cgi?src=Data_providers_and_dates (accessed 09/10/2024).

21. Kurki MI, Karjalainen J, Palta P et al. FinnGen provides genetic insights from a well-phenotyped isolated population. *Nature* 2023; **613**: 508–18. https://doi.org/10.1038/s41586-022-05473-8.

22. Bycroft C, Freeman C, Petkova D et al. The UK Biobank resource with deep phenotyping and genomic data. *Nature* 2018; **562**: 203–9. https://doi.org/10.1038/s41586-018-0579-z.

23. Abbott TEF, Fowler AJ, Dobbs TD, Harrison EM, Gillies MA, Pearse RM. Frequency of surgical treatment and related hospital procedures in the UK: a national ecological study using hospital episode statistics. *BJA Br. J. Anaesth.* 2017; **119**: 249–57. https://doi.org/10.1093/bja/aex137.

24. Bupa - Code Search main. https://codes.bupa.co.uk/procedures (accessed 12/05/2021).

25. Ruth Mitchell GH. UK Biobank Genetic Data: MRC-IEU Quality Control, version 2. *data.bris*, 2019. https://data.bris.ac.uk/data/dataset/1ovaau5sxunp2cv8rcy88688v (accessed 09/10/2024).

26. Watanabe K, Taskesen E, van Bochoven A, Posthuma D. Functional mapping and annotation of genetic associations with FUMA. *Nat. Commun.* 2017; **8**: 1826. https://doi.org/10.1038/s41467-017-01261-5.

27. Leeuw CA de, Mooij JM, Heskes T, Posthuma D. MAGMA: Generalized Gene-Set Analysis of GWAS Data. *PLOS Comput. Biol.* 2015; **11**: e1004219. https://doi.org/10.1371/journal.pcbi.1004219.

28. Chen Z, Aziz PF, Greenland P, Passman R, Gordon AS, Webster G. Age-Dependent Contributions of Rare and Common Genetic Variation in Atrial Fibrillation. *Circ. Genomic Precis. Med.* 2025; **18**: e004958. https://doi.org/10.1161/CIRCGEN.124.004958.

29. Bulik-Sullivan B, Finucane HK, Anttila V et al. An atlas of genetic correlations across human diseases and traits. *Nat. Genet.* 2015; **47**: 1236–41. https://doi.org/10.1038/ng.3406.

30. Bulik-Sullivan BK, Loh P-R, Finucane HK et al. LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nat. Genet.* 2015; **47**: 291–5. https://doi.org/10.1038/ng.3211.

31. Werme J, van der Sluis S, Posthuma D, de Leeuw CA. An integrated framework for local genetic correlation analysis. *Nat. Genet.* 2022; **54**: 274–82. https://doi.org/10.1038/s41588-022-01017-y.

32. Lambert SA, Wingfield B, Gibson JT et al. The Polygenic Score Catalog: new functionality and tools to enable FAIR research. 2024. .

33. Collister JA, Liu X, Clifton L. Calculating Polygenic Risk Scores (PRS) in UK Biobank: A Practical Guide for Epidemiologists. *Front. Genet.* 2022; **13**. https://www.frontiersin.org/journals/genetics/articles/10.3389/fgene.2022.818574.

34. Kertai MD, Li Y-J, Ji Y et al. Genome-wide association study of new-onset atrial fibrillation after coronary artery bypass grafting surgery. *Am. Heart J.* 2015; **170**: 580-590.e28. https://doi.org/10.1016/j.ahj.2015.06.009.

35. Westphal S, Stoppe C, Gruenewald M et al. Genome-wide association study of myocardial infarction, atrial fibrillation, acute stroke, acute kidney injury and delirium after cardiac surgery - a sub-analysis of the RIPHeart-Study. *BMC Cardiovasc. Disord.* 2019; **19**: 26. https://doi.org/10.1186/s12872-019-1002-x.

36. Body SC, Collard CD, Shernan SK et al. Variation in the 4q25 Chromosomal Locus Predicts Atrial Fibrillation After Coronary Artery Bypass Graft Surgery. *Circ. Cardiovasc. Genet.* 2009; **2**: 499–506. https://doi.org/10.1161/CIRCGENETICS.109.849075.

37. Sigurdsson MI, Muehlschlegel JD, Fox AA et al. Genetic Variants Associated With Atrial Fibrillation and PR Interval Following Cardiac Surgery. *J. Cardiothorac. Vasc. Anesth.* 2015; **29**: 605–10. https://doi.org/10.1053/j.jvca.2014.10.028.

38. Ellinor PT, Lunetta KL, Albert CM et al. Meta-analysis identifies six new susceptibility loci for atrial fibrillation. *Nat. Genet.* 2012; **44**: 670. https://doi.org/10.1038/ng.2261.

39. Weng L-C, Choi SH, Klarin D et al. Heritability of Atrial Fibrillation. *Circ. Cardiovasc. Genet.* 2017; **10**: e001838. https://doi.org/10.1161/CIRCGENETICS.117.001838.

40. Larach DB, Lewis A, Bastarache L et al. Limited clinical utility for GWAS or polygenic risk score for postoperative acute kidney injury in non-cardiac surgery in European-ancestry patients. *BMC Nephrol.* 2022; **23**: 339. https://doi.org/10.1186/s12882-022-02964-8.

41. Douville NJ, Larach DB, Lewis A et al. Genetic predisposition may not improve prediction of cardiac surgery-associated acute kidney injury. *Front. Genet.* 2023; **14**: 1094908. https://doi.org/10.3389/fgene.2023.1094908.

42. Gibbison B, Murphy G, O'Brien B, Pufulete M. An Update on Guidelines to Prevent and Manage Atrial Fibrillation After Cardiac Surgery and a Survey of Practice in the UK. *J. Cardiothorac. Vasc. Anesth.* 2024; **38**: 2307–13. https://doi.org/10.1053/j.jvca.2024.07.043.

43. Fry A, Littlejohns TJ, Sudlow C et al. Comparison of Sociodemographic and Health-Related Characteristics of UK Biobank Participants With Those of the General Population. *Am. J. Epidemiol.* 2017; **186**: 1026–34. https://doi.org/10.1093/aje/kwx246.