Title: Long-term Outcomes of Prophylactic Total Gastrectomy in Patients with

Hereditary Diffuse Gastric Cancer

Authors: Hui Jun Lim[1], BMed/MD, Maria O'Donovan[2], FRCPath, Claire Lamb[3], MNutr,

Sophie Samra[1], BSc Adult Nursing Nyarai Chinyama[3], MSc Advanced Clinical Practice, Paul

C Fletcher[4], MD, PhD, Hisham Ziauddeen[4], MRCPsych, PhD, Ju Yi Tai[5], Patrick Murphy[5],

Richard H Hardwick[3], Massimiliano Di Pietro[1], MD, PhD, Rebecca C Fitzgerald[1], MD, PhD,

J Robert O'Neill, FRCSEd, PhD[3, 6*]

[1]Early Cancer Institute, University of Cambridge, Cambridge, United Kingdom

[2]Department of Histopathology, Cambridge University Hospitals NHS Foundation Trust,

Cambridge, United Kingdom

[3]Cambridge Oesophago-gastric Centre, Cambridge University Hospitals NHS Foundation

Trust, Cambridge, United Kingdom

[4]Department of Psychiatry, University of Cambridge, Herchel Smith Building,

Addenbrooke's Hospital, Cambridge, CB2 0SZ, United Kingdom

[5]University of Cambridge, Cambridge, United Kingdom

[6]University of Cambridge Department of Surgery, Box 202, Addenbrookes Hospital, Hills

Road, Cambridge, CB2 0QQ, United Kingdom

*Corresponding author

Correspondence to:

Dr J Robert O'Neill

University of Cambridge Department of Surgery

Box 202, Addenbrookes Hospital, Hills Road, Cambridge, CB2 0QQ

ro317@cam.ac.uk (ORCID 0000-0003-2230-7137)

Word count: 3000 words

Declarations

- **Ethics approval and consent to participate**

  Consent was obtained prospectively from all and this study was carried out under the approval of Anglia and Oxford Medical Research Ethics Committee with reference number 97/5/32 and any amendments approved by the Cambridgeshire Research Ethics Committee

- **Consent for publication**

  Not applicable.

- **Availability of data and materials**

  Data are available from the corresponding author on reasonable request.

- **Competing interests**

  The authors declare that they have no competing interests.

- **Funding**

  Lim HJ is a PhD student on the Research Training Fellowship funded by National Medical Research Council, Singapore. Di Pietro M is funded by the UK Medical Research Council and received additional funds from the Cancer Research UK Cambridge Centre (grant number CTRQQR-2021\100012). O'Neill JR acknowledges the support of the Cancer Research UK Cambridge Centre Thoracic Cancer programme (CTRQQR-2021\100012). This work was supported by the NIHR Cambridge Biomedical Research Centre (NIHR203312). The views expressed are those of the authors and not necessarily those of the NIHR or the Department of Health and Social Care.

- **Authors' contributions**

  Hui Jun Lim, Claire Lamb, Sophie Samra, Nyarai Chinyama, Maria O'Donovan, Richard H Hardwick and J Robert O'Neill assisted with participant recruitment and data collection. Paul C Fletcher and Hisham Ziauddeen facilitated the set-up of data collection. Ju Yi Tai and Patrick Murphy aided in data collection. Lim HJ and O'Neill JR performed data analyses and wrote the manuscript. Di Pietro M and Fitzgerald R contributed intellectual guidance and examined the manuscript critically for publication. All authors have reviewed and approved the final manuscript.

- **Acknowledgements**

  We would like to thank Marc Tischkowtiz and Samantha Grimes for their contributions and involvement in the care and follow-up of patients. In addition, we would like to thank Nikos Demiris for his guidance in the statistical analyses.

Key Points

Question: What are the clinicopathological and long-term impact on weight and health-

related quality of life outcomes for patients with hereditary diffuse gastric cancer who

undergo prophylactic total gastrectomy?

Findings: In this study which included 60 patients with HDGC who underwent PTG, there

was a lower major complication rate at <2% and the majority of patients at 88.0% had pT1a

on gastrectomy specimens. Partial weight regain was seen in those with normal pre-operative

BMI between 2 to 5 years and QOL worsened markedly at 1 to 3 months but largely returned

to baseline by 12 to 24 months and remained stable at baseline levels beyond 10 years.

Meaning: PTG can be undertaken with a low major complication rate and most QOL

domains return to baseline within 24 months and sustained in the long-term within the

context of a specialist multidisciplinary service.

Abstract

Importance:

Hereditary diffuse gastric cancer (HDGC) is predominantly due to germline *CDH1* variants and confers a high lifetime risk of diffuse gastric cancer (DGC). Prophylactic total gastrectomy (PTG) is thought to prevent DGC but has potential long-term functional, nutritional and quality-of-life (QOL) sequelae.

Objective:

We report the clinical, pathological and long-term weight and QOL outcomes from a prospective cohort of patients with HDGC who underwent PTG.

Design and Setting:

This is a single-centre prospective cohort study (Familial Gastric Cancer Study) of *CDH1* pathogenic variant carriers managed at Addenbrooke's Hospital (2005–2024).

Participants:

Patients with *CDH1* pathogenic variants and completed □2 QOL questionnaires were included.

Intervention:

Patients who underwent PTG or endoscopic surveillance were included.

Main Outcomes and Measures:

The clinical, pathological, weight and QOL outcomes for patients undergoing PTG (n=60) are reported. The QOL of patients undergoing endoscopic surveillance only (n=25) and surveillance followed by PTG were also reported. Longitudinal changes in QOL were analysed using linear mixed models with Bonferroni-Holm correction.

Results:

Median age at PTG was 31 years (IQR 25–41); 75% underwent vagal-sparing PTG. The

major complication rate was <2% with no operative mortalities and a median hospital stay of

8 days. Foci of intramucosal signet ring cell (SRC) carcinoma (pT1a) were seen in 88%, pT0

in 10% with a median number of SRC foci of 8. The disease-specific and overall-survivals

was 100% with median follow-up of 9.7 years. Weight loss was universal, reaching ~15% at

3 months and extending to 20% by 24 months. Partial weight regain was seen in those with

normal pre-operative BMI between 2–5 years. QOL worsened markedly at 1–3 months but

largely returned to baseline by 12–24 months and remained stable at baseline levels beyond

10 years. Younger patients (<35) had better role functioning (p=0.038) and men reported

greater persistent appetite loss (p=0.030) following PTG.

Conclusions and Relevance:

PTG can be undertaken with a low major complication rate, and is a curative treatment for

HDGC. Although substantial early functional and weight effects occur, most QOL domains

recover to baseline by 12–24 months and are sustained long-term with regular follow-up

within a specialist multidisciplinary service.

Keywords: Hereditary diffuse gastric cancer; prophylactic total gastrectomy; quality of life;

weight outcomes

Background

Hereditary diffuse gastric cancer (HDGC) syndrome is predominantly attributed to germline *CDH1* mutations.[1, 2] Individuals with pathogenic germline *CDH1* mutations have a 33.5 fold increased risk of developing diffuse gastric cancer by the age of 30 and women with HDGC carry a 6 fold higher risk of lobular breast cancer.[3] Although advances in genetic testing have improved identification of *CDH1* mutation carriers, the mechanisms by which *CDH1* and other genes drive HDGC remain poorly understood, limiting preventative strategies. Given the insidious onset and poor prognosis of advanced DGC, prophylactic total gastrectomy (PTG) remains the only definitive risk-reducing option.[4-7] Current guidelines recommend PTG for pathogenic *CDH1* carriers, with endoscopic surveillance allowing individualised timing.[7, 8] However, PTG may result in long-term functional and nutritional sequelae affecting health-related quality of life (QOL).[9]

As many individuals with HDGC are asymptomatic and may never develop cancer, understanding surgical risks and long-term outcomes is essential to informed decision-making.[10, 11] Most QOL data following total gastrectomy are derived from patients treated for invasive disease, with limited follow-up or confounding cancer therapies.[12, 13] Long-term studies in sporadic gastric cancer demonstrate persistent QOL impairment up to five years post-operatively, underscoring the need for extended assessment.[14]

To date, only a small number of retrospective studies have evaluated outcomes following PTG in HDGC, mostly limited to two years of follow-up.[15-19] These suggest PTG is safe, with low major complication rates and resolution of most symptoms by two years. Only 1 study has assessed QOL beyond five years, reporting no significant differences over time.[19] This highlights the need for long-term evaluation.

We therefore examined clinicopathological and long-term QOL outcomes in patients with

HDGC participating in the prospective Familial Gastric Cancer cohort study and undergoing

PTG at a national referral centre.

Materials and Methods

Patient Cohort

All patients with HDGC managed as part of a national referral service at Cambridge

University Hospitals from 1st January 2005 to 31st January 2025 and who were prospectively

enrolled in the longitudinal Familial Gastric Cancer study (FGCS) were included. Patients in

the FGCS provided written informed consent to the use of clinicopathological data and

weight outcomes as well as completion of serial questionnaires. Research ethical approval for

the FGCS was provided by the Anglia and Oxford Medical Research Ethics Committee

(reference number 97/5/32) and subsequent amendments approved by the Cambridgeshire

Research Ethics Committee. All patients consented to the study received a baseline

endoscopy with high resolution white light endoscopy and narrow band imaging with

magnification followed by targeted biopsies of visible lesions, with particular focus on pale

areas, and random gastric biopsies were taken according to the Cambridge protocol. For

patients who opted to undergo endoscopic surveillance, 6-monthly or annual surveillance was

performed depending on endoscopy findings. Patients diagnosed with invasive disease on

baseline endoscopy were excluded.

There were several factors considered in the decision-making for PTG, including age, co-

morbidities, family history, burden of signet ring cell (SRC) foci on endoscopy and patient

choice. All patients were discussed at a multidisciplinary meeting before proceeding with

PTG. All patients underwent baseline endoscopy and extensive pre-operative counselling and

prehabilitation with a multi-disciplinary team (MDT). Selective use of pre-operative CT was

applied based on patient factors. Following PTG, regular assessments of patients' nutritional

status, diet, weight and symptoms were performed in conjunction with dietician input and

oral supplementation or rarely nasojejunal enteral feeding if required due to nutritional

failure. Surgical feeding jejunostomies were not placed at PTG.

Patients who continued surveillance underwent 6-monthly, annual or 2-yearly surveillance

depending on various factors, including age, endoscopic findings and patients' preference,

following discussion with their gastroenterologist and with input as needed from the MDT.

Surgical Approach and Peri-operative Protocol

All patients underwent PTG by the open approach via either midline or rooftop incision with

1 of 2 specialised upper gastrointestinal surgeons. Lymphadenectomy was D1 as per

international guidance with some D0 cases in the early cohort and D2 selectively based on

patient factors. The posterior vagal trunk and coeliac innervation was preserved where

possible, but the anterior vagal trunk was ligated at the hiatus. An end-to-side circular stapled

oesophagojejunostomy and side to side semi-mechanical jejuno-jejunostomy was performed

in all cases with intra-operative confirmation of complete gastric mucosal resection before

anastomosis. Cholecystectomy was undertaken if gallstones were identified pre-operatively.

Analgesia was via epidural or surgically placed rectus-sheath catheters with intravenous

patient-controlled opiate analgesia. Patients followed a standardised enhanced recovery

pathway with a planned level 2 ward stay for 1 to 2 days and then managed on a level 0 ward.

Nasogastric tubes were not used and surgical drains placed very selectively.

Surgical and Post-operative Outcomes Assessment

Prospectively collected clinical, pathological and post-operative outcomes were maintained

using a bespoke online database. Peri-operative outcomes included operative approach,

duration, lymphadenectomy, duration of high dependency and total inpatient stay and post-

operative complications. Complications within 30 post-operative days were reported

according to the Gastrectomy Complications Consensus Group classification and using the

Clavien-Dindo system with major complications defined as ≥Clavien-Dindo IIIa.[20, 21] Post-

operative mortality was regarded as within in-patient stay or 90 days. The interval from date

of surgery to date of disease recurrence was defined as the disease-free interval and date of

surgery to death from any cause as the overall survival interval, censoring at last follow-up

date.

Processing and Histopathological Assessment of Gastrectomy Specimen

Gastrectomy specimens were received fresh the stomach was opened along the greater curve

and examined for macroscopically visible abnormalities or lesions. The proximal and distal

resection margins were removed and fixed in 10% formalin. The anterior and posterior walls

of the stomach were separated and pinned onto corkboards. The anterior and posterior walls

were fixed for 24 hours in PAXgene® tissue FIX and 10% formalin respectively then

unpinned and placed in PAXgene® tissue stabilizer for the anterior wall. The anterior and

posterior stomach walls were divided into cardia, fundus, body, transition zone, antrum and

pylorus. These were blocked entirely into cassettes for paraffin embedding. Alternate blocks

were used for diagnostic and research purposes. Sections from diagnostic blocks were stained

with haematoxylin and eosin to assess for in-situ, pagetoid signet ring lesions and SRC

carcinoma foci. All cases were reviewed by upper gastrointestinal specialist pathologists.

Histopathological data included size, location, number of in-situ SRC lesions and SRC foci,

proximal and distal resection margins, completeness of excision of gastric mucosa, total

number of lymph nodes harvested, number of positive lymph nodes and tumour-node-

metastasis (TNM) staging summarised according to TNM staging of 8th edition of American

Joint Committee on Cancer.[22] Histological types were classified according to 2019 World

Health Organization classification of tumours of the digestive system.[23]

Weight Outcomes Assessment

Pre-operative weight was prospectively collected when patients were scheduled for surgery

and used as the baseline to derive post-operative changes in weight and body mass index

(BMI; defined as weight (kg)/ height^2 (m)). Standard post-operative follow-up intervals

included 2 and 6 weeks, 3, 6, 9 and 12 months and then 6 monthly intervals until the third

post-operative year and annually following this or at sooner intervals if needed.

Health-related Quality of Life Assessment

We prospectively assessed impact of PTG on QOL using European Organisation for

Research and Treatment of Cancer (EORTC) QLQ-C30 and QLQ-STO22 questionnaires at

post-operative months 1, 3, 12 and annually thereafter.[24, 25] Raw questionnaire responses were

used to derive scores using the scoring manuals.[26, 27] For patients continuing surveillance,

prospective questionnaire responses were requested at baseline endoscopy and annually at

subsequent endoscopies. For analysis of QOL changes, the first questionnaire response

during surveillance was taken to be the baseline (dynamic baseline). The QOL assessment

immediately prior to surgery was taken as the baseline for relative analysis of changes

following PTG.

Statistical Methods

Continuous variables were summarised using median (interquartile or total range as

described), and categorical variables by frequency (%). The scores were summarised using

mean (95% confidence intervals). To account for inter-subject variance in baseline and

subsequent QOL measures, the percentage change from baseline was used for subsequent

statistical analyses. Longitudinal QOL changes were analysed using linear mixed models

with time, age grouped as <35 or >35, sex and vagal☐sparing status as fixed effects and

patient as a random effect.[28] First☐order autoregressive covariance was selected for residuals.

Bonferroni-Holm correction accounted for multiple comparisons. A random intercept for

center was included to account for within-center clustering. The model with the lowest BIC

value was used to ensure robustness of the model. A two-sided P<0.05 indicated statistically

significant. Analyses were performed in the R statistical environment, version 4.4.1 (R

Foundation) and Jamovi, version 2.7.[29, 30]

Results

Patient Characteristics and Clinicopathological Outcomes
267
The patient characteristics, clinical and histopathological outcomes of patients who

underwent endoscopic surveillance and PTG are shown in Table 1. An overview of the

clinical pathway is shown in Figure 1. A total of 60 patients underwent PTG with median

age at surgery of 31 years old (25, 41) and 45% (27/60) were female. Within the PTG

cohort, there were 24 patients who initially opted for endoscopic surveillance and

underwent at least 1 surveillance endoscopy and proceeded to PTG with a median

number of 3 surveillance endoscopies over a median of 18 months. All patients having

PTG had pathogenic *CDH1* variants. There were 3 patients who underwent endoscopic

surveillance followed by PTG who completed at least 2 pre-operative and 2 post-

operative QOL questionnaires.

The median operative time for PTG was 149 minutes (IQR 131, 234), and 45 patients

(75.0%) underwent posterior vagal-sparing total gastrectomy. A total of 44 (73.3%)

underwent D1 lymphadenectomy. A patient had a major complication with an

anastomotic leak from the jejuno-jejunal anastomosis diagnosed on post-operative day 9

requiring re-operation (Clavien-Dindo IIIb). There were no post-operative mortalities.

The median length of high dependency stay was 2 days (range 0 to 5) with a median

inpatient stay of 8 days (range 5 to 17).

Histopathological Outcomes

All patients had complete resection of gastric mucosa and R0 resection. The final stage

was pT1bN0M0 in 1 patient (1.7%) which was due to superficial submucosal invasion in

1 lesion in the PTG specimen. This represents 0.05% (1/1697) of all histopathological

lesions diagnosed on gastrectomy specimens. The final pathological stage was

pT1aN0M0 in 53 patients (88%) and pT0N0M0 in 6 (10%) reflecting the high rate of

SRC foci in patients with pathogenic *CDH1* mutations. The median number of SRC foci

was 8 but there was a wide range from 0 to 273 reflecting a wide heterogeneity in this

group and the choice of timing for PTG. The median number of lymph nodes harvested

was 12 (range 0 to 54). No peri-neural or lympho-vascular invasion was reported.

Disease-specific survival (DSS) and overall survival were 100% with median follow-up

of 9.7 years (Supplementary Figure 1).

Assessment of Long-term Weight Changes Following PTG

The post-operative weight and BMI of 57 patients was collected during follow-up. Post-

operative weight loss was rapid and universal reaching 15% of baseline at 3 post-operative

months. This reached a plateau of 15% to 20% total body weight loss between 12 and 24

post-operative months (Figure 2a). When stratified by pre-operative BMI, all patients lost

weight by 12 post-operative months, but the greatest absolute loss occurred in patients with a

higher pre-operative BMI (Figure 2b). Furthermore, in patients with a normal pre-operative

BMI, weight gain was seen between 2 and 5 years post-operatively with most in a BMI range

20 to 25 kg/m^2.

Health-related Quality of Life Outcomes

A summary of the scores in various functional domains and symptom scales in the

surveillance cohort is shown in Figure 3. As anticipated, beyond small variances in symptom

and functional domain scores, all outcomes were stable over 3 years.

Changes in QOL from pre-operative baseline until after the first post-operative decade are

shown in Figure 5. Significant changes were seen with a high rate of most symptoms and

lower functional status across most domains at 1 and 3 post-operative months. Most changes

had resolved to baseline levels by 12 or up to 24 months. Strikingly, in long-term follow-up

from 2 to beyond 10 years, patients did not report significant different functional levels from

baseline. Similarly, in symptom scores, recovery was close to baseline at 12 or up to 24

months. A trend was noted in more prevalent symptoms relating to appetite loss, eating

difficulties, diarrhoea, reflux and fatigue although no significant difference from baseline

could be observed at 1 years onwards to >10 years (Figure 3). A trend was also noted in

reduced anxiety over time although this did not reach significance compared to baseline

levels.

The changes in QOL outcomes for patients undergoing PTG were compared according to age

at time of surgery (Supplementary Figure 2), gender (Supplementary Figure 3) and posterior

vagal-sparing or non vagal-sparing surgery (Supplementary Figure 4). There were no

significant differences in majority of functioning domains and symptoms within these groups.

However, younger patients (<35) had significantly better role functioning **(p-value = 0.038)**.

Most symptom changes were similar between genders however, men experienced

significantly more appetite loss compared to women and this persisted in the long-term (**p-**

value = 0.030). A summary of key findings is shown in Supplementary Figure 5.

We next sought to examine absolute QOL score changes in the same patients directly from

surveillance through PTG and then into follow-up where 3 patients were included

(Supplementary Figure 6). The short-term impact of PTG and resolution of symptoms and

functional impairment can be seen at 12 to 24 post-operative months.

Discussion

In this study, PTG performed within the context of a specialist service was associated with

low peri-operative morbidity and excellent long-term oncologic outcomes. Major post-

operative complications occurred in <2%, there were no postoperative mortalities and median

length of stay was 8 days. The peri-operative outcomes compare favourably with previously

reported series from high-volume centres, including those from National Institutes of Health

and international multi-centre analyses, which reported major complication rates >10% and

longer hospital stays.[17, 21] The majority of patients had a pathological stage of pT1aN0M0

(88.0%), no lymph node metastases and complete gastric mucosal resection achieved for all.

Our follow-up demonstrated DSS of 100% with median follow-up of 9.7 years. [33] These

results demonstrated PTG undertaken within a specialist programme with expert endoscopic

surveillance can be a safe curative treatment.

Weight loss was universal following PTG, with the most rapid decline occurring within the

first 3 postoperative months and a nadir between 12 and 24 months. Total weight loss ranged

from 15% to 20% of pre-operative body weight, with greater losses observed among pre-

operative BMI >30 group. Partial weight re-gain occurred 2 to 5 years post-operatively

among patients with normal pre-operative BMI. These findings are similar to those from

Memorial Sloan Kettering Cancer Center cohort with the greatest loss seen in the pre-

operatively obese group at a median of 28% at 2 years followed by 18% and 14% amongst

overweight and normal weight groups respectively.[18] We predict these observations may in

part be attributed to the specialist dietetic input and regular follow-up of their nutritional

status and weight with intervention for those with persistent nutritional compromise.

Health-related QOL declined across multiple symptom and functional domains in the early

postoperative period, particularly within the first 3 months. Appetite loss, eating difficulties,

fatigue, nausea, vomiting, and taste disturbance were most prominent. QOL measures

returned to baseline within 6 to 12 months and remained stable for more than a decade,

providing reassurance regarding long-term functional recovery. Patients managed with

endoscopic surveillance alone demonstrated stable QOL over time. Among patients who

underwent surveillance followed by PTG, serial assessments showed recovery of symptoms

and function to near-baseline levels. Our finding that the recovery of symptoms and

functional outcomes is preserved over a decade is novel and should provide some reassurance

to patients with HDGC contemplating PTG. Notably, anxiety remained prominent among

patients undergoing surveillance whereas improvement was observed following PTG. This

finding may reflect reassurance associated with elimination of gastric cancer risk and

highlights the need for psychological support for individuals undergoing surveillance.

We next assessed contribution of gender, age at surgery and preservation of the posterior

vagal nerve on outcomes. Patients aged 35 or younger had better role functioning in line with

the expectation that younger patients generally can recover over a shorter duration and return

to daily activities. It is possible these patients were inclined to undergo PTG and it was based

on patient choice. We demonstrated men experienced greater loss of appetite compared to

women. Our results did not show a significant difference in QOL outcomes between vagal-

preserving and non vagal-preserving surgery. The small numbers of patients with a complete

vagotomy in our study may limit the power to detect a clinical difference and we did not

routinely use intra-operative nerve monitoring to confirm posterior trunk was functional.[31]

Future work will be needed to explore the neurohormonal consequences of posterior

vagotomy on patients.

Our study is limited by a relatively small sample size. Nonetheless, HDGC is a rare condition

and larger cohorts are from MSKCC and NIH groups[17, 18] A further limitation is voluntary

reporting and it is possible we missed an important group with a negative experience who did

not complete the questionnaires. Regular contact and close follow-up suggest this is less

likely. We also facilitated questionnaire completion by online and paper formats. Lastly, our

outcomes are in the context of a specialised service which may limit generalisability.

The authors propose this study improves our understanding of the long-term consequences of

PTG. A strength of this study is that the effect of total gastrectomy on symptoms and function

has been determined independent of a confounding disease process or concurrent systemic

treatment for cancer. Identification of individual domains which had worse outcomes in

specific sub-groups of patients will allow them to be targeted specifically to facilitate

recovery and understand the physiological basis for this in future work. These findings are

relevant to the wider literature as with improving oncological treatments and earlier detection

of gastric cancer, the functional impact and long-term consequences become more important.

In conclusion, our study has shown PTG to be safe with excellent clinical and pathological

outcomes. Despite the potential morbidity, we have shown that patients are able to achieve

comparable outcomes to their baseline and to patients in surveillance within 12 to 24 months

maintained in the long-term. These findings form a useful resource for patients contemplating

PTG to understand medium and long-term consequences of PTG.

References

1. Corso G, Pedrazzani C, Pinheiro H, Fernandes E, Marrelli D, Rinnovati A, et al. E-
cadherin genetic screening and clinico-pathologic characteristics of early onset gastric
cancer. Eur J Cancer. 2011;47(4):631–9.

2. Schrader KA, Masciari S, Boyd N, Wiyrick S, Kaurah P, Senz J, et al. Hereditary
diffuse gastric cancer: association with lobular breast cancer. Familial Cancer.
2008;7:73–82.

3. Ryan CE, Fasaye GA, Gallanis AF, Gamble LA, McClelland PH, Duemler A,
Samaranayake SG, Blakely AM, Drogan CM, Kingham K, Patel D, Rodgers-Fouche
L, Siegel A, Kupfer SS, Ford JM, Chung DC, Dowty JG, Sampson J, Davis JL.
Germline CDH1 Variants and Lifetime Cancer Risk. JAMA. 2024;332(9):722-729.

4. van der Sluis L, van Dieren JM, van der Post RS, Bisseling TM. Current advances and
challenges in Managing Hereditary Diffuse Gastric Cancer (HDGC): a narrative
review. Hered Cancer Clin Pract. 2024;22(1):21.

5. Cosma LS, Schlosser S, Tews HC, Müller M, Kandulski A. Hereditary Diffuse
Gastric Cancer: Molecular Genetics, Biological Mechanisms and Current Therapeutic
Approaches. Int J Mol Sci. 2022;23(14):7821.

6. Monster JL, Kemp LJS, Gloerich M, van der Post RS. Diffuse gastric cancer:
Emerging mechanisms of tumor initiation and progression. Biochim Biophys Acta
Rev Cancer. 2022;1877(3):188719.

7. Blair VR, McLeod M, Carneiro F, Coit DG, D'Addario JL, van Dieren JM, et al.
Hereditary difuse gastric cancer: updated clinical practice guidelines. Lancet Oncol.
2020;21(8):e386–97.

8. Lee CYC, Olivier A, Honing J, Lydon AM, Richardson S, O'Donovan M,
Tischkowitz M, Fitzgerald RC, di Pietro M. Endoscopic surveillance with systematic

random biopsy for the early diagnosis of hereditary diffuse gastric cancer: a prospective 16-year longitudinal cohort study. Lancet Oncol. 2023;24(1):107-116.

9. van der Kaaij RT, van Kessel JP, van Dieren JM, Snaebjornsson P, Balagué O, van Coevorden F, et al. Outcomes after prophylactic gastrectomy for hereditary diffuse gastric cancer. Br J Surg. 2018;105:e176-82.

10. Hallowell N, Badger S, Richardson S, Caldas C, Hardwick RH, Fitzgerald RC, Lawton J. An investigation of the factors effecting high-risk individuals' decision-making about prophylactic total gastrectomy and surveillance for hereditary diffuse gastric cancer (HDGC). Fam Cancer. 2016;15(4):665-676.

11. Gamble LA, Grant RRC, Samaranayake SG, Fasaye GA, Koh C, Korman L, Asif B, Heller T, Hernandez JM, Blakely AM, Davis JL. Decision-making and regret in patients with germline CDH1 variants undergoing prophylactic total gastrectomy. J Med Genet. 2023;60(3):241-246.

12. Gallanis AF, Gamble LA, Samaranayake SG, Lopez R, Rhodes A, Rajasimhan S, Fasaye GA, Juma O, Connolly M, Joyce S, Berger A, Heller T, Blakely AM, Hernandez JM, Davis JL. Costs of Cancer Prevention: Physical and Psychosocial Sequelae of Risk-Reducing Total Gastrectomy. J Clin Oncol. 2024;42(4):421-430.

13. Wang WB, Song HN, Huang DD, Luo X, Cai HY, Yan JY, Chen WZ, Xing CG, Dong QT, Chen XL. Impact of Body Composition and Physical Function on Quality of Life After Gastrectomy for Gastric Cancer. Front Surg. 2022;8:832351.

14. Lee SS, Chung HY, Kwon OK, Yu W. Quality of life in cancer survivors 5 years or more after total gastrectomy: a case-control study. Int J Surg. 2014;12(7):700-705.

15. Hayat MY, Yakubu U, Jayasinghe J, Patel B. The Psychological Impact of Prophylactic Total Gastrectomy in Patients Who Are High Risk for Hereditary Diffuse Gastric Cancer: A Review of the Literature. Cureus. 2025;17(5):e84751.

16. Worster E, Liu X, Richardson S, Hardwick RH, Dwerryhouse S, Caldas C, Fitzgerald

RC. The impact of prophylactic total gastrectomy on health-related quality of life: a

prospective cohort study. Ann Surg. 2014;260(1):87-93.

17. Gallanis AF, Bowden C, Lopez R, Gamble LA, Samaranayake SG, Payne C, Snyder

D, Fasaye GA, Joyce S, Broesamle R, Miao N, Miettinen M, Quezado M, Kim SA,

Korman L, Heller T, Blakely AM, Hernandez JM, Davis JL. Lessons learned from

150 total gastrectomies for prevention of cancer. J Gastrointest Surg.

2025;29(1):101889.

18. Vos EL, Salo-Mullen EE, Tang LH, Schattner M, Yoon SS, Gerdes H, Markowitz AJ,

Mandelker D, Janjigian Y, Offitt K, Coit DG, Stadler ZK, Strong VE. Indications for

Total Gastrectomy in CDH1 Mutation Carriers and Outcomes of Risk-Reducing

Minimally Invasive and Open Gastrectomies. JAMA Surg. 2020;155(11):1050-1057.

19. Kaurah P, Talhouk A, MacMillan A, Lewis I, Chelcun-Schreiber K, Yoon SS,

Huntsman D. Hereditary diffuse gastric cancer: cancer risk and the personal cost of

preventive surgery. Fam Cancer. 2019;18(4):429-438.

20. Dindo D, Demartines N, Clavien PA. Classification of surgical complications: a new

proposal with evaluation in a cohort of 6336 patients and results of a survey. Ann

Surg. 2004;240:205-213.

21. Schneider MA, Kim J, Berlth F, Sugita Y, Grimminger PP, Sano T, Rosati R,

Baiocchi GL, Bencivenga M, De Manzoni G, Nunobe S, Yang HK, Gutschow CA;

GastroBenchmark Consortium; GASTRODATA Consortium. Defining benchmarks

for total and distal gastrectomy: global multicentre analysis. Br J Surg. 2024

31;111(2):znad379.

22. Amin MB, Greene FL, Edge SB, Compton CC, Gershenwald JE, Brookland RK,

Meyer L, Gress DM, Byrd DR, Winchester DP. The Eighth Edition AJCC Cancer

Staging Manual: Continuing to build a bridge from a population-based to a more

"personalized" approach to cancer staging. CA Cancer J Clin. 2017;67(2):93-99.

23. Nagtegaal ID, Odze RD, Klimstra D, Paradis V, Rugge M, Schirmacher P,

Washington KM, Carneiro F, Cree IA; WHO Classification of Tumours Editorial

Board. The 2019 WHO classification of tumours of the digestive system.

Histopathology. 2020;76(2):182-188.

24. Aaronson NK, Ahmedzai S, Bergman B, Bullinger M, Cull A, Duez NJ, Filiberti A,

Flechtner H, Fleishman SB, Haes JCD, et al. The European Organization for Research

and Treatment of Cancer QLQ–C30: A quality-of-life instrument for use in

international clinical trials in oncology. J Natl Cancer Inst. 1993;85:365-376.

25. Vickery CW, Blazeby JM, Conroy T, Arraras J, Sezer O, Koller M, Rosemeyer D,

Johnson CD, Alderson D; EORTC Quality of Life Group. Development of an EORTC

disease-specific quality of life module for use in patients with gastric cancer. Eur J

Cancer. 2001;37(8):966-971.

26. Fayers P, Aaronson NK, Bjordal K, Groenvold M, Curran D, Bottomley A. EORTC

QLQ-C30 Scoring Manual. 3rd ed. Brussels: European Organisation for Research and

Treatment of Cancer. 2001.

27. Gastric Cancer (Update of QLQ-STO22). EORTC Quality of Life. Accessed 1

August, 2025. Retrieved from https://qol.eortc.org/questionnaire/gastric-cancer-

update-of-qlq-sto22/

28. Detry MA, Ma Y. Analyzing Repeated Measurements Using Mixed Models. JAMA.

2016;315(4):407-8.

29. R Core Team (2024). R: A Language and Environment for Statistical Computing. R

Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.

30. The jamovi project (2025). jamovi (Version 2.7) [Computer Software]. Retrieved from https://www.jamovi.org

31. Yan Z, Wei M, Zhang T, Guo J, Yu A, Liang Y, Huang Y, Cui X, Wang H, Zhou K, Dong Z, Yu W. Vagus Nerve Preservation for Early Distal Gastric Cancer With Monitoring and Indocyanine Green Labeling: A Randomized Clinical Trial. JAMA Surg. 2025;160(1):85-92.

Figure Legends

Figure 1: Flowchart showing Pathway of Patients with Hereditary Diffuse Gastric Cancer

who underwent Endoscopic Surveillance and/or Prophylactic Total Gastrectomy.

Figure 2a: Weight Loss from Pre-operative Baseline (%) following Prophylactic Total

Gastrectomy.

Figure 2b: Change in Body Mass Index (BMI) following Prophylactic Total Gastrectomy in

Patients Grouped by Pre-operative BMI with Normal (BMI 20 to 25), Overweight (BMI 25 to

30) and Obese (BMI >30) Groups. Lines represent best -fit lines and error bars are 95%

confidence intervals of the best-fit line.

Figure 3: Change in EORTC QLQ-C30 and EORTC QLQ-STO22 Scores (mean ± 95%

confidence interval) from Dynamic Baseline over Time (months) in Endoscopic Surveillance

Cohort. The number of patients included at the various timepoints during endoscopic

surveillance is shown below the bottom row of graphs and is representative for all the graphs.

Figure 4: Change in EORTC QLQ-C30 and EORTC QLQ-STO22 Scores (mean ± 95%

confidence interval) from Baseline (pre-surgery) over Time in Patients following

Prophylactic Total Gastrectomy (* = significant change from baseline (P<0.05)). The number

of patients included at the various timepoints following PTG is shown below the bottom row

of graphs and is representative for all the graphs.

Supplementary Figure 1: Disease-specific Survival of Patients with Hereditary Diffuse

Gastric Cancer Patients undergoing Prophylactic Total Gastrectomy.

Supplementary Figure 2: Effect of Age at Surgery on Change in Quality of Life Scores

following Prophylactic Total Gastrectomy. Points represent mean ± 95% confidence interval.

Age at surgery was grouped as ≤35 years old or >35 years old to allow two similarly sized

groups. The outcome highlighted in the red box indicates a significant difference in

percentage change from baseline scores between age-groups following surgery.

Supplementary Figure 3: Effect of Gender on Change in Quality of Life Scores following

Prophylactic Total Gastrectomy. Points represent mean ± 95% confidence interval. The

outcome highlighted in the red box indicates a significant difference in percentage change

from baseline scores between genders.

Supplementary Figure 4: The Effect of Preservation of the Posterior Vagus on Change in

Quality of Life Scores following Prophylactic Total Gastrectomy. Points represent mean ±

95% confidence interval.

Supplementary Figure 5: Summary of Impact of Prophylactic Total Gastrectomy on Health-

related Quality of Life over Time.

Supplementary Figure 6: Comparison of Percentage Change in EORTC Scores (mean ±

95% confidence interval) for 3 Patients who underwent Endoscopic Surveillance followed by

Prophylactic Total Gastrectomy.

Table 1: Summary of Patient Characteristics of Surveillance Cohort with QOL Data and Prophylactic Total Gastrectomy (PTG) Cohort, and Histopathological and Surgical Outcomes, 2005 to 2025

Characteristic	Overall N	
	60 (100%)[1]	25 (100%)
Cohort	**Surgery**	**Endoscopic Surveillance**
Age at Surgery/ Baseline Endoscopy (Years)	31 (25, 41)	36 (27, 45)
Gender		
Male	33 (55%)	12 (48.0%)
Female	27 (45%)	13 (52.0%)
Germline *CDH1* Variant		
Pathogenic	60 (100.0%)	18 (72.0%)
Truncating	19 (31.7%)	6 (24.0%)
Frameshift	13 (21.7%)	3 (12.0%)
Deletion	10 (16.6%)	3 (12.0%)
Splice site	12 (20.0%)	6 (24.0%)
Missense	6 (10.0%)	0
Likely pathogenic	0	7 (28.0%)
Truncating	0	0
Deletion	0	0
Frameshift	0	2 (8.0%)
Splice site	0	5 (20.0%)
Missense	0	0

Variant of unknown significance	0 0
BMI at Surgery (kg/m²)	25.9 (22.9, 30.4)
Operative Approach	
Open	60 (100%)
Lymphadenectomy	
D0	12 (20.0%)
D1	44 (73.3%)
D2	4 (6.7%)
Vagal-Sparing Procedure	
Yes	45 (75%)
No	15 (25%)
Maximum Clavien-Dindo Grade (30 Days)	
0	48 (80%)
I-II	12 (20%)
Pneumonia	6 (10.0%)
Urinary tract infection	2 (3.2%)
Cardiac dysrhythmia	1 (1.6%)
Epidural site catheter infection	1 (1.6%)
Lower limb deep venous thrombosis	1 (1.6%)
Post-operative bleed managed conservatively	1 (1.6%)
III-IVb	1 (1.7%)
V	0 (0%)

HDU Length of Stay (Days)

    Median (Q1, Q3)   2 (1, 3)

Total Length of Stay (Days)

    Median (Q1, Q3)   8 (7, 9)

pT

    T0   6 (10%)

    T1a   53 (88%)

    T1b   1 (1.7%)

pN

    N0   60 (100%)

Total SRC Intramucosal

    Median (Q1, Q3)   8 (1, 30)

Total SRC In Situ

    Median (Q1, Q3)   1 (0, 4)

Total Lymph Nodes Examined

    Median (Q1, Q3)   12 (7, 20)

---

[1]Median (Q1, Q3); n (%); SRC: Signet ring cell

(a)

(b)

Better Outcome

Worse Outcome

Functional Domains and Global Health Status

Worse Outcome

Better Outcome

Psychosocial Areas

Anxiety Body Image Financial Difficulties

Symptoms

Nausea and Vomiting Constipation Diarrhoea

Dysphagia Taste Eating

Reflux Dry Mouth Appetite Loss

↑ **Better Outcome**

↓ **Worse Outcome**

Functional Domains and Global Health Status

↑ **Worse Outcome**

↓ **Better Outcome**

Psychosocial Areas

Worse Outcome

Better Outcome

Symptoms

No. of Months following PTG	1/3	12	24	36	48	60	72	84	96	108	120	>120
No. of Patients	30	28	24	20	16	15	21	13	13	9	6	20