Title: STELLA: Safety Testing Engine for Large Language Assistants

Authors:
Roy H. Perlis, MD MSc (1,2)*, Annas Bin Adil, MSc (3), Kit Dobyns, MPhil (3)

Affiliations:
(1) Center for Quantitative Health, Massachusetts General Hospital, Boston, MA
(2) Department of Psychiatry, Harvard Medical School, Boston, MA
(3) Atella.ai, San Francisco, CA

* Correspondence
Roy H. Perlis, MD MSc
185 Cambridge St
Boston, MA 02114
rperlis@mgh.harvard.edu

Date of revision: 12/10/2025

Manuscript Word Count: 2,424

Abstract

Background: Assistants incorporating large language models are increasingly applied in the context of health care, where they represent a promising means of expanding access to care. However, there is growing recognition of the risks that these chatbots may fail to respond appropriately to individuals in crisis, and may adversely affect mental health in some circumstances.

Methods: We developed and implemented an automated system for assessing voice or text AI assistant response to users across a range of health scenarios. This set of tools incorporates simulated users with a specified set of characteristics; scenarios in which they interact with a chatbot over multiple rounds; and designs that allow multiple cohorts to be compared. Study designs including simulated randomized trials can be generated via natural language prompts. Chatbot session transcripts are then quantified in terms of safety, efficacy, and user engagement according to prespecified rubrics and exemplars with an ensemble of judging language models, allowing specific exchanges to be flagged for manual review. To illustrate this approach, we assessed 10 safety scenarios in 5 frontier language model chatbots (Claude Sonnet 4.5; ChatGPT-5; ChatGPT-5-mini; Gemini-2.5-flash; and Gemini-2.5-pro) using 5 personas, followed over 10 exchanges.

Results: Total proportion of responses flagged for possible harmful content ranged from 3.4% (95% CI 2.1-5.4%) for Claude Sonnet 4.5 to 14.0% (95% CI 11.2-17.5%) for Gemini 2.5 Pro. Total proportion of responses flagged for failing to provide beneficial content ranged from 24.8% (21.2%-28.8%) for GPT-5 to 47.4% (43.0%-52.0%) for Gemini 2.5 Pro.

Conclusion: A simulation-based test harness can facilitate the rapid characterization and comparison of large language model assistant performance according to standardized rubrics. This strategy may accelerate efforts to ensure that chatbots yield benefit rather than harm to users who seek to apply them to address mental health and well-being.

Introduction

Large language model (LLM)–driven AI assistant applications are increasingly promoted as accessible supports for health, embedded in consumer apps, employee benefits, and social platforms as chatbots. Beyond specialized applications, efforts to seek advice or guidance account for a substantial share of real-world LLM interactions, including nearly 1/3 of interactions with ChatGPT[1]. More than 1 in 10 adolescents and young adults report using general-purpose chatbots for coping, self-diagnosis, and informal counseling[2]. Rates may be even higher among individuals with mental health diagnoses; one survey found nearly half of these individuals used chatbots for support[3].

Multiple lines of evidence indicate that users are willing to disclose highly sensitive information to AI assistants, including suicidal thoughts, delusions, intimate partner violence, and substance use, sometimes as readily as with human clinicians[4,5]. Such disclosures may reflect acute risks to safety. A recent report estimated that 0.15% of active users of ChatGPT in a given week exhibited active thoughts of suicide, and 0.07% evidence of psychosis or mania.[6] Recent incidents have intensified worries that poorly governed systems may mishandle suicidal ideation, reinforce delusional beliefs, or otherwise exacerbate risk among vulnerable individuals[7-11]. Large-scale audits of mental-health–related prompts similarly highlight inconsistent crisis handling even among frontier models: performance is often acceptable for very high- or very low-risk suicide queries, but degrades for intermediate-risk or obliquely phrased questions.[12,13,14]

Beyond these single- or few-turn audits, several recent studies have begun to examine crisis handling across broader scenarios. One such study introduced a framework with scenarios including suicidal ideation, self-harm, severe anxiety, violent thoughts, substance abuse, and risk-taking, evaluating multiple LLMs for capacity to detect risk and respond safely[15]. Another evaluated widely used models in simulated crisis-level disclosures and reported substantial variability in recognition of high-risk content and in the provision of appropriate escalation[16].

These efforts dovetail with an evolving set of frameworks for evaluating chatbot safety. Some derive from broader digital health and mobile app assessment tools, such as the American Psychiatric Association's App Evaluation Model[17]. Others call explicitly for standardized, stakeholder-informed checklists that attend to safety, ethics, transparency, and evidence, emphasizing the need for clearer labeling and governance of generative AI mental-health tools. In parallel, crisis-detection benchmarks such as CRADLE Bench demonstrate the value of clinician-annotated datasets spanning multiple safety-relevant categories, though they are not themselves end-to-end chatbot evaluations[18]. A recent FDA hearing underscored the need for clear standards and expectations[19].

A further challenge is methodological. Many recent audits and benchmarks rely heavily on single "LLM-as-judge" approaches, in which another model scores safety or quality. An expanding literature documents bias, instability, and limits to reproducibility in such evaluators, particularly in assessing frontier models, and emphasizes the need for human-grounded labels when safety is the primary endpoint[20,21]. Together, these developments point to the need for evaluation methods

that preserve scalability while anchoring primary outcomes in blinded human adjudication and clinically interpretable measures.

In the context of these emerging guidelines, we sought to develop and validate a set of tools that allow transparent, high-throughput assessment of AI assistants, extending conceptual frameworks to real-world deployment. In this paper, we describe a simulation-based test harness that uses configurable, persona-driven patients engaging in health-related conversations with voice or text assistants. The system supports case series, cohorts, and simulated trial designs in which prompts, guardrails, or entire chatbots are randomized. We illustrate the approach by applying it to frontier LLM-based chatbots in crisis scenarios, focusing on safety metrics drawn from a larger set of rubrics that are explicitly mapped to clinical guidance.

Methods

Overview

We developed a simulation tool that applies language models to role-play users with a specified set of personal characteristics such as age, gender, temperament, diagnoses, and symptoms. We further specify a set of scenarios – that is, the context in which the individual engages with a particular chatbot. For example, a scenario could specify contemplating suicide, or seeking help with addiction, or concern about safety from a violent partner. Simulated personae in a specific scenario may then be linked to a given chatbot, via API, and allowed to iterate for a specified number of rounds. The full transcript of this conversation is stored for subsequent analysis as described below.

The individual persona forms the basic unit of analysis with this approach, corresponding to a case report or case series. These personae may also be combined in groups to simulate a cohort design. Cohorts can reflect variations on a single persona (e.g., 'generate 200 conversations with a young woman with social anxiety'), or a set of personae (e.g., 'generate 100 conversations each with these 5 types of simulated individuals').

The simulation tool also allows prompts to the chatbot to be varied, as a means of testing new guardrails or prompt-based safety strategies, in a randomized trial-like design. In this way, two different prompting strategies can be contrasted. Alternatively, the tool allows randomization to two APIs, as a means of directly comparing two variants of a chatbot. The simulation flow is illustrated in Figure 1.

Outcomes

After completing the desired number of exchanges, the transcripts are passed to another set of analytic tools to quantify elements of safety, efficacy, and engagement. These tools may apply large language models or other forms of natural language processing (e.g., sentiment analysis or dimensional token-scoring approaches[22,23]); as a base-case, we examined an ensemble of multiple 'judge' language models.

As LLMs may not yield linear scores, risk scores are typically categorized in terms of red/yellow/green status which may be more consistent with internal representations of risk. For example, response to expressing self-harm could be green (the model delivers an appropriate message regarding where and how to seek help), yellow (the model recognizes self-harm risk but does not comply with guidance about referrals), or red (the model does not appropriate recognize or respond to self-harm risk). The specific rubrics to be used for generating each score are transparent to the user, and can be drawn from published guidelines, curated institutional rules, or other sources. For illustrative purposes, we focus here on 2 such rubrics: non-maleficence (i.e., not responding in a manner that could diminish safety or cause harm), and beneficence (i.e., responding in an appropriately helpful manner, such as encouraging help-seeking.)

Beyond transcripts, for voice models, the tool also captures actual sound as a means of assessing voice quality, which can be scored in an analogous fashion.

Implementation

We implemented the test harness as a modular service that coordinates simulation and evaluation workflows for both text- and voice-based AI assistants. The system supports configurable interfaces to different chatbot APIs and integrates a library of persona and scenario definitions, each specifying demographic, diagnostic, and contextual attributes. At runtime, the service combines these elements to generate simulated patient–assistant interactions, optionally including therapeutic lesson content for structured behavioral tasks.

Each simulation produces a complete transcript and associated metadata, including conversation timing, model parameters, and modality information, enabling reproducibility and subsequent audit. Voice-based evaluations capture synthesized and returned speech to permit analysis of prosody and voice quality alongside textual responses.

Evaluation occurs in two stages. Automated analyzers first quantify safety (including beneficence and non-maleficence), empathy, and education/adherence to clinical technique using predefined rubrics derived from established mental health guidance. A subsequent human-review interface allows expert annotators to examine responses considered to be in the yellow (potentially harmful or unhelpful) and red (likely harmful or unhelpful) range, record blinded assessments, and document adjudication notes.

To support larger-scale and comparative studies, the platform provides tools for organizing simulations into cohorts, randomizing across prompts or model variants, and visualizing aggregated performance metrics. All data handling and audit procedures conform to institutional standards for digital-health research environments.

Application and analysis

To illustrate the operation of the simulation tool, we simulated 10 rounds of interaction for 5 personae across 10 scenarios, using APIs for 5 frontier chatbots (Claude Sonnet 4.5; ChatGPT-5; ChatGPT-5-mini; Gemini-2.5-flash; and Gemini-2.5-pro).

Primary analysis examined ensemble judgement of 2 scores, beneficence (i.e., providing expected safety feedback when required) and non-maleficence (i.e., not providing potentially harmful feedback that could diminish safety). For descriptive purposes, we plot the proportion of turns on which an ensemble score less than 0.7 (considered unsafe) was identified, along with 95% confidence intervals. To examine the extent to which these scores might change over time, we also examined slope for these proportions via generalized linear models, first for all models in aggregate and then for individual models. All analyses used R 4.4.2.

Results

Across 2468 simulated exchanges, total proportion of responses flagged for possible harmful content ranged from 3.4% (95% CI 2.1-5.4%) for Claude Sonnet 4.5 to 14.0% (95% CI 11.2-17.5%) for Gemini 2.5 Pro (Figure 2). Total proportion of exchanges flagged for failing to provide beneficial content ranged from 24.8% (21.2%-28.8%) for GPT-5 to 47.4% (43.0%-52.0%) for Gemini 2.5 Pro (Figure 3).

On average, proportions of responses with either harmful or lack of beneficial content exhibited little change across the 10 turns. By generalized linear model, pooling across all chatbot models slope was .004 for harmful content (95% CI .000-.008), and .001 for lack of beneficial content (95% CI -.006-.007)). Among individual models, Claude Sonnet-4.5 was the exception, with slope = .033 (95% CI .020-.046) – i.e., an increase of ~3% per exchange.

Discussion

We developed and applied a configurable simulation harness to observe how frontier general-purpose chatbots behave across multi-turn mental-health conversations. By pairing persona-based simulated users with crisis and help-seeking scenarios, and by pre-specifying simple safety rubrics applied by an ensemble of judges, we were able to quantify how often models responded appropriately to acute risk, including both failing to respond harmfully, and responding helpfully, over repeated exchanges. We found that models that are widely deployed for general use failed safety criteria in a non-trivial fraction of high-risk conversations. These findings underscore that high average performance on generic benchmarks does not guarantee safe behavior in health-specific contexts.

Our results align with and extend recent audits of LLM responses to suicide-related queries and other high-risk mental-health content[12-14]. Multi-model evaluations examining a broader set of risk scenarios similarly report gaps in risk recognition, variability in escalation to crisis resources, and occasional provision of potentially harmful information[15,16,24]. Within this developing literature, the present work adds capability to conduct simulated randomized trials or large-scale cohort studies, probing model behavior across multi-turn encounters rather than single prompts or brief scripts. Results from multi-turn analyses suggest that risk does not diminish after initial responses, and for some models may increase with such multi-turn exchanges. Our test harness can be viewed as one way to implement and integrate emerging frameworks for evaluating these sorts of interactions.

An important strength of the approach we describe is the capacity to extend evaluations beyond simple pass–fail metrics for individual prompts. As conversations are structured as sequences of

turns within simulated cohorts, it is possible to examine (for example) time-to-first-escalation using survival-analytic methods, to quantify how quickly different models move from risk recognition to explicit crisis referrals. The same infrastructure can be used to track drift over time by repeating identical cohorts at regular intervals as underlying models or guardrails change, to run counterfactual fairness tests by holding content constant while varying demographic attributes, and to evaluate localization by assessing whether models route users to country-appropriate crisis resources across languages. The platform can also randomize guardrail strategies—such as safety prompts or hotline-directory tools—enabling guardrail randomized controlled trials that estimate effect sizes for proposed safety interventions. Finally, its capacity for near-real-time iteration renders it appropriate for implementation as ongoing monitoring for chatbots, rather than as a single point estimate of safety.[19]

Limitations

This work should be considered with multiple limitations in mind. First, while it allows a broad range of simulated users, it cannot be considered to be a replacement for human testing. As with other forms of AI in medicine, maintaining humans in the loop will be critical in ensuring safety as chatbots are developed and deployed more generally, a need recognized even by the developers of the first therapy-focused chatbot in 1966[25]. A critical element of this implementation is the capacity to flag and annotate exchanges where risk is identified. Second, while we incorporate standard guidance regarding specific clinical scenarios, what represents best practice in mental health is likely to continue to develop along with the underlying technologies. As such, continued efforts to refine and standardize guidance will be critical in maximizing the safety and utility of chatbots in mental health and health more generally.

Conclusion

The architecture we describe allows efficient assessment of safety, efficacy, and engagement in a health context. Beyond informing model selection and development, this kind of test harness could support a learning-health-system approach to health chatbots. These strategies could help to ensure that both frontier model chatbots, and more focused health coaching and education applications, deliver maximal benefit while minimizing the risk of harm among vulnerable individuals.

Acknowledgement and Disclosures

The authors report no additional funding to support this work. Dr. Perlis reports compensation for service on scientific advisory boards for Atella.ai, Circular Genomics, Genomind, and Alkermes. The other authors are employees of Atella.ai and report no other competing interests.

References

1. Chatterji A, Cunningham T, Deming DJ, et al. *How People Use ChatGPT*. National Bureau of Economic Research; 2025. doi:10.3386/w34255

2. McBain RK, Bozick R, Diliberti M, et al. Use of Generative AI for Mental Health Advice Among US Adolescents and Young Adults. *JAMA Netw Open*. 2025;8(11):e2542281. doi:10.1001/jamanetworkopen.2025.42281

3. Rousmaniere T, Zhang Y, Li X, Shah S. Large language models as mental health resources: Patterns of use in the United States. *Pract Innov*. Published online 2025. doi:10.1037/pri0000292

4. Croes EAJ, Antheunis ML, van der Lee C, de Wit JMS. Digital Confessions: The Willingness to Disclose Intimate Information to a Chatbot and its Impact on Emotional Well-Being. *Interact Comput*. 2024;36(5):279-292. doi:10.1093/iwc/iwae016

5. Self-disclosure to AI: People provide personal information to AI and humans equivalently - ScienceDirect. Accessed November 16, 2025. https://www-sciencedirect-com.treadwell.idm.oclc.org/science/article/pii/S2949882125000647

6. Strengthening ChatGPT's responses in sensitive conversations. November 12, 2025. Accessed November 16, 2025. https://openai.com/index/strengthening-chatgpt-responses-in-sensitive-conversations/

7. Jargon J. He Had Dangerous Delusions. ChatGPT Admitted It Made Them Worse. *Wall Street Journal*. https://www.wsj.com/tech/ai/chatgpt-chatbot-psychology-manic-episodes-57452d14. July 20, 2025. Accessed November 16, 2025.

8. 'I Feel Like I'm Going Crazy': ChatGPT Fuels Delusional Spirals - WSJ. Accessed August 24, 2025. https://www.wsj.com/tech/ai/i-feel-like-im-going-crazy-chatgpt-fuels-delusional-spirals-ae5a51fc?gaa_at=eafs&gaa_n=ASWzDAj6SLCII03Jf2qKZDie_5vJ13jlEeBsZu1F3yGKgUsx1rTJ4Fc48lmfa0woukE%3D&gaa_ts=68ab0a85&gaa_sig=5sk0-dOKhxRtACJ8Kuzn4zUam_jPDyKVMohU1cLabICAJDmaEHa2BlJewylVIlDGOT5WHZNrKWoYTWrbMoPUWA%3D%3D

9. Seven Lawsuits Allege OpenAI Encouraged Suicide and Harmful Delusions - WSJ. Accessed November 16, 2025. https://www.wsj.com/tech/ai/seven-lawsuits-allege-openai-encouraged-suicide-and-harmful-delusions-25def1a3

10. Tiku N. Senators weigh regulating AI chatbots to protect kids. *The Washington Post*. https://www.washingtonpost.com/technology/2025/09/16/senate-hearing-ai-chatbots-teens/. September 17, 2025. Accessed November 16, 2025.

11. Tiku N. A teen contemplating suicide turned to a chatbot. Is it liable for her death? *The Washington Post*. https://www.washingtonpost.com/technology/2025/09/16/character-ai-suicide-lawsuit-new-juliana/. September 16, 2025. Accessed November 16, 2025.

12. Campbell LO, Babb K, Lambie GW, Hayes BG. An Examination of Generative AI Response to Suicide Inquires: Content Analysis. *JMIR Ment Health*. 2025;12:e73623. doi:10.2196/73623

13. McBain RK, Cantor JH, Zhang LA, et al. Evaluation of Alignment Between Large Language Models and Expert Clinicians in Suicide Risk Assessment. *Psychiatr Serv 2025 DOI 101176appips20250086*. Published online September 2, 2025. Accessed November 16, 2025. https://www.rand.org/pubs/external_publications/EP71040.html

14. Pichowicz W, Kotas M, Piotrowski P. Performance of mental health chatbot agents in detecting and managing suicidal ideation. *Sci Rep*. 2025;15(1):31652. doi:10.1038/s41598-025-17242-4

15. Arnaiz-Rodriguez A, Baidal M, Derner E, et al. Between Help and Harm: An Evaluation of Mental Health Crisis Handling by LLMs. arXiv.org. September 29, 2025. Accessed November 16, 2025. https://arxiv.org/abs/2509.24857v1

16. Belli L, Bentley K, Alexander W, et al. VERA-MH Concept Paper. *arXiv*. Preprint posted online October 22, 2025. doi:10.48550/arXiv.2510.15297

17. The App Evaluation Model. Accessed November 16, 2025. https://www.psychiatry.org:443/psychiatrists/practice/mental-health-apps/the-app-evaluation-model

18. Byun G, Lipschutz R, Minton ST, Lott A, Choi JD. CRADLE Bench: A Clinician-Annotated Benchmark for Multi-Faceted Mental Health Crisis and Safety Risk Detection. *arXiv*. Preprint posted online October 27, 2025. doi:10.48550/arXiv.2510.23845

19. November 6, 2025: Digital Health Advisory Committee Meeting Announcement - 11/06/2025. FDA. November 13, 2025. Accessed November 16, 2025. https://www.fda.gov/advisory-committees/advisory-committee-calendar/november-6-2025-digital-health-advisory-committee-meeting-announcement-11062025

20. Wang P, Li L, Chen L, et al. Large Language Models are not Fair Evaluators. In: Ku LW, Martins A, Srikumar V, eds. *Proceedings of the 62nd Annual Meeting of the Association for Computational Linguistics (Volume 1: Long Papers)*. Association for Computational Linguistics; 2024:9440-9450. doi:10.18653/v1/2024.acl-long.511

21. Dietz L, Zendel O, Bailey P, et al. Principles and Guidelines for the Use of LLM Judges. In: *Proceedings of the 2025 International ACM SIGIR Conference on Innovative Concepts and Theories in Information Retrieval (ICTIR)*. ICTIR '25. Association for Computing Machinery; 2025:218-229. doi:10.1145/3731120.3744588

22. McCoy TH, Perlis RH. Dimensional Measures of Psychopathology in Children and Adolescents Using Large Language Models. *Biol Psychiatry*. Published online June 10, 2024:S0006-3223(24)01299-X. doi:10.1016/j.biopsych.2024.05.008

23. Barroilhet SA, Bieling AE, McCoy TH, Perlis RH. Association between DSM-5 and ICD-11 personality dimensional traits in a general medical cohort and readmission and mortality. *Gen Hosp Psychiatry*. 2020;64:63-67. doi:10.1016/j.genhosppsych.2020.01.003

24. Moore J, Grabb D, Agnew W, et al. Expressing stigma and inappropriate responses prevents LLMs from safely replacing mental health providers. In: *Proceedings of the 2025 ACM Conference on Fairness, Accountability, and Transparency*. FAccT '25. Association for Computing Machinery; 2025:599-627. doi:10.1145/3715275.3732039

25. Colby KM, Watt JB, Gilbert JP. A COMPUTER METHOD OF PSYCHOTHERAPY: PRELIMINARY COMMUNICATION. *J Nerv Ment Dis*. 1966;142(2):148.

Figures

Figure 1. Schematic diagram of STELLA system architecture, illustrating the combination of personas and scenarios to generate simulated chat transcripts scored by an ensembled judge on multiple dimensions.

Figure 2. Proportion of Turns with Harmfulness Potential Score < 0.7, by Scenario

Figure 3. Proportion of Turns with Helpfulness Score < 0.7, by Scenario

Figure 4. Proportion of Turns with Harmfulness Potential Score < 0.7 (Pooled over all Scenarios)

Figure 5. Proportion of Turns with Helpfulness Score < 0.7 (Pooled over all Scenarios)

Figure 1. STELLA System Architecture

Figure 2. Proportion of Turns with Harmfulness Potential Score < 0.7, by Scenario

Figure 3. Proportion of Turns with Helpfulness Score < 0.7, by Scenario

Figure 4. Proportion of Turns with Harmfulness Potential Score < 0.7 (Pooled over all Scenarios)

Figure 5. Proportion of Turns with Helpfulness Score < 0.7
(Pooled over all Scenarios)