Host Genetic Architecture between Epstein–Barr Virus Activity and Multiple Sclerosis Reveals Shared Pathways

Yoshiaki Yasumizu[1,†,*], Namkwon Kim[1,†], Cyprien A. Rivier[1,2,†], Jeonghyeon Moon[1,†,‡], Shohei Kojima[3,†], Heng-Le Chen[1], Nicholas Buitrago-Pocasangre[1], Elizabeth Quinn[1], Stephen Vaughn[1], Annalisa Morgan[1], Shufan Huo[1,2], Andrew Silberfeld[1,2], Tomokazu S. Sumida[1], Kazuyoshi Ishigaki[3,4,5], Erin E. Longbrake[1], Guido J. Falcone[1,2,*†], David A. Hafler[1,6,7,*†]

[1] Department of Neurology, Yale School of Medicine, New Haven, CT, 06510, USA

[2] Yale Center for Brain and Mind Health, Yale School of Medicine, New Haven, CT, 06510, USA

[3] Keio University Human Biology-Microbiome-Quantum Research Center (WPI-Bio2Q), Tokyo, 160-8582, Japan

[4] Department of Microbiology and Immunology, Keio University School of Medicine, Tokyo, 160-8582, Japan

[5] Laboratory for Human Immunogenetics, Riken Center for Integrative Medical Sciences, Kanagawa, 230-0045, Japan

[6] Department of Immunobiology, Yale School of Medicine, New Haven, CT, 06510, USA

[7] Broad Institute of MIT and Harvard, Cambridge, MA, 02142, USA

† Equal Contributions

‡Current address: Dental and Life Science Institute, School of Dentistry, Pusan National University, Yangsan, 50612, Republic of Korea.

* Correspondence

Abstract

Epstein–Barr virus (EBV) is strongly implicated as an essential environmental trigger of multiple sclerosis (MS), yet the host genetic mechanisms governing EBV activity and how infection triggers the disease are not known. We developed a pipeline to quantify EBV DNA from whole-genome sequencing data and applied it to population-scale cohorts. Using this pipeline, we performed a cross-ancestry genome-wide association study (GWAS) of EBV DNA positivity in 617,186 individuals and identified 39 independent susceptibility risk loci, with the strongest associations in the HLA region. We validated this finding in our independent cohort (N=94) and found that quantitative PCR (qPCR)-confirmed EBV DNA positive individuals were enriched in the top decile of EBV polygenic risk scores (PRS) containing newly discovered loci. A significant overlap with genetic variants associated with MS risk was observed. PRS and Mendelian randomization analyses further supported a causal role of EBV activity on MS risk, which was also seen in other autoimmune diseases. A meta-analysis of qPCR based case–control studies showed elevated EBV DNA positivity in MS. By establishing a single-cell RNA-seq method optimized for EBV detection, we identified EBV-infected B cells, primarily in memory B cells, atypical B cells and antibody-secreting cells from MS and healthy individuals. Notably, EBV-infected memory B cells and atypical B cells showed strong upregulation of cytokines and costimulatory signals that influence T cell activation, IFNg secreting Tregs, and regulators of B cell differentiation and survival. EBV-infected memory B cells also upregulated risk genes from both the EBV and MS GWAS, suggesting that EBV-infected B cells constitute a critical hub that modulates T cell responses while simultaneously activating MS susceptibility pathways within the B cell compartment. Together, these findings define a genetic and cellular framework linking EBV infection to the initiation of MS.

Introduction

Multiple sclerosis (MS) is the most common central nervous system (CNS) disease of young adults. The disease most commonly begins with an autoimmune, relapsing remitting phase mediated by circulating autoreactive T cells[1] that migrate to the central nervous system (CNS) and induce a marked inflammatory response both in the white matter and cortical grey matter. A secondary, neurodegenerative disease course occurs in a significant number of patients, presumably triggered by the initial T cell-mediated autoimmune process. As with other autoimmune disorders, MS is triggered by unfavorable environmental interactions in genetically susceptible individuals. Indeed, we have identified 233 common genome loci associated with disease risk and one variant associated with disease progression.[2–4] The majority of these map to the immune system, targeting T cells, regulatory T cells, and, in particular, B cells.[2,5,6] As a result, MS patients exhibit a spectrum of immunopathology, including a skewing of T cells toward pro-inflammatory Th1 and Th17 programs,[6–8] loss of regulatory T-cell function,[9,10] and activation of B cells.[11]

Compelling epidemiologic data implicates Epstein–Barr virus (EBV) as the initiating event breaking immune tolerance in genetically susceptible individuals. A large longitudinal study of U.S. military personnel reported that EBV infection occurs before MS development[12] , indicating that EBV is a necessary causal factor for MS. Serum neurofilament light chain, a marker of axon injury, begins to rise only after EBV seroconversion. Moreover, MS patients exhibit higher EBV activity,[13–15] increased frequencies of EBV-reactive T cells[16,17], and EBV-derived antigens that cross-react with myelin antigens [18,19]. EBV primarily infects B cells, directly influencing their function[20], and we have observed distinctive alterations in B-cell phenotypes in MS (*manuscript in preparation*). Importantly, B-cell depletion therapy is a highly effective treatment for the early relapsing remitting, autoimmune phase of MS,[21] underscoring the central role of B cells in disease pathogenesis.

EBV plays a causal role in several other human diseases. Epidemiologic and immunologic evidence links EBV to non-Hodgkin lymphoma, including Burkitt lymphoma, nasopharyngeal carcinoma, gastric carcinoma, and, as discussed above, to MS, rheumatoid arthritis (RA), and systemic lupus erythematosus (SLE).[22] Although EBV infection is ubiquitous affecting more than 90% of the global population, its clinical manifestations vary widely. In most individuals, EBV infection remains asymptomatic or latent, whereas a subset of subjects develop infectious mononucleosis or chronic active EBV infection. Moreover, infectious mononucleosis is not only a symptomatic form of EBV infection but also a strong epidemiological risk factor for MS.[23] However, few individuals develop malignancies or

autoimmune diseases after EBV infection. This variability suggests that the host's ability to
control EBV reactivation differs among individuals, potentially due to underlying genetic
factors.[24]

EBV activity can be assessed through two major biomarkers: anti-EBV antibodies and EBV
DNA levels. Immunoglobulin (Ig) G targeting EBV antigens reflects prior exposure. Over
90% of healthy adults and nearly 100% of patients with MS are seropositive.[15,25] In contrast,
EBV DNA directly measures viral activity; blood EBV DNA is increased in patients with MS
and SLE.[14,26] Circulating EBV DNA is also clinically valuable for nasopharyngeal carcinoma
screening.[27] In aggregate, these data suggest that circulating EBV DNA load would be a
novel tool to investigate viral activity as linked to the genetic background.

Genome-wide association studies (GWAS) have allowed identification of genetic variants
associated with various traits such as diseases or viral infections. For example,
measurement of HIV viral load overcame the limited resolution of infection-status GWAS and
successfully identified key loci in the HLA and *CCR5* regions.[28] Recent advances in
sequencing technologies now allow simultaneous quantification of host and viral genomes
from the same dataset. Our group previously developed *VIRTUS*,[29] a pipeline for detecting
viral transcripts from RNA-seq data, and EBV detection from bulk RNA-seq has also been
reported by others.[30] Similar approaches have recently been applied to whole-genome
sequencing (WGS) data, leading to a GWAS of human herpesvirus 6 (HHV-6) viral load that
revealed integration sites and links to SLE disease activity.[31] For EBV, however, prior GWAS
have focused almost exclusively on serologic antibody titers,[32] which capture infection
history rather than active viral replication. Only one prior EBV DNA GWAS, conducted in
10,585 Chinese participants, has been published;[33] limited sample sizes precluded the
identification of genome-wide significant associations.

Here, we analyzed 617,186 participants from the All of Us Research Program[34] and the UK
Biobank,[35] applying harmonized viral-detection and quality-control pipelines across
ancestries. By performing meta-analysis across these cohorts, we effectively increased the
sample size, enabling the largest EBV DNA GWAS (N=617,186) to date (Extended Data Fig.
1). We further integrated phenome-wide association studies (PheWAS), Mendelian
randomization (MR), and polygenic risk score (PRS) analysis to systematically assess the
causal relationships between EBV activity and human diseases. Combining GWAS with
single-cell RNA-seq (scRNAseq) and spatial transcriptomic analyses, we identified specific
cell types and anatomical niches involved in EBV control. Notably, using *VIRTUS3*, we

successfully identified EBV-infected cells from people with MS and characterized their
genetic effects.

Together, these findings provide a large-scale, multi-ancestry genetic map of host control
over EBV activity and EBV infection and transcriptome dynamics in people with MS at
single-cell resolution, offering mechanistic insight into EBV-associated autoimmunity,
establishing a foundation for the discovery of therapeutic targets aimed at modulating EBV
reactivation, and elucidating a mechanism by which genetic variants associated with MS risk
drive EBV infection in B cells, potentially breaking immune tolerance.

Results

EBV DNA detection from WGS

To detect EBV reads in WGS datasets, we developed an analysis pipeline that filters and
counts reads mapped to the EBV decoy sequence in the human reference genome. We
applied our pipeline to the diverse cohort of study participants enrolled in the All of Us
Research Program. After quality control, 409,331 individuals were included. DNA information
was obtained from two sources: whole blood (N=359,796) and saliva (N=48,086). EBV DNA
was detected in 20.8% of blood samples and 52.7% of saliva samples. Age, sex, and
ancestry were significantly associated with EBV positivity. In both blood and saliva, higher
age was strongly associated with higher detection rates (Fig. 1a, blood: OR=1.16 per 10
years, 95% CI=1.15–1.16, $p<1\times10^{-100}$; saliva: OR=1.10 per 10 years, $95\%\ CI$=1.09–1.11,
$p<1\times10^{-65}$). Female sex was associated with a lower rate of EBV positivity compared with
males (Fig. 1a, blood: OR=0.83, $95\%\ CI$=0.81–0.84, $p<1\times10^{-100}$; saliva: OR=0.78, 95%
CI=0.75–0.81, $p<1\times10^{-35}$).

When examining ancestry, Europeans had the lowest EBV positivity, whereas all other
ancestries showed significantly higher odd ratios of EBV expression. In blood, African
(OR=2.17, $p<1\times10^{-100}$), Latino/admixed American (OR=2.02, $p<1\times10^{-100}$), East Asian
(OR=1.47, p=9.2×10^{-43}), Middle Eastern (OR=1.31, p=1.2×10^{-4}), and South Asians
(OR=1.29, p=6.1×10^{-11}) ancestries had significantly higher odds ratios relative to Europeans
(Fig. 1b). In saliva, African (OR=1.61, p=4.1×10^{-65}), Latino/admixed American (OR=2.02,
$p<1\times10^{-100}$), East Asian (OR=1.94, p=5.8×10^{-32}) and South Asian ancestries also showed
increased odds (OR=1.42, p=1.4×10^{-5}) (Fig. 1b). Middle Eastern ancestry showed a trend
for higher EBV positivity in saliva but did not reach statistical significance (OR=1.32,
p=0.078).

Next, we examined the distribution of EBV DNA load. When normalized by the number of
reads mapped to the human genome, the EBV DNA load was on average 3,826-fold higher
in saliva compared to blood (Fig. 1c). Moreover, a subset of individuals carried viral loads
that far exceeded the population average. Indeed, in blood, the top 1% of samples
accounted for 64% of the total EBV DNA load, with the top 5% and 10% contributing 83%
and 91%, respectively. In saliva, the skew was even more pronounced: the top 1%
accounted for 80%, the top 5% for 94%, and the top 10% for 97%. To statistically evaluate
this trend, we approximated the distribution with a generalized Pareto distribution (GPD). A
positive shape parameter (ξ) indicates a heavy-tailed distribution (Fig. 1c). In blood, ξ was
estimated as 0.92 (95% *CI*:0.86–0.98) at the 95th percentile threshold, while in saliva, ξ was
even larger at 1.15 (95% *CI*:1.04–1.25). Varying the threshold from the 90th to 99th
percentiles yielded consistently positive ξ values, confirming the robustness of these results.
These findings demonstrate that the distribution of EBV DNA load is heavy-tailed in both
blood and saliva, suggesting the presence of a small number of super-carriers who
contribute disproportionately to the overall viral burden.

To comprehensively investigate diseases co-occurring with EBV infection, we conducted a
Phenome-wide Association Study (PheWAS). Using European-ancestry participants from All
of Us, we performed a PheWAS of EBV DNA positivity and viral load (Fig. 1d and Extended
Data Fig.2, Supplementary Tables 1 and 2). Both EBV DNA positivity and viral load showed
strong associations with immunosuppressive conditions such as HIV infection, post-
transplant immunosuppression, and malnutrition. Notably, there were also strong
associations with tobacco use and chronic obstructive pulmonary disease (COPD).[36]
Furthermore, the analysis identified a significant association with infectious mononucleosis,
a manifestation of primary EBV infection in adolescents. Unexpectedly, neither MS nor SLE
emerged as positively associated traits (MS: EBV positivity, *OR*=0.83, *FDR*=0.034; viral load,
FDR=0.78. SLE: EBV positivity, *FDR*=0.27; viral load, *FDR*=0.96). In addition to confirming
known associations with rheumatoid arthritis, we observed correlations with encephalitis,
encephalopathy, and polyneuropathies, suggesting links to other inflammatory neurological
diseases. Although more traits were significantly associated with EBV DNA positivity
(*FDR*<0.05, 475 traits) than with EBV viral load (270 traits), the viral load analysis
specifically highlighted associations with virus-related malignancies such as non-Hodgkin
lymphoma and T cell lymphoma, as well as myopathy. The PheWAS results not only
reproduced well-established associations between EBV and related diseases, validating our
analytical pipeline, but also revealed broader links to autoimmune and inflammatory
disorders, underscoring the extensive disease spectrum influenced by EBV infection.

Multi-Ancestry genome-wide association study of EBV DNA

We applied a similar pipeline to the UK Biobank, in which WGS was performed using DNA

derived from whole blood. EBV-derived DNA was detected in 15.5% of UK Biobank samples.

We then performed a cross-ancestry GWAS combining UK Biobank samples with blood-

derived samples from the All of Us cohort. We selected 5,020,877 SNPs with a minor allele

frequency (MAF) of ≥1% in All of Us (Methods). Association tests were performed using

Regenie[37] in all ancestry groups from the All of Us Research Program and in European

participants from the UK Biobank. A European meta-analysis was conducted using METAL[38]

(N=493,312), and a cross-ancestry meta-analysis (European from UK Biobank and

European, African, and Latino/Admixed American from All of Us) was performed using MR-

MEGA[39] (N=617,408) (Supplementary Table 3). In the cross-ancestry analysis, we identified

39 genome-wide significant loci associated with EBV positivity and 16 loci associated with

viral load (Fig. 2a and Extended Data Fig. 4a, Supplementary Tables 4-6). Using LD score

regression (LDSC),[40] the SNP-based heritability (h2) of EBV positivity was estimated at

1.77% (*SE*=0.22%), while that of viral load was 1.24% (*SE*=0.18%). There was a positive

genetic correlation between EBV positivity and EBV viral load (*rg*=0.71, *p*=4.08x10^{-24}).

We observed a slight inflation of test statistics in the cross-ancestry meta-analysis,

particularly for the binary trait (Extended Data Fig. 3a, EBV positivity, λ_{GC}=1.15; viral load,

λ_{GC}=1.07). Importantly, the λ_{GC} values from each individual GWAS were well controlled

(λ_{GC}<1.08 in all cases, Supplementary Table 3). We evaluated whether the MHC contributed

to the inflation; removing this region yielded comparable values (EBV positivity, λ_{GC}=1.14;

viral load, λ_{GC}=1.06). Of note, in the LDSC analysis, intercepts were low (EBV positivity,

intercept=1.04; viral load, *intercept*=1.03), indicating that the observed inflation is likely due

to polygenicity rather than residual confounding.

Next, we examined the lead SNPs within each locus in detail. Outside of the HLA region,

approximately half of the associated variants were intronic (Supplementary Table 4,

positivity:63.1%; viral load:53.3%), and roughly half were located in cis-eQTLs

(Supplementary Table 4, positivity:55%; viral load:60%). The majority of these loci

(Supplementary Table 4, positivity: 65.8%; viral load: 60%) have been previously implicated

in immune-related traits, including MS, SLE, RA, IgM levels, and blood cell counts in prior

GWAS (Fig. 2a). Remarkably, there was a significant overlap with genetic variants

associated with MS risk (Fig. 2B, *OR*=32.1, *p*=1.3x10^{-12}) and severity (Fig. 2B, *OR*=180.6,

p=7.54x10^{-3}) including MS susceptibility alleles, *HLA-DRB1**04:04, *HLA-A**02:01 and to a

lesser extent, *HLA-DRB*1*15:01. Specifically, among the top hits, 29.7% of the EBV

positivity-associated genes and 33% of the viral load-associated genes overlapped with

genes reported in MS GWAS, including *DYSF*, which has been associated with MS severity[4]
(Fig. 3b and Extended Data Fig.4b). When we examined the direction of effects for EBV
DNA positivity across autoimmune diseases, we found that *RSBN1*, *SLAMF7*, *CTLA4*,
EOMES, *CD86*, *TP63*, *TRAF3*, *CLEC16A*, and *IKZF3* showed concordant effects, with
increased EBV DNA positivity associated with increased disease risk, while *SP140*, *SH2B3*,
and *PTPN11* exhibited discordant directions of effect with less EBV DNA expression (Fig.
2a). The genes highlighted by the GWAS provided biological insights into the regulation of
EBV activity. For example, *TERT*, encoding the catalytic subunit of telomerase, emerged as
one of the top hits. Indeed, the top variant within this locus, rs7726159, is associated with
telomere length[41] (Supplementary Table 5, Extended Data Fig. 4c). Given that EBV is known
to drive telomerase regulation to promote immortalization of infected cells,[42] this finding may
highlight a mechanistic link between EBV replication and telomere biology. We also
observed strong support for biologically plausible genes, including *IFI16*, a nuclear DNA
sensor that restricts herpesvirus reactivation, and *SP140*, a PML-nuclear body component
involved in antiviral chromatin regulation (Extended Data Fig. 4d).[43,45,47]

Stratified LD score regression[44] revealed that genetic heritability for both EBV DNA positivity
and EBV DNA load was concentrated in transcriptionally active genomic regions, particularly
at transcription start sites (TSS; enrichment=23.4±4.1 [positivity], 31.3±5.3 [load]),
FANTOM5 enhancers (19.9±10.3 [positivity], 32.7±13.8 [load]), and coding regions (17.2±5.2
[positivity], 17.0±6.3 [load]), indicating that genetic variants influencing EBV control tend to
localize to regulatory and coding elements with high transcriptional activity (Extended Data
Fig. 3B). Pathway enrichment analysis revealed that EBV susceptibility loci were primarily
concentrated in immune response–related pathways (Fig. 2c, Extended Data Fig. 4e,
Supplementary Tables 7 and 8). Both EBV positivity and EBV DNA load showed significant
enrichment for T cell activation, CD8[+] αβ T cell activation, and T cell proliferation, reflecting a
shared involvement of cytotoxic T cell–mediated immune responses. In contrast, cell cycle
G1/S phase transition was specifically enriched for EBV positivity, while EBV DNA load
additionally involved epithelial-associated pathways such as positive regulation of
keratinocyte proliferation and multiciliated epithelial cell differentiation, indicating potential
contributions of epithelial cell proliferation and differentiation to viral load regulation.

Tissue-, Cell-, and Niche-Specific Heritability Analyses
To elucidate the biological contexts underlying host control of EBV activity, we performed
partitioned heritability analysis using stratified LD score regression (S-LDSC),[46] which infers
tissue-level enrichment based on external functional annotations. Both EBV positivity and

EBV DNA load exhibited marked enrichment in immune-related tissues (Fig. 2d and
Extended Data Fig. 4f). For EBV positivity, significant heritability enrichment was observed in
spleen, peripheral blood, lymph nodes, synovial fluid, and lung, suggesting that systemic and
mucosal immune environments contribute to host susceptibility to EBV activity. In contrast,
EBV DNA load showed its strongest enrichment in EBV-transformed lymphocytes, followed
by spleen, lymphoid tissue, and blood, indicating a link between viral load regulation and B
cell transformation processes.

We next sought to interpret the polygenic signals identified in the EBV GWAS in the context
of specific cell types and spatial niches using integrated single-cell and spatial transcriptomic
analyses. We first applied single-cell disease relevance score (scDRS)[48] to single-cell RNA-
seq data from PBMCs of EBV-positive and control individuals[49], analyzing all cell types
jointly. This analysis revealed enrichment of EBV-associated polygenic signals in cytotoxic
lymphocyte populations, including CD8 effector memory T cells, NK cells, and gd T cells
(Extended Data Fig. 5a).

Using the same dataset, we then focused the analysis on B-cell lineages that are directly
involved in EBV infection. The dataset included publicly available single-cell RNA-seq
profiles from patients with infectious mononucleosis and hemophagocytic
lymphohistiocytosis (HLH). Existing pipelines based on Cell Ranger remove overlapping
reads, resulting in substantial loss of reads from the compact EBV genome, and they may
misalign reads with sequence similarity to human transcripts.[50] To overcome these
limitations, we developed VIRTUS3, an analysis framework optimized for viral detection in
single-cell data (https://github.com/yyoshiaki/VIRTUS3). In this workflow, reads are first
aligned to the human genome using Cell Ranger. Unmapped reads are then recovered and
re-aligned to viral genomes using Alevin. This pipeline can filter out reads with stringent
similarity to the human genome, and also manipulate the reads aligned to multiple genes on
viral genomes. This approach enabled the detection of 1.84-fold more EBV-positive cells
compared with conventional methods (Figs. 3a-c). EBV-positive cells were predominantly
detected among plasmablasts, followed by B memory and B intermediate cells, whereas
naïve B cells showed almost no EBV positivity (Fig. 3d). Using scDRS, we further evaluated
the cell-type–specific distribution of EBV DNA-associated polygenic scores. In memory B
cells, polygenic scores were significantly elevated in EBV-positive compared with EBV-
negative cells ($p=3.0\times10^{-\square}$), while no difference was observed in plasmablasts (Fig. 3e and
Extended Data Fig. 5b). These findings suggest that host genetic control of EBV activity is
exerted primarily within the memory B-cell compartment rather than at the terminally
differentiated plasmablast stage.

We next investigated the immune microenvironments in which host genetic factors regulating
EBV activity are concentrated by applying scDRS-spatial, a spatial heritability framework we
previously developed[51]. At the tissue level, polygenic signals showed the strongest
enrichment in lymph nodes and tonsils, confirming that these lymphoid tissues represent
major sites of EBV regulation (Extended Data Fig. 5c). We then analyzed the spatial
localization of genetic signals within tonsillar tissue, a key reservoir for EBV infection, latency,
and reactivation. Polygenic signals for both EBV DNA positivity and viral load were highly
enriched within the germinal center, T:B border, and proliferating follicle regions, which are
immune niches characterized by active B cell responses and dynamic B–T cell interactions
(Figs. 3f–h and Extended Data Fig. 5d). These findings indicate that host genetic factors
controlling EBV activity act predominantly within lymphoid microenvironments supporting T
cell–dependent B cell activation. In particular, the germinal center and T:B border emerge as
key immunological niches where genetic variation may influence EBV persistence and
immune control.

Antigen presentation by HLA and EBV activity

The strongest associations in both GWASs were located within the HLA locus. In addition, a
secondary genome-wide significant signal was identified at rs27300, a variant residing within
a linkage disequilibrium block encompassing the *LNPEP* and *ERAP1/2* genes (Fig. 4a and
Extended Data Fig. 6a). This variant exhibited a significant eQTL effect on *ERAP2*
expression in lymphoblastoid cell lines (LCLs) (p=2.33x10^{-118}, Fig. 4b), but not with *LNPEP*
(p=7.07x10^{-6}, Fig. 4b). *ERAP2* encodes an aminopeptidase essential for peptide trimming
selectively during antigen presentation via MHC class I molecules. This allelic directionality
provides additional support for a functional role of *ERAP2* in EBV control. The *ERAP2*-
increasing C allele corresponds to the protective direction in the EBV GWAS, suggesting
that more efficient peptide trimming enhances antigen presentation and improves the
clearance of EBV-infected cells.

We also investigated the associations with HLA alleles and amino acid usage and EBV
susceptibility. Using All of Us participants of European ancestry, classical HLA alleles were
imputed with SNP2HLA,[52] and their associations with EBV activity (DNA positivity and DNA
viral load) were examined. In both traits, *HLA-DRB1**04:04 showed the most significant
association. After conditioning analysis, the top alleles associated with EBV DNA positivity
were *HLA-DRB1**04:04, *B**14:02, *A**02:01, *DQA1**05, and *B**35:01, while those for EBV DNA
load were *HLA-DRB1**04:04, *A**02:01, *B**35:01, *DQA1**05:01, and *B**14, showing a broadly
consistent pattern between the two traits (Fig. 4C, Extended Data Figs. 6B, 7, 8,
Supplementary Tables 9, 10). While to a lesser degree, EBV DNA positivity was also seen

with the strongest MHC associated allele, *DRB1**15:01. Furthermore, an amino acid–level
association analysis identified multiple residues significantly correlated with EBV activity (Fig.
4c, Extended Data Figs. 6b, 7, 8, and Supplementary Tables 11, 12). Notably, valine at
position 95 of *HLA-A**02:01 (A95V), detected in both positivity and viral load analyses, was
located within the F pocket of the peptide-binding groove, while position 30 of HLA-DRB1
(L/R/Y variants), detected in the viral load analysis, was located in the P7 pocket (Fig. 4d).
Both residues are situated within key structural regions that determine peptide-binding
specificity, suggesting that EBV-associated HLA signals may reflect functional variation in
peptide-binding pockets influencing viral antigen presentation and immune recognition.
These findings suggested that antigen presentation is a key component of EBV regulation
and a hotspot of susceptibility.

Causal inference of EBV activity on autoimmune diseases through Mendelian randomization

To test whether EBV DNA positivity was causally associated with risk of MS and other major
autoimmune diseases, we performed bidirectional Mendelian randomization (MR). For this
purpose, we leveraged a set of genetic instruments derived from the European meta-
analysis of EBV DNA positivity (27 independent SNPs after accounting for linkage
disequilibrium). We observed significant associations between genetically-determined EBV
DNA positivity and MS (*OR*=1.45, *95% CI*=1.18–1.80, *p*=5.9×10⁻☐), RA (*OR*=1.27, *95%*
CI=1.05–1.54, *p*=0.015), and SLE (*OR*=1.44, *95% CI*=1.08–1.91, *p*=0.012) using MR-
PRESSO[53]-corrected IVW estimates, with sensitivity analyses using the Weighted Median
method confirming the directionality of these associations (MS: *OR*=1.41, *95% CI*=1.08–1.83,
p=0.011; RA: *OR*=1.28, *95% CI*=1.06–1.56, *p*=0.012; SLE: *OR*=1.65, *95% CI*=1.37–1.99,
p=1.2×10⁻⁷) (Fig. 5a). Ischemic stroke was included as a negative control and showed no
association with EBV DNA positivity (*OR*=0.97, *95% CI*=0.93–1.02, *p*=0.31). Stratification of
instruments showed that robust associations were observed for MHC variants (IVW: MS
OR=1.69, 95% *CI*=1.57–1.82, *p*=3.5×10⁻⁴⁶; RA *OR*=1.30, *95% CI*=1.23–1.37, *p*=4.7×10⁻²¹;
SLE *OR*=1.74, *95% CI*=1.55–1.94, *p*=1.9×10⁻²²), while non-MHC variants showed no
significant causal effects on any autoimmune disease, indicating that the MHC explains a
large proportion of the inferred causality for these autoimmune diseases (Extended Data Fig.
9a). In the reverse direction, genetic liability to MS, RA, and SLE was not significantly
associated with EBV DNA levels (all *p*>0.05 for IVW and Weighted Median), supporting a
unidirectional causal effect of EBV activity on autoimmune disease risk rather than reverse
causation.

Polygenic risk scoring and EBV DNA measured by quantitative PCR (qPCR)

To evaluate whether identified genome-wide significant risk loci for EBV DNA capture biologically meaningful variation in viral load, we assessed polygenic risk scores (PRSs) in an independent cohort with qPCR-based measurement of EBV DNA. We calculated PRSs based on risk variants identified in the meta-analysis of European ancestry using PRScs.[54] We evaluated association with EBV DNA positivity measured by qPCR in an independent Yale cohort of European ancestry (*N*=94; mean age 40.2±14.7 years; 71% female). For PRS based on 729,766 variants across the whole genome, EBV DNA positivity was 2/10 (20%) in the top decile, 1/74 (1.4%) in the middle group, and 0/10 (0%) in the bottom decile for both positivity and viral load PRS (Fig.5b). Ordinal trend tests remained significant after covariate adjustment (positivity: *p*=0.024; viral load: *p*=0.016). Consistently, a 1 standard deviation increase in the PRS was associated with a higher EBV DNA load after adjustment (log10 EBNA1 copies/mL) (positivity: *β*=0.21, *SE*=0.076, *p*=0.006; load: *β=0.26*, *SE*=0.075, *p*=0.0009). When we stratified variants by their location inside or outside the MHC region, only the PRSs derived from the MHC region showed significant associations with qPCR-confirmed EBV DNA positivity and load (adjusted trend tests: positivity *p*=0.006; load *p*=0.014; effect sizes for 1 SD increase: positivity *β*=0.24, *SE*=0.074, *p*=0.001; load *β*=0.27, *SE*=0.074, *p*=0.0003) (Extended Data Fig. 9b). In contrast, PRSs constructed from variants outside the MHC region showed no significant association (adjusted trend *p*>0.65 for both positivity and load), underscoring the contribution of HLA variation. Taken together, these findings demonstrate that qPCR-measured EBV DNA levels in our independent cohort closely track with the polygenic architecture identified by our EBV DNA GWAS, supporting the validity and biological relevance of the GWAS results.

Association of PRS of EBV DNA with multiple sclerosis

Next, to investigate how the polygenic signal of EBV activity contributes to MS onset, we analyzed PRSs derived from EBV DNA GWAS. We first constructed PRSs using the EBV DNA GWAS performed in the European ancestry subset of the All of Us cohort and applied them to individuals of European ancestry in the UK Biobank (*N*=487,181; 2,423 MS cases; 484,758 controls). A 1 standard deviation increase in the EBV DNA positivity PRS was associated with a significant increase in MS risk (*OR*=1.11, *95% CI*=1.07–1.16, *p*<0.0001) after covariate adjustments. Stratification of individuals by PRS percentiles (bottom 10%, middle 80%, top 10%) revealed a significant ordinal association between higher EBV DNA PRS and increased risk of MS (Fig. 5c). MS prevalence increased from 0.52% in the bottom decile to 0.68% in the top decile, with individuals in the top decile showing a 33% increased odds of MS compared with those in the bottom decile (*OR*=1.33, *95% CI*=1.13–1.57,

p=0.0006). When we further separated PRSs into variants inside versus outside the MHC
region, the association remained robust for MHC-based scores (*OR per SD*=1.12, *95%*
CI=1.07–1.16, p<0.0001; top vs bottom decile *OR*=1.44, 95% *CI*=1.22–1.70, p<0.0001) but
not for scores derived from non-MHC variants (*OR per SD*=0.99, 95% *CI*=0.95–1.03,
p=0.71), indicating that MHC variation largely accounts for the shared genetic architecture
between EBV activity and MS susceptibility. Quintile analyses confirmed this pattern, with
individuals in the top quintile of the whole-genome EBV DNA positivity PRS showing 43%
higher odds of MS compared with the bottom quintile (*OR*=1.43, *95% CI*=1.26–1.61,
p<0.0001), an effect entirely attributable to MHC variants (MHC-only Q5 vs Q1: *OR*=1.45,
95% *CI*=1.28–1.64, p<0.0001; non-MHC Q5 vs Q1: *OR*=1.01, *95% CI*=0.89–1.14, p=0.91).
Taken together, these results suggest that the polygenic burden associated with EBV activity
meaningfully contributes to MS risk.

EBV DNA detection in blood from MS patients
We next examined whether EBV DNA levels are directly associated with MS by qPCR. We
compared whole-blood EBV DNA positivity between MS patients and non-MS individuals
collected at Yale. EBV DNA was detected in 10.9 percent of MS patients (24/220) and 7.65
percent of controls (13/170), showing a higher proportion in MS, although the difference did
not reach statistical significance (p=0.301, tested by Fisher's Exact Test).

To more rigorously assess this association, we conducted a meta-analysis of 11 studies
encompassing 1,725 individuals and 526 EBV DNA–positive events (Fig. 6a, Extended Data
Fig. 10, Table S13). The fixed-effect model yielded an odds ratio of 1.77 (95% *CI*=1.41–2.22,
p<0.0001), and the random-effects model produced a nearly identical estimate (*OR*=1.76,
95 % *CI*=1.34–2.33, p=0.0010). Heterogeneity was low (I^2 = 9.8 percent), indicating
consistent effects across studies. Although the effect size in MS is more modest than in
other autoimmune diseases such as SLE or RA, where odds ratios typically range from 3.5
to 5,[26,55] well-controlled case–control comparisons consistently demonstrate higher
circulating EBV DNA in MS. This pattern reinforces the concept that EBV activity is elevated
around the time of MS onset.

Single-cell resolution cellular dynamics in EBV-infected B cells in people with
MS
We next sought to directly characterize the interaction between EBV infection and MS-
related genetic programs by performing single-cell analyses of PBMCs from MS and healthy
individuals with EBV detection at single-cell resolution. A comprehensive survey of public
scRNA-seq datasets revealed virtually no EBV-positive cells.[56] This was unlikely to reflect a

true biological absence but rather a technical limitation of conventional single-cell RNA-seq,
which captures only polyadenylated transcripts and therefore fails to detect the major latent
EBV noncoding RNA *EBER1*, which lacks a polyA tail. To overcome this limitation, we
introduced a spike-in probe enabling specific capture of *EBER1*, which is a similar approach
to EBV-seq.[57] Yet even with this improved method, EBV-infected cells remained exceedingly
rare, with none detected in PBMCs (0 of 51,950 cells from 3 donors) and only three cells
detected in tonsils (3 of 134,073 cells from 12 donors) from both healthy donors and MS
patients, underscoring a fundamental limitation of single-cell approaches when the infected
population is extremely sparse (Extended Data Fig. 11).

Previous studies have shown that blood EBV DNA load strongly correlates with the number
of *EBER1*-positive cells,[58] suggesting that EBV DNA–positive samples may enable the
recovery of EBV-infected cells at single-cell resolution. We therefore analyzed banked frozen
PBMCs from EBV–positive individuals screened by qPCR analysis. We performed single-cell
RNAseq with the *EBER1* spike-in probe, prioritizing EBV DNA–positive MS and non-MS
samples. With this technique, we could detect 247 DNA-positive cells from 201,398 B cells
(0.12%) (Figs. 6b and c). The EBV–positive cells were predominantly memory B cells and
plasma cells (Fig. 6c and Extended Data Fig. 12a), and the number of EBV–positive cells
were positively correlated with qPCR-based EBV DNA load (r=0.56, p=0.035, Fig. 6d). While
we did not find a significant difference in terms of somatic hyper mutation (p=0.0879,
Extended Data Fig. 12b), *IGHM* were more frequent and *IGHA1*, and *IGHD* were less
frequent (Extended Data Fig. 12c). Most of the EBV–positive cells expressed latent EBV
genes including *EBER1*, while some atypical B cells, naïve B cells, intermediate memory B
cells, and antibody-secreting B cells expressed lytic genes (Fig. 6e).

Next, we examined how EBV-positive B cells alter their transcriptional programs. EBV
infection induced two broad patterns of gene expression changes, separating mature
populations including antibody-secreting cells, switched memory B cells from more immature
populations naïve (naïve, unswitched-memory, intermediate memory, and atypical B cells
(Fig. 6f, Table S15). Pathway analysis revealed marked upregulation of BCR signaling,
antigen receptor–mediated signaling, B-cell activation, and RNA splicing in memory B cells
and antibody-secreting cells, whereas antigen presentation via MHC class I was particularly
enriched in naive B cells (Fig. 6g, Table S16). A focused analysis of cytokines and receptors
involved in immune interactions (Fig. 6h) showed increased expression of *IL12A* and *IL23A*,
in EBV infected B cells which promote Th1 and Th17 differentiation, together with reduced
expression of the immunoregulatory cytokine *IL10* and its receptor *IL10RA*. EBV-positive
cells also upregulated *IL16*, *IL6ST* (gp130), and *IL6R*, molecules implicated in MS

pathogenesis. While *CXCR4* was downregulated, molecules involved in recruiting and
activating pathogenic T-cell subsets, including *CCR6* (Th17-associated) and *CXCR3* (Th1-
associated), were also increased. In addition, EBV-positive cells showed elevated
expression of receptors crucial for B-cell differentiation and survival, including *CXCR5*,
TNFRSF13B (TACI), *TNFRSF13C* (BAFF-R), *TNFRSF14* (HVEM), and *TNFRSF17* (BCMA).

Finally, we investigated how host genetic factors interact with EBV infection across B-cell
subsets. Using scDRS to evaluate EBV GWAS signals, we found that the EBV DNA
positivity–associated polygenic signal was significantly enriched in EBV-positive B naïve and
switched memory B cells, whereas no meaningful enrichment was observed in antibody-
secreting cells (Fig. 6i). These findings suggest that early infection responses in naïve B
cells and persistent infection in switched memory B cells represent key cellular contexts in
which EBV regulation is exerted. We then performed a similar analysis using MS onset–
associated gene sets. Polygenic scores for MS were markedly elevated in EBV-positive
unswitched memory B cells and switched memory B cells, while antibody-secreting cells
showed a reduction in MS-related signal (Fig. 6i). Specifically, MS risk gene expression was
increased specifically within EBV-infected cells in atypical B cells and unswitched-memory B
cells subsets, indicating that EBV-infected B cells constitute a biological context where MS
risk genes are preferentially engaged. Notably, many MS susceptibility genes, EBV
susceptibility genes, and shared loci between the two GWAS, including *IKZF3*, *SP140*,
CD86, *NFKB1*, *TRAF3*, and *LPP*, showed increased expression in EBV-positive cells (Fig.
6j). These observations indicate that EBV infection activates a gene network that intersects
with MS susceptibility specifically within defined B-cell subsets. Taken together, these results
demonstrate that EBV infection enhances both antiviral polygenic signals and MS
susceptibility gene expression in select B-cell subsets, revealing a cellular setting in which
viral regulation and autoimmune genetic risk converge.

Discussion

Here, we developed a pipeline to quantify EBV DNA from whole-genome sequencing data
and applied it to population-scale cohorts. A cross-ancestry GWAS of EBV DNA positivity in
617,186 individuals identified 39 susceptibility risk loci. We validated this finding in our
independent cohort and found that qPCR-confirmed EBV DNA-positive individuals were
markedly enriched in the top decile of EBV polygenic risk scores containing these newly
discovered loci. Surprisingly, there was a significant overlap with genetic variants associated
with MS, RA, and SLE susceptibility alleles, and Mendelian randomization analyses further
supported a causal influence of EBV activity on MS, largely driven by HLA variation. By

establishing a single-cell RNA-seq method optimized for *EBER1* detection, we found that EBV DNA expression was primarily in memory, atypical, and antibody-secreting B cells and that EBV-infected memory B cells strongly upregulated cytokines and risk genes from both the EBV and MS GWAS. These data demonstrate that EBV–infected B cells are a key modulator of T cells and that MS susceptibility genes are preferentially activated within EBV-infected B cells, defining a genetic and cellular framework linking EBV infection to MS and potentially other autoimmune diseases.

MS is the autoimmune disease for which EBV is most strongly implicated as a necessary environmental factor, and both tissue and cell specificity of EBV regulation further align with MS-specific immunobiology in which memory B cells drive CNS infiltration.[59] Moreover, it has been clearly demonstrated that the EBV virus harbors sequences that are cross-reactive with myelin epitopes including GlialCAM that are likely also playing a critical role in the pathogensis of MS[18,19]. Thus, EBV uniquely may drive activation of myelin reactive T cells both by breaking immune tolerance and by presenting crossreactive antigens.

Although EBV DNA showed no positive association with MS in our PheWAS,[60] well-controlled case-control meta-analyses have consistently reported elevated EBV DNA levels in MS patients. This discrepancy is likely explained by the time course of EBV involvement in the onset of MS. Specifically, EBV seroconversion occurs an average of 7.5 years before clinical onset,[12] MS diagnosis requires CNS lesions disseminated in time and space,[61] and a prodromal period precedes diagnosis.[62] As a result, EBV load at the time of diagnosis is likely to be partially deflated. Furthermore, individuals with MS in cohorts such as All of Us include patients receiving long-term immunotherapy and those many years after diagnosis, making it difficult to capture EBV activity during the true initiation phase. These considerations highlight the importance of examining newly diagnosed, treatment-naïve patients or MS high-risk individuals prior to disease onset to accurately evaluate EBV activity in early MS.

Our large-scale WGS analysis also clarified the broader epidemiology and biology of EBV infection across populations. In the United States, individuals of African, Latino/admixed American, and Asian ancestry showed higher EBV positivity rates, consistent with serology-based findings.[63] Female individuals showed lower EBV DNA positivity despite a higher prevalence of autoimmunity. This pattern suggests that sex differences in EBV susceptibility and the host immune response to EBV may shape autoimmune risk. We also found that a small subset of "super-carriers" accounted for a disproportionate fraction of EBV DNA detected in the population.

Cross-ancestry GWAS clarified the host genetic architecture underlying EBV activity. Strong
enrichment of associations in the HLA region, in T-cell–related pathways including
association in *CTLA4*, *EOMES*, and *CD86*, and at the T–B border of secondary lymphoid
tissues, underscores the central role of B cells in antigen presentation to T cells in EBV
regulation. EBV employs multiple immune-evasion strategies, including BNLF2a-mediated
inhibition of TAP1.[64] Here, we showed that HLA polymorphisms shape antigen presentation
efficiency and directly influence EBV activity. *HLA-DRB1**04:04, a known risk allele for RA
and MS,[65,66] and an allele that presents myelin-specific TCRs in MS patients,[67] was
associated with higher EBV activity. The risk allele *HLA-DRB1**15:01 was also associated
with higher EBV presentation, though to a lesser extent than the *HLA-DRB1**04:04 allele,
suggesting other factors contribute to the genetic risk associated with these MHC haplotypes.
In contrast, *HLA-A**02:01 was associated with reduced EBV activity and is protective against
MS[68] and EBV⁺ Hodgkin lymphoma.[69] HLA molecules shape antigen-presentation
preferences and thymic TCR repertoire formation.[70] In addition, components of the antigen-
processing pathway, including *ERAP2*, emerged as important regulators of EBV control. Our
findings indicate that HLA diversity and related genes exert strong effects on EBV
susceptibility and that several EBV-associated alleles overlap with autoimmune risk alleles,
confirming suggested shared genetic pathways linking EBV control and autoimmunity.[71]
Moreover, PRS and MR analysis highlighted the HLA contribution to autoimmune diseases
including MS.

Non-HLA loci further underscore that EBV regulation relies heavily on intrinsic programs of
the host B cell. Autoimmune diseases and infections exhibit broad sharing of genetic risk
variants across conditions.[72] Genes in which the direction of effect was concordant between
EBV DNA positivity and autoimmune disease risk, including *RSBN1*, *SLAMF7*, *CTLA4*,
EOMES, *CD86*, *TP63*, *TRAF3*, *CLEC16A*, and *IKZF3*, converged on pathways governing B-
cell differentiation, germinal-center dynamics,[73] NF-κB signaling,[74] T cell effector function,
and antigen presentation. This pattern suggests that B-cell differentiation and activation
programs provide a shared biological foundation that simultaneously promotes EBV
persistence and increases autoimmune susceptibility. In contrast, genes showing discordant
effects, such as *SP140*, *SH2B3* and *PTPN11*, map to pathways involved in
immunoregulatory thresholds and negative feedback, raising the possibility that trade-offs
between antiviral responsiveness and autoimmune risk underlie their opposing directions of
effect. *SP140* localizes to promyelocytic leukemia protein nuclear bodies, a key site of viral
genome surveillance, and can repress viral gene activation.[75,76] There were also loci whose
directional relationship with autoimmunity was unclear but nonetheless pointed to plausible

mechanisms of EBV regulation. For example, *IFI16* functions as an intrinsic viral DNA
sensor that can detect EBV genomes in the nucleus, restrict lytic reactivation, and participate
in epigenetic silencing of viral DNA.[77] Associations in the *TERT* locus further suggests that
variation in telomere biology may influence EBV replication and the long-term survival of
infected B cells, consistent with the ability of EBV to co-opt telomerase activity during
immortalization. Together, these loci highlight a second layer of EBV control, the intrinsic
programs of the infected B cell, which operate in parallel with HLA-mediated antigen
presentation. This integrated view suggests that host genetic variation shapes not only
antiviral immunity but also the cellular context that EBV exploits to sustain lifelong infection
and influence on autoimmunity.

Building on these findings, we further leveraged three key advances, improved single-cell
detection of viral transcripts, an optimized bioinformatics pipeline, and sample prioritization
based on EBV DNA by qPCR, to achieve the direct identification of EBV-infected cells in
patients with MS at single-cell resolution. By analyzing a large cohort enriched for untreated,
newly diagnosed MS cases, we were able to capture EBV-positive cells during the clinically
relevant early phase of disease. These infected cells were concentrated within memory B
cells, ABCs and ASCs.

EBV-infected B cells showed a coordinated activation of immune communication pathways,
including upregulation of *IL12A* and *IL23A*, which drive Th1 and Th17 differentiation. IL-12
has been shown to diminish Treg suppressive capacity in inflammatory settings, shifting the
balance toward effector responses.[78] This pro-inflammatory shift was further reinforced by
reduced expression of *IL10* and its receptor *IL10R*, consistent with impaired regulatory B-cell
function described in MS.[79] EBV-positive cells also increased *IL16*, *IL6ST* (gp130), and IL6R,
molecules implicated in MS pathogenesis, and upregulated the chemokine receptors *CCR6*
and *CXCR3*, which position B cells within Th17- and Th1-rich inflammatory niches rather
than homeostatic environments. In parallel, EBV-positive B cells showed elevated
expression of receptors crucial for B-cell differentiation and survival, including *CXCR5*,
TNFRSF13B (TACI), *TNFRSF13C* (BAFF-R), *TNFRSF14* (HVEM), and *TNFRSF17* (BCMA).
Together, these findings indicate that EBV infection simultaneously activates programs that
potentiate pathogenic T-cell–B-cell interactions, promote trafficking to inflamed sites, and
reinforce intrinsic B-cell maturation and survival, revealing a broad rewiring of B-cell
immunobiology relevant to MS.

Notably, both MS susceptibility genes and EBV activity–associated genes were highly
expressed within these EBV-infected memory B cells. This represents the first direct

evidence that host genetic risk for MS and genetic determinants of EBV activity converge
within EBV-infected memory B cells. Given that memory B cells serve as potent, antigen-
specific professional antigen-presenting cells, targeting this compartment provides a
compelling biological rationale for the remarkable efficacy of B-cell–depleting therapies in
MS.

Taken together, our work identifies the epidemiology of EBV across ancestries and
elucidates the genetic basis of EBV B cell expression. By integrating multiple omics data, we
mapped the key immunological "hotspots" where this regulation occurs. These findings
support the existence of shared biological pathways linking EBV to autoimmune diseases
including MS. As the genetic variants associated with MS are linked to EBV DNA expression,
our results are also consistent with the higher EBV DNA expression in MS B cells. Future
studies using Massive Parallel Reporter Assays of autoimmune genetic variants will allow
the determination of how EBV gene products interact with disease associated haplotypes in
driving B cells to a state of activated, antigen presentation state potentially breaking immune
tolerance triggering autoreactive T cells.

Data and code availability

GWAS summary statistics will be available at the GWAS Catalog and PGS Catalog upon
acceptance. Locusplot for all lead SNPs and the adjacent region is available at
10.6084/m9.figshare.30712496. Single-cell RNAseq data generated in this study will be
deposited in GEO and CZ CELLxGENE upon acceptance. Code generated for this work will
be publicly accessible on GitHub upon acceptance. The pipeline used for EBV DNA
detection from WGS is available at https://github.com/shohei-kojima/WGS_EBV_search.
VIRTUS3 is available at https://github.com/yyoshiaki/VIRTUS3. All codes used for the
analysis will be available on github
(https://github.com/yyoshiaki/EBV_MS_2025_Manuscript) upon acceptance.

Author contributions

622 Conceptualization, Y.Y., S.K., K.I., C.R., and D.A.H; data analysis, Y.Y., C.R., S.K., N.K.,
623 and H.L.C.; experiment, N.K. and J.M.; data curation and sample collection, L.Q., S.H., A.S.,
624 N.B.P, A.M. and E.E.L.; writing, Y.Y., N.K., C.R., S.K., E.E.L., G.J.F. and D.A.H.; supervision,
625 Y.Y., T.S.S., K.I., E.E.L., G.J.F., and D.A.H.; funding acquisition, Y.Y., D.A.H., and G.J.F.

Competing interests

Conflict of interest: DAH has received research funding from Bristol Myers Squibb, Novartis,
Sanofi, and Genentech. He has been a consultant for Bayer Pharmaceuticals, Repertoire

Inc., Bristol Myers Squibb, Compass Therapeutics, EMD Serono, Genentech, Juno

Therapeutics, Novartis Pharmaceuticals, Proclara Biosciences, Sage Therapeutics, and

Sanofi Genzyme. EEL has received research support from Biogen, LabCorp, Intus, and

Genentech. She has been a consultant for Bristol Myers Squibb, EMD Serono, Genentech,

Sanofi Genzyme, and NGM Bio. AM has been a consultant for Genentech.

634   Acknowledgement

This work was supported by NIH grants (P01 AI073748, U19 AI089992 U24 AI11867, R01

AI22220, UM 1HG009390, P01 AI039671, P50 CA121974, and R01 CA227473); the Race

to Erase MS (to DAH); the National MS Society (to DAH and EEL); Department of Defense

(HT9425-24-1-0522 to EEL); the Common Mechanisms in Autoimmunity Insight Award

jointly funded by Breakthrough T1D, the Lupus Research Alliance, and the National MS

Society (to YY), and Robert E. Leet and Clara Guthrie Patterson Trust Mentored Research

Award, Bank of America, Private Bank, Trustee (to YY). YY is supported by JSPS Overseas

Research Fellowships. NK is supported by the Sejong Science Fellowship (Overseas) from

the National Research Foundation of Korea (RS-2024-00342077). CR is supported by an

AAN/AHA Ralph L. Sacco Scholars Fellowship (AHA.24RSSPOST1328228), a Pilot Grant

from the Claude D. Pepper Older Americans Independence Center at Yale School of

Medicine (P30AG021342), and a Pilot Award from the Yale Center for Brain and Mind Health.

The illustrations were generated using BioRender.com.

648   **Methods**

649   **Human Subjects**

In this study, we analyzed whole-genome sequencing (WGS) data and linked electronic

health record information from participants enrolled in the All of Us Research Program and

the UK Biobank. All analyses were conducted in accordance with each data resource's

governance framework (All of Us: Registered Tier; UK Biobank: Application 58743), and only

authorized researchers accessed the data within secure computing environments. For All of

Us, WGS data derived from both blood and saliva were used, whereas UK Biobank provided

WGS data derived from blood DNA. In both cohorts, all datasets were de-identified, and no

personally identifiable information was available to researchers. Clinical information included

ICD-9/10 diagnosis codes, age, sex, self-reported ancestry, and laboratory measurements,

limited to variables approved for research use within each biobank.

A cohort of individuals at increased risk for MS was recruited at Yale University. This cohort

study consists of (1) family members of MS patients and (2) asymptomatic individuals

incidentally discovered to have morphologically suspicious lesions on brain MRI. Participants

donate biospecimens and are prospectively followed. European-ancestry cohort members

were selected for PRS analysis. A cohort of patients with MS used for qPCR and scRNAseq

was also recruited at Yale School of Medicine.

EBV DNA quantification from WGS

The human reference genome, GRCh38, which is used in AoU and UKB, contains the EBV

decoy sequence, named chrEBV. To detect EBV reads in WGS samples, we investigated

reads mapped to chrEBV stored in the CRAM files. A WGS sample can sometimes contain

very abundant EBV reads due to viremia and have the potential to cause a heavy

computational burden. To decrease the computational cost, samples with more than 350,000

reads mapped to chrEBV were randomly downsampled using samtools to 350,000 reads.

Reads were then filtered to retain only proper pairs, passing standard quality control metrics,

with duplicates, secondary, and supplementary alignments removed. Mapping length was

calculated from the CIGAR string, and only reads with an aligned length of ≥145 bp were

retained. The EBV genome carries a repetitive region, named BamHI W repeat (positions

12,000–35,355 bp). EBV reads derived from this region can be multi-mapping due to the

repetitive nature of the BamHI W repeat region in the EBV reference genome. Our pipeline

accounts for this repetitive nature and avoids counting such multi-mapping reads repeatedly.

Finally, paired-end reads passing all filters were defined as EBV reads, and counts were

summarized separately inside and outside the BamHI W repeat. In the case of analysis with

downsampling, it estimates the number of EBV reads in the original CRAM file by accounting

for the downsampling ratio.

Association analysis of EBV DNA

We analyzed the association of EBV positivity with age_at_collection, genetically predicted

ancestry, which is provided by All of US, and sex. Observations with missing values were

excluded. Logistic regression (generalized linear model with binomial distribution and logit

link) was fitted separately for blood samples and saliva samples. The model was specified

as:

$$\text{logit}\{P(\text{EBVpos} = 1)\} = \beta_0 + \beta_1 \cdot \text{age\_scaled} + \beta_2 \cdot \text{ancestry\_pred} + \beta_3 \cdot \text{sex}$$

To facilitate interpretation, age was scaled in 10-year units (age_scaled = age_at_collection /

10). European ancestry (eur) and male sex were used as reference categories. Robust

(HC3) standard errors were applied. Coefficients were reported as odds ratios (ORs) with

95% confidence intervals. Analyses were conducted in Python using the package

statsmodels. Logistic regression models were estimated using statsmodels GLM (Binomial,
logit) with robust HC3 standard errors.

Generalized Pareto distribution (GPD) analysis

To assess the tail behavior of EBV DNA distributions, we applied a peaks-over-threshold
(POT) approach using the Generalized Pareto distribution (GPD). Normalized EBV DNA
reads (normalized by the number of human-genome coverage) were used. For each sample
type (blood, saliva), the 95th percentile of the distribution was chosen as the threshold u,
and exceedances above u were modeled with the GPD parameterized by shape (ξ) and
scale (β). Parameter estimation was performed by maximum likelihood, and 95% confidence
intervals of ξ were obtained via profile likelihood. For conceptual illustration, simulated GPD
curves with ξ = -0.3, 0, 1.0, and a common scale were generated to highlight the difference
between finite, exponential, and heavy-tailed distributions.

Genotype Data Processing and Quality Control

Genotype data were processed independently for the All of Us Research Program[34] and the
UK Biobank,[35] following harmonized quality control procedures to enable cross-cohort
analyses. Unless otherwise noted, equivalent filters and thresholds were applied across both
datasets. The variant set for downstream analyses was defined based on the All of Us ACAF
PGEN callset and applied consistently to both cohorts.

For the analysis of All of Us, we utilized the All of Us ACAF PGEN callset, which includes
variants with "AF>1% or AC>100 in any ancestry," filtered using the FT flag. Participants
were excluded if DNA was derived from saliva, retaining only blood-derived samples.
Following previous studies, we removed individuals with immunodeficiency disorders (ICD-
10 codes D80–D89) or those with 'ID_057.1' (Human immunodeficiency virus infection) in
PhecodeX. Related and flagged individuals were excluded using the All of Us auxiliary files
(relatedness_flagged_samples.tsv and flagged_samples.tsv). We retained only samples with
inferred sex chromosomes XX or XY, sex at birth recorded as male or female, and available
age-at-collection information.

Variant-level quality control was performed per ancestry with the following parameters: MAF
between 0.01 and 0.99, Hardy–Weinberg equilibrium (HWE) $p>1\times10^{-\square}$ (using keep-fewhet),
biallelic variants only, and a genotype missingness rate <5%. An ancestry-specific variant list
was generated for each group, and the union across all ancestries and cohorts was used for
subsequent analyses.

For the UK Biobank cohort, EBV DNA was quantified from WGS data using the same

pipeline as All of Us, counting reads mapping to the EBV reference genome (chrEBV). EBV

read counts were normalized by mean sequencing coverage and log10-transformed

(log10[normalized reads + 1]) for the quantitative trait analysis; a binary EBV DNA positivity

variable was defined as having at least one EBV-mapped read. Sample-level quality control

included: retention of individuals of European genetic ancestry based on the UK Biobank

genetic ethnicity field (Data-Field 22006); exclusion of one member from each pair of related

individuals (*kinship coefficient* $\geq$ 0.0442) using the UK Biobank relatedness file; exclusion of

individuals with sex chromosome aneuploidy or discordance between reported and genetic

sex; removal of WGS preparation plates with unusually high EBV positivity rates (>28%);

and exclusion of individuals in the top 1% of normalized EBV reads. As in All of Us, we

excluded individuals with immunodeficiency disorders (ICD-10 codes D80–D89) or HIV

infection (ICD-10 codes B20–B24), as well as those taking immunosuppressant medications

at baseline (Data-Field 20003). Principal components were computed on unrelated

European individuals and projected onto all European samples. For GWAS, we used

imputed genotype data from UK Biobank. For REGENIE Step 1, we used directly genotyped

variants after quality control (*MAF* $\geq$ 0.05, *genotype missingness* <5%, *MAC* $\geq$ 40, HWE

p>1×10^{-2}, biallelic SNPs only) and LD pruning (window 200 kb, step 50 variants, r^2<0.2),

yielding 346,104 variants. For REGENIE Step 2 association testing, we applied filters of

MAC ≥ 40 and imputation INFO ≥ 0.8. For the meta-analysis, we restricted to the same set

of variants defined by the All of Us ACAF callset by matching on chromosome, position, and

alleles.

749 Genome-Wide Association Analysis and meta-analysis

Genome-wide association analyses were performed using REGENIE[37] with comparable

parameters across cohorts/population. In step 1, LD-pruned variants were selected using

ancestry-specific thresholds.[80] For individuals of African ancestry, we used --indep-pairwise

1000 100 0.15; for non-African ancestries, --indep-pairwise 1000 100 0.3. Covariates

included the first 20 principal components, sex at birth, and age at collection (including

quadratic and age×sex interaction terms), with additional cohort-specific covariates such as

sequencing batch metrics, fasting time, lymphocyte percentage, smoking status, time of

blood draw, and month of collection to account for technical and seasonal variation. Step 2

analyses were restricted to SNPs that passed QC in at least four of the six ancestry groups.

For the cross ancestry meta analysis, cohort/population with the participants more than

10,000 (European from UK Biobank and European, African, and Latino/Admixed American

from All of Us) were used. We performed the meta analysis using MR-MEGA, which

accounts for ancestry heterogeneity through genetic principal components.[39] For the meta-

analysis restricted to Europeans (UK Biobank + All of Us), METAL[38] was used with an

inverse-variance weighted fixed-effects model. The visualization and the selection of lead

SNPs were performed using gwaslab.[81] The annotation of lead SNPs was performed with

VEP,[82] OpenTable[83] and ImmuNexUT platform.[84]

LD Score Regression and Tissue-specific Heritability Analyses

We performed LDSC analyses to estimate SNP-based heritability and partition it across

tissue- and =cell-type–specific annotations.[40,44,46] GWAS summary statistics were derived

from a meta-analysis of the All of Us and UK Biobank European ancestry cohorts, combined

using METAL as described above. To ensure consistency, only HapMap3 (HM3) variants

were retained during harmonization. Reference LD scores and regression weights for

European populations were obtained from the Broad Institute's public Google Cloud

repository   (https://console.cloud.google.com/storage/browser/broad-alkesgroup-public-

requester-pays). Baseline model v1.2 annotations were used to account for known genomic

features.[44] For each trait, we estimated SNP heritability and genetic correlations using LDSC.

Tissue- and cell-type–specific heritability enrichment was assessed using LDSC (--h2-cts)

with Multi_tissue_gene_expr annotations.[46]

Pathway analysis

First,   we   performed   SNP   annotation   with   gene   locations   (NCBI37.3,

https://ctg.cncr.nl/software/MAGMA/aux_files/NCBI37.3.zip) and the reference data created

from   1000   genomics   Phase3   (g1000_eur,

https://ctg.cncr.nl/software/MAGMA/ref_data/g1000_eur.zip) using magma --annotate (with

the option, window=10,10).[85] For the pathway-level analyses, we used MAGMA v1.10 on the

same European-ancestry METAL meta-analysis summary statistics used for LDSC. SNPs

were first annotated to genes using NCBI37.3 gene locations with a ± 10 kb window and the

1000 Genomes Phase 3 European reference panel (g1000_eur) as the linkage

disequilibrium (LD) reference, and variants within the extended MHC region (chr 6: 25-34

Mb) were excluded. Gene-based association tests were then performed with MAGMA using

SNP-level p values and sample sizes while accounting for LD from g1000_eur, yielding

gene-level z statistics. These results were subsequently used for competitive gene-set

analyses that were carried out using MSigDB v2025.1 gene sets, including Hallmark, KEGG

canonical pathways, and Gene Ontology (biological process, molecular function, and cellular

component) collections.

PheWAS

796 We conducted analyses in individuals of European ancestry from the All of Us cohort. ICD-9

797 and ICD-10 codes were harmonized using PhecodeX.[86] The R package SPAtest was used

798 to perform score tests based on the saddlepoint approximation, and beta coefficients were

799 estimated using Firth's method. Visualization was performed with matplotlib.

EBV detection from scRNAseq data.

We developed a pipeline, VIRTUS3, to detect EBV transcripts from 10x Genomics scRNA-

seq data. First, we performed alignment to the human reference (GRCh38, refdata-gex-

GRCh38-2020-A) and cell barcode calling using Cell Ranger count. The resulting barcodes

were used as a whitelist for viral quantification. Unmapped reads were extracted from the

position-sorted BAM file with samtools view -f 4 and converted back to FASTQ with

cellranger bamtofastq. These unmapped reads were then quantified against an EBV

reference (NC_007605.1, B95-8 strain, CDS plus EBER1/2) using Salmon/Alevin [87] with the

barcode whitelist. The pipeline was applied to the 10x 3' data of PBMC from EBV-HLH, EBV-

IM, and healthy individuals.[49] We compared the number of EBV-positive cells by VIRTUS3

with the reported number in Suzuki *et al.*[49] In their analysis, EBV detection was performed

using Cell Ranger v5.0 with the custom reference made by the human genome (GRCh38)

and the genome of the EBV representative strain (NC_007605).

Polygenic signal enrichment analysis using scDRS and scDRS-spatial

The polygenic signals were calculated using scDRS[48] and scDRS-spatial.[51] The gene scores

derived from MAGMA were used. For PBMC scRNAseq data and the visualization of spatial

transcriptome data, we used imputed gene expression calculated by MAGIC.[88] For tissue-

and region-specific analysis, we used raw counts for scDRS. The validation of the usage of

MAGIC is described in our previous publication.[51] Polygenic enrichment for each cell was

calculated using scDRS (v1.0.3) with --n-ctrl 1000. We used the number of detected genes

per cell as a covariate. Group-level statistics were summarized using scdrs perform-

downstream and visualized with scdrs.util.plot_group_stats.

HLA typing and single-marker association testing

We followed the protocol described by Sakaue et al.[89] Briefly, we used the ACAF PGEN call

set of European participants from the All of Us cohort. Sample quality control was performed

as in the GWAS. Variant filtering was conducted using PLINK2[90] with the following options: --

maf 0.005 --chr 6 --geno 0.1 --hwe 1e-10 --snps-only just-acgt --min-alleles 2 --max-alleles 2.

Variants were lifted over to hg19 using CrossMap, and phasing was performed with EAGLE

(v2.4.1).[91] HLA imputation was carried out using Minimac4 (v4.1.6)[92] with the publicly

available 1KGP HLA panel (https://github.com/immunogenomics/HLA_analyses_tutorial).
Single-marker association testing was performed with PLINK2 --glm, adjusting for age, sex,
PC1, and PC2 as covariates. We also performed conditioning using PLINK2. The
visualization of HLA structure was performed using PyMOL.

Genotyping using SNP microarray

We genotyped SNPs from individuals collected at Yale University using the Illumina Global
Diversity Array. The genotyping was performed at the Keck Microarray Shared Resource
core laboratory at Yale University. We performed standard quality control procedures;
variant-level filters included genotyping missingness <5%, MAF > 0.05, HWE p > 1×10^{-6}.
Genetic ancestry was determined with principal component analysis using 1000 Genomes
as a reference panel. Samples of European genetic ancestry were imputed using IMPUTE 2,
and strand alignment was performed using SHAPEIT. SNPs with INFO > 0.7 and MAF >
0.01 were retained in the final genomic data.

Mendelian Randomization Analysis

Two-sample Mendelian randomization (MR) was performed to assess potential causal
relationships between EBV DNA positivity and autoimmune diseases. Genetic instruments
for EBV DNA positivity were derived from the European-ancestry meta-analysis of EBV DNA
GWAS. Instruments were selected at genome-wide significance ($p<5\times10^{-}$) and clumped
using PLINK-based LD clumping ($r^2<0.0001$, *window*=1000kb) with the European 1000
Genomes Phase 3 reference panel. SNPs with significant heterogeneity in the exposure
GWAS (HetPVal<0.05) were excluded prior to instrument selection. Outcome GWAS
summary statistics were obtained from publicly available sources: MS from the International
Multiple Sclerosis Genetics Consortium (IMSGC), RA from the European-ancestry GWAS,
SLE from published meta-analyses, and ischemic stroke from the MEGASTROKE
consortium (used as a negative control). Harmonization of exposure and outcome data was
performed using the genal Python package, with palindromic SNPs retained (action=1).
When exposure SNPs were not directly available in outcome datasets, proxy SNPs were
identified using LD-based searching (r²≥0.6, ±10kb window).MR analyses were conducted
using the inverse-variance weighted (IVW) method with random effects as the primary
analysis. Sensitivity analyses included the Weighted Median method, and MR-PRESSO for
outlier detection and correction. MR-PRESSO[53] was applied when evidence of pleiotropy or
heterogeneity was detected (Cochran's Q p<0.0001, or discordant IVW and Weighted
Median estimates). When MR-PRESSO identified significant outliers, results were reported
after outlier removal. Causal estimates are reported as odds ratios with 95% confidence

intervals. To assess the contribution of the MHC region, instruments were stratified into
MHC-only (chromosome 6: 25–34Mb) and non-MHC variants, and MR analyses were
performed separately for each subset. Bidirectional MR was performed to test for potential
reverse causation, using genetic instruments for MS, RA, and SLE as exposures and EBV
DNA as the outcome.

Polygenic risk scoring

PRSs were constructed using PRScs[54] a Bayesian continuous shrinkage method that infers
posterior effect sizes by integrating GWAS summary statistics with an external LD reference
panel. The final SNP set for each PRS was determined by the intersection of GWAS variants
with the reference panel, yielding 729,766 variants for the genome-wide scores. For stratified
analyses, MHC-only and no-MHC PRSs were derived by respectively retaining or excluding
variants within the HLA region on chromosome 6 (chr6:25–34 Mb, GRCh38).

For validation in the independent Yale cohort, we used GWAS summary statistics from the
European-ancestry meta-analysis (All of Us + UK Biobank) for EBV DNA positivity (binary
trait) and EBV DNA load (quantitative trait). The Yale cohort comprises European-ancestry
individuals at increased risk of MS (relatives of MS patients and those with morphologically
suggestive brain MRI lesions; N=94; mean age 40.2±14.7 years; 71% female) with
measured EBV DNA load by qPCR. PRS were calculated for each participant, then
standardized to mean 0 and SD 1. Participants were grouped into three PRS categories
based on cohort-specific percentiles (≤10%, 10–90%, and ≥90%). EBV DNA positivity was
modeled using Firth bias-reduced logistic regression with the PRS as either a continuous
predictor (per SD increase) or categorical variable (percentile groups), adjusting for age, sex,
and the first 10 genotype principal components. Ordinal trend tests across PRS categories
were obtained from models including an ordinal coding of the three groups. EBV DNA load
was analyzed using linear regression of log10-transformed EBV DNA copies/mL with the
same covariate sets.

For the UK Biobank MS association analysis, we used GWAS summary statistics from the
European-ancestry subset of All of Us only (to avoid sample overlap) to construct PRSs that
were then applied to European-ancestry individuals in the UK Biobank (N=487,181; 2,423
MS cases; 484,758 controls). PRS values were standardized and participants were grouped
by percentiles (≤10%, 10–90%, and ≥90%, or quintiles). Associations with MS were
assessed using logistic regression adjusted for age, sex, and the first 10 principal
components.

EBV detection by qPCR

Genomic DNA was extracted from whole blood (1-2mL) using the QIAamp DNA Blood Midi kit (Qiagen, 51183), following the manufacturer's spin-column protocol. Briefly, blood was incubated with a protease and lysis buffer at 70°C for 10 minutes, mixed with 100% ethanol, and loaded onto the spin column. After centrifugation, columns were washed with AW1 and AW2 buffers, and DNA was eluted with 300µL of elution buffer. DNA was stored at –80 °C before analysis.

EBV genomic DNA was quantified by TaqMan qPCR targeting *BamH1W*. Reactions contained DNA (5µL), Fast Advanced qPCR master mix (Applied Biosystems, 444457), primers, and a FAM-labeled probe. qPCR was performed on a StepOnePlus (Applied Biosystems) with the following cycling conditions: 95 °C for 20 s, followed by 40 cycles of 95 °C for 1 s and 60°C for 20 seconds. An EBV DNA standard (ATCC, VR-3247SD: $6.6{\times}10^5$ copies/µL; Lot: 70063234). Ten-fold serial dilutions of the standard (10^5 to 10^1 copies) were prepared and run with each qPCR experiment to define assay sensitivity and ensure plate-to-plate consistency. For each run, a standard curve was generated by plotting Ct values against the log-transformed input copy numbers, followed by calculation of the correlation coefficient (R^2) from the resulting linear regression.

Meta-analysis of EBV DNA in patient with MS and controls

We searched PubMed from inception to 2025 using predefined keywords ("multiple sclerosis", "Epstein–Barr virus", "PCR", "viral load"). Only peer-reviewed original research articles that examined EBV DNA in peripheral blood from both MS patients and healthy controls using qPCR were included. Because the sensitivity of EBV DNA detection varies across laboratories and assays, we restricted the analysis to studies in which EBV DNA positivity in either group fell within the range of 5–75 percent to minimize the influence of extreme outliers. When raw counts were not reported, the number of EBV-positive subjects was estimated by multiplying the total sample size by the reported percentage and rounding to the nearest integer. Meta-analysis of odds ratios was performed using the metabin function implemented in the meta package in R.

Single-cell RNA-seq optimized for EBV detection

PBMCs were isolated from whole blood by density-gradient centrifugation using Lymphoprep (STEMCELL, 18061) and cryopreserved in Bambanker Freezing Media (Bulldog-Bio, BB01) for storage in liquid nitrogen until use. PBMCs were thawed rapidly, and B cells were enriched from PBMCs using a pan-B cell isolation kit (STEMCELL, 19554) or B cell isolation kit (STEMCELL, 19054) according to the manufacturer's protocols. Enriched B cells were

counted and adjusted to a final concentration corresponding to a target loading of 20,000
cells.

Tonsil cells were collected by needle aspiration. Collected samples were centrifuged at
500×g for 5 minutes to remove the supernatant, and the cell pellet was resuspended in 1mL
PBS. Total and leukocyte counts were determined using trypan blue and methylene blue,
respectively. After counting cells were centrifuged at 500×g for 5 minutes, the supernatant
was discarded, and the pellet was cryopreserved in Bambanker Freezing Media for storage
in liquid nitrogen until use.

To detect the EBV transcript, a custom EBV-specific primer was added to the cell
suspension immediately prior to loading onto the Chromium GEM-X Single Cell 5' v3 Gene
Expression platform (10x Genomics). The primer (sequence: 5'-AAG CAG TGG TAT CAA
CGC AGA GTA CAA AAC ATG CGG ACC ACC AGC-3'; 10 µM stock, 1 µL added per
sample) was synthesized by Integrated DNA Technologies. Libraries were sequenced on an
Illumina NovaSeqXP.

Single-cell RNAseq analysis
The fastq files are processed with cellranger Sequenced reads were processed using Cell
Ranger (v8.0.1) with pre-built reference refdata-gex-GRCh38-2024-A and refdata-cellranger-
vdj-GRCh38-alts-ensembl-7.1.0 downloaded at 10x Genomics's Website. EBV transcripts
were detected by VIRTUS3. For the pooled samples, donor assignment was performed
using the Vireo pipeline[93] in Demuxafy.[94] For each sample, cell barcodes and the aligned
BAM file generated by Cell Ranger were used as input to generate pileups of exonic SNPs.
SNP pileup was performed using cellsnp-lite with the 1000 Genomes Project–derived
genotype reference panel (GRCh38; MAF>0.01), applying filters of minimum allele count
(minCOUNT≥20) and minor allele frequency (minMAF≥0.1). To restrict genotyping to the
relevant donors, we subset the donor VCF to the SNPs detected in the sample-level pileup
using bcftools. Vireo was run with an expected donor number of 8 and enabling ambient
RNA modeling (--callAmbientRNAs). The analysis was performed using scanpy 1.10.2.[95]
The embedding and cluster assignment was performed using screfmapping pipeline with
pre-defined B cell reference.[6] Briefly, we performed label transfer using CellTypist[96] with the
model Immune_All_Low. B cell population was extracted based on the predicted cluster.
Label transfer and embedding was performed by symphonypy with pre-defined B cell
reference (the dataset will be published on Figshare).[97] BCR were analyzed using airrflow
and scirpy.[98,99] DEG analysis was performed using sc.tl.rank_genes_groups, and pathway

analysis was performed ClusterProfiler.[100] For tonsil data analysis, we used Azimuth[101] with
the tonsil reference dataset.[102]

Figure legends

Figure 1 EBV DNA characteristics

(a) EBV DNA positivity across predicted ancestors, gender, and age in blood (left) and saliva
(right). (b) Forest plots showing odds ratios (points) and 95% confidence intervals (horizontal
lines) from logistic regression of EBV DNA positivity for blood samples (left) and saliva
samples (right). The age effect represents the change per 10-year increase. European
ancestry and male sex were used as references. The dashed line indicates *Odds ratio*
(OR)=1. (c) Distribution of EBV DNA fitted with Generalized Pareto distribution (GPD). Left:
Simulated examples of GPD probability density functions illustrating the effect of the shape
parameter ξ on the tail behavior: finite tail (ξ < 0), exponential tail (ξ = 0), and heavy tail (ξ >
0). Middle and Right: Histograms of normalized EBV DNA reads in blood (middle) and saliva
(right). Orange curves represent GPD fits above the 95th percentile threshold (red dashed
line). Estimated ξ values are shown for each sample type. (d) Phenome-wide association
study (PheWAS) of EBV DNA positivity in European participants from the All of Us cohort.
Each point represents a clinical phenotype, colored by disease category. The x-axis denotes
phenotype groups and the y-axis shows the $-log_{10}$*(FDR-adjusted P value)*. The dashed line
indicates the FDR threshold of 0.05. Details are available in Supplementary Table 1. AFR,
African ancestry; AMR, Latino/admixed American ancestry; EAS, East Asian ancestry; EUR,
European ancestry; SAS, South Asian ancestry

Figure 2 EBV DNA GWAS

(a) Manhattan plots showing genome-wide association statistics of EBV DNA expression.
The y-axis shows $-log_{10}$P of each variant, and lead SNPs and annotated genes were
highlighted. Heatmap shows the genes previously reported in MS,[2,4,103] SLE,[104] and RA[105]
GWAS. The dashed line indicates a genome-wide significant level (p=5.0x10[-8]). Symbols
indicate effect direction relative to the EBV GWAS (+ concordant, – discordant). Directional
symbols are displayed only for loci reaching P<5×10⁻□ in the respective MS/SLE/RA GWAS,
while nonsignificant or unavailable associations are shown as blank cells. Genes not
reported in the GWASs with P<5×10⁻□ in the most significant loci of EBV DNA positivity
GWAS were highlighted in light blue. Details are also available in Table S4. (b) Overlap of
genes highlighted in MS onset GWAS,[2] MS severity GWAS[4] and EBV positivity GWAS.
Statistical significance was evaluated using a two-tailed Fisher's exact test on a 2×2
contingency table, with all protein-coding genes (Gencode v47) used as the background (a

and b). (c) Gene ontology biological process pathways enriched in GWAS of EBV DNA
Positivity. (d) Partitioned heritability was estimated using LD score regression for EBV
positivity. Each dot represents a tissue or cell-type annotation, and the size indicates
whether the enrichment passed the FDR < 0.2 threshold.

Figure 3 Tissue- and cell-type–specific heritability enrichment for EBV
positivity and viral load.

(a) UMAP visualization of PBMC single-cell RNA-seq data from EBV-positive and control
individuals (DRA017005),[49] showing major immune cell clusters. (b) Cells with detected EBV
transcripts identified by VIRTUS3 (blue) overlaid on the UMAP plot. (c) Comparison of the
number of EBV-positive cells detected by VIRTUS3 versus Cell Ranger across samples. (d)
Distribution of predicted cell types among EBV-positive cells, highlighting enrichment in
plasmablasts and memory B cells. (e) Polygenic scores derived from the EBV DNA positivity
GWAS in B memory and plasmablast populations. The Mann-Whitney U test was applied. (f)
Region-level enrichment of EBV DNA positivity polygenic signals in tonsils[102] estimated by
scDRS-spatial. Heatmap colors depict the proportion of significant cells (*FDR*<0.2) evaluated
using scDRS[48] Squares denote significant disease associations (*FDR*<0.05). (g and h)
Spatial transcriptomic map of human tonsil tissue showing annotated microanatomical
regions and EBV DNA positivity polygenic scores across the same tissue section.

Figure 4 Association of HLA alleles and amino acid variants with EBV DNA
positivity and viral load.

(a) Locus plot of rs27300 in EBV Positivity GWAS. (b) eQTL plot of rs27300 in *ERAP2* and
LNPEP in EBV transformed lymphocytes. The eQTL data was downloaded from GTEx
Portal.[106] (c) Associations between HLA alleles and EBV DNA positivity. The size of each
point corresponds to the allele frequency, and the top associated alleles are labeled. The
alleles highlighted by conditioning were also shown. HLA association analysis using All of Us
European participants was performed by imputing classical HLA alleles with SNP2HLA. (d)
Amino acid residue associated with EBV DNA positivity/viral load in HLA-A*02:01 (top) and
HLA-DRB1*04:04 (bottom). The structure was downloaded from PDB (3MRF for HLA-
A*02:01 and 7nzf for HLA-DRB1*04:04).

Figure 5 Mendelian randomization and polygenic risk score analyses of EBV
DNA positivity.

(a) Forest plot of Mendelian randomization (MR) estimates assessing the causal effect of
genetically predicted EBV DNA positivity on autoimmune diseases (multiple sclerosis [MS],
rheumatoid arthritis [RA], and systemic lupus erythematosus [SLE]) and stroke (negative

control). Points represent odds ratios; horizontal lines indicate 95% confidence intervals. The dashed vertical line indicates the null effect (OR = 1). (b) EBV DNA positivity measured by qPCR across categories of the EBV DNA positivity PRS in an independent Yale cohort of European-ancestry individuals without MS (N = 94; 3 EBV DNA+ cases). (c) MS prevalence across EBV DNA positivity PRS categories in the UK Biobank European-ancestry cohort (N = 487,181; 2,423 MS cases). For panels b and c, whole-genome PRSs were constructed using PRScs with the European-ancestry meta-analysis GWAS (b) or All of Us European-ancestry GWAS (c, to avoid sample overlap). Individuals were stratified into bottom 10%, middle 80%, and top 10% of the PRS distribution. Percentages indicate the proportion of EBV DNA-positive individuals (b) or MS cases (c) within each category.

Figure 6 Single-cell EBV detection from people with MS and healthy individuals

(a) Forest plot showing OR and 95 percent confidence intervals for EBV DNA positivity comparing patients with multiple sclerosis (MS) and non-MS controls across published studies and our cohort. Squares indicate study-specific ORs with sizes proportional to study weights, and horizontal lines represent 95 percent CIs. The diamonds indicate the pooled effect estimates under fixed-effect and random-effects models. Quantification of heterogeneity: τ^2=0.0169, τ=0.1301, I^2=9.8 percent [0.0 percent; 49.4 percent], H=1.05 [1.00; 1.41]. A funnel plot is shown in Extended Data Fig. 10. (b and c) UMAP embedding of B-cell subsets from PBMCs (b) and EBV–positive cells (c). (d) Relationship between EBV copies per mL (qPCR) and the percentage of EBV$^+$ B cells, stratified by disease status (MS vs non-MS). The shaded region denotes a 95 percent confidence interval. Pearson's *r* and *p-value* are shown in the figure. (e) Heatmap of viral transcript detection across B-cell subsets, displaying number of cells expressing canonical EBV latent and lytic transcripts. (f) Pearson's correlation of differentially expressed genes in EBV–positive cells compared to EBV–negative cells across the B cell population. (g) Gene Ontology (GO) enrichment analysis comparing EBV–positive versus EBV–negative cells for each B cell population. Dot size represents gene ratio, and color indicates adjusted *p*-value. (h) Expression heatmap of cytokines, chemokine receptors, and immunomodulatory molecules differentially expressed in EBV–positive cells. Expression was normalized for each gene. (i) scDRS-based polygenic score enrichment for EBV DNA positivity (left) and MS onset genes (right) across B-cell subsets, comparing EBV–positive and EBV–negative cells. Statistical test was performed by the Mann-Whitney U test. (j) Volcano plot of differentially expressed genes in EBV–positive versus EBV–negative memory B cells (Busm, Bmem int, Bsm, Bsm *FCER2*, and ABC are pooled). Genes highlighted in the EBV and MS GWAS are annotated.

Extended Data Figure 1 Study design of cross-ancestry meta-analysis of EBV DNA GWAS.

Extended Data Figure 2 Phenome-wide association study (PheWAS) of DNA EBV viral load in European participants from the All of Us cohort.

Phenome-wide association study (PheWAS) of normalized EBV DNA load in European participants from the All of Us cohort. Each point represents a clinical phenotype, colored by disease category. The x-axis denotes phenotype groups, and the y-axis shows the $-log_{10}(FDR)$. The dashed line indicates the FDR threshold of 0.05. Details are available in Supplementary Table 2.

Extended Data Figure 3 QC plots and annotation of GWAS hits.

(a) QQ plot of GWAS for EBV DNA positivity (left) and EBV DNA load (right). (b) Partitioned heritability enrichment in various genomic categories calculated by S-LDSC. EBV DNA positivity (left) and EBV DNA load (right).

Extended Data Figure 4 Supportive data for GWAS.

(a) Manhattan plots showing genome-wide association statistics of EBV DNA load. log_{10}(Normalized EBV read) The y-axis shows $-log_{10}P$ of each variant, and lead SNPs and annotated genes were highlighted. The dashed line indicates a genome-wide significant level (p=5.0x10-8). Details are also available in Supplementary Table 4. (b) Overlap of genes highlighted in MS onset GWAS, MS severity GWAS, and EBV load GWAS. Statistical significance was assessed using a two-tailed Fisher's exact test. (c and d) Locusplot of rs7726159 (*TERT2*) and rs2814778 (*ACKR1*, located near *IFI16*) in EBV Positivity GWAS (e) Gene ontology biological process pathways enriched in GWAS of EBV DNA Load. (f) Partitioned heritability was estimated using LD score regression for EBV DNA load. Each dot represents a tissue or cell-type annotation, and the size indicates whether the enrichment passed the *FDR*<0.2 threshold.

Extended Data Figure 5 Biological interpretation of GWAS results.

(a) Proportion of scDRS-significant cells (*FDR*<0.2) across immune cell types for EBV DNA positivity and viral load. (b) Polygenic scores derived from the EBV load GWAS in B memory and plasmablast populations. The Mann-Whitney U test was applied. (c and d) Enrichment in tissues (c) and anatomical regions in tonsils (d) of EBV polygenic signals estimated by scDRS-spatial. Heatmap colors depict the proportion of significant cells (*FDR*<0.2) evaluated using scDRS. Squares denote significant disease associations (*FDR*<0.05), and cross symbols denote significant heterogeneity in association (*FDR*<0.05).

Extended Data Figure 6 HLA association with EBV DNA load.

(a) Locus plot of rs27300 in EBV viral load GWAS. (b) Associations between HLA alleles and
EBV DNA load. The size of each point corresponds to the allele frequency, and the top
associated alleles are labeled. The alleles highlighted by conditioning were also shown.

Extended Data Figure 7 Conditioning of HLA allele and amino acids
association with EBV DNA Positivity.

Association results for classical HLA alleles across loci (left) and amino acids (right) shown
as $-\log_{10}$(P values). The most strongly associated alleles in each conditioning round are
labeled. HLA amino acid association results for EBV DNA positivity across iterative
conditional analyses (Rounds 0–5).

Extended Data Figure 8 Conditioning of HLA allele and amino acids
association with EBV DNA load.

Association results for classical HLA alleles across loci (left) and amino acids (right) shown
as $-\log_{10}$(P values). The most strongly associated alleles in each conditioning round are
labeled. HLA amino acid association results for EBV DNA load across iterative conditional
analyses (Rounds 0–5).

Extended Data Figure 9 Mendelian randomization and polygenic risk score
stratified by HLA region.

(a) Forest plots of Mendelian randomization (MR) estimates assessing the causal effect of
genetically predicted EBV DNA positivity on autoimmune diseases (multiple sclerosis [MS],
rheumatoid arthritis [RA], and systemic lupus erythematosus [SLE]) and stroke (negative
control). Three instrument sets were analyzed: whole genome, MHC-only, and genome
without the MHC region. Points represent odds ratios; horizontal lines indicate 95%
confidence intervals. The dashed vertical line indicates the null effect (OR = 1). (b) EBV DNA
positivity measured by qPCR across categories of the EBV DNA positivity PRS in an
independent Yale cohort of European-ancestry individuals without MS (N = 94; 3 EBV DNA+
cases). (c) MS prevalence across EBV DNA positivity PRS categories in the UK Biobank
European-ancestry cohort (N = 487,181; 2,423 MS cases). For panels b and c, PRSs were
constructed using PRScs with the European-ancestry meta-analysis GWAS (b) or All of Us
European-ancestry GWAS (c) for three variant sets: whole genome, MHC-only, and non-
MHC. Individuals were stratified into bottom 10%, middle 80%, and top 10% of each PRS
distribution. Percentages indicate the proportion of EBV DNA-positive individuals (b) or MS
cases (c) within each category.

Extended Data Figure 10 Funnel plot assessing publication bias in studies
comparing EBV DNA positivity between MS and controls.
Each point represents an individual study included in the meta-analysis. The vertical dashed
line indicates the pooled odds ratio under the common-effect model (*OR*=1.77). The dashed
diagonal lines denote the 95 percent confidence limits of the expected distribution in the
absence of publication bias.

Extended Data Figure 11 EBV profiling of tonsils from MS and nonMS
individuals
(a and b) UMAP plot showing predicted clusters (a) and detected EBV–positive cells (b).

Extended Data Figure 12 EBV profiling of blood B cells from MS and nonMS
individuals
(a) Percentage of EBV–positive cells across B cell populations per samples. Rows indicate
each donor. The heatmap in orange shows qPCR based copy number of EBV DNA in blood.
(b) Forest plot showing the association of isotype with EBV DNA positivity. Clusters were
treated as a covariete to extract global isotype preference. IGHE, IGHG3, IGHG4 were
removed from the plot because no EBV–positive cells were detected. (c) Somatic hyper
mutation frequency distribution across clusters between EBV–positive and –negative cells.

Supplementary Tables

Supplementary Table 1 PheWAS of Whole-Blood EBV Positivity in AoU European Participants

Supplementary Table 2 PheWAS of Whole-Blood EBV DNA Load in AoU European Participants

Supplementary Table 3 Number individuals and λ_{GC} of individual GWAS

Supplementary Table 4 Lead SNPs of cross-ancestry meta-analysis

Supplementary Table 5 Statistics of EBV Positivity GWAS top hits in all single GWAS

Supplementary Table 6 Statistics of EBV viral load GWAS top hits in all single GWAS

Supplementary Table 7 MAGMA Gene-Set Associations with EBV DNA Positivity

Supplementary Table 8 MAGMA Gene-Set Associations with EBV DNA Load

Supplementary Table 9 HLA allele associated with EBV Positivity

Supplementary Table 10 HLA allele associated with EBV DNA Load

Supplementary Table 11 HLA amino acids associated with EBV Positivity

Supplementary Table 12 HLA amino acids associated with EBV DNA Load

Supplementary Table 13 meta analysis of EBV DNA Positivity

Supplementary Table 14 Sample information for scRNAseq

Supplementary Table 15 Differentially expressed genes in EBV–positive cells

Supplementary Table 16 pathway enriched in EBV–positive cells

Reference

1.   Hafler, D.A., and Weiner, H.L. (1989). MS: a CNS and systematic autoimmune disease.
Immunol. Today *10*, 104–107.

2.   Patsopoulos, N.A., Baranzini, S.E., Santaniello, A., Shoostari, P., Cotsapas, C., Wong,
G., Beecham, A.H., James, T., Replogle, J., Vlachos, I.S., et al. (2019). Multiple
sclerosis genomic map implicates peripheral immune cells and microglia in
susceptibility. Science *365*, eaav7188.

3.   International Multiple Sclerosis Genetics Consortium, Hafler, D.A., Compston, A.,
Sawcer, S., Lander, E.S., Daly, M.J., De Jager, P.L., de Bakker, P.I.W., Gabriel, S.B.,
Mirel, D.B., et al. (2007). Risk alleles for multiple sclerosis identified by a genomewide
study. N. Engl. J. Med. *357*, 851–862.

4.   International Multiple Sclerosis Genetics Consortium, and MultipleMS Consortium
(2023). Locus for severity implicates CNS resilience in progression of multiple sclerosis.
Nature *619*, 323–331.

5.   Farh, K.K.-H., Marson, A., Zhu, J., Kleinewietfeld, M., Housley, W.J., Beik, S., Shoresh,
N., Whitton, H., Ryan, R.J.H., Shishkin, A.A., et al. (2015). Genetic and epigenetic fine
mapping of causal autoimmune disease variants. Nature *518*, 337–343.

6.   Yasumizu, Y., Takeuchi, D., Morimoto, R., Takeshima, Y., Okuno, T., Kinoshita, M.,
Morita, T., Kato, Y., Wang, M., Motooka, D., et al. (2024). Single-cell transcriptome
landscape of circulating CD4+ T cell populations in autoimmune diseases. Cell Genom
4, 100473.

7.   Pappalardo, J.L., Zhang, L., Pecsok, M.K., Perlman, K., Zografou, C., Raddassi, K.,
Abulaban, A., Krishnaswamy, S., Antel, J., van Dijk, D., et al. (2020). Transcriptomic
and clonal characterization of T cells in the human central nervous system. Sci.
Immunol. *5*. https://doi.org/10.1126/sciimmunol.abb8786.

8.   Cao, Y., Goods, B.A., Raddassi, K., Nepom, G.T., Kwok, W.W., Love, J.C., and Hafler,
D.A. (2015). Functional inflammatory profiles distinguish myelin-reactive T cells from
patients with multiple sclerosis. Sci. Transl. Med. *7*, 287ra74.

9.   Sumida, T.S., Lincoln, M.R., He, L., Park, Y., Ota, M., Oguchi, A., Son, R., Yi, A.,
Stillwell, H.A., Leissa, G.A., et al. (2024). An autoimmune transcriptional circuit drives
FOXP3+ regulatory T cell dysfunction. Sci. Transl. Med. *16*, eadp1720.

10.   Viglietta, V., Baecher-Allan, C., Weiner, H.L., and Hafler, D.A. (2004). Loss of functional
suppression by CD4+CD25+ regulatory T cells in patients with multiple sclerosis. J. Exp.
Med. *199*, 971–979.

11.   Schafflick, D., Xu, C.A., Hartlehnert, M., Cole, M., Schulte-Mecklenbeck, A., Lautwein,
T., Wolbert, J., Heming, M., Meuth, S.G., Kuhlmann, T., et al. (2020). Integrated single
cell analysis of blood and cerebrospinal fluid leukocytes in multiple sclerosis. Nat.
Commun. *11*, 247.

12.   Bjornevik, K., Cortese, M., Healy, B.C., Kuhle, J., Mina, M.J., Leng, Y., Elledge, S.J.,
Niebuhr, D.W., Scher, A.I., Munger, K.L., et al. (2022). Longitudinal analysis reveals
high prevalence of Epstein-Barr virus associated with multiple sclerosis. Science *375*,
296–301.

13.   Soldan, S.S., Su, C., Monaco, M.C., Yoon, L., Kannan, T., Zankharia, U., Patel, R.J.,
Dheekollu, J., Vladimirova, O., Dowling, J.W., et al. (2024). Multiple sclerosis patient-

derived spontaneous B cells have distinct EBV and host gene expression profiles in
active disease. Nat. Microbiol. *9*, 1540–1554.

14. Agostini, S., Mancuso, R., Guerini, F.R., D'Alfonso, S., Agliardi, C., Hernis, A.,
Zanzottera, M., Barizzone, N., Leone, M.A., Caputo, D., et al. (2018). HLA alleles
modulate EBV viral load in multiple sclerosis. J. Transl. Med. *16*, 80.

15. Bray, P.F., Bloomer, L.C., Salmon, V.C., Bagley, M.H., and Larsen, P.D. (1983).
Epstein-Barr virus infection and antibody synthesis in patients with multiple sclerosis.
Arch. Neurol. *40*, 406–408.

16. Schneider-Hohendorf, T., Wünsch, C., Falk, S., Raposo, C., Rubelt, F., Mirebrahim, H.,
Asgharian, H., Schlecht, U., Mattox, D., Zhou, W., et al. (2025). Broader anti-EBV TCR
repertoire in multiple sclerosis: disease specificity and treatment modulation. Brain *148*,
933–940.

17. Schneider-Hohendorf, T., Gerdes, L.A., Pignolet, B., Gittelman, R., Ostkamp, P., Rubelt,
F., Raposo, C., Tackenberg, B., Riepenhausen, M., Janoschka, C., et al. (2022).
Broader Epstein-Barr virus-specific T cell receptor repertoire in patients with multiple
sclerosis. J. Exp. Med. *219*. https://doi.org/10.1084/jem.20220650.

18. Lanz, T.V., Brewer, R.C., Ho, P.P., Moon, J.-S., Jude, K.M., Fernandez, D., Fernandes,
R.A., Gomez, A.M., Nadj, G.-S., Bartley, C.M., et al. (2022). Clonally expanded B cells
in multiple sclerosis bind EBV EBNA1 and GlialCAM. Nature *603*, 321–327.

19. Wucherpfennig, K.W., and Strominger, J.L. (1995). Molecular mimicry in T cell-mediated
autoimmunity: viral peptides activate human T cell clones specific for myelin basic
protein. Cell *80*, 695–705.

20. Küppers, R. (2003). B cells under influence: transformation of B cells by Epstein-Barr
virus. Nat. Rev. Immunol. *3*, 801–812.

21. Wei, J., Moon, J., Yasumizu, Y., Zhang, L., Radassi, K., Buitrago-Pocasangre, N.,
Deerhake, M.E., Strauli, N., Chen, C.-W., Herman, A., et al. (2025). Transcriptomic
profiling after B cell depletion reveals central and peripheral immune cell changes in
multiple sclerosis. J. Clin. Invest. *135*. https://doi.org/10.1172/JCI182790.

22. Damania, B., Kenney, S.C., and Raab-Traub, N. (2022). Epstein-Barr virus: Biology and
clinical disease. Cell *185*, 3652–3670.

23. Xu, Y., Hiyoshi, A., Smith, K.A., Piehl, F., Olsson, T., Fall, K., and Montgomery, S.
(2021). Association of infectious mononucleosis in childhood and adolescence with risk
for a subsequent multiple sclerosis diagnosis among siblings. JAMA Netw. Open *4*,
e2124932.

24. Hwang, A.E., Hamilton, A.S., Cockburn, M.G., Ambinder, R., Zadnick, J., Brown, E.E.,
Mack, T.M., and Cozen, W. (2012). Evidence of genetic susceptibility to infectious
mononucleosis: a twin study. Epidemiol. Infect. *140*, 2089–2095.

25. Larsen, P.D., Bloomer, L.C., and Bray, P.F. (1985). Epstein-Barr nuclear antigen and
viral capsid antigen antibody titers in multiple sclerosis. Neurology *35*, 435–438.

26. Li, Z.-X., Zeng, S., Wu, H.-X., and Zhou, Y. (2019). The risk of systemic lupus
erythematosus associated with Epstein-Barr virus infection: a systematic review and
meta-analysis. Clin. Exp. Med. *19*, 23–36.

27. Chan, K.C.A., Woo, J.K.S., King, A., Zee, B.C.Y., Lam, W.K.J., Chan, S.L., Chu, S.W.I.,
Mak, C., Tse, I.O.L., Leung, S.Y.M., et al. (2017). Analysis of plasma Epstein-Barr virus
DNA to screen for nasopharyngeal cancer. N. Engl. J. Med. *377*, 513–522.

28. McLaren, P.J., Coulonges, C., Bartha, I., Lenz, T.L., Deutsch, A.J., Bashirova, A.,
Buchbinder, S., Carrington, M.N., Cossarizza, A., Dalmau, J., et al. (2015).
Polymorphisms of large effect explain the majority of the host genetic contribution to
variation of HIV-1 virus load. Proc. Natl. Acad. Sci. U. S. A. *112*, 14658–14663.

29. Yasumizu, Y., Hara, A., Sakaguchi, S., and Ohkura, N. (2021). VIRTUS: a pipeline for
comprehensive virus analysis from conventional RNA-seq data. Bioinformatics *37*,
1465–1467.

30. Kumata, R., Ito, J., Takahashi, K., Suzuki, T., and Sato, K. (2020). A tissue level atlas of
the healthy human virome. BMC Biol. *18*, 55.

31. Sasa, N., Kojima, S., Koide, R., Hasegawa, T., Namkoong, H., Hirota, T., Watanabe, R.,
Nakamura, Y., Oguro-Igashira, E., Ogawa, K., et al. (2025). Blood DNA virome
associates with autoimmune diseases and COVID-19. Nat. Genet. *57*, 65–79.

32. Kachuri, L., Francis, S.S., Morrison, M.L., Wendt, G.A., Bossé, Y., Cavazos, T.B.,
Rashkin, S.R., Ziv, E., and Witte, J.S. (2020). The landscape of host genetic factors
involved in immune response to common viral infections. Genome Med. *12*, 93.

33. Guo, J., Huang, X., Zhang, C., Huang, P., Li, Y., Wen, F., Wang, X., Yang, N., Xu, M.,
Bi, Y., et al. (2022). The blood virome of 10,585 individuals from the ChinaMAP. Cell
Discov. *8*, 113.

34. All of Us Research Program Genomics Investigators (2024). Genomic data in the All of
Us Research Program. Nature *627*, 340–346.

35. UK Biobank Whole-Genome Sequencing Consortium (2025). Whole-genome
sequencing of 490,640 UK Biobank participants. Nature *645*, 692–701.

36. McManus, T.E., Marley, A.-M., Baxter, N., Christie, S.N., Elborn, J.S., O'Neill, H.J.,
Coyle, P.V., and Kidney, J.C. (2008). High levels of Epstein-Barr virus in COPD. Eur.
Respir. J. *31*, 1221–1226.

37. Mbatchou, J., Barnard, L., Backman, J., Marcketta, A., Kosmicki, J.A., Ziyatdinov, A.,
Benner, C., O'Dushlaine, C., Barber, M., Boutkov, B., et al. (2021). Computationally
efficient whole-genome regression for quantitative and binary traits. Nat. Genet. *53*,
1097–1103.

38. Willer, C.J., Li, Y., and Abecasis, G.R. (2010). METAL: fast and efficient meta-analysis
of genomewide association scans. Bioinformatics *26*, 2190–2191.

39. Mägi, R., Horikoshi, M., Sofer, T., Mahajan, A., Kitajima, H., Franceschini, N., McCarthy,
M.I., COGENT-Kidney Consortium, T2D-GENES Consortium, and Morris, A.P. (2017).
Trans-ethnic meta-regression of genome-wide association studies accounting for
ancestry increases power for discovery and improves fine-mapping resolution. Hum.
Mol. Genet. *26*, 3639–3650.

40. Bulik-Sullivan, B.K., Loh, P.-R., Finucane, H.K., Ripke, S., Yang, J., Schizophrenia
Working Group of the Psychiatric Genomics Consortium, Patterson, N., Daly, M.J.,
Price, A.L., and Neale, B.M. (2015). LD Score regression distinguishes confounding

from polygenicity in genome-wide association studies. Nat. Genet. *47*, 291–295.

41. Wen, S., Kuri-Morales, P., Hu, F., Nag, A., Tachmazidou, I., Deevi, S.V.V., Taiy, H.,
Smith, K.R., Loesch, D.P., Burren, O.S., et al. (2025). Comparative analysis of the
Mexico City Prospective Study and the UK Biobank identifies ancestry-specific effects
on clonal hematopoiesis. Nat. Genet. *57*, 572–582.

42. Dolcetti, R., Giunco, S., Dal Col, J., Celeghin, A., Mastorci, K., and De Rossi, A. (2014).
Epstein-Barr virus and telomerase: from cell immortalization to therapy. Infect. Agent.
Cancer *9*, 8.

43. Witt, K.C., Dziulko, A., An, J., Pekovic, F., Cheng, A.X., Liu, G.Y., Lee, O.V., Turner,
D.J., Lari, A., Gaidt, M.M., et al. (2025). SP140-RESIST pathway regulates interferon
mRNA stability and antiviral immunity. Nature *643*, 1372–1380.

44. Finucane, H.K., Bulik-Sullivan, B., Gusev, A., Trynka, G., Reshef, Y., Loh, P.-R., Anttila,
V., Xu, H., Zang, C., Farh, K., et al. (2015). Partitioning heritability by functional
annotation using genome-wide association summary statistics. Nat. Genet. *47*, 1228–
1235.

45. Amatullah, H., Fraschilla, I., Digumarthi, S., Huang, J., Adiliaghdam, F., Bonilla, G.,
Wong, L.P., Rivard, M.-E., Beauchamp, C., Mercier, V., et al. (2022). Epigenetic reader
SP140 loss of function drives Crohn's disease due to uncontrolled macrophage
topoisomerases. Cell *185*, 3232-3247.e18.

46. Finucane, H.K., Reshef, Y.A., Anttila, V., Slowikowski, K., Gusev, A., Byrnes, A., Gazal,
S., Loh, P.-R., Lareau, C., Shoresh, N., et al. (2018). Heritability enrichment of
specifically expressed genes identifies disease-relevant tissues and cell types. Nat.
Genet. *50*, 621–629.

47. Sumida, T.S., Dulberg, S., Schupp, J.C., Lincoln, M.R., Stillwell, H.A., Axisa, P.-P.,
Comi, M., Unterman, A., Kaminski, N., Madi, A., et al. (2022). Type I interferon
transcriptional network regulates expression of coinhibitory receptors in human T cells.
Nat. Immunol. *23*, 632–642.

48. Zhang, M.J., Hou, K., Dey, K.K., Sakaue, S., Jagadeesh, K.A., Weinand, K.,
Taychameekiatchai, A., Rao, P., Pisco, A.O., Zou, J., et al. (2022). Polygenic
enrichment distinguishes disease associations of individual cells in single-cell RNA-seq
data. Nat. Genet., 1–9.

49. Suzuki, T., Sato, Y., Okuno, Y., Torii, Y., Fukuda, Y., Haruta, K., Yamaguchi, M.,
Kawamura, Y., Hama, A., Narita, A., et al. (2024). Single-Cell Transcriptomic Analysis of
Epstein-Barr Virus-Associated Hemophagocytic Lymphohistiocytosis. J. Clin. Immunol.
44, 103.

50. Cell   Ranger's   Gene   Expression   Algorithm   10x   Genomics.
https://www.10xgenomics.com/support/software/cell-ranger/latest/algorithms-
overview/cr-gex-algorithm.

51. Yasumizu, Y., Kinoshita, M., Zhang, M.J., Motooka, D., Suzuki, K., Nojima, S., Koizumi,
N., Okuzaki, D., Funaki, S., Shintani, Y., et al. (2024). Spatial transcriptomics elucidates
medulla niche supporting germinal center response in myasthenia gravis-associated
thymoma. Cell Reports *43*. https://doi.org/10.1016/j.celrep.2024.114677.

52. Jia, X., Han, B., Onengut-Gumuscu, S., Chen, W.-M., Concannon, P.J., Rich, S.S.,

Raychaudhuri, S., and de Bakker, P.I.W. (2013). Imputing amino acid polymorphisms in
human leukocyte antigens. PLoS One *8*, e64683.

53. Verbanck, M., Chen, C.-Y.Y., Neale, B., and Do, R. (2018). Detection of widespread
horizontal pleiotropy in causal relationships inferred from Mendelian randomization
between complex traits and diseases. Nat. Genet. *50*, 693–698.

54. Ge, T., Chen, C.-Y., Ni, Y., Feng, Y.-C.A., and Smoller, J.W. (2019). Polygenic
prediction via Bayesian regression and continuous shrinkage priors. Nat. Commun. *10*,
1776.

55. Sorgato, C.C., Lins-E-Silva, M., Leão, J.C., Vasconcelos, L.R., Romão, T.P., Duarte,
A.L., and Gueiros, L.A. (2020). EBV and CMV viral load in rheumatoid arthritis and their
role in associated Sjögren's syndrome. J. Oral Pathol. Med. *49*, 693–700.

56. Bäckerholm, A., Tian, Y., Holmqvist, I., Xie, G., Vracar, D., Abrahamsson, S., Guibentif,
C., and Tang, K.-W. (2022). Detection of latent Epstein-Barr virus gene expression in
single-cell sequencing of peripheral blood mononuclear cells. Microbiology.

57. Younis, S., Moutusy, S.I., Rasouli, S., Jahanbani, S., Pandit, M., Wu, X., Acharya, S.,
Sharpe, O., Wijeratne, T.U., Harris, M.L., et al. (2025). Epstein-Barr virus reprograms
autoreactive B cells as antigen-presenting cells in systemic lupus erythematosus. Sci.
Transl. Med. *17*, eady0210.

58. Fournier, B., Boutboul, D., Bruneau, J., Miot, C., Boulanger, C., Malphettes, M., Pellier,
I., Dunogué, B., Terrier, B., Suarez, F., et al. (2020). Rapid identification and
characterization of infected cells in blood during chronic active Epstein-Barr virus
infection. J. Exp. Med. *217*. https://doi.org/10.1084/jem.20192262.

59. Läderach, F., Piteros, I., Fennell, É., Bremer, E., Last, M., Schmid, S., Rieble, L.,
Campbell, C., Ludwig-Portugall, I., Bornemann, L., et al. (2025). EBV induces CNS
homing of B cells attracting inflammatory T cells. Nature *646*, 171–179.

60. Nyeo, S.S., Cumming, E.M., Burren, O.S., Pagadala, M.S., Gutierrez, J.C., Ali, T.A.,
Kida, L.C., Chen, Y., Hu, F., Hollis, B., et al. (2025). Population-scale sequencing
resolves correlates and determinants of latent Epstein-Barr Virus infection. Genetics.

61. Montalban, X., Lebrun-Frénay, C., Oh, J., Arrambide, G., Moccia, M., Pia Amato, M.,
Amezcua, L., Banwell, B., Bar-Or, A., Barkhof, F., et al. (2025). Diagnosis of multiple
sclerosis: 2024 revisions of the McDonald criteria. Lancet Neurol. *24*, 850–865.

62. Makhani, N., and Tremlett, H. (2021). The multiple sclerosis prodrome. Nat. Rev.
Neurol. *17*, 515–521.

63. Ford, J.L., and Stowe, R.P. (2013). Racial-ethnic differences in Epstein-Barr virus
antibody titers among U.S. children and adolescents. Ann. Epidemiol. *23*, 275–280.

64. Lee, J., Manon, V., and Chen, J. (2025). Structurally diverse viral inhibitors converge on
a shared mechanism to stall the antigen transporter TAP. Proc. Natl. Acad. Sci. U. S. A.
122, e2516676122.

65. Raychaudhuri, S., Sandor, C., Stahl, E.A., Freudenberg, J., Lee, H.-S., Jia, X.,
Alfredsson, L., Padyukov, L., Klareskog, L., Worthington, J., et al. (2012). Five amino
acids in three HLA proteins explain most of the association between MHC and
seropositive rheumatoid arthritis. Nat. Genet. *44*, 291–296.

66. Patsopoulos, N.A., Barcellos, L.F., Hintzen, R.Q., Schaefer, C., van Duijn, C.M., Noble,
J.A., Raj, T., IMSGC, ANZgene, Gourraud, P.-A., et al. (2013). Fine-mapping the
genetic association of the major histocompatibility complex in multiple sclerosis: HLA
and non-HLA effects. PLoS Genet. *9*, e1003926.

67. Mycko, M.P., Waldner, H., Anderson, D.E., Bourcier, K.D., Wucherpfennig, K.W.,
Kuchroo, V.K., and Hafler, D.A. (2004). Cross-reactive TCR responses to self antigens
presented by different MHC class II molecules. J. Immunol. *173*, 1689–1698.

68. Moutsianas, L., Jostins, L., Beecham, A.H., Dilthey, A.T., Xifara, D.K., Ban, M., Shah,
T.S., Patsopoulos, N.A., Alfredsson, L., Anderson, C.A., et al. (2015). Class II HLA
interactions modulate genetic risk for multiple sclerosis. Nat. Genet. *47*, 1107–1113.

69. Niens, M., Jarrett, R.F., Hepkema, B., Nolte, I.M., Diepstra, A., Platteel, M., Kouprie, N.,
Delury, C.P., Gallagher, A., Visser, L., et al. (2007). HLA-A*02 is associated with a
reduced risk and HLA-A*01 with an increased risk of developing EBV+ Hodgkin
lymphoma. Blood *110*, 3310–3315.

70. Ishigaki, K., Lagattuta, K.A., Luo, Y., James, E.A., Buckner, J.H., and Raychaudhuri, S.
(2022). HLA autoimmune risk alleles restrict the hypervariable region of T cell receptors.
Nat. Genet. *54*, 393–402.

71. Schmidt, A., Alawathurage, T.M., Frach, L., Richter, S., David, F.S., Schaefer, M.,
Henne, S.K., Boztug, K., Proebstel, A.-K., Noethen, M.M., et al. (2025). Host control of
latent Epstein-Barr virus infection. medRxiv, 2025.07.19.25331823.
https://doi.org/10.1101/2025.07.19.25331823.

72. Harroud, A., and Hafler, D.A. (2023). Common genetic factors among autoimmune
diseases. Science *380*, 485–490.

73. Lazarian, G., Yin, S., Ten Hacken, E., Sewastianik, T., Uduman, M., Font-Tello, A.,
Gohil, S.H., Li, S., Kim, E., Joyal, H., et al. (2021). A hotspot mutation in transcription
factor IKZF3 drives B cell neoplasia via transcriptional dysregulation. Cancer Cell *39*,
380-393.e8.

74. Devergne, O., Hatzivassiliou, E., Izumi, K.M., Kaye, K.M., Kleijnen, M.F., Kieff, E., and
Mosialos, G. (1996). Association of TRAF1, TRAF2, and TRAF3 with an Epstein-Barr
virus LMP1 domain important for B-lymphocyte transformation: role in NF-kappaB
activation. Mol. Cell. Biol. *16*, 7098–7108.

75. Cable, J.M., Wongwiwat, W., Grabowski, J.C., White, R.E., and Luftig, M.A. (2025).
Sp140L functions as a herpesvirus restriction factor suppressing viral transcription and
activating interferon-stimulated genes. Proc. Natl. Acad. Sci. U. S. A. *122*,
e2426339122.

76. Boisvert, F.M., Hendzel, M.J., and Bazett-Jones, D.P. (2000). Promyelocytic leukemia
(PML) nuclear bodies are protein structures that do not accumulate RNA. J. Cell Biol.
148, 283–292.

77. Xu, H., Li, X., Rousseau, B.A., Akinyemi, I.A., Frey, T.R., Zhou, K., Droske, L.E.,
Mitchell, J.A., McIntosh, M.T., and Bhaduri-McIntosh, S. (2022). IFI16 partners with
KAP1 to maintain Epstein-Barr virus latency. J. Virol. *96*, e0102822.

78. Koch, M.A., Thomas, K.R., Perdue, N.R., Smigiel, K.S., Srivastava, S., and Campbell,
D.J. (2012). T-bet(+) Treg cells undergo abortive Th1 cell differentiation due to impaired

expression of IL-12 receptor β2. Immunity *37*, 501–510.

79. Knippenberg, S., Peelen, E., Smolders, J., Thewissen, M., Menheere, P., Cohen
Tervaert, J.W., Hupperts, R., and Damoiseaux, J. (2011). Reduction in IL-10 producing
B cells (Breg) in multiple sclerosis is accompanied by a reduced naïve/memory Breg
ratio during a relapse but not in remission. J. Neuroimmunol. *239*, 80–86.

80. Li, B., Veturi, Y., Lucas, A., Bradford, Y., Verma, S.S., Verma, A., Park, J., Wei, W.-Q.,
Feng, Q., Namjou, B., et al. (2021). Multi-ancestry gene-trait connection landscape
using electronic health record (EHR) linked biobank data. bioRxiv.
https://doi.org/10.1101/2021.10.21.21265225.

81. He, Y., Koido, M., Shimmori, Y., and Kamatani, Y. (2023). GWASLab: a Python
package for processing and visualizing GWAS summary statistics.

82. McLaren, W., Gil, L., Hunt, S.E., Riat, H.S., Ritchie, G.R.S., Thormann, A., Flicek, P.,
and Cunningham, F. (2016). The Ensembl Variant Effect Predictor. Genome Biol. *17*,
122.

83. Ochoa, D., Hercules, A., Carmona, M., Suveges, D., Baker, J., Malangone, C., Lopez,
I., Miranda, A., Cruz-Castillo, C., Fumis, L., et al. (2023). The next-generation Open
Targets Platform: reimagined, redesigned, rebuilt. Nucleic Acids Res. *51*, D1353–
D1359.

84. Ota, M., Nagafuchi, Y., Hatano, H., Ishigaki, K., Terao, C., Takeshima, Y., Yanaoka, H.,
Kobayashi, S., Okubo, M., Shirai, H., et al. (2021). Dynamic landscape of immune cell-
specific gene regulation in immune-mediated diseases. Cell *184*, 3006-3021.e17.

85. de Leeuw, C.A., Mooij, J.M., Heskes, T., and Posthuma, D. (2015). MAGMA:
generalized gene-set analysis of GWAS data. PLoS Comput. Biol. *11*, e1004219.

86. Shuey, M.M., Stead, W.W., Aka, I., Barnado, A.L., Bastarache, J.A., Brokamp, E.,
Campbell, M., Carroll, R.J., Goldstein, J.A., Lewis, A., et al. (2023). Next-generation
phenotyping: introducing phecodeX for enhanced discovery research in medical
phenomics. Bioinformatics *39*. https://doi.org/10.1093/bioinformatics/btad655.

87. Srivastava, A., Malik, L., Smith, T., Sudbery, I., and Patro, R. (2019). Alevin efficiently
estimates accurate gene abundances from dscRNA-seq data. Genome Biol. *20*, 65.

88. van Dijk, D., Sharma, R., Nainys, J., Yim, K., Kathail, P., Carr, A.J., Burdziak, C., Moon,
K.R., Chaffer, C.L., Pattabiraman, D., et al. (2018). Recovering Gene Interactions from
Single-Cell Data Using Data Diffusion. Cell *174*, 716-729.e27.

89. Sakaue, S., Gurajala, S., Curtis, M., Luo, Y., Choi, W., Ishigaki, K., Kang, J.B., Rumker,
L., Deutsch, A.J., Schönherr, S., et al. (2023). Tutorial: a statistical genetics guide to
identifying HLA alleles driving complex disease. Nat. Protoc. *18*, 2625–2641.

90. Chang, C.C., Chow, C.C., Tellier, L.C., Vattikuti, S., Purcell, S.M., and Lee, J.J. (2015).
Second-generation PLINK: rising to the challenge of larger and richer datasets.
Gigascience *4*, 7.

91. Loh, P.R., Danecek, P., Palamara, P.F., Fuchsberger, C., Reshef, Y.A., Finucane, H.K.,
Schoenherr, S., Forer, L., McCarthy, S., Abecasis, G.R., et al. (2016). Reference-based
phasing using the Haplotype Reference Consortium panel. Nat. Genet. *48*, 1443–1448.

92. Fuchsberger, C., Abecasis, G.R., and Hinds, D.A. (2015). Minimac2: Faster genotype
imputation. Bioinformatics *31*, 782–784.

93. Huang, Y., McCarthy, D.J., and Stegle, O. (2019). Vireo: Bayesian demultiplexing of
pooled single-cell RNA-seq data without genotype reference. Genome Biol. *20*, 273.

94. Neavin, D., Senabouth, A., Arora, H., Lee, J.T.H., Ripoll-Cladellas, A., sc-eQTLGen
Consortium, Franke, L., Prabhakar, S., Ye, C.J., McCarthy, D.J., et al. (2024).
Demuxafy: improvement in droplet assignment by integrating multiple single-cell
demultiplexing and doublet detection methods. Genome Biol. *25*, 94.

95. Wolf, F.A., Angerer, P., and Theis, F.J. (2018). SCANPY: large-scale single-cell gene
expression data analysis. Genome Biol. *19*, 15.

96. Domínguez Conde, C., Xu, C., Jarvis, L.B., Rainbow, D.B., Wells, S.B., Gomes, T.,
Howlett, S.K., Suchanek, O., Polanski, K., King, H.W., et al. (2022). Cross-tissue
immune cell analysis reveals tissue-specific features in humans. Science *376*.
https://doi.org/10.1126/science.abl5197.

97. Kang, J.B., Nathan, A., Weinand, K., Zhang, F., Millard, N., Rumker, L., Moody, D.B.,
Korsunsky, I., and Raychaudhuri, S. (2021). Efficient and precise single-cell reference
atlas mapping with Symphony. Nat. Commun. *12*, 5890.

98. Gabernet, G., Marquez, S., Bjornson, R., Peltzer, A., Meng, H., Aron, E., Lee, N.Y.,
Jensen, C.G., Ladd, D., Polster, M., et al. (2024). nf-core/airrflow: An adaptive immune
receptor repertoire analysis workflow employing the Immcantation framework. PLoS
Comput. Biol. *20*, e1012265.

99. Sturm, G., Szabo, T., Fotakis, G., Haider, M., Rieder, D., Trajanoski, Z., and Finotello,
F. (2020). Scirpy: a Scanpy extension for analyzing single-cell T-cell receptor-
sequencing data. Bioinformatics *36*, 4817–4818.

100.Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan,
L., et al. (2021). clusterProfiler 4.0: A universal enrichment tool for interpreting omics
data. Innovation (Camb.) *2*, 100141.

101.Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., Zheng, S., Butler, A., Lee, M.J.,
Wilk, A.J., Darby, C., Zager, M., et al. (2021). Integrated analysis of multimodal single-
cell data. Cell *184*, 3573-3587.e29.

102.Massoni-Badosa, R., Aguilar-Fernández, S., Nieto, J.C., Soler-Vila, P., Elosua-Bayes,
M., Marchese, D., Kulis, M., Vilas-Zornoza, A., Bühler, M.M., Rashmi, S., et al. (2024).
An atlas of cells in the human tonsil. Immunity.
https://doi.org/10.1016/j.immuni.2024.01.006.

103.Housley, W.J., Fernandez, S.D., Vera, K., Murikinati, S.R., Grutzendler, J., Cuerdon, N.,
Glick, L., De Jager, P.L., Mitrovic, M., Cotsapas, C., et al. (2015). Genetic variants
associated with autoimmunity drive NFκB signaling and responses to inflammatory
stimuli. Sci. Transl. Med. *7*, 291ra93.

104.Wang, Y.-F., Zhang, Y., Lin, Z., Zhang, H., Wang, T.-Y., Cao, Y., Morris, D.L., Sheng,
Y., Yin, X., Zhong, S.-L., et al. (2021). Identification of 38 novel loci for systemic lupus
erythematosus and genetic heterogeneity between ancestral groups. Nat. Commun. *12*,
772.

105. Ishigaki, K., Sakaue, S., Terao, C., Luo, Y., Sonehara, K., Yamaguchi, K., Amariuta, T.,
Too, C.L., Laufer, V.A., Scott, I.C., et al. (2022). Multi-ancestry genome-wide
association analyses identify novel genetic mechanisms in rheumatoid arthritis. Nat.
Genet., 1–12.

106. Consortium, T.G. (2020). The GTEx Consortium atlas of genetic regulatory effects
across human tissues. Science *369*, 1318–1330.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

EBV Detection and GWAS

409,331 participants across the US
(Multi-ancestry)

313,774 participants across the UK
(Europeans)

Whole Blood
(723,105)

Saliva
(48,899)
(not used for GWAS)

Short Read Whole
Genome sequence

SNV/indel call

Quantification of
reads mapped on
EBV genome

Association test
Ancestry/Age/Gender

PheWAS

EBV DNA GWAS
(DNA positivity/DNA load)
  Europeans (METAL)
  Multi-ancestry (MR-MEGA)

Extended Data Figure 1

Extended Data Figure 2

Extended Data Figure 3

Extended Data Figure 4

a

Prop. of sig. cells (FDR < 0.20)

0.0   0.1   0.2

EBV DNA Positivity

EBV viral load

ASDC
B intermediate
B memory
B naïve
CD14 Mono
CD16 Mono
CD4 Naive
CD4 Proliferating
CD4 TCM
CD4 TEM
CD8 Naive
CD8 Proliferating
CD8 TCM
CD8 TEM
Eryth
HSPC
ILC
MAIT
NK
NK Proliferating
NK_CD56bright
Plasmablast
Platelet
Treg
cDC1
cDC2
dnT
gdT
pDC

b

B memory
p=4.5x10⁻³

Plasmablast
N.S.

Polygenic Score

−1

EBV+ EBV-
Viral Load

0.0

−0.5

−1.0

EBV+ EBV-
Viral Load

c

Prop. of sig. cells (FDR < 0.20)

0.0   0.1   0.2

EBV Positivity

EBV Load

brain
cerebellum
heart
kidney
lymphnode
prostate
spinalcord
tonsil

d

Prop. of sig. cells (FDR < 0.20)

0.0   0.1   0.2

EBV Positivity

EBV Load

Epithelial-1
Epithelial-2
Germinal Center
Inter-Follicular Zone
Muscle
Proliferating Follicle
Subepithelial
T cell zone
T:B Border

Extended Data Figure 5

Extended Data Figure 6

Extended Data Figure 7

Extended Data Figure 8

a

b

c

Extended Data Figure 9

Extended Data Figure 10

a

predicted.celltype.l1

- ● Activated Naive B
- ● CD4 T
- ● CD8 T
- ● Cycling B
- ● Cycling T
- ● Cycling follicular dendritic
- ● Cycling myeloid
- ● Dendritic
- ● Double negative T
- ● Epithelial
- ● Follicular dendritic
- ● Germinal center B
- ● Granulocytes
- ● Innate lymphoid
- ● Mast
- ● Memory B
- ● Monocyte/macrophage
- ● Naive B
- ● Naive CD4 T
- ● Naive CD8 T
- ● Natural killer
- ● Plasma
- ● Plasmacytoid dendritic

b

total_counts_EBV

- ● Activated Naive B
- ● CD4 T
- ● CD8 T
- ● Cycling B
- ● Dendritic
- ● Double negative T
- ● Granulocytes
- ● Innate lymphoid
- ● Mast

Extended Data Figure 11

Extended Data Figure 12