The feasibility and efficacy of a virtual, symptom-guided aerobic exercise intervention to

improve cognition in mild traumatic brain injury: A single-blind pilot randomized control trial

with an active comparator group.

Emma M. Tinney[1,2], Mark C. Nwakamma[2,3], Goretti España-Irla[2,3], Lauren Kong[2], Colette

Chen[2], Jeremy Hwang[2], Amanda O'Brien[2], Madeleine Perko[2,3], Ryan Luke Sodemann[2],

Jacqueline Caefer[2,3], Julia Manczurowsky[3], Alexandra Stillman[4], Charles H. Hillman[1,2,3],

Timothy P. Morris[2,3,5]

[1]Department of Psychology, Northeastern University, Boston, MA, USA

[2]Institute for Cognitive & Brain Health, Northeastern University, Boston, MA, USA

[3]Department of Physical Therapy, Movement, & Rehabilitation Sciences, Northeastern

University, Boston, MA, USA

[4]Department of Neurology, Beth Israel Deaconess Medical Center, Boston, Massachusetts, USA.

[5]Department of Applied Psychology, Northeastern University, Boston, MA, USA

Corresponding Author: Emma M. Tinney, tinney.e@northeastern.edu

Abstract

Background: Mild traumatic brain injury (mTBI) affects millions worldwide, with cognitive

impairment substantially impacting daily functioning. Despite this burden, evidence-based non-

pharmaceutical interventions remain lacking in clinical practice. Emerging evidence suggests

aerobic exercise may improve post-mTBI cognition; however, the methodological limitations,

particularly inadequate control groups, prevent definitive conclusions.

Objective: This pilot randomized controlled trial examined the feasibility and preliminary

efficacy of a 12-week virtual exercise intervention for community-dwelling adults aged 18-55

years with mTBI within one year of injury.

Methods: Thirty-seven participants were randomized to either symptom-guided aerobic exercise

or active balance control; both delivered virtually three times weekly for 30 minutes over 12

weeks. Primary outcomes assessed feasibility metrics; secondary outcomes examined cognitive

function.

Results: Of enrolled participants, 75% completed the intervention with 94.2% session adherence

and zero adverse events, demonstrating excellent feasibility and safety. The aerobic group

demonstrated greater improvements in executive function compared to balance controls, with

large effect sizes for TMT B-A difference scores in both post-intervention comparisons (Hedges'

$g = 1.20$, 95% CI [0.00, 2.41]) and Group $\times$ Time interactions (Hedges' $g = 1.38$, 95% CI [0.27,

2.49]). Additionally, the aerobic group reported fewer sleep disturbances post intervention ($g =$

1.65, 95% CI [0.22, 3.09]).

Conclusion: These findings establish that virtual, supervised, symptom-guided exercise

interventions are feasible and safe for mTBI populations, with preliminary evidence suggesting

aerobic exercise specifically benefits cognitive flexibility and sleep quality following mTBI. A

fully powered randomized controlled trial is warranted to confirm these effects

Key words: mild TBI, Cognition, aerobic exercise, rehabilitation,

Introduction

Mild traumatic brain injury (mTBI) affects an estimated 60 million individuals worldwide

annually, with a substantial proportion experiencing persistent cognitive impairment that

significantly impacts return to work and daily functioning[1]. Despite this considerable public

health burden, there are no FDA-approved treatments to directly address cognitive dysfunction

following mTBI[2], leaving clinicians with limited options beyond symptom monitoring and

activity modification.

Aerobic exercise has emerged as a promising therapeutic approach, with preclinical evidence

demonstrating neuroprotective effects, including reduced neuroinflammation, enhanced

neurogenesis, and improved synaptic plasticity[3]. Previous human studies examining exercise

interventions for cognitive recovery after mTBI have been limited by significant heterogeneity,

including inadequate control groups, unsupervised interventions, and lack of individualized

symptom-guided prescriptions[4–12]. These limitations have prevented definitive conclusions about

exercise efficacy and optimal implementation approaches for mTBI populations.

The current study addresses critical gaps in prior studies through three key innovations. First, we

implemented a virtual yet supervised exercise intervention that overcomes barriers to exercise

participation in mTBI populations, such as transportation challenges, symptom exacerbation in

gym environments, and cognitive demands of navigating unfamiliar spaces and scheduling

[13]. Second, we implemented a symptom-threshold approach using metrics from validated

exercise tolerance testing protocols[14], allowing for real time modification of intensity based on

individual symptoms. This personalized approach minimizes risk of symptom exacerbation while

promoting self-efficacy and adherence. Third, we included an active balance control group to

isolate aerobic exercise effects from standard practice of care and general physical activity

benefits, a methodological control absent from prior studies. Therefore, the purpose of this study

was to test the feasibility of an individualized, virtual, supervised exercise intervention in mTBI.

The primary aim of this pilot study was to establish feasibility of delivering a 12-week

supervised virtual exercise intervention in community-dwelling adults with a recent mTBI

(within one year of injury), including recruitment rates, adherence, retention, and safety

parameters. Secondary aims examined precision estimates of the effect of the intervention versus

the active comparator group (means, variances, and effect sizes with 95% confidence

intervals)[15,16] for changes in cognitive function and a battery of psychosocial questionnaires.

Methods

Details of all study protocols are described in the protocol paper[17] and below. This study received

ethical approval from the Northeastern University Institutional Review Board (#24-10-30).

Study design

This was a two-arm randomized single-blind control trial. After baseline testing, participants

were randomized after the baseline session using a block randomization in block sizes of four

and six with equal allocation to each group, with allocation concealed in REDCap. Participants

were blinded to study hypotheses, but investigators were not blinded. Participants engaged in

supervised exercise sessions three-times per week for twelve weeks with outcome assessments

obtained before and after the intervention.

Participants

Adults aged 18-55 years with a diagnosis of mTBI within the past year were recruited for

participation in an exercise intervention (The Exercise and Concussion Health Study (TECHS))

(ClinicalTrials.gov: NCT06494592) via advertisements around the Greater Boston area.

Eligibility required that all participants received a formal concussion or mTBI diagnosis by a

physician (AS). Diagnoses were confirmed using OSU-TBI-ID questionnaire[18], requiring a direct

blow or impact to the head with either a loss of consciousness of <5 minutes, dazed, or a gap in

memory for <5 minutes. Participants were excluded if they had a skull breach, subdural

hematoma, prior diagnosis of cognitive or physical disability, clinical diagnosis of neurological

or neuropsychiatric disorder, undergoing treatment for cardiovascular events, not fluent in

English, not medically cleared for exercise, use of an assisted walking device, or not magnetic

resonance imaging compatible.

Between January 2023 and June 2025 216 inquiries were received, 83 individuals were assessed

for eligibility. 37 were consented and enrolled into the study. See the consort diagram in Figure 1

for details. An a-priori sample size of 12 per arm was chosen based on Julious et al., 2005,

considering the feasibility of patient recruitment and the optimal precision and variance about the

effect estimates to inform future studies.

Cognitive Testing

Trail Making Test[19], Verbal Fluency[20], and Hopkins Verbal Learning Test[21] were administered at

baseline and endpoint. These were chosen to provide measures of domains with known deficits

after a mTBI and are commonly used in clinical practice. All cognitive testing was administered

by two trained research staff (EMT and MCN).

Symptom Threshold and Cardiorespiratory Fitness Testing

We utilized an in person hybrid modified exertional bike test, similar to the Buffalo concussion

treadmill test and others[14,22] to objectively determine exercise intensity (via heart rate) at which

concussion symptoms increase ≥3 units above baseline, combined with sub-maximal VO□max

estimation using the Astrand-Rhyming protocol[23] (Figure 2). The assessment consisted of three

phases on a cycle ergometer: (1) 7-minute self-paced warm-up, (2) 6-minute Astrand-Rhyming

test at 50 RPM with predetermined workload to estimate VO□max, and (3) symptom threshold

test with resistance increasing every 30 seconds until meeting termination criteria (≥3-point

symptom increase, age-predicted maximum heart rate, or RPE ≥18). Heart rate (Polar H7), RPE

(Borg scale[24]), and symptoms were recorded every two minutes. The symptom threshold heart

rate was used to prescribe aerobic exercise intensity for intervention sessions. Two trained staff

(EMT, MCN) administered all testing at baseline and post-intervention (see Nwakamma et al.,

for details).

Balance Test

The Balance Error Scoring System (BESS)[25] was used to evaluate postural stability, consisting of

three stance positions performed on two different surfaces (firm and foam), yielding six total

testing conditions held for 20 seconds. The BESS was administered and scored by two trained

research staff (EMT and MCN).

Psychosocial Assessments

A battery of questionnaires was administered at baseline and endpoint including the Pittsburgh

Sleep Quality Index (PSQI)[26], Patient-Reported Outcomes Measurement Information System

(PROMIS)[27–29], and International Physical Activity Questionnaire (IPAQ)[30], asking specifically

about pre-injury physical activity engagement.

Intervention

All participants completed supervised one-on-one virtual exercise sessions three times weekly

for 30 minutes over 12 weeks (36 total sessions). Weekly texts confirmed scheduling, adherence,

and remaining sessions. Each intervention group had three sets of five exercises with

modifications to maintain engagement and progression. Two trained staff led each virtual

session: one exercise leader and one safety observer. Heart rate (HR) was monitored via

hyperate4health (https://hyperate4health.netlify.app/) and manually recorded every 5 minutes

(continuous recording worked for only 54% of participants for ≥18 sessions). Participants rated

symptoms and exertion every five minutes. Aerobic sessions targeted 20 minutes at 65-80% of

symptom threshold HR. If symptoms increased by ≥3 points, participants took a one-minute

break and reduced intensity to 60% of threshold HR. If symptoms persisted, another break

occurred and intensity decreased to 40% of threshold HR (Figure 2).

Endpoint satisfaction survey

Participants were asked to complete an optional satisfaction survey, answering Likert scale

questions about scheduling, communication, difficulty, and open-ended responses about

satisfaction with the study.

Statistical Analysis

Descriptive statistics are provided for all feasibility metrics. To calculate demographic

differences between groups, we performed analyses of variance (ANOVA), reporting sum of

squares, and 95% confidence intervals. For categorical variables, we performed chi-squared

tests, reporting χ^2 and 95% confidence intervals.

Statistical models to test for differences in outcomes at end point and rate of change over time

between groups were used. However, inferences were not made based on p-values as sample size

was chosen to demonstrate feasibility, precision, and variance around the mean difference.

Therefore, all model results are reported as effect sizes and 95% confidence intervals around the

point estimate.

We used two complementary approaches to evaluate treatment effects. All models controlled for

four fixed covariates: age, sex, education, and days since injury. Forward selection with AIC

identified additional covariates improving model fit (previous injuries, adherence rate, baseline

VO☐max, baseline symptoms, total METs, time in heart rate zone). VO☐max analyses excluded

baseline VO☐max, and symptom analyses excluded baseline symptoms to avoid circularity.

We conducted Analysis of Covariance (ANCOVA) to compare post-intervention scores between

groups while controlling for baseline performance. Assumptions were verified using Shapiro-

Wilk, Levene's, and Group × Pre-Score interaction tests. Outliers were identified but not

removed. We also fitted linear mixed-effects models to assess group differences in change over

time. Models used restricted maximum likelihood estimation (*lme4* package in R). Full details

are in supplementary material 1.

Results

Participants

Thirty-seven participants signed informed consent, underwent baseline testing and were

randomized to one of the two study arms. Nine participants withdrew from the study before

completing the intervention (24%) and four participants were excluded from the final analysis

for varying reasons (Figure 1). The final sample included 24 participants, 12 in each group.

Groups did not differ in demographics except for biological sex (Table 1).

Adverse Events

No participant experienced any adverse events during the baseline, endpoint, or intervention.

However, several minor events happened outside the study that affected participation in the

study. One participant was forced to withdraw due to diagnosis of postural orthostatic

tachycardia syndrome, but this was not due to the intervention. One participant developed

pneumonia but received a doctor's clearance to continue participating and resumed full protocol.

One participant dropped out due to receiving unspecified inpatient care.

Intervention feasibility

In the aerobic exercise intervention (n=12), participants were exercised at 65-80% of their

symptom-threshold heart rate for 36 sessions with no adverse events. Seven of the twelve

participants exceeded their prescribed HR zones for at least 50% of the intervention, but none

experienced clinically significant symptom exacerbation (mean three-unit increase). Notably, 10

of 12 participants demonstrated decreased correlations between heart rate and symptom severity

over time, indicating improved exercise tolerance (supplementary material 10). The balance

intervention successfully minimized increases in HR with clear numerical differences in HR

between intervention arms (Supplementary Material 14). An average adherence rate of 96% was

achieved for the aerobic condition and 92.5% for balance. Participants' reported a three-unit

exercise-induced symptom severity increase in 3.8% of all completed sessions. Twenty-one out

of 24 (87.5%) participants reported at least one symptom on the CIF. (Supplementary Material

3).

Endpoint Differences of Primary outcomes

Pre-post changes in primary cognitive and behavioral outcomes are reported as mean and

standard deviation (Table 2).The aerobic group reported fewer sleep disturbances compared to

the balance group after controlling for baseline sleep quality and total physical activity levels

(β=3.75,SE=1.46,g=1.65,95%CI[0.22, 3.09]) (Figure 3c). Specifically, the aerobic group

displayed a 3.75-unit greater reduction in sleep disturbance scores. The aerobic group completed

the TMT-B an average of 13.15 seconds faster than the balance group after controlling for

baseline performance and number of prior injuries (β=13.15,SE= 6.05,g=1.24,95%CI[0.04,2.45])

(Figure 3c). The aerobic group demonstrated greater improvement in the TMT B-A difference

score compared to the balance group after controlling for baseline performance and number of

prior injuries (β=13.36,SE=6.31,g=1.20,95% CI[0.00,2.41]). Specifically, the aerobic group

showed a 13.36-second greater reduction in the time difference between TMT-B and TMT-A

(Figure 3c). The balance group demonstrated higher PROMIS global physical health compared

to the aerobic group after controlling for baseline performance, number of prior injuries, and

time in heart rate zone (β = -7.97, SE = 3.67, g = -1.33, 95% CI [-2.63, -0.03]) (Figure 3c).

Specifically, the aerobic group scored 7.97 points lower on PROMIS GPH at post-intervention

compared to the balance group, representing a large effect size (g = -1.33).

No further large effect sizes were observed in post-test differences between groups. AIC-guided

covariate selection revealed that time in heart rate zone, adherence rate, and number of prior

injuries consistently improved model fit across multiple cognitive and health outcomes.

However, neither time in heart rate zone nor adherence rate correlated with magnitude of

cognitive or sleep improvements, suggesting that exceeding prescribed intensity thresholds may

not confer additional benefits and that high adherence rates in both groups contributed equally to

positive outcomes. Notably, number of prior injuries emerged as a potential moderator of

treatment response, particularly for the balance group's delayed recall performance, though small

subgroup sizes warrant cautious interpretation. See Supplementary Material for complete

ANCOVA results (12), outcome visualizations (4), expanded exploratory covariate analyses (5-

7), and complimentary results of the linear mixed effects models assessing differences in change

over time between groups (13).

Study Feedback

27 participants (72%) completed the endpoint satisfaction survey collected via email. Generally.

participants were satisfied with the study but recommended improved scheduling. Many

participants expressed gratitude for the study and that participating helped in their subjective

recovery (Supplementary Material 9).

Discussion

This pilot study tested the feasibility of a single-blinded, supervised, virtual exercise intervention

for community-dwelling individuals within one year of mTBI. Results demonstrated that this

approach is feasible and safe, with high adherence and no adverse events. Participants reported

high satisfaction with study procedures, and preliminary evidence suggests aerobic exercise may

improve cognitive flexibility compared to an active balance control group mimicking typical

vestibular physical therapy.

These findings demonstrate that virtual aerobic exercise can safely and feasibly be utilized as a

rehabilitative intervention following mTBI, effectively overcoming traditional barriers to

exercise participation in mTBI populations. The supervised format with continuous symptom

monitoring enabled real-time intensity adjustments while maintaining safety and engagement.

Although the dropout rate was slightly higher than other exercises studies in patient populations

(between 15-20%[31,32]), this was improved throughout the study with enhanced participant

communication. Zero adverse events, high adherence, and symptom exacerbation in only 3.8%

of sessions demonstrate adequate safety protocols. Participants frequently exceeded prescribed

heart rate zones suggesting difficulty of floor-based exercises (compared to running or cycling)

to manipulate HR. However, exercising above the symptom-threshold did not increase

symptoms, and the heart rate-symptom correlations decreased over time with continued exercise

exposure (Supplementary Material 10). These results support that the intervention is feasible,

acceptable, and safe for mTBI populations, warranting progression to a fully powered

randomized controlled trial.

Improvement in TMT-B and TMT B-As in the aerobic group align with several prior studies

demonstrating TMT improvements in individuals with TBI of varying severities following

aerobic exercise [12,33], though both studies lacked a control group. This holds clinical significance

given that TMT-B measures executive function, a cognitive domain typically impaired in

TBI[34,35] and corresponds with functional recovery[36,37]. The mechanisms underlying exercise

related improvements in executive function may be driven by improved synaptic plasticity and

associated downstream cellular and molecular mechanisms in prefrontal and subcortical

networks that support executive control[39,40]. Given our balance intervention consisted of non-

aerobic physical activity, which itself has shown improvement in cognitive function in other

populations[41], it is unclear why TMT-B appears particularly sensitive to aerobic exercise post-

TBI. Notably, here, 83% of aerobic participants achieved clinically meaningful improvement

(≥10% faster completion times) on TMT-B, compared to 42% of balance (Figure 3C). The

improvements in TMT-B following aerobic exercise may have meaningful implications for real-

world functional outcomes, including return to work, academic performance, and independent

living skills. Prior work has shown improvement in TMT-B using deep brain stimulation in

TBI[42], while our study shows that aerobic exercise may provide a non-invasive and accessible

therapeutic option. Further discussion of cognitive changes is found in supplementary material

13.

The aerobic group demonstrated fewer sleep disturbances compared to the balance group at post-

intervention. This finding holds clinical significance given the persistent nature of sleep

dysfunction following TBI. Sleep deficits after TBI can persist for years post-injury, representing

one of the most common and debilitating complaints in TBI[43]. Beyond their immediate impact

on quality of life, sleep disturbances are strongly associated with poorer cognitive outcomes and

compromised brain health trajectories after TBI[44], positioning sleep as a critical therapeutic

target for recovery. The mechanisms underlying exercise-related improvements in sleep may

operate through both direct and indirect pathways. Exercise directly influences circadian rhythm

regulation, adenosine accumulation, and thermoregulatory processes that facilitate sleep onset

and maintenance[52]. Indirectly, exercise may improve sleep by reducing anxiety and depressive

symptoms, enhancing mood regulation, and promoting daytime wakefulness, all factors

commonly disrupted after TBI. These converging mechanisms position aerobic exercise as a

particularly promising therapeutic intervention for treating post-TBI sleep disturbances. Despite

this theoretical foundation, empirical evidence supporting exercise interventions for sleep in TBI

populations remains limited. In one of the few studies, Damanio and colleagues[47] demonstrated

that sleep quality improved following 8 weeks of daily aerobic exercise in chronic TBI.

However, the absence of a control group limits causal inference from those findings. The current

study addresses this critical gap with an active control group, providing robust evidence that

aerobic exercise drives improvements in sleep.

Neither group demonstrated changes to cardiorespiratory fitness (CRF), likely because

participants exercised at 65-80% of symptom threshold HR, which was lower than HR max for

most participants. Importantly though, CRF increases appear unnecessary for beneficial effects

on executive function in mTBI, suggesting exercise influences brain health through pathways

independent of cardiovascular conditioning. Indeed, prior work has shown that light intensity

exercise can increase executive function in non-injured adults [49]. Alternative mechanisms of

action may include acute exerc*ise-induced increases in cerebral blood flow and BDNF, reduced

neuroinflammation, enhanced neurogenesis, and improved sleep quality, all achievable without

cardiovascular conditioning[50]. These findings have significant clinical implications: symptom-

threshold guided exercise can yield meaningful cognitive benefits at intensities below those

needed for CRF improvements. This supports individualized, moderate-intensity exercise

protocols that are safer, more tolerable, and reduce barriers to mTBI rehabilitation by minimizing

concerns about symptom exacerbation.

Limitations

Our study has several limitations. The small sample size limits generalizability, but was chosen

a-priori to provide adequate precision and variance around the mean effect[15]. Combining acute,

subacute, and chronic injury phases limits our ability to determine optimal timing of aerobic

exercise delivery; future studies should examine these phases separately. Inter-trainer variability

in rapport and instruction may have introduced inconsistencies. The unbalanced sex distribution

in the aerobic group precluded sex-difference analyses; future studies should balance groups to

test sex-specific effects. Additionally, while our three cognitive tests represent standard clinical

markers, a more comprehensive battery may capture additional cognitive changes. Finally, the

interpretation of the PROMIS Global Physical Health result warrants caution, as baseline

differences between groups may have contributed to regression to the mean.

Conclusion

A virtual, supervised, and individualized exercise intervention is both feasible and acceptable in

community-based individuals with mTBI. Supervised aerobic exercise is potentially effective at

improving executive function deficits observed post-mTBI. A larger randomized controlled trial

is necessary to definitively assess the efficacy of the study.

Acknowledgements

We would like to thank the participants for their participation in this study. We would also like to

thank the TBI Lab undergraduate research assistants for their help in administering and

monitoring the interventions. Additionally, this work was done in part with the Discovery and

Explorer Cluster, supported by Northeastern University's Research Computing team. The data

was partially collected using REDCap, supported by NIH UL1TR002544.

Data will be available upon request and uploaded to fitbir repository.

Study was funded by Northeastern University, specifically TPM start-up funding and the Institute

for Cognitive and Brain Health Fellowship (ET).

Bibliography

1. Maas AIR, Menon DK, Manley GT, et al. Traumatic brain injury: progress and
challenges in prevention, clinical care, and research. *Lancet Neurol*.
2022;21(11):1004-1060. doi:10.1016/S1474-4422(22)00309-X

2. Hiskens MI. Targets of Neuroprotection and Review of Pharmacological
Interventions in Traumatic Brain Injury. *J Pharmacol Exp Ther*. 2022;382(2):149-166.
doi:10.1124/jpet.121.001023

3. Cabral DF, Rice J, Morris TP, Rundek T, Pascual-Leone A, Gomes-Osman J.
Exercise for Brain Health: An Investigation into the Underlying Mechanisms Guided
by Dose. *Neurotherapeutics*. 2019;16(3):580-599. doi:10.1007/s13311-019-00749-w

4. Alarie C, Gagnon I, Quilico E, Teel E, Swaine B. Physical Activity Interventions for
Individuals With a Mild Traumatic Brain Injury:: A Scoping Review. *J Head Trauma*
Rehabil. 2021;36(3):205. doi:10.1097/HTR.0000000000000639

5. Morris T, Gomes Osman J, Tormos Muñoz JM, Costa Miserachs D, Pascual Leone
A. The role of physical exercise in cognitive recovery after traumatic brain injury:
A systematic review. *Restor Neurol Neurosci*. 2016;34(6):977-988.
doi:10.3233/RNN-160687

6. Pawlowski J, Dixon-Ibarra A, Driver S. Review of the Status of Physical Activity
Research for Individuals With Traumatic Brain Injury. *Arch Phys Med Rehabil*.
2013;94(6):1184-1189. doi:10.1016/j.apmr.2013.01.005

7. Sharma B, Allison D, Tucker P, Mabbott D, Timmons BW. Cognitive and neural
effects of exercise following traumatic brain injury: A systematic review of
randomized and controlled clinical trials. *Brain Inj*. 2020;34(2):149-159.
doi:10.1080/02699052.2019.1683892

8. Snowden T, Morrison J, Boerstra M, et al. Brain changes: aerobic exercise for
traumatic brain injury rehabilitation. *Front Hum Neurosci*. 2023;17.
doi:10.3389/fnhum.2023.1307507

9. Larson-Dupuis C, Léveillé E, Desjardins M, et al. Subtle long-term cognitive effects
of a single mild traumatic brain injury and the impact of a three-month aerobic
exercise intervention. *J Sports Med Phys Fitness*. 2021;61(1):87-95.
doi:10.23736/S0022-4707.20.10918-6

10. Snyder AR, Greif SM, Clugston JR, et al. The Effect of Aerobic Exercise on
Concussion Recovery: A Pilot Clinical Trial. *J Int Neuropsychol Soc*. 2021;27(8):790-
804. doi:10.1017/S1355617721000886

11. Ding K, Tarumi T, Tomoto T, et al. A proof-of-concept trial of a community-based
aerobic exercise program for individuals with traumatic brain injury. *Brain Inj*.
2021;35(2):233-240. doi:10.1080/02699052.2020.1865569

12. Chin LM, Keyser RE, Dsurney J, Chan L. Improved Cognitive Performance
Following Aerobic Exercise Training in People with Traumatic Brain Injury. *Arch Phys*
Med Rehabil. 2015;96(4):754. doi:10.1016/J.APMR.2014.11.009

13. Driver S, Ede A, Dodd Z, Stevens L, Warren AM. What barriers to physical activity
do individuals with a recent brain injury face? *Disabil Health J.* 2012;5(2):117-125.
doi:10.1016/j.dhjo.2011.11.002

14. Leddy JJ, Willer B. Use of Graded Exercise Testing in Concussion and Return-to-
Activity Management. *Curr Sports Med Rep.* 2013;12(6):370.
doi:10.1249/JSR.0000000000000008

15. Julious SA. Sample size of 12 per group rule of thumb for a pilot study. *Pharm Stat.*
2005;4(4):287-291. doi:10.1002/pst.185

16. Teresi JA, Yu X, Stewart AL, Hays RD. Guidelines for Designing and Evaluating
Feasibility Pilot Studies. *Med Care.* 2022;60(1):95.
doi:10.1097/MLR.0000000000001664

17. Tinney EM, Nwakamma MC, España-Irla G, et al. The Exercise and Concussion
Health Study (TECHS): Pilot and Feasibility Protocol. *Contemp Clin Trials Commun.*
Published online January 29, 2026:101608. doi:10.1016/j.conctc.2026.101608

18. Corrigan JD, Bogner J. Initial reliability and validity of the Ohio State University TBI
Identification Method. *J Head Trauma Rehabil.* 2007;22(6):318-329.
doi:10.1097/01.HTR.0000300227.67748.77

19. Bowie CR, Harvey PD. Administration and interpretation of the Trail Making Test. *Nat*
Protoc. 2006;1(5):2277-2281. doi:10.1038/nprot.2006.390

20. Henry JD, Crawford JR. A Meta-Analytic Review of Verbal Fluency Performance in
Patients With Traumatic Brain Injury. *Neuropsychology.* 2004;18(4):621-628.
doi:10.1037/0894-4105.18.4.621

21. Brandt J. The hopkins verbal learning test: Development of a new memory test with
six equivalent forms. *Clin Neuropsychol.* 1991;5(2):125-142.
doi:10.1080/13854049108403297

22. Miutz LN, Burma JS, Brassard P, Phillips AA, Emery CA, Smirl JD. Comparison of
the Buffalo Concussion Treadmill Test With a Physiologically Informed Cycle Test:
Calgary Concussion Cycle Test. *Sports Health.* 2024;16(5):837-850.
doi:10.1177/19417381231217744

23. Astrand PO, Ryhming I. A nomogram for calculation of aerobic capacity (physical
fitness) from pulse rate during sub-maximal work. *J Appl Physiol.* 1954;7(2):218-
221. doi:10.1152/jappl.1954.7.2.218

24. Borg G. *Borg's Perceived Exertion and Pain Scales.* Human Kinetics; 1998:viii, 104.

25. Bell DR, Guskiewicz KM, Clark MA, Padua DA. Systematic Review of the Balance
Error Scoring System. *Sports Health*. 2011;3(3):287-295.
doi:10.1177/1941738111403122

26. Buysse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep
Quality Index: a new instrument for psychiatric practice and research. *Psychiatry*
Res. 1989;28(2):193-213. doi:10.1016/0165-1781(89)90047-4

27. Cella D, Riley W, Stone A, et al. The Patient-Reported Outcomes Measurement
Information System (PROMIS) developed and tested its first wave of adult self-
reported health outcome item banks: 2005–2008. *J Clin Epidemiol*.
2010;63(11):1179-1194. doi:10.1016/j.jclinepi.2010.04.011

28. Cella D, Yount S, Rothrock N, et al. The Patient-Reported Outcomes Measurement
Information System (PROMIS): Progress of an NIH Roadmap Cooperative Group
During its First Two Years. *Med Care*. 2007;45(5):S3.
doi:10.1097/01.mlr.0000258615.42478.55

29. Cella D, Choi SW, Condon DM, et al. PROMIS® Adult Health Profiles: Efficient
Short-Form Measures of Seven Health Domains. *Value Health J Int Soc*
Pharmacoeconomics Outcomes Res. 2019;22(5):537-544.
doi:10.1016/j.jval.2019.02.004

30. Bassett DR. International physical activity questionnaire: 12-country reliability and
validity. *Med Sci Sports Exerc*. 2003;35(8):1396.
doi:10.1249/01.MSS.0000078923.96621.1D

31. Stubbs B, Vancampfort D, Rosenbaum S, et al. Dropout from exercise randomized
controlled trials among people with depression: A meta-analysis and meta
regression. *J Affect Disord*. 2016;190:457-466. doi:10.1016/j.jad.2015.10.019

32. Vancampfort D, Van Damme T, Brunner E, et al. Dropout From Exercise
Interventions in Adults With Fibromyalgia: A Systematic Review and Meta-analysis.
Arch Phys Med Rehabil. 2024;105(3):571-579. doi:10.1016/j.apmr.2023.06.002

33. López LP, Coll-Andreu M, Torras-Garcia M, et al. Aerobic exercise and cognitive
function in chronic severe traumatic brain injury survivors: a within-subject A-B-A
intervention study. *BMC Sports Sci Med Rehabil*. 2024;16(1):201.
doi:10.1186/s13102-024-00993-4

34. Lange RT, Iverson GL, Zakrzewski MJ, Ethel-King PE, Franzen MD. Interpreting the
Trail Making Test Following Traumatic Brain Injury: Comparison of Traditional Time
Scores and Derived Indices. *J Clin Exp Neuropsychol*. 2005;27(7):897-906.
doi:10.1080/1380339049091290

35. Iverson GL, Lange RT, Green P, Franzen MD. Detecting Exaggeration and
Malingering With the Trail Making Test. *Clin Neuropsychol*. 2002;16(3):398-406.
doi:10.1076/clin.16.3.398.13861

36. Ross SR, Millis SR, Rosenthal M. Neuropsychological prediction of psychosocial
outcome after traumatic brain injury. *Appl Neuropsychol*. 1997;4(3):165-170.
doi:10.1207/s15324826an0403_4

37. Ip RY, Dornan J, Schentag C. Traumatic brain injury: Factors predicting return to
work or school. *Brain Inj*. 1995;9(5):517-532. doi:10.3109/02699059509008211

38. Heyder K, Suchan B, Daum I. Cortico-subcortical contributions to executive control.
Acta Psychol (Amst). 2004;115(2):271-289. doi:10.1016/j.actpsy.2003.12.010

39. Tinney EM, Ai M, España-Irla G, Hillman CH, Morris TP. Physical activity and
frontoparietal network connectivity in traumatic brain injury. *Brain Behav*.
2024;14(9):e70022. doi:10.1002/brb3.70022

40. España-Irla G, Tinney EM, Ai M, Nwakamma M, Morris TP. Functional Connectivity
Patterns Following Mild Traumatic Brain Injury and the Association With Longitudinal
Cognitive Function. *Hum Brain Mapp*. 2025;46(8):e70237. doi:10.1002/hbm.70237

41. Rogge AK, Röder B, Zech A, et al. Balance training improves memory and spatial
cognition in healthy adults. *Sci Rep*. 2017;7(1):5661. doi:10.1038/s41598-017-
06071-9

42. Schiff ND, Giacino JT, Butson CR, et al. Thalamic deep brain stimulation in traumatic
brain injury: a phase 1, randomized feasibility study. *Nat Med*. 2023;29(12):12.
doi:10.1038/s41591-023-02638-4

43. Mathias JL, Alvaro PK. Prevalence of sleep disturbances, disorders, and problems
following traumatic brain injury: A meta-analysis. *Sleep Med*. 2012;13(7):898-905.
doi:10.1016/j.sleep.2012.04.006

44. Tinney EM, España-Irla G, Warren AEL, et al. Axonal injury, sleep disturbances, and
memory following traumatic brain injury. *Ann Clin Transl Neurol*. n/a(n/a).
doi:10.1002/acn3.52145

45. Fiorin F da S, Godinho DB, dos Santos EB, et al. Relationship among depression,
fatigue, and sleep after traumatic brain injury: The role of physical exercise as a non-
pharmacological therapy. *Exp Neurol*. 2025;386:115156.
doi:10.1016/j.expneurol.2025.115156

46. Korkutata A, Korkutata M, Lazarus M. The impact of exercise on sleep and sleep
disorders. *Npj Biol Timing Sleep*. 2025;2(1):5. doi:10.1038/s44323-024-00018-w

47. Damiano DL, Zampieri C, Ge J, Acevedo A, Dsurney J. Effects of a rapid-resisted
elliptical training program on motor, cognitive and neurobehavioral functioning in
adults with chronic traumatic brain injury. *Exp Brain Res*. 2016;234(8):2245-2252.
doi:10.1007/s00221-016-4630-8

48. Terwee CB, Peipert JD, Chapman R, et al. Minimal important change (MIC): a
conceptual clarification and systematic review of MIC estimates of PROMIS
measures. *Qual Life Res Int J Qual Life Asp Treat Care Rehabil*. 2021;30(10):2729-
2754. doi:10.1007/s11136-021-02925-y

49. Morris TP, Fried PJ, Macone J, et al. Light aerobic exercise modulates executive
function and cortical excitability. *Eur J Neurosci*. 2020;51(7):1723-1734.
doi:10.1111/ejn.14593

50. Mansoor M, Ibrahim A, Hamide A, Tran T, Candreva E, Baltaji J. Exercise-Induced
Neuroplasticity: Adaptive Mechanisms and Preventive Potential in
Neurodegenerative Disorders. *Physiologia*. 2025;5(2):13.
doi:10.3390/physiologia5020013

 Table 1 Participant Demographics

	Aerobic (n=12)	Balance (n=12)	Excluded/Dropped (n=13)	Overall (n=37)	Difference between aerobic vs balance (SS/χ²), [95% CI]
Age (years)					(84.4) [-12.98, 5.48]
Mean (SD)	28.2 (8.67)	31.9 (12.6)	29.8 (10.8)	29.9 (10.6)	
Median [Min, Max]	25.5 [19.0, 47.0]	32.0 [18.0, 55.0]	25.0 [19.0, 50.0]	25.0 [18.0, 55.0]	
Ethnicity					(1.1) [-0.47, 0.30]
Hispanic/Latino	2 (16.7%)	2 (16.7%)	0 (0%)	4 (10.8%)	
Not Hispanic/Latino	10 (83.3%)	9 (75.0%)	11 (84.6%)	30 (81.1%)	
Unknown/Not Reported	0 (0%)	1 (8.4%)	2 (15.4%)	3 (8.1%)	
Race					(0) [-0.79, 0.79]
Asian	1 (8.3%)	1 (8.3%)	3 (23.1%)	5 (13.5%)	
Black/African American	0 (0%)	0 (0%)	1 (7.7%)	1 (2.7%)	

White	10 (83.3%)	10 (83.3%)	9 (69.2%)	29 (78.4%)	
More than One Race	1 (8.3%)	1 (8.3%)	0 (0%)	2 (5.4%)	
Biological Sex					(4.69)
Female	11 (91.7%)	5 (41.7%)	12 (92.3%)	28 (75.7%)	[27.8, 52.2]
Male	1 (8.3%)	7 (58.3)	1 (7.7%)	9 (24.3%)	
Gender Identity					(7.64) [-1.14, 0.81]
Woman	9 (75.0%)	5 (41.7%)	11 (84.6%)	25 (67.6%)	
Man	1 (8.3%)	7 (58.3)	1 (7.7%)	9 (24.3%)	
Non-Binary	2 (16.7%)	0 (0%)	1 (7.7%)	3 (8.1%)	
Education (years)					(3.37) [-1.62, 3.12]
Mean (SD)	15.8 (2.86)	15.0 (2.73)	15.5 (2.15)	15.4 (2.53)	
Median [Min, Max]	15.0 [12.0, 21.0]	15.0 [11.0, 21.0]	16.0 [12.0, 21.0]	15.0 [11.0, 21.0]	
Total MET minutes per					(1341665) [-4466,

					3520]
week (International Physical Activity Questionnaire)					
Mean (SD)	4650 (4830)	4180 (4600)	3350 (2350)	4040 (3970)	
Median [Min, Max]	2960 [320, 18200]	2350 [372, 13800]	2840 [924, 7930]	2610 [320, 18200]	
Time since Injury (days)					(22694) [-25.58, 148.58]
Mean (SD)	133 (129)	69 (60.4)	85.3 (77.7)	94.7 (94.1)	
Median [Min, Max]	54.0 [16.0, 355]	39.0 [25, 200]	52.0 [9, 245]	49.0 [9.0, 355]	
Number of prior injuries					(15.04) [-0.61, 3.77]
Mean (SD)	3 (3.05)	1.42 (1.98)	1.62 (1.61)	1 (2.32)	
Median [Min, Max]	2 [0, 9]	0.5 [0, 6]	2 [0, 4]	1 [0, 9]	
Intervention Adherence					(72.2) [-2.63, 9.57]

Rate (% of sessions attended)					
Mean (SD)	96% (6.66)	92.5% (7.70)	64.5% (33.1)	83.8 (24.6)	
Median [Min, Max]	100 [81.8, 100]	94.9 [78.4, 100]	67.5 [0, 100]	92.5 [0, 100]	
Time in Heart Rate Zone					(0.24) [-0.14, 0.54]
Mean (SD)	57.29% (45.3%)	77.08% (35.7%)		67.19% (41.18%)	
Median [Min, Max]	75% [0, 100]	100% [0, 100]		93.7 [0, 100]	
Mechanism of Injury					(0.17) [-1.11, 1.44]
Bicycle Accident	2 (16.7%)	0 (0%)	1 (7.7%)	3 (8.1%)	
Direct Blow to the head	5 (41.7%)	6 (50.0%)	4 (30.8%)	15 (40.5%)	
Fall	0 (0%)	3 (25.0%)	1 (7.7%)	4 (10.8%)	
Motor Vehicle Accident	2 (16.7%)	0 (0%)	4 (30.8%)	6 (16.2%)	
Recreation/Sport	3 (25%)	2 (16.7%	2 (15.4%)	7 (18.9%)	

Measure	Aerobic Pre M (SD)	Aerobic Post M (SD)	Balance Pre M (SD)	Balance Post M (SD)	Post-test Differences (Hedges' g (95% CI))
Whiplash	0 (0%)	1 (8.3%)	0 (0%)	1 (2.7%)	
Violence	0 (0%)	0 (0%)	1 (7.7%)	1 (2.7%)	

483   *Table 2 Group Means at Baseline and Endpoint*

Measure	Aerobic Pre M (SD)	Aerobic Post M (SD)	Balance Pre M (SD)	Balance Post M (SD)	Post-test Differences (Hedges' g (95% CI))
Hopkin Verbal Learning Test Total Recall	29.67 (3.31)	29.58 (2.54)	25.33 (6.47)	28.42 (3.15)	0.01 [-1.25, 1.26]
Hopkin Verbal Learning Test Delayed Recall	10.75 (1.60)	10.25 (1.86)	8.92 (2.35)	9.92 (1.44)	1.25 [-0.05, 2.55]
Hopkin Verbal Learning Test Retention	0.96 (0.12)	0.93 (0.19)	0.91 (0.15)	0.92 (0.11)	0.69 [-0.51, 1.89]
Hopkin Verbal Learning Test Recognition Discrimination	23.75 (0.87)	22.92 (1.24)	23.25 (0.75)	23.00 (0.74)	-0.05 [-1.27, 1.17]

Index					
Verbal Fluency Letter	39.83 (10.92)	47.17 (12.71)	42.17 (10.35)	44.75 (10.42)	0.26 [-0.93, 1.44]
Verbal Fluency Animals	25.25 (4.92)	24.58 (3.70)	23.00 (3.57)	22.08 (3.65)	-1.01 [-2.35, 0.34]
Trail Making A	19.67 (6.15)	18.25 (4.63)	24.33 (8.72)	20.42 (6.44)	0.28 [-0.95, 1.51]
Trail Making B	52.67 (16.08)	40.75 (10.06)	58.42 (30.52)	56.00 (30.08)	1.24 [0.04, 2.45]
Trail Making B-A	33.00 (15.63)	22.50 (8.72)	34.08 (24.50)	35.58 (28.24)	1.20 [0.00, 2.41]
Pittsburg Sleep Quality Index Global	7.50 (2.35)	7.20 (3.43)	7.67 (3.17)	8.11 (2.98)	1.65 [0.22, 3.09]
PROMIS Global Physical Health T-Score	45.61 (4.59)	48.93 (6.18)	48.98 (4.92)	50.81 (8.02)	-1.33 [-2.63, -0.03]
PROMIS Global Mental Health T-Score	46.61 (5.17)	47.58 (9.73)	48.18 (5.17)	51.14 (9.43)	-0.09 [-1.38, 1.21]

PROMIS Cognitive					0.254 [-0.937, 1.445
T-Score	37.82 (7.81)	44.55 (10.25)	43.51 (9.66)	49.86 (8.48)	]
Balance Error					-0.27 [-1.52,
Scoring System	14.75 (7.47)	11.25 (6.59)	9.75 (2.63)	10.75 (8.54)	0.98]
					-0.24 [-1.50,
Resting Heart Rate	78.25 (16.93)	75.00 (19.98)	73.67 (14.79)	70.50 (12.27)	1.01]
Cardiorespiratory					-0.77 [-2.06,
Fitness	38.64 (10.85)	42.08 (11.90)	37.09 (6.85)	40.63 (9.29)	0.52]
Concussion					-0.64 [-1.85,
Symptoms	13.50 (19.60)	1.42 (1.73)	5.50 (6.22)	0.17 (0.39)	0.56]

Figure 1: Consort diagram

Figure 2: Modified Exertional Bike Test and Intervention. a) Modified Exertional Bike Test and decision tree. b) virtual

intervention set up. Study staff led exercise sessions from a dedicated room in the research institute. The safety officer monitored

the session for several criteria including flushed face, increased perfusion, pale features and followed an emergency protocol if

signs were observed.

Figure 3. Forrest plots and Interaction plots A a) Forest Plot of Post Intervention Group Differences b) Percent change in TMT B

test, with 10% change showing clinically significant improvement. Negative percentages show faster TMT B times at endpoint. c)

Pittsburg Sleep Quality Index (PSQI), Trail making test (TMT) B, Trail making test B-A score, and PROMIS Global Physical

Health T Score.

497