Clinical and Epidemiological Characteristics of Pelvic Organ Prolapse: A Retrospective Study

Author

Mohamed Boulahia, MD

Department of Obstetrics and Gynecology, Tertiary Care Hospital, Algiers, Algeria

Corresponding Author: Dr. Mohamed Boulahia — dr.mohamedboulahia@gmail.com

ORCID iD: https://orcid.org/0009-0006-2117-2479

Abstract

Background

Pelvic organ prolapse (POP) is a common condition among multiparous and postmenopausal women. In North Africa, delayed consultation often leads to advanced stage presentation.

Objective

To describe the epidemiological profile, clinical characteristics, and surgical outcomes of women treated for POP at a Tertiary Care Hospital, Algiers, Algeria.

Methods

We conducted a retrospective descriptive study of 31 patients who underwent surgical management for POP between January 2022 and December 2023. Data collected included age, parity, menopausal status, type and stage of prolapse, associated symptoms, and surgical procedures performed. Descriptive statistics were applied.

Results

The mean age was 63.2 years (range 47 to 80), with mean parity 5.7; 87% of patients were postmenopausal. Multicompartment prolapse predominated (87%), with stage III disease in 90% of cases. The most frequent procedure was vaginal hysterectomy with anterior and posterior colporrhaphy (VH + APR), performed in 58% of patients. Early postoperative outcomes were favorable: 90% recovered without major complications. Minor issues included transient

urinary retention and mild infection, managed conservatively. Average hospital stay was 3 to 5 days.

Conclusion

POP in this population was strongly associated with high parity and menopause, typically presenting at advanced stages. Vaginal surgical management, particularly VH + APR, proved effective and safe. Preventive strategies such as pelvic floor rehabilitation, family planning education, and earlier gynecological consultation are essential to reduce disease burden in resource limited settings.

Keywords: pelvic organ prolapse, pop-q, multiparity, vaginal hysterectomy, colporrhaphy, algeria, retrospective study

Introduction

Pelvic organ prolapse (POP) is a frequent and disabling condition in women, particularly in multiparous and postmenopausal patients. It results from weakening of the pelvic floor muscles and connective tissue that support the uterus, bladder, and rectum.

Global prevalence varies depending on definitions and populations studied, but POP is estimated to affect up to 50% of women over 50 years old [1,2]. Risk factors include high parity, aging, vaginal childbirth, menopause, obesity, chronic cough, and heavy physical labor [3,4]. In North African and South Asian populations, delayed consultation and sociocultural factors contribute to advanced-stage presentation, as reported in studies from Tunisia [5,6] and India [7].

This study aimed to describe the epidemiological and clinical characteristics of women treated for POP at a Tertiary Care Hospital, Algiers, Algeria, and to assess the surgical procedures performed and their short-term outcomes.

Materials and Methods

Study Design and Setting

Retrospective descriptive study conducted in the Department of Obstetrics and Gynecology, Tertiary Care Hospital, Algiers, Algeria, from January 2022 to December 2023.

Study Population

31 patients who underwent surgical management for POP.

- **Inclusion criteria**

  Clinical/paraclinical diagnosis of POP confirmed by a specialist, surgical treatment during the study period.

- **Exclusion criteria**

  Incomplete medical records, non-surgical management, associated gynecologic malignancy.

Data Collection

Medical records and operative reports were reviewed. Variables included age, parity, menopausal status, occupation, type and degree of prolapse, associated symptoms, surgical procedure, and short-term outcomes.

Surgical Management

All patients underwent vaginal surgery. The most frequent procedure was vaginal hysterectomy with anterior and posterior colporrhaphy (VH + APR). Additional procedures included perineorrhaphy and vault repair.

Data Analysis

Descriptive statistics were used. Quantitative variables expressed as means, ranges, and percentages.

Results

Epidemiological Characteristics

The mean age was 63.2 years (range 47-80), with nearly half of the patients between 60 and 69 years old. Multiparity was the most consistent risk factor, with a mean parity of 5.7 and more than two-thirds of patients reporting three or more deliveries. Obesity and menopausal status were also frequent contributors, with 87% of patients being postmenopausal. The age distribution of patients is summarized in Table 1, and the prevalence of major risk factors is illustrated in Figure 1.

Age group (years)	Patients (n)	Percentage (%)
40-49	1	3.2%
50-59	8	25.8%
60-69	14	45.2%
70-79	7	22.6%
≥80	1	3.2%
Total	**31**	**100%**

Table 1. Age distribution of women treated for pelvic organ prolapse in Algiers (2022–2023)

Tabulated data showing the frequency and percentage of cases across age groups among 31 patients. The table demonstrates that the majority of cases occurred in women aged 60–69 years, with smaller proportions in younger and older age groups.

Epidemiology

Common urogenital prolapse

Predominant in women

Prevalence increases with age

25-50% in multiparous women

Risk factors
- Parity
- Obesity
- Menopause

Figure 1. Epidemiology of pelvic organ prolapse in Algerian women.

Infographic summarizing the epidemiological profile of 31 women treated surgically for pelvic organ prolapse in Algiers (2022–2023). The diagram highlights the predominance of POP in women, its increasing prevalence with age, and a rate of 25–50% among multiparous women. Major risk factors identified include parity, obesity, and menopause. Created by the author based on study data.

Clinical Characteristics

Multi-compartment prolapse predominated, affecting 87% of patients, while isolated cystocele and rectocele were less frequent. Most cases presented at stage III according to the POP-Q classification, reflecting delayed consultation. Common symptoms included pelvic heaviness (71%), urinary disturbances (52%), and sexual discomfort (35%).

Surgical Management

All patients underwent vaginal surgery. The most frequent procedure was vaginal hysterectomy with anterior and posterior colporrhaphy (VH + APR), performed in 58% of cases. Other procedures included anterior colporrhaphy (19%), posterior colporrhaphy (13%), and perineorrhaphy/vault repair (10%). The distribution of surgical techniques is detailed in Table 2.

Procedure	Patients (n)	Percentage (%)
VH + APR	18	58%
Anterior colporrhaphy	6	19%
Posterior colporrhaphy	4	13%
Perineorrhaphy / vault repair	3	10%

Table 2. Vaginal surgical procedures performed for pelvic organ prolapse in 31 women in Algiers (2022–2023)

Summary of surgical techniques used, including vaginal hysterectomy with anterior and posterior colporrhaphy (VH + APR), anterior colporrhaphy, posterior colporrhaphy, and perineorrhaphy/vault repair. Frequencies and percentages are provided to illustrate the predominance of VH + APR in this cohort.

Postoperative Outcomes

Early outcomes were favorable, with 90% of patients recovering without major complications. Minor issues included transient urinary retention and mild infection, managed conservatively. No major hemorrhage, visceral injury, or recurrence was observed in the immediate postoperative period. The average hospital stay was 3–5 days.

Discussion

This study demonstrates that pelvic organ prolapse (POP) in Algeria predominantly affects older, multiparous, postmenopausal women, which is consistent with epidemiological patterns reported in other North African countries such as Tunisia and Morocco, as well as in South Asian populations like India [5-7]. The predominance of advanced-stage, multi-compartment prolapse in our cohort reflects delayed consultation, a finding that highlights the impact of sociocultural factors and limited awareness on disease progression. The epidemiological trends summarized

in Figure 1 reinforce the strong association between parity, obesity, and menopause as major risk factors.

International comparisons provide further context. In Europe and North America, earlier detection of POP is more common, largely due to preventive screening programs and greater patient education [8]. These differences emphasize the importance of awareness campaigns and accessible gynecological services in resource-limited settings such as Algeria, where patients often present only when symptoms become disabling.

Surgical management remains the cornerstone of treatment for advanced POP. In our series, vaginal hysterectomy with anterior and posterior colporrhaphy (VH + APR) was the most frequently performed procedure, accounting for more than half of all cases. The distribution of surgical techniques, summarized in Table 2, demonstrates the predominance of vaginal approaches, which are favored for their effectiveness and safety. Early postoperative outcomes were favorable, with low complication rates and no immediate recurrences, aligning with international literature that supports VH + APR as a reliable option when uterine preservation is not prioritized [9].

From a public health perspective, these findings underscore the need for preventive strategies. Pelvic floor muscle training, improved obstetric care, and family planning education could reduce the incidence of advanced POP. Encouraging earlier gynecological consultation may also help identify cases at earlier stages, improving outcomes and reducing the burden of complex surgery. Future research should focus on community-based prevalence studies and long-term surgical outcomes to guide national health policy and optimize patient care.

Conclusion

This retrospective study of 31 women treated surgically for pelvic organ prolapse in Algiers highlights the strong association between the condition and multiparity, menopause, and advancing age. The epidemiological profile, summarized in Figure 1 and Table 1, confirms that most patients presented at advanced stages, reflecting delayed consultation and limited awareness in this population. Vaginal surgical management, particularly vaginal hysterectomy with anterior and posterior colporrhaphy, was the predominant intervention, as shown in Table 2, and proved both effective and safe, with favorable short-term outcomes and minimal complications.

These findings emphasize the need for preventive strategies in Algeria and similar resource-limited settings. Pelvic floor rehabilitation, improved obstetric care, family planning education, and earlier gynecological consultation could reduce the burden of advanced prolapse and improve women's quality of life. Future research should focus on community-based

prevalence studies and long-term surgical outcomes to inform national health policy and optimize patient care.

References

1. Maher C, Feiner B, Baessler K, Schmid C. Surgical management of pelvic organ prolapse in women. *Cochrane Database Syst Rev.* 2013;2013(4):CD004014. doi:10.1002/14651858.CD004014.pub5

2. Swift SE. The distribution of pelvic organ support in a population of female subjects seen for routine gynecologic health care. *Am J Obstet Gynecol.* 2000;183(2):277-285. doi:10.1067/mob.2000.107583

3. Hendrix SL, Clark A, Nygaard I, Aragaki A, Barnabei V, McTiernan A. Pelvic organ prolapse in the Women's Health Initiative: gravity and gravidity. *Am J Obstet Gynecol.* 2002;186(6):1160-1166. doi:10.1067/mob.2002.123819

4. Jelovsek JE, Maher C, Barber MD. Pelvic organ prolapse. *Lancet.* 2007;369(9566):1027-1038. doi:10.1016/S0140-6736(07)60462-0

5. Ben Miled A, Chelli D, Boudaya F, et al. Pelvic organ prolapse: epidemiological and clinical study in Tunisian women. *Tunis Med.* 2015;93(4):234-239

6. El Fekih C, Ben Amara F, Ben Hmid A, et al. Pelvic organ prolapse: clinical and therapeutic aspects in 100 cases. *Pan Afr Med J.* 2014;17:234

7. Bhatia N, Bhatia K. Pelvic organ prolapse in Indian women: prevalence and risk factors. *Int J Reprod Contracept Obstet Gynecol.* 2016;5(7):2200-2204

8. Olsen AL, Smith VJ, Bergstrom JO, Colling JC, Clark AL. Epidemiology of surgically managed pelvic organ prolapse and urinary incontinence. *Obstet Gynecol.* 1997;89(4):501-506. doi:10.1016/S0029-7844(97)00058-6

9. Dietz HP, Haylen BT, Broome J. Ultrasound in the quantification of female pelvic organ prolapse. *Ultrasound Obstet Gynecol.* 2001;18(5):511-514. doi:10.1046/j.0960-7692.2001.00494.x