Perceptions of COVID-19 Vaccination During Pregnancy in a Cohort of US Adults (CHASING COVID) in 2023

Rachael Piltch-Loeb[1,2,3]*, PhD MSPH; Subha Balasubramanian[1], MPH; McKaylee Robertson[1,3], PhD MPH; Chloe Teasdale[1,3], PhD MPH; Sasha Fleary[1,4], PhD MS; Josefina Nuñez Sahr[1], MPH; Denis Nash[1,3], PhD MPH; Kate Penrose[1], MSPH; Bai Xi Jasmine Chan[1], MSPH; Angela Parcesepe[1,5,6], PhD MPH

Affiliations:

1. City University of New York (CUNY) Institute for Implementation Science in Population Health, New York, NY, USA.
2. Department of Environmental, Occupational, Geospatial Health Sciences, City University of New York (CUNY), New York, NY, USA
3. Epidemiology and Biostatistics, Graduate School of Public Health and Health Policy, City University of New York (CUNY), New York, NY, USA.
4. Department of Community Health and Social Sciences, Graduate School of Public Health and Health Policy, City University of New York (CUNY), New York, NY, USA
5. Maternal and Child Health, University of North Carolina at Chapel Hill Gillings School of Global Public Health, Chapel Hill, North Carolina, USA.
6. Carolina Population Center, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

Corresponding Author
Email: rachael.piltch-loeb@sph.cuny.edu (RP)

Abstract

Objective

Pregnant women are at higher risk of severe COVID-19, and vaccination significantly reduces the risk of severe infection. Despite its benefits, only 13% of pregnant women in the U.S. had received the updated 2024–25 vaccine by December 2024, with uptake varying across sociodemographic groups. This study examines perceptions on COVID-19 vaccination during pregnancy among U.S. adults enrolled in the Chasing Covid Cohort, analyzing responses across 16 surveys between March 2020 and December 2023 (N=4488).

Methods

Key variables included sociodemographic characteristics, susceptibility to severe COVID-19 disease, perceived worry about COVID-19, individual and household vaccination status, symptoms of anxiety and depression, trusted information sources, and having a regular healthcare provider. Perceptions of vaccine safety and efficacy during pregnancy were measured using five Likert-scale statements, categorized into agreement, uncertainty, and disagreement. Exploratory factor analysis identified two constructs—safety and efficacy—which were analyzed in relation to participant characteristics using bivariate analysis and chi-square tests, and multivariable robust Poisson regression models.

Results

Among all respondents and women of reproductive age, less than half (40%) perceived the COVID-19 vaccine as safe during pregnancy, and just over half recognized its efficacy. Individuals with a personal physician and those who trusted public health institutions or healthcare providers were more inclined to agree with the vaccine's safety and efficacy.

Conclusions

These findings highlight the influence of demographic factors on vaccine perceptions, the potential impact of social networks during pregnancy, and the critical role of trust in public health institutions in promoting vaccine uptake.

1. Background

Pregnant women and birthing people ("pregnant women") are at higher risk of severe COVID-19[1,2]. Pregnant women with COVID-19 were more likely to be admitted to an intensive care unit (ICU), require invasive ventilation, require extracorporeal membrane oxygenation, and die when compared to nonpregnant women of reproductive age with COVID-19, according to the United States Centers for Disease Control and Prevention (CDC) COVID-19 surveillance system[1]. SARS-CoV-2 infection during pregnancy may also be associated with several adverse birth outcomes including preeclampsia, preterm birth, and stillbirth, especially among those with severe COVID-19 disease[2].

COVID-19 vaccination significantly reduces the risk of severe SARS-CoV-2 infection in the general population and during pregnancy[3]. Despite being largely excluded from the initial clinical trials, pregnant women have since been included in routine roll-out and follow-up studies focusing on the safety and efficacy of COVID-19 vaccination[4]. A 2021 cohort study of over 10,000 vaccinated pregnant women found the vaccine 89% effective preventing COVID-19-related hospitalization[5]. A 2022 global review found no major mRNA vaccine adverse events among pregnant women and confirmed their effectiveness preventing severe disease[3]. Additionally, maternal mRNA COVID-19 vaccination during pregnancy was associated with lower risks of severe neonatal morbidity, death, and ICU admission and no increase in neonatal readmission or hospital admission up to age 6 months[6]. Until infants are age-eligible for vaccination (six months), maternal vaccination provides passive protection against symptomatic infection[7].

The CDC and the American College of Obstetrics and Gynecology (ACOG) recommend COVID-19 vaccination during pregnancy [8,9]. However, as of December 2024, only 13% of pregnant women received an updated 2024-25 COVID-19 vaccine[10]. Uptake varies significantly by race/ethnicity - highest amongst non-Hispanic Asian (22%) and non-Hispanic White women (18%), with lower uptake observed among Hispanic (8%) and Black (6%) women[10], age (lower in younger women)[11], and education (highest with college education), urban residence and higher income [12,11]. Access to healthcare, including insurance, a primary care provider, or vaccination counseling from a healthcare provider, has been associated with higher odds of COVID-19 vaccination during pregnancy[13].

Despite ongoing efforts, significant gaps remain in understanding COVID-19 vaccine hesitancy during pregnancy. Addressing these gaps is crucial given the heightened risk of severe complications from COVID-19 for pregnant women and evolving vaccine recommendations. To gain a broader understanding of attitudes toward COVID-19 vaccination during pregnancy, we examined acceptability for pregnant women and its relationship with sociodemographics, healthcare experiences, risk perception, and trust in public health institutions in a cohort of US adults >=18 years and among women of reproductive age (18-49 years). Focusing on women of reproductive age highlights unique immunization concerns during/after pregnancy, while including all adults allows exploration of generational implications on vaccine attitudes. By deconstructing these diverse perspectives, this study hopes to provide significant insights for developing effective nuanced health communication that connects with a wide audience and addresses specific concerns of key subpopulations.

2. Methods

2.1 Study Population

The Communities, Households, and SARS-CoV-2 Epidemiology (CHASING) COVID Cohort study is a national prospective study initiated in March 2020, during the onset of the COVID-19 pandemic in the United States. We employed internet-based recruitment strategies to assemble a geographically and socio-demographically diverse cohort of participants aged ≥18 residing in the U.S. or its territories. Follow-up surveys were conducted approximately quarterly from March 2020 through December 2023. Further details on recruitment and procedures are available elsewhere[14]. Recruitment of the original study was completed between 28/03/2020 and 21/08/2020.This study was approved by the Institutional Review Boards of the City University of New York (CUNY) (New York, NY, USA) (protocol 2020-0256).

COVID-19 Vaccine Safety and Efficacy Perceptions During Pregnancy

We assessed perceptions of the COVID-19 vaccine's safety and efficacy during pregnancy in December 2023 using responses to five specific statements: a) Not enough is known about the long-term side effects of receiving the COVID-19 vaccine during pregnancy; b) There is not enough research to support getting the COVID-19 vaccine during pregnancy; c) The COVID-19 vaccine is safe to receive during pregnancy; d) The COVID-19 vaccine reduces the risk of severe disease from COVID-19 during pregnancy; e) Receiving the COVID-19 vaccine while pregnant helps protect infants until they are old enough to be vaccinated.

Data were collected using a 5-point Likert scale, which was collapsed into three categories for analysis: "Agreement" ("strongly agree" or "agree"), "Uncertainty" ("neither agree

115 nor disagree"), and "Disagreement" ("strongly disagree" or "disagree"). For two statements,

116 responses were reverse-coded to align with the direction of the other statements.

2.2 Participant Characteristics

Age, gender, race/ethnicity, education, and household annual income were collected

at cohort enrollment. Participants were classified as pregnant or recently pregnant if they

reported being pregnant at any time between March 2020 and December 2023. Both cisgender

women and non-binary individuals reporting a pregnancy were included. No transgender men

reported a pregnancy. We identified participants with any children <18 years in the household as

of October 2023.

a) Susceptibility to severe COVID-19:

Susceptibility to severe COVID-19 disease was assessed at cohort enrollment

using a composite variable based on factors the CDC identified as increasing risk for

COVID-19 complications given SARS-CoV-2 infection[15]. These included age ≥ 60

years, self-report of chronic lung disease (CLD) or chronic obstructive pulmonary disease

(COPD), current asthma, diabetes, serious heart conditions (including heart attack, high

blood pressure, angina), kidney disease, immunocompromised status, HIV, or daily

smoking. Each factor was coded as present (1) or absent (0), and a cumulative score was

calculated by summing these binary variables, resulting in a score from 0-9. This score

was then dichotomized based on the median as: more (score >1) or less (score ≤ 1)

susceptible to severe COVID-19.

b) Vaccination status - individual and household:

In over 16 rounds of follow-up surveys, participants were asked to report their

personal and household members' COVID-19 vaccination status. Participants were

considered vaccinated if they reported completing their primary vaccination series (2

doses of a two-dose vaccine [e.g., mRNA vaccines] or 1 dose of a single-dose vaccine

[e.g., J&J]). Participants were considered boosted in 2023 if they reported receiving a

booster dose in 2023.

As of October 2023, if at least one COVID-19 vaccine-eligible person, including

the enrolled participant, in a household, was COVID-19 vaccinated, the household was

considered vaccinated.

c) Trusted information sources:

In October 2023, participants were asked to identify various entities, groups, or

individuals they trusted to provide reliable information about the COVID-19 vaccine (see

Appendix). Trust in public health institutions was defined as endorsing trust in any of the

following national or international public health agencies: CDC, the World Health

Organization (WHO), or the Food and Drug Administration (FDA). Participants' trust in

healthcare providers was defined as endorsing personal physicians or other healthcare

professionals as providers of reliable information about the COVID-19 vaccine.

d) Regular healthcare provider:

Participants were categorized as having a personal doctor if they indicated they

had one person they think of as their personal doctor or healthcare provider, as assessed

in October 2023.

e) Symptoms of anxiety and depression:

In December 2023, participants completed the seven-item Generalized Anxiety

Disorder (GAD-7) scale and the eight-item Patient Health Questionnaire (PHQ-8). Those

who scored ≥10 on the GAD-7 were classified as having moderate to severe anxiety

symptoms, and those who scored ≥10 on the PHQ-8 were classified as having moderate

to severe depression symptoms [16, 17].

f) Perceived worry:

Assessed in December 2023, participants were coded as "worried" if they

responded "somewhat worried" or "very worried" to either of the following questions:

How worried are you about getting sick from COVID-19?; How worried are you about

the long-term effects of COVID-19 infection(s)?

2.3 Analytic Approach

For this analysis, we included CHASING COVID cohort participants who completed

enrollment surveys between March and July 2020, and who completed the December 2023

survey, which included questions about COVID-19 vaccination concerns during pregnancy. The

analysis was conducted first among all adults and then only among women of reproductive age

(18-49 years at enrollment). Individuals who did not answer all vaccine perception questions

were excluded from the analysis. We calculated the proportion of respondents who agreed with

each COVID-19 vaccine perception statement, indicating confidence in the vaccine.

Factor Analysis

We conducted an EFA to identify underlying constructs related to COVID-19 vaccine

perceptions during pregnancy. The iterated principal axis factoring method with varimax rotation

was used to extract two distinct factors: safety and efficacy perceptions. These categories were

then used to classify participants according to their perceptions of safety and efficacy. Bivariate

analyses examined associations between each identified factor and participant characteristics.

Chi-square tests were utilized to explore the relationships between these characteristics, and the

factors identified in the EFA.

Multivariable regression models

We used multivariable robust Poisson regression [18] to identify participant

characteristics correlated with agreement that COVID-19 vaccination during pregnancy was safe

or effective. We ran a separate model for each construct (vaccine safety and vaccine efficacy)

and for two different comparisons: 1. "any agreement" versus "some disagreement," excluding

those categorized as uncertain, and 2. "any agreement" versus "uncertainty," excluding those

categorized as having some disagreement. We ran models among all adults and among women of

reproductive age. This resulted in eight models across outcome, comparison, and population

combinations. For each predictive model, we included age, gender, education, race/ethnicity,

moderate-to-severe anxiety symptoms, moderate-to-severe depression symptoms, having a

personal doctor, susceptibility to severe COVID-19, trust in public health institutions, and trust

in healthcare providers. The model among all adults also included gender. All analyses were

performed using SAS 9.4 (Cary, NC, USA).

3. Results

Of the 4537 participants from the CHASING COVID Cohort who completed the

December 2023 survey, 4488 (99%) provided complete responses to the vaccine safety and

efficacy statements during pregnancy questions. Table 1 characterizes the study participants. Of

all participants, 36% (N=1,611) were women of reproductive age and, of those, 13% were

pregnant or had been recently pregnant. Table 2 and Table 3 present summary statistics of

COVID-19 vaccine perceptions during pregnancy within the study cohort among all adults and

women of reproductive age, respectively. Among all adults, the proportion of participants

uncertain about a statement ranged from 38% to 53%. Notably, more than half of the

participants, 53% (N=2384), disagreed with the statement, "The COVID-19 vaccine reduces the

risk of severe disease from COVID-19 during pregnancy". Among women of reproductive age,

55% (N=879) disagreed with the statement "The COVID-19 vaccine reduces the risk of severe

disease from COVID-19 during pregnancy".

Table 1. **Participant characteristics**

Characteristics	All adults	Women of reproductive age[a]
Total, N	**4488**	**1611**
Age category, N (col %)		
Age category: 18-29	985 (21.9%)	595 (36.9%)
Age category: 30-39	1299 (28.9%)	606 (37.6%)
Age category: 40-49	832 (18.5%)	410 (25.5%)
Age category: 50-64	600 (13.4%)	NA
Age category: 65+	772 (17.2%)	NA
Gender, N (col %)		
Gender - Female/Non-binary[a]	2510 (55.9%)	1611 (100%)
Gender - Male	1978 (44.1%)	NA
Race/Ethnicity, N (col %)		
Race/Ethnicity - NH White	2796 (62.3%)	856 (53.1%)
Race/Ethnicity - Hispanic	749 (16.7%)	342 (21.2%)
Race/Ethnicity - NH Asian/PI/Other	456 (10.2%)	211 (13.1%)
Race/Ethnicity - NH Black	487 (10.9%)	202 (12.5%)
Education, N (col %)		
Education - <high school/high school	507 (11.3%)	241 (15%)
Education - Some college	1176 (26.2%)	468 (29.1%)
Education - College graduate	2805 (62.5%)	902 (56%)

Household annual income, N (col %)		
Income - <$50k/Unknown	1834 (40.9%)	748 (46.4%)
Income - $50k-$100k	1424 (31.8%)	477 (29.6%)
Income - >$100k	1227 (27.4%)	386 (24%)
Any <18 y in household, N (col %)		
Any <18y kid in household - No	2543 (56.7%)	722 (44.8%)
Any <18y kid in household - Yes	1945 (43.3%)	889 (55.2%)
Pregnant, N (col %)		
Ever pregnant	210 (4.7%)	209 (13%)
Never pregnant	4278 (95.3%)	1402 (87%)
Moderate to severe symptoms of anxiety as of December 2023, N (col %)		
Moderate to severe symptoms of anxiety - No	3839 (85.5%)	1288 (80%)
Moderate to severe symptoms of anxiety - Yes	649 (14.5%)	323 (20.1%)
Depression status (PHQ) as of December 2023, N (col %)		
Depression status (PHQ) - No	3682 (82%)	1230 (76.4%)
Depression status (PHQ) - Yes	806 (18%)	381 (23.7%)
Has personal doctor as of December 2023, N (col %)		
Has personal doctor - No	1050 (24.5%)	483 (31.3%)
Has personal doctor - Yes	3237 (75.5%)	1061 (68.7%)
Susceptible to severe COVID-19 (if infected) at baseline, N (col %)		
Susceptible to severe COVID-19 (if infected) - No	3527 (78.6%)	1447 (89.8%)
Susceptible to severe COVID-19 (if infected) - Yes	961 (21.4%)	164 (10.2%)
Vaccination status as of December 2023, N (col %)		
Vaccination status - No	63 (1.5%)	34 (2.5%)
Vaccination status - Yes	4030 (98.5%)	1354 (97.6%)
Received Booster dose in 2023 N (col %)		
Boosted in 2023 - No	2653 (59.1%)	1137 (70.6%)
Boosted in 2023 - Yes	1835 (40.9%)	474 (29.4%)
Household members vaccinated as of October 2023, N (col %)		
Household members vaccinated - No	504 (11.8%)	275 (17.8%)
Household members vaccinated - Yes	3774 (88.2%)	1266 (82.2%)
Trust in government sources as of October 2023, N (col %)		
Trust in government information - No	926 (21.7%)	377 (24.5%)

Trust in government information - Yes	3339 (78.3%)	1160 (75.5%)
Trust in healthcare providers as of October 2023, N (col %)		
Trust in healthcare providers - No	1267 (29.7%)	581 (37.8%)
Trust in healthcare providers - Yes	2998 (70.3%)	956 (62.2%)
Perceived worry, as of December 2023 N (col %)		
Perceived worry - No	2372 (52.9%)	762 (47.3%)
Perceived worry - Yes	2116 (47.1%)	849 (52.7%)

[a]We classify all gender minorities (N = 129) with women due to small numbers and in recognition of the increased health risks in this population.

Table 2. Summary Statistics of COVID-19 Vaccine Perception Statements as reported by all adults >=18 years in the CHASING COVID Cohort (N=4488)

N (row%)	Agreement[b]	Uncertainity[c]	Disagreement[d]
a) Enough is known about the long-term side effects of receiving the COVID-19 vaccine during pregnancy[a]	1351 (30.1%)	1699 (37.8%)	1438 (32.0%)
b) There is enough research to support getting the COVID-19 vaccine during pregnancy[a]	1047 (23.3%)	1806 (40.2%)	1635 (36.4%)
c) The COVID-19 vaccine is safe to receive during pregnancy	468 (10.4%)	2112 (47.0%)	1908 (42.5%)
d) The COVID-19 vaccine reduces the risk of severe disease from COVID-19 during pregnancy	322 (7.2%)	1782 (39.7%)	2384 (53.1%)
e) Receiving the COVID-19 vaccine while pregnant helps protect infants until they are old enough to be vaccinated	401 (8.9%)	2386 (53.2%)	1701 (37.9%)

[a]Statements have been reworded and reverse coded to ensure consistent interpretation across all statements.
[b]Participants who expressed agreement with at least one statement within the factor were categorized as agreeing with the belief that COVID-19 vaccination during pregnancy is safe and effective.
[c]Participants who expressed uncertainty for all statements within the factor were categorized as having uncertainty
[d]Participants who expressed uncertainty or disagreement for all statements within the factor, but not exclusively uncertainty, were categorized as having some disagreement.
Note: We classify all gender minorities (N = 129) with women due to small numbers and in recognition of the increased health risks in this population.

Table 3: Summary Statistics of COVID-19 Vaccine Perception Statements as reported by women of reproductive age in the CHASING COVID Cohort (N=1611)

N (row%)	Agreement[b]	Uncertainity[c]	Disagreement[d]
a) Enough is known about the long-term side effects of receiving the COVID-19 vaccine during pregnancy[a]	584(36.2%)	536(33.2%)	491(30.4%)
b) There is enough research to support getting the COVID-19 vaccine during pregnancy[a]	457(28.3%)	555(34.4%)	599(37.2%)
c) The COVID-19 vaccine is safe to receive during pregnancy	199(12.3%)	696(43.2%)	716(44.4%)
d) The COVID-19 vaccine reduces the risk of severe disease from	152(9.4%)	580(36.0%)	879(54.5%)

COVID-19 during pregnancy			
e) Receiving the COVID-19 vaccine while pregnant helps protect infants until they are old enough to be vaccinated	183(11.4%)	756(46.9%)	672(41.7%)

[a]Statements have been reworded and reverse coded to ensure consistent interpretation across all statements.

[b]Participants who expressed agreement with at least one statement within the factor were categorized as agreeing with the belief that COVID-19 vaccination during pregnancy is safe and effective.

[c]Participants who expressed uncertainty for all statements within the factor were categorized as having uncertainty

[d]Participants who expressed uncertainty or disagreement for all statements within the factor, but not exclusively uncertainty, were categorized as having some disagreement.

Note: We classify all gender minorities (N = 129) with women due to small numbers and in recognition of the increased health risks in this population.

Factor Analysis

As seen in Table 4, the EFA identified two factors related to any agreement with vaccine

perceptions: safety and efficacy. For all adults, the items loaded significantly onto these two

factors, with statements regarding the long-term side effects, safety, and sufficient research to

support vaccination positively loading on the safety factor, while statements about the protection

of infants and reduction of risk of severe disease positively loaded on the efficacy factor. The

factor analysis for the subpopulation of women of reproductive age showed similar patterns. The

reliability analysis conducted among all adults yielded a Cronbach's alpha of 0.84 for safety

perceptions and 0.85 for efficacy perceptions, indicating good internal consistency. The

proportions of safety and efficacy perceptions were comparable between all adults and the

subpopulation of women of reproductive age. Among both groups, more than half expressed any

agreement with COVID-19 vaccine efficacy statements, with 58.2% of all adults and 59.7% of

women of reproductive age reporting any agreement (Table 5).

Table 4. **Vaccine perceptions during pregnancy scale items factor loading matrix (EFA)**

Items	All adult participants (N=4488)		Women of reproductive age (N=1611)[a]	
	Safety perceptions	Efficacy perceptions	Safety perceptions	Efficacy perceptions
a) Enough is known about the long-term side effects of receiving the COVID-19 vaccine during pregnancy[b]	**0.80147**	-0.23126	**0.79258**	-0.21322

b) There enough research to support getting the COVID-19 vaccine during pregnancy[b]	**0.82022**	-0.31582	**0.80449**	-0.34271
c) The COVID-19 vaccine is safe to receive during pregnancy	-0.35344	**0.72717**	-0.35647	**0.7449**
d) The COVID-19 vaccine reduces the risk of severe disease from COVID-19 during pregnancy	-0.2366	**0.82136**	-0.23782	**0.82952**
e) Receiving the COVID-19 vaccine while pregnant helps protect infants until they are old enough to be vaccinated	-0.21824	**0.75018**	-0.23292	**0.77648**

[a]We classify all gender minorities (N = 129) with women due to small numbers and in recognition of the increased health risks in this population.
[b]Statement were re-worded and reverse-scored to ensure consistent interpretation across all statements.
Note: Factor loadings representative of a particular factor appear in boldface.

Table 5: Alpha reliabilities and mean (SD) and median (IQR) COVID-19 perceptions during pregnancy by population

	All adult participants (N=4488)			Women of reproductive age (N=1611)[a]			
	Any agreement[b]	Uncertainty[c]	Some disagreement[d]	Any agreement[b]	Uncertainty[c]	Some disagreement[d]	Cronbach's alpha
Safety perceptions	1815 (40.4%)	1367 (30.5%)	1306 (29.1%)	655 (40.7%)	389 (24.1%)	567 (35.2%)	0.84
Efficacy perceptions	2611 (58.2%)	1390 (30.9%)	487 (10.9%)	961 (59.7%)	434 (26.9%)	216 (13.4%)	0.85

[a]We classify all gender minorities (N = 129) with women due to small numbers and in recognition of the increased health risks in this population.
[b]Participants who expressed agreement with at least one statement within the factor were categorized as agreeing with the belief that COVID-19 vaccination during pregnancy is safe and effective.
[c]Participants who expressed uncertainty for all statements within the factor were categorized as having uncertainty
[d]Participants who expressed uncertainty or disagreement for all statements within the factor, but not exclusively uncertainty, were categorized as having some disagreement.
Note: Statements a) and b) have been reworded and reverse coded to ensure consistent interpretation across all statements.

Participant characteristics associated with vaccine safety and efficacy perceptions

Among all adults, 40.4% had any agreement, 30.5% were uncertain, and 29.1% had some disagreement with the statements regarding the safety of the COVID-19 vaccine during pregnancy; 58.2% had any agreement, 30.9% were uncertain, and 10.9% had some disagreement with the statements concerning the vaccine's efficacy (Table 5).

Participants aged 30-39, non-Hispanic White individuals, and college graduates without

children were more likely to have any agreement with statements about the safety and efficacy of

the COVID-19 vaccine during pregnancy compared to those who expressed some disagreement

(Table 6). Additionally, participants who did not report moderate to severe symptoms of anxiety

or depression, and those who reported being fully vaccinated as of December 2023 or boosted in

2023, had a personal physician, and expressed trust in the public health institutions or healthcare

providers were more likely to have any agreement with statements about the COVID-19

vaccine's safety and efficacy during pregnancy than those who expressed some disagreement.

Chi-square tests indicated that all characteristics were significantly associated with safety

perceptions of the COVID-19 vaccine during pregnancy. However, moderate to severe

symptoms of anxiety and depression were not significantly associated with efficacy perceptions

of the vaccine during pregnancy (Table 6).

Table 6. Participant characteristics by perception towards COVID vaccine concerns during pregnancy among all adult participants (N=4488)

Characteristics	Safety perceptions				Efficacy perceptions			
	Any agreement[b]	Uncertainity[c]	Some disagreement[d]	p-value	Any agreement[b]	Uncertainity[c]	Some disagreement[d]	p-value
Total, N (%)	1815 (40.4%)	1367 (30.5%)	1306 (29.1%)		2611 (58.2%)	1390 (30.9%)	487 (10.9%)	
Age category, N (col %)								
Age category: 18-29	413(23%)	256(19%)	316(24%)	<0.001	632(24%)	238(17%)	115(24%)	<0.001
Age category: 30-39	572(32%)	351(26%)	376(29%)		777(30%)	378(27%)	144(30%)	
Age category: 40-49	338(19%)	256(19%)	238(18%)		468(18%)	274(20%)	90(18%)	
Age category: 50-64	219(12%)	201(15%)	180(14%)		320(12%)	223(16%)	57(12%)	
Age category: 65+	273(15%)	303(22%)	196(15%)		414(16%)	277(20%)	81(17%)	
Gender, N (col %)								
Gender - Female/Non-binary[a]	966(53%)	720(53%)	824(63%)	<0.001	1433(55%)	771(55%)	306(63%)	0.005
Gender - Male	849(47%)	647(47%)	482(37%)		1178(45%)	619(45%)	181(37%)	
Race/Ethnicity, N (col %)								
Race/Ethnicity - NH White	1214(67%)	895(65%)	687(53%)	<0.001	1735(66%)	819(59%)	242(50%)	<0.001

Race/Ethnicity - Hispanic	271(15%)	212(16%)	266(20%)		384(15%)	251(18%)	114(23%)	
Race/Ethnicity - NH Asian/PI/Other	175(10%)	135(10%)	146(11%)		273(10%)	140(10%)	43(9%)	
Race/Ethnicity - NH Black	155(9%)	125(9%)	207(16%)		219(8%)	180(13%)	88(18%)	
Education, N (col %)								
Education - <high school/high school	138(8%)	148(11%)	221(17%)	<0.001	210(8%)	206(15%)	91(19%)	<0.001
Education - Some college	367(20%)	364(27%)	445(34%)		545(21%)	436(31%)	195(40%)	
Education - College graduate	1310(72%)	855(63%)	640(49%)		1856(71%)	748(54%)	201(41%)	
Household annual income, N (col %)								
Income - <$50k/Unknown	631(35%)	572(42%)	631(48%)	<0.001	936(36%)	659(47%)	239(49%)	<0.001
Income - $50k-$100k	572(32%)	432(32%)	420(32%)		835(32%)	423(30%)	166(34%)	
Income - >$100k	612(34%)	361(26%)	254(19%)		839(32%)	306(22%)	82(17%)	
Any <18 y in household, N (col %)								
Any <18y kid in household - No	1241(68%)	1004(73%)	701(54%)	<0.001	1741(67%)	966(69%)	239(49%)	<0.001
Any <18y kid in household - Yes	574(32%)	363(27%)	605(46%)		870(33%)	424(31%)	248(51%)	
Pregnant, N (col %)								
Ever pregnant	103(6%)	29(2%)	78(6%)	<0.001	146(6%)	35(3%)	29(6%)	<0.001
Never pregnant	1712(94%)	1338(98%)	1228(94%)		2465(94%)	1355(97%)	458(94%)	
Moderate to severe symptoms of anxiety as of December 2023, N (col %)								
Moderate to severe symptoms of anxiety - No	1588(87%)	1184(87%)	1067(82%)	<0.001	2240(86%)	1193(86%)	406(83%)	0.35
Moderate to severe symptoms of anxiety - Yes	227(13%)	183(13%)	239(18%)		371(14%)	197(14%)	81(17%)	
Depression status (PHQ) as of December 2023, N (col %)								
Depression status (PHQ) - No	1531(84%)	1129(83%)	1022(78%)	<0.001	2155(83%)	1140(82%)	387(79%)	0.27
Depression status (PHQ) - Yes	284(16%)	238(17%)	284(22%)		456(17%)	250(18%)	100(21%)	
Has personal doctor as of December 2023, N (col %)				0.003				<0.001
Has personal doctor - No	397(23%)	306(24%)	347(28%)		553(22%)	343(26%)	154(33%)	

Has personal doctor - Yes	1348(77%)	993(76%)	896(72%)		1952(78%)	974(74%)	311(67%)	
Susceptible to severe COVID-19 (if infected) at baseline, N (col %)								
Susceptible to severe COVID-19 (if infected) - No	1479(81%)	1047(77%)	1001(77%)	<0.001	2109(81%)	1040(75%)	378(78%)	<0.001
Susceptible to severe COVID-19 (if infected) - Yes	336(19%)	320(23%)	305(23%)		502(19%)	350(25%)	109(22%)	
Vaccination status as of December 2023, N (col %)								
Vaccination status - No	17(1%)	11(1%)	35(4%)	<0.001	24(1%)	19(2%)	20(7%)	<0.001
Vaccination status - Yes	1771(99%)	1304(99%)	955(96%)		2533(99%)	1222(98%)	275(93%)	
Received Booster dose in 2023 N (col %)								
Boosted in 2023 - No	800(44%)	782(57%)	1071(82%)	<0.001	1284(49%)	933(67%)	436(90%)	<0.001
Boosted in 2023 - Yes	1015(56%)	585(43%)	235(18%)		1327(51%)	457(33%)	51(10%)	
Household members vaccinated as of October 2023, N (col %)								
Household members vaccinated - No	73(4%)	125(10%)	306(25%)	<0.001	131(5%)	198(15%)	175(38%)	<0.001
Household members vaccinated - Yes	1671(96%)	1169(90%)	934(75%)		2371(95%)	1115(85%)	288(62%)	
Trust in public health institutions as of October 2023, N (col %)								
Trust in public health institutions - No	177(10%)	204(16%)	545(44%)	<0.001	295(12%)	339(26%)	292(63%)	<0.001
Trust in public health institutions - Yes	1565(90%)	1082(84%)	692(56%)		2202(88%)	966(74%)	171(37%)	
Trust in healthcare providers as of October 2023, N (col %)								
Trust in healthcare providers - No	411(24%)	306(24%)	550(44%)	<0.001	588(24%)	423(32%)	256(55%)	<0.001
Trust in healthcare providers - Yes	1331(76%)	980(76%)	687(56%)		1909(76%)	882(68%)	207(45%)	
Perceived worry, as of December 2023 N (col %)								
Perceived worry - No	846(47%)	745(54%)	781(60%)	<0.001	1223(47%)	808(58%)	341(70%)	<0.001
Perceived worry - Yes	969(53%)	622(46%)	525(40%)		1388(53%)	582(42%)	146(30%)	

[a]We classify all gender minorities (N = 129) with women due to small numbers and in recognition of the increased health risks in this population.

[b]Participants who expressed agreement with at least one statement within the factor were categorized as agreeing with the belief that COVID-19 vaccination during pregnancy is safe and effective.

[c]Participants who expressed uncertainty for all statements within the factor were categorized as having uncertainty

[d]Participants who expressed uncertainty or disagreement for all statements within the factor, but not exclusively uncertainty, were

Note: Statements corresponding to safety belief have been reworded and reverse coded to ensure consistent interpretation across a statements.

While findings were similar among women of reproductive age, in that group, women

aged 18-29 and those more worried about getting sick again from COVID-19 were more likely to

have any agreement about the safety and efficacy statements of the COVID-19 vaccine during

pregnancy (Table 7). Similar to all adults, moderate to severe symptoms of anxiety and

depression were not significantly associated with perceptions of COVID-19 vaccine efficacy

(Table 7).

Table 7. Participant characteristics by perception towards COVID vaccine concerns during pregnancy among women[a] of reproductive age (N=1611)

Characteristics	Safety perceptions				Efficacy perceptions			
	Any agreement[b]	Uncertainity[c]	Some disagreement[d]	p-value	Any agreement[b]	Uncertainity[c]	Some disagreement[d]	p-value
Total, N (%)	655 (40.7%)	389 (24.1%)	567 (35.2%)		961 (59.7%)	434 (26.9%)	216 (13.4%)	
Age category, N (col %)								
Age category: 18-29	242(37%)	147(38%)	206(36%)	0.68	381(40%)	139(32%)	75(35%)	0.0
Age category: 30-39	249(38%)	135(35%)	222(39%)		341(35%)	174(40%)	91(42%)	
Age category: 40-49	164(25%)	107(28%)	139(25%)		239(25%)	121(28%)	50(23%)	
Race/Ethnicity, N (col %)								
Race/Ethnicity - NH White	399(61%)	206(53%)	251(44%)	<0.001	573(60%)	199(46%)	84(39%)	<0.00
Race/Ethnicity - Hispanic	114(17%)	87(22%)	141(25%)		172(18%)	106(24%)	64(30%)	
Race/Ethnicity - NH Asian/PI/Other	81(12%)	51(13%)	79(14%)		130(14%)	59(14%)	22(10%)	
Race/Ethnicity - NH Black	61(9%)	45(12%)	96(17%)		86(9%)	70(16%)	46(21%)	
Education, N (col %)				<0.001				<0.00
Education - <high	57(9%)	58(15%)	126(22%)		92(10%)	97(22%)	52(24%)	

school/high school								
Education - Some college	141(22%)	120(31%)	207(37%)		215(22%)	160(37%)	93(43%)	
Education - College graduate	457(70%)	211(54%)	234(41%)		654(68%)	177(41%)	71(33%)	
Household annual income, N (col %)								
Income - <$50k/Unknown	258(39%)	186(48%)	304(54%)	<0.001	371(39%)	250(58%)	127(59%)	<0.00
Income - $50k-$100k	177(27%)	120(31%)	180(32%)		298(31%)	120(28%)	59(27%)	
Income - >$100k	220(34%)	83(21%)	83(15%)		292(30%)	64(15%)	30(14%)	
Any <18 y in household, N (col %)								
Any <18y kid in household - No	365(56%)	198(51%)	204(36%)	<0.001	504(52%)	199(46%)	64(30%)	<0.00
Any <18y kid in household - Yes	290(44%)	191(49%)	363(64%)		457(48%)	235(54%)	152(70%)	
Pregnant, N (col %)								
Ever pregnant	103(16%)	29(7%)	77(14%)	<0.001	145(15%)	35(8%)	29(13%)	0.00
Never pregnant	552(84%)	360(93%)	490(86%)		816(85%)	399(92%)	187(87%)	
Moderate to severe symptoms of anxiety as of December 2023, N (col %)								
Moderate to severe symptoms of anxiety - No	546(83%)	317(81%)	425(75%)	<0.001	782(81%)	338(78%)	168(78%)	0.2
Moderate to severe symptoms of anxiety - Yes	109(17%)	72(19%)	142(25%)		179(19%)	96(22%)	48(22%)	
Depression status (PHQ) as of December 2023, N (col %)								
Depression status (PHQ) - No	522(80%)	302(78%)	406(72%)	0.003	751(78%)	320(74%)	159(74%)	0.1
Depression status (PHQ) - Yes	133(20%)	87(22%)	161(28%)		210(22%)	114(26%)	57(26%)	
Has personal doctor as of December 2023, N (col %)								
Has personal doctor - No	171(27%)	131(35%)	181(34%)	0.009	252(27%)	151(37%)	80(39%)	<0.00
Has personal doctor - Yes	463(73%)	243(65%)	355(66%)		675(73%)	261(63%)	125(61%)	
Susceptible to severe COVID-19 (if infected) at baseline, N (col %)				<0.001				<0.00

Susceptible to severe COVID-19 (if infected) - No	609(93%)	359(92%)	479(84%)		885(92%)	380(88%)	182(84%)	
Susceptible to severe COVID-19 (if infected) - Yes	46(7%)	30(8%)	88(16%)		76(8%)	54(12%)	34(16%)	
Vaccination status as of December 2023, N (col %)				<0.001				<0.00
Vaccination status - No	7(1%)	5(1%)	22(6%)		11(1%)	13(4%)	10(9%)	
Vaccination status - Yes	632(99%)	353(99%)	369(94%)		918(99%)	339(96%)	97(91%)	
Received Booster dose in 2023 N (col %)				<0.001				<0.00
Boosted in 2023 - No	352(54%)	286(74%)	499(88%)		579(60%)	355(82%)	203(94%)	
Boosted in 2023 - Yes	303(46%)	103(26%)	68(12%)		382(40%)	79(18%)	13(6%)	
Household members vaccinated as of October 2023, N (col %)				<0.001				<0.00
Household members vaccinated - No	30(5%)	67(18%)	178(33%)		71(8%)	107(26%)	97(48%)	
Household members vaccinated – Yes	603(95%)	307(82%)	356(67%)		854(92%)	305(74%)	107(52%)	
Trust in public health institutions as of October 2023, N (col %)				<0.001				<0.00
Trust in public health institutions - No	77(12%)	68(18%)	232(44%)		130(14%)	124(30%)	123(60%)	
Trust in public health institutions - Yes	554(88%)	305(82%)	301(56%)		792(86%)	287(70%)	81(40%)	
Trust in healthcare providers as of October 2023, N (col %)				<0.001				<0.00
Trust in healthcare providers - No	171(27%)	125(34%)	285(53%)		263(29%)	191(46%)	127(62%)	
Trust in healthcare providers - Yes	460(73%)	248(66%)	248(47%)		659(71%)	220(54%)	77(38%)	
Perceived worry, as of December 2023 N (col %)				<0.001				<0.00
Perceived worry - No	255(39%)	197(51%)	310(55%)		378(39%)	234(54%)	150(69%)	
Perceived worry - Yes	400(61%)	192(49%)	257(45%)		583(61%)	200(46%)	66(31%)	

Multivariate Models

All adults:

Table 8 presents unadjusted and adjusted prevalence ratios (aPRs) for any agreement vs. some disagreement with COVID-19 vaccine statements during pregnancy in the cohort (N=4,488). Participants who reported trusting public health institutions (aPR=2.48, 95% CI: 2.17–2.83) and trusting healthcare providers (aPR=1.23, 95% CI: 1.14–1.33) were more likely to have any agreement about statements related to the safety of COVID-19 vaccine during pregnancy compared to those without trust in information sources. Conversely, symptoms of anxiety (aPR=1.00, 95% CI: 0.89–1.13) and depression (aPR=0.96, 95% CI: 0.87–1.07) were not significantly associated with COVID-19 vaccine statements during pregnancy. Non-Hispanic Black participants were 19% less likely than their non-Hispanic White counterparts to have any agreement with statements related to the vaccine's safety during pregnancy (aPR=0.81, 95% CI: 0.71–0.92).

Regarding associations with vaccine efficacy, trust in public health institutions (aPR=1.73, 95% CI: 1.59–1.87), trust in healthcare providers (aPR=1.13, 95% CI: 1.08–1.17), and having a personal doctor (aPR=1.05, 95% CI: 1.01–1.09) were statistically significant. Non-Hispanic Black participants were 9% less likely to have any agreement with statements related to

268   the vaccine's efficacy (aPR=0.91, 95% CI: 0.85-0.97) during pregnancy compared to their non-

269   Hispanic White counterparts.

Table 8. Prevalence Ratios and P-values for Any Agreement vs. Some Disagreement about COVID-19 Vaccine During Pregnancy in the CHASING COVID Cohort among all adult participants

	Safety perceptions (N = 3,121)				Efficacy perceptions (N = 3,098)			
	Prevalence Ratio	P-value	Adjusted Prevalence ratio[a]	P-value	Prevalence Ratio	P-value	Adjusted Prevalence ratio[a]	P-value
Age 18-29 vs 30-39	0.94 (0.87 - 1.02)	0.13	1.03 (0.96 - 1.12)	0.41	1.00 (0.96 - 1.05)	0.89	1.05 (1.01 - 1.09)	0.03
Age 40-49 vs 30-39	0.97 (0.89 - 1.06)	0.53	0.97 (0.89 - 1.05)	0.41	0.99 (0.95 - 1.04)	0.80	0.98 (0.94 - 1.02)	0.43
Age 50-64 vs 30-39	0.91 (0.82 - 1.01)	0.07	0.89 (0.81 - 0.97)	0.01	1.01 (0.96 - 1.06)	0.81	0.98 (0.94 - 1.02)	0.41
Age 65+ vs 30-39	0.96 (0.88 - 1.06)	0.45	0.98 (0.89 - 1.07)	0.61	0.99 (0.95 - 1.04)	0.72	0.97 (0.92 - 1.02)	0.26
Gender Male vs Female/Non-binary[b]	1.18 (1.11 - 1.25)	<0.001	1.11 (1.05 - 1.17)	<0.001	1.05 (1.02 - 1.08)	0.001	1.03 (1.00 - 1.06)	0.02
Race/Ethnicity Hispanic vs NH White	0.79 (0.72 - 0.86)	<0.001	0.94 (0.86 - 1.02)	0.13	0.88 (0.84 - 0.92)	<0.001	0.95 (0.90 - 0.99)	0.03
Race/Ethnicity NH Black vs NH White	0.67 (0.59 - 0.76)	<0.001	0.81 (0.71 - 0.92)	<0.001	0.81 (0.76 - 0.87)	<0.001	0.91 (0.85 - 0.97)	0.006
Race/Ethnicity NH Asian/PI/Other vs NH White	0.85 (0.77 - 0.95)	0.003	0.89 (0.81 - 0.99)	0.02	0.98 (0.94 - 1.03)	0.51	1.00 (0.97 - 1.05)	0.82
Income $50k-$100k vs <$50k/Unknown	1.15 (1.07 - 1.25)	0.003	NA		1.05 (1.01 - 1.09)	0.02	NA	
Income >$100k vs <$50k/Unknown	1.41 (1.32 - 1.52)	<0.001	NA		1.14 (1.10 - 1.18)	<0.001	NA	
Education <high school/high school vs College graduate	0.57 (0.50 - 0.65)	<0.001	0.73 (0.64 - 0.84)	<0.001	0.77 (0.72 - 0.83)	<0.001	0.89 (0.82 - 0.95)	<0.001
Education Some college vs College graduate	0.67 (0.62 - 0.73)	<0.001	0.82 (0.76 - 0.89)	<0.001	0.82 (0.78 - 0.85)	<0.001	0.91 (0.87 - 0.95)	<0.001
Anxiety status moderate/sev vs none to mild	0.81 (0.74 - 0.90)	<0.001	1.00 (0.89 - 1.13)	0.96	0.97 (0.93 - 1.02)	0.19	1.06 (1.01 - 1.12)	0.02
Depression status moderate/sev vs none to mild	0.83 (0.76 - 0.91)	<0.001	0.96 (0.87 - 1.07)	0.49	0.97 (0.93 - 1.01)	0.12	1.00 (0.95 - 1.05)	0.89
Any <18y kid in household Yes vs No	0.76 (0.71 - 0.82)	<0.001	NA		0.89 (0.85 - 0.92)	<0.001	NA	

Pregnancy Status Ever pregnant vs Never pregnant	0.98 (0.86 - 1.11)	0.73	NA		0.99 (0.92 - 1.06)	0.76	NA	
Personal doctor Yes vs No	1.13 (1.04 - 1.21)	0.002	1.00 (0.94 - 1.08)	0.90	1.10 (1.06 - 1.15)	<0.001	1.05 (1.01 - 1.09)	0.01
Susceptibility to severe COVID-19 (if infected) at baseline above vs below	0.88 (0.81 - 0.95)	0.002	0.94 (0.86 - 1.02)	0.14	0.97 (0.93 - 1.01)	0.13	1.00 (0.96 - 1.05)	0.96
Vaccination status Yes vs No	1.99 (1.34 - 2.94)	0.001	NA		1.65 (1.26 - 2.17)	<0.001	NA	
Boosted in 2023 Yes vs No	1.90 (1.79 - 2.01)	<0.001	NA		1.29 (1.25 - 1.33)	<0.001	NA	
Household members vaccinated Yes vs No	3.33 (2.70 - 4.10)	<0.001	NA		2.08 (1.83 - 2.37)	<0.001	NA	
Trust in public health institutions Yes vs No	2.83 (2.48 - 3.22)	<0.001	2.48 (2.17 - 2.83)	<0.001	1.85 (1.70 - 2.00)	<0.001	1.73 (1.59 - 1.87)	<0.001
Trust in healthcare providers Yes vs No	1.54 (1.42 - 1.67)	<0.001	1.23 (1.14 - 1.33)	<0.001	1.29 (1.24 - 1.36)	<0.001	1.13 (1.08 - 1.17)	<0.001
Perceived worry as of December 2023 Yes vs No	1.20 (1.13 - 1.28)	<0.001	NA		1.14 (1.11 - 1.17)	<0.001	NA	

[a]Fully adjusted model, including age, race/ethnicity, education, anxiety status, depression status, personal doctor, susceptibility to severe COVID-19, trust in public health institutions and trust in healthcare providers.
[b]We classify all gender minorities (N = 129) with women due to small numbers and in recognition of the increased health risks in this population.

Note: Statements corresponding to safety belief have been re-worded and reverse coded to ensure consistent interpretation across all statements.

Women of reproductive age

Table 9 presents unadjusted and adjusted prevalence ratios (aPRs) for any agreement vs.

some disagreement with COVID-19 vaccine statements during pregnancy among women of

reproductive age in the cohort (N=1,611). Participants who trusted healthcare providers

(aPR=1.34, 95% CI: 1.18–1.52) were 34% more likely to agree with statements related to the

vaccine's safety during pregnancy compared to those who did not trust healthcare providers.

Similarly, participants who trusted public health institutions (aPR=2.15, 95% CI: 1.76–2.63)

were more than twice as likely to agree with these statements compared to those who did not

trust public health institutions. However, while the relationship between having a personal

physician (aPR=1.03, 95% CI: 0.92–1.15) and perceptions about the vaccine's safety suggests a

positive association, it is not statistically significant at the 0.05 level. Women more susceptible to

severe COVID-19 if infected with SARS-CoV-2 were 25% less likely to have any agreement

with statements about the vaccine's safety during pregnancy than those less susceptible

(aPR=0.75, 95% CI: 0.59–0.95).

Table 9. Prevalence Ratios and P-values for Any Agreement vs. Some Disagreement about COVID-19 Vaccine During Pregnancy Among Women[b] of Reproductive Age in the CHASING COVID Cohort

	Safety perceptions (N = 1,222)				Efficacy perceptions (N = 1,177)			
	Prevalence Ratio	P-value	Adjusted Prevalence ratio[a]	P-value	Prevalence Ratio	P-value	Adjusted Prevalence ratio[a]	P-value
Age 18-29 vs 30-39	1.02 (0.91 - 1.15)	0.73	1.10 (0.98 - 1.23)	0.09	1.06 (0.99 - 1.13)	0.08	1.07 (1.01 - 1.14)	0.02
Age 40-49 vs 30-39	1.02 (0.90 - 1.17)	0.73	1.02 (0.90 - 1.15)	0.78	1.05 (0.98 - 1.13)	0.20	1.02 (0.95 - 1.08)	0.63
Race/Ethnicity Hispanic vs NH White	0.73 (0.63 - 0.85)	<0.001	0.93 (0.81 - 1.07)	0.31	0.84 (0.77 - 0.91)	<0.001	0.93 (0.86 - 1.00)	0.06
Race/Ethnicity NH Black vs NH White	0.63 (0.52 - 0.78)	<0.001	0.83 (0.67 - 1.03)	0.09	0.75 (0.66 - 0.85)	<0.001	0.85 (0.75 - 0.97)	0.01
Race/Ethnicity NH Asian/PI/Other vs NH White	0.82 (0.70 - 0.97)	0.02	0.84 (0.73 - 0.98)	0.02	0.98 (0.91 - 1.05)	0.60	0.97 (0.91 - 1.04)	0.38
Income $50k-$100k vs <$50k/Unknown	1.08 (0.94 - 1.24)	0.27	NA		1.12 (1.05 - 1.20)	0.001	NA	
Income >$100k vs <$50k/Unknown	1.58 (1.41 - 1.77)	<0.001	NA		1.22 (1.14 - 1.30)	<0.001	NA	
Education <high school/high school vs College graduate	0.47 (0.38 - 0.59)	<0.001	0.65 (0.51 - 0.81)	<0.001	0.71 (0.62 - 0.80)	<0.001	0.85 (0.75 - 0.95)	0.005
Education Some college vs College graduate	0.61 (0.53 - 0.70)	<0.001	0.76 (0.66 - 0.87)	<0.001	0.77 (0.72 - 0.84)	<0.001	0.86 (0.80 - 0.93)	<0.001
Anxiety status moderate/sev vs none to mild	0.77 (0.66 - 0.90)	0.001	0.97 (0.82 - 1.16)	0.78	0.96 (0.89 - 1.03)	0.25	1.07 (0.99 - 1.15)	0.08
Depression status moderate/sev vs none to mild	0.80 (0.70 - 0.92)	0.002	1.00 (0.85 - 1.17)	0.99	0.95 (0.89 - 1.02)	0.17	1.00 (0.93 - 1.08)	0.98
Any <18y kid in	0.69	<0.001	NA		0.85	<0.001	NA	

	PR (95% CI)	p	PR (95% CI)	p	PR (95% CI)	p	PR (95% CI)	p
household Yes vs No	(0.62 - 0.77)				(0.80 - 0.89)			
Pregnancy Status Ever pregnant vs Never pregnant	1.08 (0.94 - 1.24)	0.28	NA		1.02 (0.95 - 1.10)	0.52	NA	
Personal doctor Yes vs No	1.17 (1.03 - 1.32)	0.02	1.03 (0.92 - 1.15)	0.60	1.11 (1.04 - 1.19)	0.002	1.07 (1.00 - 1.13)	0.03
Susceptibility to severe COVID-19 (if infected) at baseline above vs below	0.61 (0.48 - 0.78)	<0.001	0.75 (0.59 - 0.95)	0.02	0.83 (0.73 - 0.95)	0.005	0.92 (0.81 - 1.04)	0.20
Vaccination status Yes vs No	2.62 (1.37 - 5.00)	0.004	NA		1.73 (1.15 - 2.60)	0.008	NA	
Boosted in 2023 Yes vs No	1.97 (1.80 - 2.17)	<0.001	NA		1.31 (1.25 - 1.37)	<0.001	NA	
Household members vaccinated Yes vs No	4.36 (3.12 - 6.09)	<0.001	NA		2.10 (1.76 - 2.51)	<0.001	NA	
Trust in public health institutions Yes vs No	2.60 (2.13 - 3.18)	<0.001	2.15 (1.76 - 2.63)	<0.001	1.77 (1.56 - 1.99)	<0.001	1.60 (1.42 - 1.81)	<0.001
Trust in healthcare providers Yes vs No	1.73 (1.52 - 1.97)	<0.001	1.34 (1.18 - 1.52)	<0.001	1.33 (1.23 - 1.43)	<0.001	1.13 (1.06 - 1.21)	<0.001
Perceived worry Yes vs No	1.35 (1.21 - 1.51)	<0.001	NA		1.25 (1.18 - 1.33)	<0.001	NA	

[a]Fully adjusted model, including age, race/ethnicity, education, anxiety status, depression status, personal doctor, susceptibility to severe COVID-19, trust in public health institutions and trust in healthcare providers.
[b]We classify all gender minorities
(N = 129) with women due to small numbers and in recognition of the increased health risks in this population.
Note: Statements corresponding to safety belief have been re-worded and reverse coded to ensure consistent interpretation across all statements.

285  Any agreement with statements related to the efficacy of the COVID-19 vaccine during

286 pregnancy was positively associated with trust in healthcare providers (aPR=1.13, 95% CI: 1.06–

287 1.21), trust in public health institutions (aPR=1.60, 95% CI: 1.42–1.81), and having a personal

288 doctor (aPR=1.07, 95% CI: 1.00–1.13).

289 **Any agreement vs Uncertainty model**

290  Among all adults (S1 Table), having a personal doctor and trust in healthcare providers

291 were significantly associated with agreement on the efficacy of the COVID-19 vaccine during

292 pregnancy. Trust in public health institutions was strongly associated with both safety and

293 efficacy perceptions, but it was the only significant predictor for safety-related statements.

Among women of reproductive age (S2 Table), having a personal doctor was significantly associated with agreement on both the safety and efficacy of the COVID-19 vaccine during pregnancy. While trust in public health institutions and healthcare providers was significant for efficacy perceptions, they were not significant predictors for safety perceptions in this subgroup.

4. Conclusion

This study revealed that less than half of adults (40%) perceived the COVID-19 vaccine as safe during pregnancy, with just over half considering it effective. Perceptions were influenced by demographics, with middle-aged, non-Hispanic White individuals, women, and those with a college education more likely to view the vaccine as safe and effective. Low uptake reasons often cited fears of maternal sickness or harm, insufficient research, and potential harm to the fetus[19], or broader concerns about side effects, infertility, and mortality[20]. Conversely, having vaccinated family/friends or prior Tdap/influenza vaccination was associated with less hesitancy[21].

A key finding was the strong association between trust and positive vaccine perceptions. Individuals who trusted healthcare providers were significantly more likely to regard the vaccine as safe (1.44 times) and effective (1.24 times) during pregnancy. Similarly, trust in public health institutions for vaccine information correlated with perceiving the vaccine as safe (almost three times more likely) and effective (almost two times more likely). These results underscore the vital role of healthcare providers and governmental risk communicators in building trust.

Limitations include a restricted number of pregnant individuals, preventing assumptions of causality between beliefs and actual vaccine uptake. Women of reproductive age were thus included to address unique immunization concerns during/after pregnancy as they may become

pregnant. Future research should further explore the relationship between vaccine beliefs and

actual vaccination behavior during pregnancy, and consider the influence of various healthcare

providers.

In conclusion, this study highlights the complex interplay of demographic factors,

healthcare experiences, and trust in shaping perceptions of COVID-19 vaccination during

pregnancy. These findings are consistent with prior literature that highlights the significant

influence of physician recommendations and trust in healthcare providers on vaccination

decisions during pregnancy [23,24], including both general providers and

obstetricians/gynecologists[25]. Given suboptimal vaccination rates, current strategies may be

insufficient. Efforts to improve vaccine uptake must prioritize fostering trust, both at the

individual level through provider-patient relationships and at the institutional level via

transparent, community-centered communication strategies.

Acknowledgments

Support for this project was provided by the CUNY Institute for Implementation Science

in Population Health and the CUNY Graduate School of Public Health and Health Policy.

Support for the CHASING COVID Cohort Study was provided by the National Institute

of Allergy and Infectious Diseases (NIAID), award number UH3AI133675 (MPIs: D Nash and C

Grov), NIMH award RF1MH132360 (MPIs: D Nash and A Parcesepe), National Institute of

Child Health and Human Development grant P2CHD050924 (Carolina Population Center),

Pfizer Inc., the CUNY Institute for Implementation Science in Population Health (cunyisph.org),

and the COVID-19 Grant Program of the CUNY Graduate School of Public Health and Health

Policy. The funders played no role in the production of this manuscript nor necessarily endorse

the findings.

The City University of New York (CUNY) School of Public Health received research

support from Pfizer during the conduct of this study that was paid directly to CUNY, with Dr.

Nash serving as the Principal Investigator. All authors declare no conflicts of interest related to

this work.

References

1. Jamieson DJ, Rasmussen SA. An update on COVID-19 and pregnancy. Am J Obstet Gynecol. 2022 Feb;226(2):177-186. doi: 10.1016/j.ajog.2021.08.054. Epub 2021 Sep 14. PMID: 34534497; PMCID: PMC8438995.

2. Wei SQ, Bilodeau-Bertrand M, Liu S, Auger N. The impact of COVID-19 on pregnancy outcomes: a systematic review and meta-analysis. CMAJ. 2021 Apr 19;193(16):E540-E548. doi: 10.1503/cmaj.202604. Epub 2021 Mar 19. PMID: 33741725; PMCID: PMC8084555.

3. Nunes MC, Madhi SA. COVID-19 vaccines in pregnancy. Trends Mol Med. 2022 Aug;28(8):662-680. doi: 10.1016/j.molmed.2022.04.012. Epub 2022 May 3. PMID: 35610123; PMCID: PMC9061132.

4. Kons KM, Wood ML, Peck LC, Hershberger SM, Kunselman AR, Stetter C, Legro RS, Deimling TA. Exclusion of Reproductive-aged Women in COVID-19 Vaccination and Clinical Trials. Womens Health Issues. 2022 Nov-Dec;32(6):557-563. doi: 10.1016/j.whi.2022.06.004. Epub 2022 Jun 15. PMID: 36075817; PMCID: PMC9197956.

5. Dagan, N., Barda, N., Biron-Shental, T. et al. Effectiveness of the BNT162b2 mRNA COVID-19 vaccine in pregnancy. Nat Med 27, 1693–1695 (2021). https://doi.org/10.1038/s41591-021-01490-8

6. Jorgensen SCJ, Drover SSM, Fell DB, et al. Newborn and Early Infant Outcomes Following Maternal COVID-19 Vaccination During Pregnancy. JAMA Pediatr. 2023;177(12):1314–1323. doi:10.1001/jamapediatrics.2023.4499

7. Cardemil CV, Cao Y, Posavad CM, Badell ML, Bunge K, Mulligan MJ, Parameswaran L, Olson-Chen C, Novak RM, Brady RC, DeFranco E, Gerber JS, Pasetti M, Shriver M, Coler R, Berube B, Suthar MS, Moreno A, Gao F, Richardson BA, Beigi R, Brown E, Neuzil KM, Munoz FM; MOMI-Vax Study Group. Maternal COVID-19 vaccination and prevention of symptomatic infection in infants. Pediatrics. 2024;153(3):e2023064252. doi:10.1542/peds.2023-064252

8. Centers for Disease Control and Prevention. COVID-19 vaccines while pregnant or breastfeeding. CDC. https://www.cdc.gov/covid/vaccines/pregnant-or-breastfeeding.html. Updated September 22, 2023. Accessed on September 4, 2024.

9. American College of Obstetricians and Gynecologists. COVID-19 vaccines and pregnancy: conversation guide for clinicians. ACOG. https://www.acog.org/covid-19/covid-19-vaccines-and-pregnancy-conversation-guide-for-clinicians. Published September 2023. Accessed September 4, 2024.

10. Centers for Disease Control and Prevention. COVID-19 vaccination dashboard. https://www.cdc.gov/covidvaxview/weekly-dashboard. Accessed July 25, 2024.

11. Rawal S, Tackett RL, Stone RH, Young HN. COVID-19 vaccination among pregnant people in the United States: a systematic review. Am J Obstet Gynecol MFM. 2022 Jul;4(4):100616. doi: 10.1016/j.ajogmf.2022.100616. Epub 2022 Mar 10. PMID: 35283351; PMCID: PMC8908633.

12. Patel A, Puglisi JL, Patel S, Tarn DM. COVID-19 Vaccine Acceptance in Pregnant Women in the United States: A Systematic Review and Meta-Analysis. J Womens Health (Larchmt). 2024;33(4):453-466. doi:10.1089/jwh.2023.0498

13. Huddleston HG, Jaswa EG, Lindquist KJ, Kaing A, Morris JR, Hariton E, Corley J, Hoskin E, Gaw SL, Cedars MI. COVID-19 vaccination patterns and attitudes among

American pregnant individuals. Am J Obstet Gynecol MFM. 2022 Jan;4(1):100507. doi: 10.1016/j.ajogmf.2021.100507. Epub 2021 Oct 11. PMID: 34649015; PMCID: PMC8502724.

14. Robertson MM, Kulkarni SG, Rane M, et al. Cohort profile: a national, community-based prospective cohort study of SARS-CoV-2 pandemic outcomes in the USA-the CHASING COVID Cohort study. BMJ Open. 2021;11(9):e048778. Published 2021 Sep 21. doi:10.1136/bmjopen-2021-048778

15. Robertson MM, Shamsunder MG, Brazier E, Mantravadi M, Zimba R, Rane MS, Westmoreland DA, Parcesepe AM, Maroko AR, Kulkarni SG, Grov C, Nash D. Racial/Ethnic Disparities in Exposure, Disease Susceptibility, and Clinical Outcomes during COVID-19 Pandemic in National Cohort of Adults, United States. Emerg Infect Dis. 2022 Nov;28(11):2171-2180. doi: 10.3201/eid2811.220072. Epub 2022 Oct 3. PMID: 36191624; PMCID: PMC9622253.

16. Spitzer, R. L., Kroenke, K., Williams, J. B. W., & Löwe, B. (2006). Generalized Anxiety Disorder 7 (GAD-7) [Database record]. APA PsycTests. https://doi.org/10.1037/t02591-000

17. Kroenke, K., & Spitzer, R. L. (2002). *Patient Health Questionnaire-8 (PHQ-8)* [Database record]. APA PsycTests. https://doi.org/10.1037/t64702-000

18. Zou G. A modified Poisson regression approach to prospective studies with binary data. Am J Epidemiol. 2004 Apr 1;159(7):702-706. doi: 10.1093/aje/kwh090.4o

19. Sutton D, D'Alton M, Zhang Y, Kahe K, Cepin A, Goffman D, Staniczenko A, Yates H, Burgansky A, Coletta J, Williams Z, Gyamfi-Bannerman C. COVID-19 vaccine acceptance among pregnant, breastfeeding, and nonpregnant reproductive-aged women. Am J Obstet Gynecol MFM. 2021 Sep;3(5):100403. doi: 10.1016/j.ajogmf.2021.100403. Epub 2021 May 25. PMID: 34048965; PMCID: PMC8146275.

20. Gianfredi V, Berti A, Stefanizzi P, D'Amico M, De Lorenzo V, Moscara L, Di Lorenzo A, Venerito V, Castaldi S. COVID-19 Vaccine Knowledge, Attitude, Acceptance and Hesitancy among Pregnancy and Breastfeeding: Systematic Review of Hospital-Based Studies. Vaccines (Basel). 2023 Nov 7;11(11):1697. doi: 10.3390/vaccines11111697. PMID: 38006029; PMCID: PMC10675759.

21. Kiefer MK, Mehl R, Costantine MM, Johnson A, Cohen J, Summerfield TL, Landon MB, Rood KM, Venkatesh KK. (2022). Characteristics and perceptions associated with COVID-19 vaccination hesitancy among pregnant and postpartum individuals: A cross-sectional study. BJOG: Int J Obstet Gy. 2022; 129: 1342–1351. https://doi.org/10.1111/1471-0528.17110

22. Schiavo R, Eyal G, Obregon R, Quinn SC, Riess H, Boston-Fisher N. The science of trust: future directions, research gaps, and implications for health and risk communication. J Commun Healthc. 2022;15(4):245-259. doi: 10.1080/17538068.2022.2121199.

23. Chervenak FA, McCullough LB, Grünebaum A. Reversing physician hesitancy to recommend COVID-19 vaccination for pregnant patients. Am J Obstet Gynecol. 2022 Jun;226(6):805-812. doi: 10.1016/j.ajog.2021.11.017. Epub 2021 Nov 8. PMID: 34762864; PMCID: PMC8572733.

24. Sienicka A, Pisula A, Pawlik KK, Dobrowolska-Redo A, Kacperczyk-Bartnik J, Romejko-Wolniewicz E. The Obstetrician's Role in Pregnant Women's Decision-Making Process Regarding Influenza and COVID-19 Vaccination. Vaccines

(Basel). 2023 Oct 18;11(10):1608. doi: 10.3390/vaccines11101608. PMID: 37897010;
PMCID: PMC10610802.
25. Miraglia del Giudice, Grazia, Giorgia Della Polla, Lucio Folcarelli, Annalisa Napoli,
Raffaella Punzo, Martina Peracchini, and Italo Francesco Angelillo. "Midwives'
Knowledge, Attitudes, and Practice Regarding COVID-19 Vaccination for Pregnant
Women: A Nationwide Web-Based Survey in Italy." Vaccines 11, no. 2
(January 19, 2023): 222. https://doi.org/10.3390/vaccines11020222. PMCID:
PMC9961665 PMID: 36851098

443
444

Supporting information

S1 Table. Prevalence Ratios and P-values for Any Agreement vs. Uncertainty about

COVID-19 Vaccine During Pregnancy in the CHASING COVID Cohort among all adult

participants

S2 Table. Prevalence Ratios and P-values for Any Agreement vs. Uncertainty about

COVID-19 Vaccine During Pregnancy in the CHASING COVID Cohort among all adult

participants