1

Co-Designing HPV Vaccine Programs with Girls and Caregivers: Insights from a Human-

Centered Design Approach

Eliza Fishman, MPH[1]*, Nicole Castle, MSc[1], Munyaradzi Joel Chinguwa, BSc[2], Amenze

Eguavoen, MBBS, MPH[3], Temitope Adebola Alfred, MPH[], Steven Ellis, MSc[], Yeni Indra,

BSc[], Paul Bassi Amos, MBBS, FMCPH (Nig.)[], Brittany Iskarpatyoti, MPH[]

1. JSI Research & Training Institute, Arlington, Virginia, United States of America

2. JSI Research & Training Institute, Harare, Zimbabwe

3. Lluvia Health Organization, Lagos, Nigeria

4. Independent Consultant, Abuja, Nigeria

5. Empatika, Jakarta, Indonesia

6. JSI Research & Training Institute, Abuja, Nigeria

7. JSI Research & Training Institute, North Carolina, United States of America

* Corresponding author

E-mail: eliza.fishman@jsi.org (EF)

Abstract

Cervical cancer is a significant global health issue, ranking as the fourth highest cause of cancer-related deaths among women globally. HPV vaccination for girls aged 9–14 offers a powerful tool for cervical cancer prevention, yet global coverage remains low, particularly in low- and middle-income countries. While research has documented parental and stakeholder perspectives, few studies center on girls' own experiences and how gender norms shape their access and decision-making. This study applied a human-centered design (HCD) approach in Nigeria and Indonesia to explore barriers to HPV vaccination and co-develop context-specific solutions with girls, caregivers, and immunization stakeholders.

Eight co-creation workshops were conducted with adolescent girls (ages 10–17) and caregivers in Nigeria (n=64) and Indonesia (n=66). Using HCD techniques—including persona development, journey mapping, and prototyping—participants identified barriers, enablers, and potential solutions. Stakeholder workshops in both countries validated and refined these community-generated ideas. Data were analyzed thematically using UNICEF's Journey to Health and Immunization framework.

Across both countries, girls and caregivers reported limited awareness and inconsistent information about HPV vaccination. Misinformation, unclear communication, and insufficient preparation were common. Gender norms strongly shaped decision-making: while mothers typically led discussions, fathers or male community leaders often held final authority, particularly in Nigeria. Access challenges included transportation costs, distance to facilities, missed opportunities during school-based delivery, stockouts, and limited guidance on managing adverse events, with out-of-school girls experiencing the greatest barriers.

Co-designed solutions emphasized improving communication through trusted messengers,

strengthening coordination between schools and health facilities, increasing engagement with

fathers and community leaders, and expanding flexible, community-based delivery models to

reach out-of-school girls. Findings underscore the importance of centering girls in HPV program

design and demonstrate the value of HCD in generating actionable, locally relevant strategies.

Strengthening communication, addressing gender norms, and improving equitable service

delivery are critical to increasing HPV vaccination uptake and advancing cervical cancer

prevention.

52   Introduction

Cervical cancer is a preventable and treatable disease, yet it ranks as the fourth highest cause of

cancer-related deaths among women globally, disproportionately affecting women in low- and

middle-income countries (LMICs) where access to primary and secondary prevention and

treatment are constrained (1,2). In 2023, only 27 percent of girls globally received a single dose

of HPV vaccine. There are large disparities in the distribution of national vaccination programs

and vaccination coverage rates between high-income countries (HICs) and LMICs, with 88

percent of HICs having included HPV vaccination in their national immunization program,

compared to 40 percent of LMICs (3,4).

There have been many challenges to introducing and increasing HPV vaccination coverage in

LMICs, including supply and logistical constraints, COVID-19 pandemic disruptions, high

vaccine costs, global and national funding for vaccination programs, vaccine prioritization, and

health system capacity (1,4,5). Parents cited specific barriers to HPV vaccinations, such as cost,

transport, and time constraints; low levels of knowledge about cervical cancer and HPV

vaccination; and negative experiences with other vaccinations (6–11). Vaccine access and

coverage are also challenged by misinformation, socioeconomic constraints, and gender

inequities within and across countries. It is well known that gender and social norms affect

decision-making, access, and uptake of HPV vaccination (12–15). These barriers are particularly

pronounced in efforts to reach out-of-school girls, ensure completion of the vaccine series,

accurately estimate target populations, monitor program performance, and sustain vaccination

efforts (1,6). However, new opportunities are emerging to address these challenges, including the

entry of additional vaccine manufacturers and the WHO recommendation for alternative single

dose scheduling (16,17).

Evidence suggests that social norms related to peer approval and trust in the government and

health professionals affect parental acceptance of and decision-making around HPV vaccination

for their daughters (11,18). In many LMICs, mothers, typically the primary caregivers, often

need to seek permission from their husbands to vaccinate their children. Women's daily

responsibilities, along with financial and mobility constraints, can further limit vaccine access

and use (7,8,19).

While implementation strategies for HPV vaccination and studies on parental perspective have

been documented, few studies explore the role of girls themselves in identifying key challenges

or shaping programs designed to boost vaccine uptake (20). One study on adolescent

involvement in HPV vaccine decision-making found that most had little to no influence, with

over half remaining silent during the decision phase. This lack of influence was linked to their

late or absent participation, underscoring the importance of engaging adolescents earlier in the

decision-making process (21). Less is known about the challenges girls face and their priorities,

as well as the social and gender-related factors that influence their uptake of the HPV vaccine.

These factors are often overlooked in the design and implementation of vaccination programs.

Engaging girls early in HPV program planning and decision-making may deepen our

understanding of the factors influencing vaccination and help design solutions that strengthen

and sustain high coverage.

Country Context: Nigeria

In Nigeria, cervical cancer is the second most common cancer among women and the second

most prevalent cause of cancer-related mortality among women aged 15–44 years (22). In 2022,

there were more than 13,600 new cases of cervical cancer and more than 7,000 deaths from the

disease (23). The Government of Nigeria introduced the HPV vaccine nationwide for girls aged

9–14 years in October 2023. The first phase of the introduction covered 16 states, and more than

4.8 million girls were vaccinated (24). In May 2024, the government implemented the second

phase of the introduction for the remaining 21 states. As of December 2024, more than 12

million eligible girls have been vaccinated, bringing vaccination coverage to 71 percent (25).

Following the campaign, Nigeria incorporated the HPV vaccine into its routine immunization

schedule. The vaccine is delivered through school- and health facility-based programs. While the

introduction campaign was successful, increasing vaccination coverage through the routine

immunization system has its own set of challenges. To ensure program sustainability in Nigeria,

a deeper understanding of the social and gender-related factors influencing HPV vaccine access

and uptake from girls' perspectives is important for designing tailored interventions that can

improve and sustain high vaccination coverage nationwide.

Country Context: Indonesia

Cervical cancer is the second most common cancer among women in Indonesia. In 2023, there

were almost 37,000 new cases of cervical cancer, and more than 20,000 women died from the

disease (26,27). In November of the same year, the president announced the National Plan for

Cervical Cancer Elimination in Indonesia (2023–2030) (28). The Indonesian government

introduced the HPV vaccine in Jakarta in 2016 to girls in school years five and six through a free,

school-based program. Since its introduction in Jakarta, the government has been scaling

vaccination nationwide through schools and health facilities; as of 2023, 132 municipalities are

delivering the program to in-school and out-of-school girls aged 11 and 12 years (9,29,30).

According to the WHO, HPV vaccination coverage reached 95.9 percent for the final dose in

2023. However, it is unclear if that data includes out-of-school girls (31). Reducing barriers to

vaccination and more effectively reaching out-of-school girls with solutions that center girls in

program planning will be important to maintaining high levels of coverage as the Indonesian

government expands the HPV vaccination program and advances its cervical cancer elimination

goals.

Study Objectives

Addressing the barriers girls and women face and increasing global HPV vaccination coverage

requires the development and implementation of tailored, multi-level vaccination programs that

meet the needs and preferences of girls and caregivers. This is true in both Indonesia and Nigeria

as they integrate the HPV vaccine into their national immunization schedules. This study

explores girls' and their caregivers' experiences, needs, and perspectives around improving HPV

vaccination programs in both countries. The primary objective of this study was to explore both

gender-related and non-gender-related barriers to the delivery and uptake of HPV immunization,

while identifying potential solutions to address these barriers, as proposed by girls and women

themselves and vaccine program stakeholders.

Materials and Methods

Study Design

This study employed a qualitative research design, using human-centered design (HCD)

approaches to capture the experiences and perspectives of research participants. HCD is an

iterative, solution-focused methodology that emphasizes understanding and prioritizing the needs,

challenges, and experiences of end users and others directly involved in a system—in this case,

girls, their caregivers, and vaccine program stakeholders (32,33). The study was approved by JSI

Institutional Review Board (IRB) under the IRB reference (#24-37). The JSI IRB is registered

with the Office of Human Research Protections (OHRP) as IRB00009069. JSI has a Federal-

wide Assurance (FWA) on file [FWA00000218] and is a recognized IRB organization

[IORG0007551]. Local ethics approval was received in both Indonesia and Nigeria. Written

consent was obtained by all study participants.

Study Population and Sample Selection

The study involved girls (ages 10–17) and their caregivers from four selected communities—two

in Nigeria and two in Indonesia. Communities were purposively selected to capture high- and

low-rates of out-of-school girls, urban/rural dispersion, and, where possible, religious diversity.

Participants included equal distribution of vaccinated and unvaccinated girls, along with their

caregivers. Caregivers were predominantly mothers; however, in both Indonesia and Nigeria,

there were a few instances where non-maternal caregivers—such as grandmothers or female

boarding school caretakers—were involved.

This purposive sampling approach enabled the collection of rich, relevant insights from those

with direct experience of or influence over the vaccination process, including vaccinated and

unvaccinated caregivers and girls, EPI managers, and representatives from implementing NGOs,

multilateral agencies, and local partners. Results were shared with and refined by local vaccine

program stakeholders, ensuring that the insights gained were actionable and relevant to program

implementation.

Sample Size

A total of 130 participants were included across eight workshops, with eight girls and their

female caregivers (16 participants) in each workshop; one workshop in Indonesia included nine

girls and their caregivers. Seventy stakeholders participated in the dissemination workshops held

in Nigeria and Indonesia—45 in Nigeria (across two sub-national workshops) and 25 in

Indonesia (in one national workshop). See Table 1 for sample characteristics.

Table 1: Sample Characteristics

Country	Nigeria				Indonesia	
States	**Benue**		**Adamawa**		**East Java**	**West Java**
Religion	Christianity		Islam		Islam	Islam
Study Site	Gwer-West	Makurdi	Fufore	Toungo	Bekasi City	Kediri Regency

Setting		Rural	Urban	Semi-urban	Rural	Urban		Rural	
School Status Sampled		In-school	Out-of-school	In-school	Out-of-school	In-school	Out-of-school	In-school	Out-of-school
Number of girls sampled	Vaccinated	4	4	4	4	5	5	4	3
	Unvaccinated	4	4	4	4	3	4	4	5
Number of caregivers sampled		8	8	8	8	8	9	8	8
Number of Stakeholders		20		25		25			

Data Collection and Analysis

Data were collected at each site through two-day co-creation workshops, where participants—

girls and caregivers—worked collaboratively to identify barriers to HPV vaccination and

generate locally feasible solutions. Co-creation workshops are one method that can be used as

part of the HCD process to conduct participatory activities to identify insights and generate

solutions (34–36). During the workshops, participants developed personas, a tool used to create

fictional characters that reflect user group characteristics and needs (32,37). Personas represented

girls and their caregivers from their community who either had or had not been vaccinated.

Participants explored girls' and caregivers' characteristics, influences, experiences with health

and HPV vaccination, values and aspirations, and gender and community norms. Following this,

journey maps visually documented each persona's experiences accessing and taking (or not) the

HPV vaccine. Journey maps are used to understand key steps an end user or other stakeholder

takes as they experience a product or service (32,38). UNICEF's Journey to Health and

Immunization was used as the journey map framework (39). It was developed to follow a

caregiver's journey to vaccinating a child while taking into consideration the health provider's

journey, and it has been used by other programs applying an HCD approach (35,36). It includes

six domains that represent steps on the journey: 1) knowledge and awareness, 2) intent, 3)

preparation, cost, and access, 4) point of service, 5) experience of care, and 6) after service (40).

The journey maps supported caregivers and girls to identify key barriers to HPV vaccination

based on their experiences at each step. After identifying key challenges, participants

brainstormed ideas for potential solutions and developed prototypes, a version of an idea that can

help communicate a concept and be tested (32). Following the community-level workshops, the

personas, journey maps, and solutions were summarized and shared with relevant immunization

stakeholders at the national (Indonesia) and sub-national (Nigeria) levels. A one-day workshop

(two state-level workshops in Nigeria) was conducted to review outputs from community

workshops, refine solutions, and develop action plans.

These interactive sessions served as the main method of data collection, capturing qualitative

insights and actionable recommendations. Data were analyzed thematically based on the domains

of the UNICEF Journey to Health and Immunization framework. The study lasted approximately

10 months, from June 2024 to April 2025, which included the planning and execution of

workshops, data collection, and analysis. Recruitment began and ended at different times at each

of the study sites. In Nigeria, data collection started in 01/09/2024 and ended in 28/02/2025. In

Indonesia, data collection started in 01/10/2024 and ended in 09/04/2025.

Results

The results are organized according to the domains of the UNICEF Journey to Health and Immunization framework, exploring the barriers and enablers girls and their caregivers encountered across each stage of the vaccination journey and follows with the solutions proposed by participants and reflected upon by stakeholders to address these challenges.

Knowledge and awareness

Overall, in both countries and regardless of vaccination status, girls and their caregivers had varying but generally limited awareness and understanding of HPV vaccination. The ways they received information differed by country and vaccination status, but a similar pattern emerged: some unvaccinated caregivers in both Indonesia and Nigeria first heard about the vaccine through neighbors or social media.

Indonesia

In Indonesia, vaccinated girls reported receiving information primarily from their mothers. Most mothers received their information from formal school communications delivered through letters, WhatsApp messages, or community sessions. While these communication mechanisms afforded more systematic outreach, participants expressed that information frequently arrived late or lacked sufficient detail to meet caregivers' needs or support them to prepare their daughters for vaccination. Caregivers of vaccinated girls in Indonesia specifically ranked social media as the least trusted platform for HPV information, as it causes confusion and misunderstanding.

Nigeria

In contrast, girls in Nigeria accessed information from a wider array of sources, including peers, radio, town announcers, and health workers. Many out-of-school girls reported being exposed to a variety of informal sources that often conveyed conflicting or inconsistent information. Caregivers reported a similar pattern, drawing on diverse informal channels such as friends, daughters, religious leaders, and radio broadcasts. While this broad network enabled widespread dissemination, the compilation of information shared was described as often insufficient or unclear, limiting individuals' ability to make informed decisions. As a result, caregivers' access to accurate information varied widely, with some unaware of vaccination campaigns until after they had concluded. Girls also noted that women generally do not attend mosque and therefore miss out on important health messages that men receive there, as the messages are not further relayed.

Intent and decision-making

Across both countries, female caregivers (often mothers) consistently confirmed their active role in decisions about HPV vaccination. Mothers were viewed as the primary caregiver responsible for their daughters' health, and they typically guided vaccination choices. However, girls in both settings often had limited visibility into how decisions were made within the household, reflecting broader norms around obedience, modesty, and deference to parental authority.

Indonesia

In Indonesia, mothers overwhelmingly described themselves as the main decision-makers regarding HPV vaccination, noting that fathers usually deferred to them. Girls also tended to rely almost exclusively on their mothers for guidance, with little mention of fathers' involvement.

Despite their central role, many Indonesian mothers reported feeling under-informed and pressed for time when making vaccination decisions. They expressed concerns about misinformation and anti-vaccine narratives circulating in their communities and, in some cases, wished fathers were more engaged in discussions related to health-related decision-making processes.

Nigeria

In Nigeria, mothers were also responsible for overseeing vaccination, but decision making was more collaborative than in Indonesia. Fathers and extended family members—particularly men—played a significant role, especially for unvaccinated girls. Girls noted that male approval was often required, and mothers frequently consulted husbands, relatives, community members, and religious leaders before deciding. Among caregivers of vaccinated girls, most sought information from health professionals and supported the vaccine, though several expressed frustrations with unclear or absent consent procedures. For unvaccinated girls, the need for male approval and financial support emerged as a key barrier, and women who made decisions independently were sometimes perceived as "rebellious" or faced social backlash.

Preparation, cost, and access

Across Indonesia and Nigeria, both vaccinated and unvaccinated girls reported preparation, cost, and access as consistent barriers to HPV vaccination. Girls in both contexts described general challenges in accessing health services, including long distances to facilities, transportation-related costs, and limited availability of health workers. Caregivers in both countries echoed these issues, noting that insufficient information, unclear guidance, and logistical hurdles made it difficult for girls, particularly those out of school, to obtain the vaccine.

Indonesia

In Indonesia, unvaccinated girls showed strong interest in receiving the HPV vaccine, partly because some believed it was free, though they were often unsure about the actual cost. Their primary barrier was a lack of information on how to access the vaccine—such as cost, eligibility requirements, or necessary documentation. While parental permission was a concern, it was secondary; some out-of-school girls even noted they might be able to persuade their parents to agree. For in-school girls, a major concern was not knowing where to go if they missed school-based vaccination sessions. Caregivers of in-school girls wanted greater involvement in the school vaccination process, citing insufficient notice and unclear instructions. Caregivers of out-of-school girls highlighted potential administrative barriers, such as the need for ID cards and interactions with unwelcoming facility staff.

Nigeria

In Nigeria, girls—especially the vaccinated ones—identified barriers their peers might face, including long distances to schools where vaccination occurred and social pressures, such as losing friends who opposed the vaccine. Unvaccinated out-of-school girls emphasized two main obstacles: the distance to health facilities and the limited availability of parents, particularly mothers, to accompany them due to work, chores, fatigue, or travel. Some girls were also discouraged by peers and family members, including older sisters. For caregivers of both vaccinated and unvaccinated girls, transportation costs were a dominant barrier. Caregivers of vaccinated girls often accompanied their daughters but worried about lost income or business. Those whose daughters missed the in-school campaign were directed to health facilities but

284 encountered vague instructions, long travel distances, and inconsistent vaccine availability,

285 frequently arriving to find the vaccine unavailable or the campaign already over.

Point of service & Experience of Care

Across both Indonesia and Nigeria, girls generally viewed health workers positively, though their

experiences varied widely depending on the setting and the individual provider. In both countries,

girls and caregivers described logistical and communication challenges that affected the

vaccination experience, such as limited information about the vaccine, insufficient guidance on

managing potential adverse events, and procedural gaps during vaccination sessions. Participants

in both countries also faced system constraints, including long wait times, crowding, and

inconsistent availability of vaccines, which created barriers for girls and caregivers. For out-of-

school girls in both Indonesia and Nigeria, accessing services at health facilities often involved

delays, confusion, or stockouts, making the process more difficult than for in-school girls.

Indonesia

In Indonesia, vaccination primarily occurred within school settings, which shaped most girls'

experiences. Girls reported a mix of positive and less positive interactions with health workers,

but several noted feeling shy, uncomfortable, or excluded, especially those choosing not to get

vaccinated. Some girls were separated from their peers during vaccination activities, which

created confusion and social discomfort. Communication gaps were pronounced—many girls felt

unable to ask questions or seek clarification during the session. The school-based model also

limited access for out-of-school girls, who had to visit health facilities instead. These girls

described long waits of up to an hour before receiving the vaccine. Caregivers in Indonesia

generally understood the steps involved in school-based vaccination—registration, checks, and

instructions for adverse event management—but felt that teachers and parents needed to provide

more support. Some caregivers said teachers were unprepared to answer questions, and

caregivers of unvaccinated girls who opted out reported being treated poorly during the process.

Nigeria

In Nigeria, girls accessed vaccination both in schools and at health facilities, giving rise to more

diverse experiences. While health workers were generally perceived positively, systemic issues,

such as wait times and vaccine shortages, and poor crowd control were common. Some girls

were unable to get vaccinated due to stockouts at health facilities while other in-school girls ran

off during the campaign out of fear. Caregivers in Nigeria described generally positive

vaccination experiences regardless of the delivery point. However, they noted that neither girls

nor caregivers were adequately informed about potential adverse events or how to manage them.

Echoing the girls' accounts, caregivers highlighted logistical frustrations—including long delays,

disrupted class schedules (for in-school girls), and the challenge of traveling to health facilities

where vaccines were sometimes unavailable or health workers were unwelcoming (for out-of-

school or unvaccinated girls).

Post-vaccination experience

Post-vaccination experiences were shared only by vaccinated girls, as unvaccinated girls had

little to report. Across both Indonesia and Nigeria, post-vaccination feedback from caregivers

was generally positive, with caregivers pleased that their daughters were vaccinated and

reporting few side effects. In both countries, neither girls nor caregivers described any severe

adverse reactions, and overall satisfaction with having completed vaccination was high.

Indonesia

328 In Indonesia, vaccinated girls reported minor side effects and noted a lack of guidance on post-

329 vaccination care. Some caregivers, particularly in Kediri, shared perceptions that differed from

330 the girls' experiences; they were pleased with the vaccination outcome, while girls expressed

331 frustration about returning to schoolwork immediately after vaccination and a lack of guidance

332 on post-vaccination care.

Nigeria

334 In Nigeria, vaccinated girls generally expressed happiness and enthusiasm, often encouraging

335 peers to get vaccinated. They reported no significant barriers or negative post-vaccination

336 experiences. Caregivers in Nigeria did raise some concerns about side effects and limited

337 information on post-vaccination care, but, like Indonesia, no severe adverse reactions were

338 reported.

Proposed Solutions

Girls and caregivers developed solutions to address the challenges they had identified. State- and

national-level stakeholders were engaged to reflect on the findings, share their own experiences,

and collaboratively identify and prioritize actionable solutions. Each group's solutions are

summarized in Table 2 and Table 3.

Table 2: Summary of Solutions Developed by Girls, Caregivers, and Stakeholders in

Nigeria

Subtheme	Girls	Caregivers	Stakeholders
Communication/Awareness	· School programs: Information should be delivered by the teachers or health educators during free periods and through peer-to-peer education to share information on vaccine safety, side effects, and benefits. Include motivators/incentives like provision of school supplies, etc. · Community awareness: Messages should be shared through town announcers in the mornings and evenings. Comprehensive health education should be delivered by qualified health care personnel to ensure credibility and engagement.	· Use multiple communication channels including town announcers, community leaders, radio, tv, teachers, school authorities, religious houses, traditional leaders, women groups, Facebook, etc. to inform the public about the vaccines; if one channel doesn't serve them, another will. ·Vaccine champions should be identified in the communities to share their experiences and raise awareness about the vaccine in the community.	· Use multi-channel communication strategies, including town announcers, community leaders, and structured outreach programs to enhance knowledge and acceptance.
Coordination/Outreach	· School peer to peer education: train in-school girls to train or educate other out-of-school girls.	· Men should be engaged to obtain their support and buy-in as influencers and decision makers.	· Closer collaboration with local security forces and community networks can help ensure safe delivery. · Increase targeted engagement with men.
Service Delivery	· Out-of-school peer-to-peer education enhances confidence and decision-making skills as well as increases girls'	· House-to-house vaccination should be considered because of its potential to reach	

Subtheme			
	happiness. Peers can refer or accompany others to the health facility for vaccine uptake.	both in school and out-of-school girls.	
System Strengthening/Supply Chain		· Improve school vaccination program by including proper planning to prevent stock-outs, more vaccinators, class-to-class vaccination instead of an assembly of all eligible girls, lesson free vaccination days where possible, a system to identify missed opportunities, and bidirectional communication with health care workers.	· Improve supply chain management and real-time tracking system.

Table 3: Summary of Solutions Developed by Girls, Caregivers, and Stakeholders in

Indonesia

Subtheme	Girls	Caregivers	Stakeholders
Communication/Awareness	· Display a poster with information on how to search for trusted information on HPV and HPV vaccination, including how to find information about HPV vaccination on TikTok.

· During in-school vaccination, use storytelling to pass the time and provide information about health and the vaccine for those waiting to get | ·Organize educational activities at schools in coordination with puskesmas (public health clinics) to equip girls with information about HPV vaccination.

· Provide one-page leaflets from doctors/nurses for girls to show their parents.

· Provide socialization at schools two days before vaccination day to | · Develop engaging educational materials on HPV and the vaccine, including videos and interactive storybooks.

· Increase strategic collaboration with religious leaders to disseminate accurate information and address faith-based concerns. |

	vaccinated. · Provide a fun, informative video before vaccination.	provide more information about HPV (with video and printed materials). · Organize FAQ meetings for parents led by the health department.	· Create a comprehensive FAQ resource on HPV vaccination.
Service Delivery	· Provide education sessions at schools for girls and their parents that are facilitated by teachers and health workers. · Organize outreach led by the puskesmas in coordination with the community and pesantren (Islamic boarding schools).	· Organize community socialization about HPV vaccination in communities residing around landfills. · Teachers should include parents in WhatsApp groups to inform them of the vaccination process (at least three days before). · Establish a coordination and collaboration process for schools and health centers, including a coordination system connecting the puskesmas with parents/caregivers.	
System Strengthening/Supply Chain			· Implement standardized, child-friendly service delivery protocols.

Girls in both countries suggested solutions centered on improving awareness and knowledge. In

Indonesia, proposed strategies often focused on improving individual knowledge—such as

making information clearer and more accessible for girls—but also reflected group-oriented

elements, including peer support among girls and a desire among caregivers for community-

based information sharing and collaboration with schools. Nigerian girls, on the other hand,

predominantly recommended community-level interventions involving health workers,

community leaders, and peers to improve vaccine uptake. Across both countries, girls

emphasized the importance of timely, consistent information and improved access to vaccination

services. Additional solutions focused on enhancing the vaccination experience and improving

access.

Caregivers in Nigeria focused their suggestions on community-driven solutions, including using

local leaders, town announcers, media campaigns, and home visits to raise awareness and combat

misinformation. Incentives and logistical support (e.g., transportation vouchers or providing food

during information sessions) were also recommended to improve access. Indonesian caregivers

for in-school girls emphasized school-centered approaches and saw girls themselves as effective

messengers in convincing parents about the value of vaccination. They also highlighted the need

for better coordination among schools, teachers, and health workers, as well as improved

communication to prepare families for vaccination. These proposed solutions closely aligned

with those expressed by the girls, reinforcing the shared priorities of improved knowledge,

support, and access.

Similarly, stakeholders in both Nigeria and Indonesia noted low public awareness and

widespread misinformation as a critical challenge to their programming. In Nigeria, stakeholders

emphasized the importance of multi-channel communication strategies, proposing the use of

town announcers, community leaders, and structured outreach programs to enhance knowledge

and acceptance. In Indonesia, communication between schools and parents is channeled through

schools, but there is a lack of clear processes and standardized guidance on how schools should

engage with caregivers, including what information to share, when to share it, and in what format.

To address this, stakeholders proposed the development of educational and informational

materials for both girls and their caregivers.

Stakeholders in both countries also focused heavily on logistical and programmatic operation

challenges. Operational hurdles in Nigeria—such as vaccine stock-outs, delays in distribution,

and incomplete data reporting—were seen as major impediments to effective rollout.

Stakeholders underscored the need for improved supply chain management and real-time

tracking systems. They also advocated for closer collaboration with local security forces and

community networks to ensure safe delivery in conflict-affected regions. In Indonesia, access-

related issues were compounded by the lack of engagement with marginalized groups and data

collection gaps, particularly for unregistered pesantren (Islamic boarding schools). At the point

of service delivery, stakeholders noted the unclear delineation of responsibilities between schools

and health services and weak cross-sectoral coordination.

Stakeholders in both Nigeria and Indonesia identified that greater engagement is needed with

men, as fathers, decision makers, and religious and community leaders. However, stakeholders

also stressed the importance of promoting gender inclusivity within HPV vaccination programs

to ensure that strategies are responsive to the needs of girls and their caregivers.

Discussion and recommendations

Gender Norms and Decision-Making

The findings underscore the significant influence of gender norms on decision-making dynamics

and access to HPV vaccination in both Indonesia and Nigeria. Immunization programs in LMICs

have traditionally focused on women as the primary caregivers for children's health (8,41).

However, existing literature also emphasizes men's dominant role as household decision-makers

(42–47). This study found this duality plays an important role in overall acceptance of the

vaccine. While caregivers and girls in both contexts frequently identified mothers as the primary

decision-makers for their daughters' health, men may hold the final say and power in

operationalizing that decision. In Indonesia, several mothers expressed a desire for greater

involvement from fathers, reflecting a perceived gap in shared responsibility. In contrast, in

Nigeria, fathers and male community or religious leaders often play a more overt and

authoritative role in decisions concerning girls' health. Some Nigerian mothers described

scenarios in which men held the final authority over health matters, and women who acted

independently risked social criticism—highlighting the persistent influence of traditional gender

expectations. These findings illustrate the complex and context-specific nature of gender norms,

particularly as they relate to intra-household and health-related power dynamics (48,49). This

study, in which girls and women voiced a desire for greater involvement from fathers and male

figures, highlights the importance of addressing gender norms and underscores the need to

actively engage male decision-makers in HPV program planning. Recent studies have

increasingly recognized that men play influential roles as decision-makers, household heads, and

community leaders—and that their support can significantly influence vaccine uptake (50–53).

Developing targeted outreach strategies to engage men as supportive partners in their daughters'

health and HPV vaccination efforts is essential (54). In Indonesia, this could include father-

focused education sessions at schools and community events. In Nigeria, outreach through

mosques, churches, and traditional gatherings may be more effective in increasing fathers'

awareness and support for vaccination.

Gender norms and expectations shape not only girls' and their caregivers' attitudes and access to

the vaccine but can influence the effectiveness of vaccination campaigns themselves. This study

highlights how social stigma and gender norms in Indonesia discourage girls from actively

seeking information. In Nigeria, concerns about vaccine side effects and persistent myths about

infertility remain major barriers, especially among unvaccinated girls and their caregivers. To

effectively address these gender-related challenges, vaccination campaigns should consider the

unique ways in which both girls and caregivers navigate health systems (55). For instance,

stigma in Indonesia and fears of infertility in Nigeria require context-specific, culturally relevant

messaging. Approaches like HCD offer value by engaging girls and their caregivers directly,

ensuring that solutions are rooted in local realities and shaped by the voices of those most

impacted (36). For HPV program planning specifically, HCD approaches can potentially

improve uptake and sustainability (35,37,56).

432 Limited Awareness and Information Gaps

Our findings reveal that both adolescent girls and their caregivers lack sufficient, consistent, and

accurate information about HPV vaccination. Misinformation remains a significant barrier to

uptake. While this information gap has been well-documented from the perspectives of parents

and stakeholders, it has rarely been examined from the perspective of girls themselves (6–

8,10,11,18,57). Our study makes clear that there is a widespread desire for more information;

however, the content, messenger, and mode of delivery must be better aligned with the needs of

the end users—namely, adolescent girls and their caregivers. Participants across both countries

emphasized this point, with all proposing solutions centered on improving knowledge and

awareness of the HPV vaccine. This underscores both the urgency of the issue and the need for a

shift in communication approaches. In Nigeria, participants strongly recommended peer-to-peer

education as an effective way to reach girls with accurate, relatable information. In Indonesia,

participants stressed the need for more structured and engaging content, along with better

training for teachers to deliver key messages effectively, as teachers were considered the most

trusted source. The literature highlights that strengthening communication strategies to address

mothers' perceptions of HPV can significantly improve vaccine uptake among both mothers and

their daughters (58–60). Furthermore, enhancing communication about the HPV vaccine may

empower young people to take a more active role in their health care decisions, ultimately

contributing to higher vaccination coverage. These insights highlight the critical value of

involving end users in the planning and design of HPV communication strategies (37).

Ultimately, the challenge is not only about what messages are delivered but also who delivers

them and how (61,62). Our findings suggest that current messengers are not always reaching or

resonating with their intended audiences. Greater alignment between messenger and audience is

needed to ensure communication efforts are trusted, accessible, and impactful. Specifically,

strengthening vaccine education using trusted messengers (such as teachers, community leaders,

and health workers) who deliver accurate, culturally appropriate information tailored to diverse

audience needs is strongly recommended (54,56).

Community Influence and Social Norms

This study highlights important cross-country differences in how social norms and decision-

making structures influence HPV vaccination uptake in Indonesia and Nigeria. These findings

underscore the importance of tailoring vaccination strategies to the local sociocultural context

(63–65). In Indonesia, decision-making was primarily individualized and centered within the

household. Mothers and daughters were often the primary decision-makers, and concerns about

vaccination reflected personal or family-level considerations. This aligns with a systematic

review on health-seeking behavior in Indonesia, which found that individualized circumstances

were more influential than social factors (66). This suggests that in Indonesia, program strategies

should focus on empowering households—particularly mothers—as key agents of change.

Campaigns may be most effective when they emphasize the benefits of HPV vaccination for a

girl's health, future well-being, and academic or career success, using messaging that resonates at

the family level.

In contrast, in Nigeria, decision-making is more collective and embedded within broader

community dynamics. Religious and traditional leaders, along with peer networks, serve as

influential gatekeepers. Drawing on the socio-ecological model, health choices are not simply

personal— they are communal decisions, shaped by shared beliefs and group trust (67–69). Thus,

HPV programs in Nigeria should prioritize community-based approaches, leveraging respected

community figures to champion vaccination, combat misinformation, and foster a culture of

acceptance. Establishing peer-to-peer networks can also help reinforce accurate knowledge and

increase collective confidence in the vaccine (70,71).

These contrasts highlight a broader implication: understanding the social ecosystem—who

makes decisions, who influences them, and what values they are anchored in—is essential for

developing effective, context-sensitive strategies that can lead to higher and more equitable HPV

vaccination coverage. In more family-oriented settings, interventions should speak directly to the

family unit; in communal settings, they must ripple outward through social structures and trusted

voices. Recognizing who the reference groups are in each setting—whether it's a mother, a

religious leader, or a peer—allows programs to align their approaches and messages with local

values and decision-making pathways.

488 Service Delivery: In-School vs. Out-of-School Girls

Effective HPV vaccination requires not only demand generation but also reliable and equitable

service delivery. Our findings point to important differences in how service delivery challenges

affect vaccine uptake in Indonesia and Nigeria, with distinct implications for reaching both in-

school and out-of-school girls.

In Indonesia, the school-based model has generally provided a consistent platform for vaccine

delivery. However, this approach leaves significant gaps for out-of-school girls, who often face

informational rather than logistical barriers (72). Moreover, the current model does not

adequately create a welcoming environment or recognize that girls may have questions and need

support during the vaccination process. Opportunities exist to improve the school experience by

reducing wait times and fostering support through storytelling activities and more engaging

interactions with teachers and health workers.

Service delivery challenges in Nigeria are more systemic, multifaceted, and align with the

existing literature on HPV barriers (73). Even when girls and caregivers were informed and

motivated to seek vaccination, they frequently encountered structural barriers: long wait times,

vaccine stockouts, transportation difficulties, and insufficient infrastructure all limited their

ability to access vaccines (1,73–75). These barriers particularly affect out-of-school girls, for

whom health services may be even more out of reach due to geographic, financial, or social

constraints (72).

These findings carry several important implications for program design. First, school-based

vaccination alone is insufficient to achieve equitable coverage (76). Out-of-school girls—who

are often among the most vulnerable—require alternative and flexible delivery mechanisms.

Mobile clinics, house-to-house outreach, and partnerships with community health centers and

religious institutions can help bridge this gap (72,76). In Indonesia, mobile outreach combined

with trusted local institutions may extend the reach of services beyond the classroom. In Nigeria,

these approaches are even more critical, given the breadth of logistical challenges that impede

access.

Second, there is an urgent need to strengthen health system capacity and planning. In Nigeria,

this includes improving supply chain management, investing in health infrastructure, and training

frontline workers to deliver vaccines more efficiently and sensitively. In Indonesia,

improvements in communication, standardization of procedures, and support and training for

health workers and teachers can improve the quality and consistency of school-based delivery.

Ultimately, the needs of in-school and out-of-school girls differ not just in terms of access points,

but also in the types of support they require. A flexible, equity-focused service delivery model

that recognizes these differences is essential to improving HPV vaccination coverage and

ensuring no girl is left behind.

Conclusion

This study is one of the few that places girls at the forefront of HPV vaccine program planning,

highlighting the value of designing interventions around the needs and lived experiences of the

end user. By centering girls, HPV vaccine programs can better tailor strategies that not only

improve uptake but also enhance long-term sustainability. Importantly, the findings point to the

potential of peer-to-peer education as a powerful strategy, leveraging girls' own social networks

to increase trust, correct misinformation, and strengthen vaccine confidence.

The findings underscore the critical need for context-specific, gender-responsive approaches to

increase HPV vaccination in both Indonesia and Nigeria. In both settings, persistent

misinformation and limited awareness call for targeted communication strategies delivered by

trusted messengers. Reaching out-of-school girls also demands flexible, equity-focused service

delivery models that extend beyond school-based platforms.

Ultimately, a one-size-fits-all approach is inadequate—HPV programs must be grounded in local

decision-making structures, gender norms, and access barriers to ensure broader and more

equitable vaccine coverage.

Acknowledgement

We want to acknowledge all the data collectors in both Indonesia and Nigeria who made the

collection of this research possible. This material was produced by the HPV Vaccine

Acceleration Program Partners Initiative (HAPPI) Consortium and may be used freely. The

HAPPI Consortium, in collaboration with our esteemed partners, is managed by JSI together

with Clinton Health Access Initiative (CHAI), the International Vaccine Access Center (IVAC)

at the Johns Hopkins Bloomberg School of Public Health, Jhpiego, and PATH. This research and

the HAPPI consortium are funded by the Gates Foundation under grant number INV-057603.

References

1.   Guillaume D, Waheed D e N, Schleiff M, Muralidharan KK, Vorsters A, Limaye RJ.
Global perspectives of determinants influencing HPV vaccine introduction and scale-up in
low- and middle-income countries. Thomas LF, editor. PLOS ONE. 2024 Jan
16;19(1):e0291990.

2.   WHO. Cervical Cancer. 2024 [cited 2025 July 14]. Cervical cancer. Available from:
https://www.who.int/news-room/fact-sheets/detail/cervical-cancer

3.   Giannone G, Giuliano AR, Bandini M, Marandino L, Raggi D, Earle W, et al. HPV
vaccination and HPV-related malignancies: impact, strategies and optimizations toward global
immunization coverage. Cancer Treat Rev. 2022 Dec;111:102467.

4.   Malvolti S, Soble A, Bloem P, LaMontagne DS, Aggarwal R, Pitisuttithum P, et al. The
Global Demand and Supply Balance of the Human Papillomavirus Vaccine: Implications for
the Global Strategy for the Elimination of Cervical Cancer. Vaccines. 2023 Dec 19;12(1):4.

5.   Branda F, Pavia G, Ciccozzi A, Quirino A, Marascio N, Gigliotti S, et al. Human
Papillomavirus (HPV) Vaccination: Progress, Challenges, and Future Directions in Global
Immunization Strategies. Vaccines. 2024 Nov 19;12(11):1293.

6.   Marshall S, Fleming A, Sahm LJ, Moore AC. Identifying intervention strategies to
improve HPV vaccine decision-making using behaviour change theory. Vaccine. 2023
Feb;41(7):1368–77.

7.   Agha S, Bernard D, Francis S, Fareed A, Nsofor I. Determinants of Human
Papillomavirus Vaccine Acceptance among Caregivers in Nigeria: A Fogg Behavior Model-
Based Approach. Vaccines. 2024 Jan 13;12(1):84.

8.   Balogun FM, Omotade OO. Parental intention to vaccinate adolescents with HPV
vaccine in selected communities in Ibadan, Southwest Nigeria: an application of Integrated
Behavioral Model. Hum Vaccines Immunother [Internet]. 2022 Nov 30 [cited 2025 July
9];18(5). Available from:
https://www.tandfonline.com/doi/full/10.1080/21645515.2022.2069959

9.   Wijayanti KE, Schütze H, MacPhail C. Parents' attitudes, beliefs and uptake of the
school-based human papillomavirus (HPV) vaccination program in Jakarta, Indonesia – A
quantitative study. Prev Med Rep. 2021 Dec;24:101651.

10.   Endarti D, Satibi -, Kristina SA, Farida MA, Rahmawanti Y, Andriani T. Knowledge,
Perception, and Acceptance of HPV Vaccination and Screening for Cervical Cancer among
Women in Yogyakarta Province, Indonesia. Asian Pac J Cancer Prev [Internet]. 2018 Apr
[cited 2025 July 9];19(4). Available from: https://doi.org/10.22034/APJCP.2018.19.4.1105

11.   Wijayanti K, Schutze H, Mac Phail C, Ivers R. Exploring Parents' Decisions Regarding
HPV Vaccination for Their Daughters in Jakarta, Indonesia: A Qualitative Study. Asian Pac J
Cancer Prev. 2023 Nov 1;24(11):3993–8.

12.   Alarcão V, Zdravkova B. Attitudes and Practices towards HPV Vaccination and Its
Social Processes in Europe: An Equity-Focused Scoping Review. Societies. 2022 Sept
18;12(5):131.

13.   Wong LP, Wong PF, Megat Hashim MMAA, Han L, Lin Y, Hu Z, et al.
588   Multidimensional social and cultural norms influencing HPV vaccine hesitancy in Asia. Hum
589   Vaccines Immunother. 2020 July 2;16(7):1611–22.

590   14.   Sundaram N, Voo TC, Tam CC. Adolescent HPV vaccination: empowerment, equity and
591   ethics. Hum Vaccines Immunother. 2020 Aug 2;16(8):1835–40.

592   15.   Sundstrom B, Smith E, Delay C, Luque JS, Davila C, Feder B, et al. A reproductive
593   justice approach to understanding women's experiences with HPV and cervical cancer
594   prevention. Soc Sci Med. 2019 July;232:289–97.

595   16.   Tsu VD, LaMontagne DS, Atuhebwe P, Bloem PN, Ndiaye C. National implementation
596   of HPV vaccination programs in low-resource countries: Lessons, challenges, and future
597   prospects. Prev Med. 2021 Mar;144:106335.

598   17.   WHO. World Health Organization. 2022 [cited 2025 Aug 5]. WHO updates
599   recommendations on HPV vaccination schedule. Available from:
600   https://www.who.int/news/item/20-12-2022-WHO-updates-recommendations-on-HPV-
601   vaccination-schedule

602   18.   Wijayanti KE, Schütze H, MacPhail C. Parents' attitudes, beliefs and uptake of the
603   school-based human papillomavirus (HPV) vaccination program in Jakarta, Indonesia – A
604   quantitative study. Prev Med Rep. 2021 Dec;24:101651.

605   19.   National Primary Health Care Developement Agency U. Gender Barriers and
606   Behavioural and Social Drivers Analysis Immunisation in Nigeria. 2024 Apr.

607   20.   Lubeya MK, Mwanahamuntu M, Chibwesha CJ, Mukosha M, Monde MW, Kawonga M.
608   Implementation Strategies Used to Increase Human Papillomavirus Vaccination Uptake by
609   Adolescent Girls in Sub-Saharan Africa: A Scoping Review. Vaccines. 2023 July
610   16;11(7):1246.

611   21.   Fenton ATHR, Eun TJ, Clark JA, Perkins RB. Calling the Shots? Adolescents' Influence
612   on Human Papillomavirus Vaccine Decision-Making During Clinical Encounters. J Adolesc
613   Health. 2020 Apr;66(4):447–54.

614   22.   ICO/IARC HPV Information Centre. Nigeria: Human Papillomavirus and Related
615   Cancers, Fact Sheet 2023. Fact Sheet. 2023 Mar 10;

616   23.   World Cancer Research Fund. Cervical cancer statistics [Internet]. Cervical Cancer
617   Statistics. [cited 2025 July 14]. Available from: https://www.wcrf.org/preventing-
618   cancer/cancer-statistics/cervical-cancer-statistics/

619   24.   Gavi, the Vaccine Alliance. Nigeria to vaccinate 7.7□million girls against leading cause
620   of cervical cancer. 2023 [cited 2025 July 14]. Nigeria to vaccinate 7.7 million girls against
621   leading cause of cervical cancer. Available from: https://www.gavi.org/news/media-
622   room/nigeria-vaccinate-77-million-girls-against-leading-cause-cervical-cancer

623   25.   Steen L. Achieving 71% HPV Vaccination Coverage in Nigeria [Internet]. Evans School
624   of Public Policy & Governance. 2025 [cited 2025 July 14]. Available from:
625   https://evans.uw.edu/achieving-71-hpv-vaccination-coverage-in-nigeria/

626   26.   ICO/IARC Information Centre on HPV and Cancer. Indonesia: Human Papillomavirus
627   and Related Cancers, Fact Sheet 2023. Fact Sheet. 2023;

628   27.   Ferlay E, Lam L, Colombet M, Pineros Z, Soerjomataram B. Global Cancer Observatory
629   Cancer Today. Lyon, France: International Agency for Research on Cancer; 2024.

630   28.   WHO. Indonesia announces declaration on cervical cancer elimination [Internet]. 2023
631   [cited 2025 July 15]. Available from: https://www.who.int/news/item/13-11-2023-indonesia-
632   announces-declaration-on-cervical-cancer-elimination

633   29.   UNICEF Indonesia. A future without cervical cancer: Protecting Indonesia's girls
634   through HPV immunization | UNICEF Indonesia [Internet]. 2025 [cited 2025 July 17].
635   Available from: https://www.unicef.org/indonesia/health/stories/future-without-cervical-
636   cancer-protecting-indonesias-girls-through-hpv-immunization

637   30.   Translation O of A to DCS for SD&. Health Ministry Expands Free HPV Vaccination
638   Coverage [Internet]. Sekretariat Kabinet Republik Indonesia. 2023 [cited 2025 July 17].
639   Available from: https://setkab.go.id/en/health-ministry-expands-free-hpv-vaccination-
640   coverage/

641   31.   WHO. Immunization Data. 2024 [cited 2025 July 17]. WHO Immunization Data portal -
642   Detail Page. Available from: https://immunizationdata.who.int/global/wiise-detail-page

643   32.   Design for Health, Gates Foundation, USAID CENTER FOR INNOVATION AND
644   IMPACT. Glossary of Design Terms.

645   33.   Holeman I, Kane D. Human-centered design for global health equity. Inf Technol Dev.
646   2020 July 2;26(3):477–505.

647   34.   Blynn E, Harris E, Wendland M, Chang C, Kasungami D, Ashok M, et al. Integrating
648   Human-Centered Design to Advance Global Health: Lessons From 3 Programs. Glob Health
649   Sci Pract. 2021 Nov 29;9(Supplement 2):S261–73.

650   35.   Nzilibili SM, Maziku H, Araya A, Kateule R, Malamla MA, Salum S, et al. Using
651   Human-Centered Design to Bridge Zero-Dose Vaccine Gap: A Case Study of Ilala District in
652   Tanzania. Vaccines. 2025 Jan 6;13(1):38.

653   36.   Shearer JC, Nava O, Prosser W, Nawaz S, Mulongo S, Mambu T, et al. Uncovering the
654   Drivers of Childhood Immunization Inequality with Caregivers, Community Members and
655   Health System Stakeholders: Results from a Human-Centered Design Study in DRC,
656   Mozambique and Nigeria. Vaccines. 2023 Mar 17;11(3):689.

37.   Reñosa MDC, Bärnighausen K, Wachinger J, Endoma V, Landicho J, Aligato MF, et al.
Messy but worth it: human-centred design as applied within a successful vaccine-promotive
campaign. BMJ Glob Health. 2024 July;9(7):e014870.

38.   Göttgens I, Oertelt-Prigione S. The Application of Human-Centered Design Approaches
in Health Research and Innovation: A Narrative Review of Current Practices. JMIR MHealth
UHealth. 2021 Dec 6;9(12):e28102.

39.   UNICEF. Resources. 2023 [cited 2025 Aug 5]. Resources | UNICEF Human Centered
Design 4 Health. Available from: https://www.hcd4health.org/resources

40.   UNICEF Health Section Implementation Research and Delivery Science Unit and the
Office of Innovation Global Innovation Centre. Demand for Health Services A Human-
Centred Field Guide for Investigating and Responding to Challenges [Internet]. 2018 Oct.
Available from: https://www.unicef.org/innovation/reports/demand-health-services

41.   Wentzell EA, Inhorn MC. Reconceiving masculinity and 'men as partners' for ICPD
Beyond 2014: Insights from a Mexican HPV study. Glob Public Health. 2014 July
3;9(6):691–705.

42.   Kwambai TK, Dellicour S, Desai M, Ameh CA, Person B, Achieng F, et al. Perspectives
of men on antenatal and delivery care service utilisation in rural western Kenya: a qualitative
study. BMC Pregnancy Childbirth [Internet]. 2013 Dec [cited 2025 July 23];13(1). Available
from: https://bmcpregnancychildbirth.biomedcentral.com/articles/10.1186/1471-2393-13-134

43.   Berenson AB, Laz ,Tabassum H, Hirth ,Jacqueline M, McGrath ,Christine J, and Rahman
M. Effect of the decision-making process in the family on HPV vaccination rates among
adolescents 9–17 years of age. Hum Vaccines Immunother. 2014 July 7;10(7):1807–11.

44.   Mboane R, Bhatta MP. Influence of a husband's healthcare decision making role on a
woman's intention to use contraceptives among Mozambican women. Reprod Health
[Internet]. 2015 Dec [cited 2025 July 21];12(1). Available from: https://reproductive-health-
journal.biomedcentral.com/articles/10.1186/s12978-015-0010-2

45.   Ngom P, Debpuur C, Akweongo P, Adongo P, Binka FN. Gate-Keeping and Women's
Health Seeking Behaviour in Navrongo, Northern Ghana. Afr J Reprod Health. 2003
Apr;7(1):17.

46.   Reuben Mahiti G, Mbekenga CK, Dennis Kiwara A, Hurtig AK, Goicolea I. Perceptions
about the cultural practices of male partners during postpartum care in rural Tanzania: a
qualitative study. Glob Health Action [Internet]. 2017 Jan [cited 2025 July 23];10(1).
Available from: https://www.tandfonline.com/doi/full/10.1080/16549716.2017.1361184

47.   Baland JM, Ziparo R. Intra-household Bargaining in Poor Countries. In: Towards Gender
Equity in Developement. Oxford, United Kingdom: Oxford University Press; 2018.

48.   Azad AD, Charles AG, Ding Q, Trickey AW, Wren SM. The gender gap and healthcare:
associations between gender roles and factors affecting healthcare access in Central Malawi,

June–August 2017. Arch Public Health [Internet]. 2020 Dec [cited 2025 July 21];78(1). Available from: https://archpublichealth.biomedcentral.com/articles/10.1186/s13690-020-00497-w

49. Kumar N, Quisumbing A, Manohar S, Banerjee A, Gupta S, Chauhan A, et al. Intrahousehold dynamics in South Asia: understanding the relationships among women's empowerment, task sharing, women's decision making, and diets [Internet]. Cold Spring Harbor Laboratory; 2025 [cited 2025 July 23]. Available from: http://medrxiv.org/lookup/doi/10.1101/2025.01.08.25320196

50. Melillo S, Strachan R, O'Brien CJ, Wonodi C, Bormet M, Fountain D. Effects of local faith-actor engagement in the uptake and coverage of immunization in low- and middle-income countries: A literature review. Christ J Glob Health. 2022 June 20;9(1):2–32.

51. Miyoshi A, Takiuchi T, Kimura T. HPV vaccination in Japan: can educational intervention promote a father's intention to encourage his daughter's vaccination? Int J Clin Oncol. 2020 Apr 1;25(4):746–54.

52. Vereinte Nationen, editor. A new role for men: partners for women's empowerment. New York: UNFPA; 1997. 16 p.

53. Baker P, Leon N, Colvin CJ, Griffith DM. Health policies must consider gender, including men. Lancet Glob Health. 2023 Dec;11(12):e1847–8.

54. JSI Research & Training Institute, PATH. HPV Evidence Brief: Communications.

55. Internation Vaccine Acess Center, Gavi. Addressing Gender Inequities to Improve Immunization Coverage for Zero-dose Childre. 2022 Dec.

56. Fisher H, Evans K, Ferrie J, Yates J, Roderick M, Audrey S. Young women's autonomy and information needs in the schools-based HPV vaccination programme: a qualitative study. BMC Public Health [Internet]. 2020 Dec [cited 2025 July 24];20(1). Available from: https://bmcpublichealth.biomedcentral.com/articles/10.1186/s12889-020-09815-x

57. Zimet GD, Rosberger Z, Fisher WA, Perez S, Stupiansky NW. Beliefs, behaviors and HPV vaccine: Correcting the myths and the misinformation. Prev Med. 2013 Nov;57(5):414–8.

58. Moradi S, Maboudi M. Enhancing human papillomavirus vaccination uptake: improving communication and countering misinformation. Evid Based Nurs. 2025 Mar 12;ebnurs-2024-104234.

59. Xu M, Choi J, Capasso A, DiClemente R. Improving HPV Vaccination Uptake Among Adolescents in Low Resource Settings: Sociocultural and Socioeconomic Barriers and Facilitators. Adolesc Health Med Ther. 2024 July;Volume 15:73–82.

60. JSI Research & Training Institute. Resource Guide for Strengthening HPV Vaccination Service Delivery. 2023.

61.   Njogu R, Deb Roy A, Bagudu Z, Borda H, Jurgensmeyer M, Muralidharan K, et al.
Effective communication by health care workers to promote HPV vaccination: Lessons
learned from Kenya, India, and Nigeria. Vaccine. 2024 July;42:S27–32.

62.   Cartmell KB, Mzik CR, Sundstrom BL, Luque JS, White A, Young-Pierce J. HPV
Vaccination Communication Messages, Messengers, and Messaging Strategies. J Cancer Educ.
2019 Oct;34(5):1014–23.

63.   Crawshaw AF, Kitoko LM, Nkembi SL, Lutumba LM, Hickey C, Deal A, et al. Co-
designing a theory-informed, multicomponent intervention to increase vaccine uptake with
Congolese migrants: A qualitative, community-based participatory research study (LISOLO
MALAMU). Health Expect Int J Public Particip Health Care Health Policy. 2024
Feb;27(1):e13884.

64.   Jalloh MF, Wilhelm E, Abad N, Prybylski D. Mobilize to vaccinate: lessons learned from
social mobilization for immunization in low and middle-income countries. Hum Vaccines
Immunother. 2020 May 3;16(5):1208–14.

65.   Habib GL, Yousuf H, Bredius L, Bindraban NR, Winter MM, Scherder EJA, et al. The
importance of cultural tailoring of communicators and media outlets in an influenza
vaccination awareness campaign: a digital randomized trial. Sci Rep [Internet]. 2023 Feb 16
[cited 2025 July 24];13(1). Available from: https://www.nature.com/articles/s41598-023-
27910-y

66.   Widayanti AW, Green JA, Heydon S, Norris P. Health-Seeking Behavior of People in
Indonesia: A Narrative Review. J Epidemiol Glob Health. 2020;10(1):6.

67.   Limaye RJ, Sara AB, Siddique AR, Vivas C, Malik S, Omonoju K. Interpersonal and
community influences affecting childhood vaccination decision-making among Nigerian
caregivers: Perceptions among frontline workers in Nigeria. J Child Health Care. 2019
Sept;23(3):403–14.

68.   Akinyemi JO, Adedini SA, Odimegwu CO. Individual versus community-level measures
of women decisionmaking involvement and child survival in Nigeria. South Afr J Child
Health. 2017 Mar 31;11(1):26.

69.   Osamor P, Grady C. FACTORS ASSOCIATED WITH WOMEN'S HEALTH CARE
DECISION-MAKING AUTONOMY: EMPIRICAL EVIDENCE FROM NIGERIA. J Biosoc
Sci. 2018 Jan;50(1):70–85.

70.   Hoffman E, Kahan T, Auerbach E, Brody H, Abramson NN, Haiken S, et al. Peer
education as a strategy to promote vaccine acceptance: A randomized controlled trial within
New York community healthcare practices. Vaccine. 2024 Nov;42:126028.

71.   Posavac EJ, Kattapong KR, Dew DE. Peer-Based Interventions to Influence Health-
Related Behaviors and Attitudes: A Meta-Analysis. Psychol Rep. 1999
Dec;85(3_suppl):1179–94.

72.   Rosser EN, Wysong MD, Rosen JG, Limaye RJ, Park S. HPV Vaccine Delivery
768   Strategies to Reach Out-of-School Girls in Low- and Middle-Income Countries: A Narrative
769   Review. Vaccines. 2025 Apr 22;13(5):433.

770   73.   JSI Research & Training Institute, PATH. HPV Evidence Brief: Pitfalls. 2025.

771   74.   Etim EOE, Odiachi A, Dougherty L, Alabi M, Adetunji A, Adedimeji A. Understanding
772   caregivers' and community influencers' perspectives on the barriers to childhood
773   immunisation in Northern Nigerian States with public-private partnerships in routine
774   immunisation programme. BMC Public Health [Internet]. 2025 Apr 21 [cited 2025 July
775   24];25(1). Available from:
776   https://bmcpublichealth.biomedcentral.com/articles/10.1186/s12889-025-22710-7

777   75.   Waheed DEN, Bolio A, Guillaume D, Sidibe A, Morgan C, Karafillakis E, et al. Planning,
778   implementation, and sustaining high coverage of human papillomavirus (HPV) vaccination
779   programs: What works in the context of low-resource countries? Front Public Health
780   [Internet]. 2023 Apr 14 [cited 2025 July 24];11. Available from:
781   https://www.frontiersin.org/articles/10.3389/fpubh.2023.1112981/full

782   76.   JSI Research & Training Institute, PATH. HPV Evidence Brief: Effective Delivery. 2025.