**CPR Preparedness Across Massachusetts Public High Schools: A Statewide Cross-**

**Sectional Study**

Authors

Meng Yang[1,†,*], Nayan L Sapers[2,†], Ivette I Chen[2,†], William A Porcaro[3,4,*]

[†]These authors contributed equally to this work and share first authorship.

[*]Correspondence: wporcaro@mah.harvard.edu (WAP); mengxxyang@gmail.com (MY)

[1]Harvard Division of Continuing Education, Cambridge, MA, USA;

[2]Harvard College, Cambridge, MA, USA;

[3]Department of Emergency Medicine, Mount Auburn Hospital, Beth Israel Lahey Health,

Cambridge, MA, USA;

[4]Harvard Medical School, Boston, MA, USA

**Abstract**

**Background:** Out-of-hospital cardiac arrest (OHCA) accounts for over 350,000 deaths

annually in the United States, and survival depends on early bystander cardiopulmonary

resuscitation (CPR). Although many cardiac arrests occur on or near school grounds,

Massachusetts has no statewide CPR graduation requirement and little current data on school

preparedness.

**Methods:** We conducted a cross-sectional electronic survey of all 413 public high schools in

Massachusetts (including charter, vocational, and technical) between September 29 and

November 17, 2025. The 14-item survey asked about enrollment, staff size, CPR and

automated external defibrillator (AED) resources, student and staff training, and the presence

of a cardiac emergency response plan (CERP). The CERP item was excluded from analysis

due to inconsistent interpretation. We summarized resources per 1,000 students or staff,

compared Title I and non–Title I schools, and explored geographic variation and

multivariable predictors of AED availability and CPR teaching.

**Results:** One hundred schools responded (24.2%), representing 13 of 14 counties, with a

median enrollment of 662 students, and 33.0% were Title I schools. Overall, 72.0% reported

teaching CPR. The median student training rate was 138.2 per 1,000 students, though only

10.0% of schools with non-missing data reported all students trained and 15.0% reported at

least 70% of students trained. Among responding schools, Title I schools had fewer trained

students than non-Title I schools (median 50.3 vs 199.8 per 1,000; Holm-Bonferroni adjusted

$p = 0.025$), despite similar AED and manikin availability. This disparity persisted in

sensitivity analyses using median imputation for missing student training data. County-level

analyses suggested geographic variation in both training rates and AED density, although

county-level estimates were based on small numbers of responding schools.

**Conclusions:** Among responding Massachusetts high schools, most reported some CPR

instruction, but only a small minority achieved broad student coverage, with particularly low

training rates in Title I schools. These exploratory findings underscore the need for policies

ensuring equitable CPR training access, particularly in Title I schools, and support targeted

investment in school-based cardiac emergency preparedness

**Key Terms**

Cardiopulmonary Resuscitation; Automated External Defibrillator; CPR in Schools; School

Emergency Preparedness; Health Equity

**Introduction**

Out-of-hospital cardiac arrest (OHCA) is a major public health concern in the United States,

occurring in more than 350,000 people each year, including an estimated 23,000 occurring in

children under eighteen.[1,2] Survival depends heavily on early bystander cardiopulmonary

resuscitation (CPR), and each minute without CPR or defibrillation is associated with a 7-

10% decrease in survival.[3,4] Prior studies estimate that one in every 250 to 600 cardiac arrests

occurs on school grounds,[5,6] which highlights the importance of CPR education and

preparedness in school settings.

Although school-based CPR instruction is widely supported as an effective approach to

increase community preparedness, implementation across the United States remains

variable.[7] Massachusetts is one of only nine states without a statewide CPR mandate for

students,[8] and current data on national CPR training and resources are limited.[9,10] Evidence

from national studies also suggests that schools in lower-income communities face greater

barriers to CPR training because of limited equipment and local resource constraints.[11–14]

Because nearly all adolescents pass through the public school system, school-based CPR

training represents a scalable mechanism for reaching an entire age cohort. Students trained

in high school may become adult bystanders and household responders, meaning training

coverage at graduation may influence long-term community readiness for cardiac arrest.

Evaluating CPR preparedness at the high school level therefore provides insight into future

population-level bystander CPR capacity. However, whether current school programs achieve

broad and equitable student training coverage remains unclear.

To address these gaps, we conducted an evaluation of CPR and automated external

defibrillator (AED) infrastructure across public Massachusetts high schools. We assessed

student and staff CPR training rates and equipment availability, with attention to

socioeconomic and regional variation. Together, these findings aim to inform efforts to

expand and improve the equity of school-based CPR education in Massachusetts.

**Methods**

This observational cross-sectional study used an electronic survey to assess CPR and AED

infrastructure in Massachusetts public high schools. Data collection occurred between

September 29, 2025, and November 17, 2025. Eligible schools included all public, charter,

vocational, and technical schools serving grades 9–12 listed in the Massachusetts Department

of Elementary and Secondary Education School and District Profiles database.[15] Nonpublic

schools were not included in the sampling frame as they operate under different funding

streams and regulatory requirements. The study was reviewed by the Harvard University

Committee on the Use of Human Subjects and determined to be exempt human subjects

research in accordance with federal regulations.

The survey consisted of fourteen items administered electronically. Respondents provided

their role, school name, and district. Schools reported student enrollment, staff size, numbers

of CPR manikins and AEDs, and estimated counts of students and staff trained in CPR. The

survey also asked whether CPR is taught and whether the school maintained a Cardiac

Emergency Response Plan (CERP). Because Massachusetts high schools are required to

maintain a Medical and Behavioral Health Emergency Response Plan (MBHERP),

qualitative review indicated that many respondents conflated CERPs with the state-mandated

MBHERP. Because responses could not be consistently interpreted, the CERP item was

excluded from all analyses. Title I status was determined using the 2024–2025 Massachusetts

Department of Elementary and Secondary Education database.[15]

The survey was intended for school personnel with direct knowledge of emergency

preparedness or health education, including school nurses, principals, athletic directors,

health and physical education teachers, and district administrators. All Massachusetts

superintendents were contacted via six rounds of email during the study period. No financial

or administrative incentives were provided for participation, and completion required

voluntary response by school personnel. Additional dissemination occurred through partner

organizations, including the Athletic Trainers of Massachusetts (ATOM), the Massachusetts

School Nurses Organization (MSNO), and Harvard CrimsonEMS and the American Heart

Association's Massachusetts High School CPR Ambassadors. Schools that responded or

opted out were removed from subsequent outreach rounds. Follow-up outreach was

conducted, when needed, to clarify implausible entries and to obtain school-specific

information when respondents represented district-level offices.

One hundred unique Massachusetts high schools (24.2% response rate) remained for analysis

following data cleaning. Most responses were submitted by school nurses or health

administrators (60.0%), followed by school administrators (23.0%), health, physical

education teachers, or athletic staff (13.0%), and science teachers (4.0%). Student-facing

metrics (AEDs, manikins, students trained) were normalized per 1,000 enrolled students; staff

training was normalized per 1,000 staff. Analyses involving student or staff CPR training

rates used complete-case samples, defined as schools with non-missing and internally

consistent values for the corresponding variable. Implausible values (e.g., staff training rates

exceeding 1,000 per 1,000 staff) were excluded a priori. Identifying information such as

respondent names and email addresses was retained only for follow-up communication and

verification and was not included in the analytic dataset. All schools verified consent for

inclusion in de-identified aggregate analysis through an opt-out email sent after the study

period.

**Statistical Analysis**

Descriptive statistics were summarized as medians with interquartile ranges (IQRs) for

continuous variables due to non-normal distributions and as frequencies with percentages for

categorical variables. Differences between Title I and non-Title I schools were assessed using

Mann-Whitney U tests for continuous variables and chi-square tests for categorical variables;

Fisher's exact tests were used when expected cell counts were <5. To account for multiple

hypothesis testing within each domain, p-values were adjusted using the Holm-Bonferroni

method. Geographic variation in CPR training and AED availability was evaluated using

Kruskal-Wallis tests. Counties with fewer than three responding schools were excluded from

these analyses because small group sizes yield unstable variance estimates and violate the

assumptions of rank-based tests.

Two multivariable regression models were fit to explore patterns in CPR and AED resources.

Model 1 used negative binomial regression to examine predictors of AED count, with log

enrollment included as an offset such that coefficients reflected AEDs per student. Predictors

included Title I status, enrollment (per one hundred students), and county (modeled as

indicator variables to adjust for geographic clustering). Model 2 used logistic regression to

examine whether Title I status and enrollment predicted whether schools reported teaching

CPR without county adjustment due to small cell sizes. Residual diagnostics (e.g. Breusch-

Pagan test for heteroscedasticity, variance inflation factors for multicollinearity, and events-

per-variable assessment for the logistic model) did not reveal major deviations from model

assumptions. Given the self-reported sample, these models were treated as exploratory and

were intended to describe associations rather than test hypotheses or generate population-

level estimates.

Sensitivity analyses assessed robustness to missing data. Twenty schools (20%) had missing

values for student CPR training counts; in sensitivity analyses, these values were imputed

using the median training rate within each Title I stratum before repeating Mann–Whitney U

tests. For Model 1, an additional linear regression model with log-transformed AEDs per one

thousand students as the outcome was fit. Because most missing values reflected "do not

know" responses, these analyses were used to evaluate directional consistency rather than

produce unbiased estimates.

All analyses were conducted in Python version 3.10 using scipy, statsmodels, and scikit-

learn. Statistical significance was defined as $\alpha = 0.05$ (two-tailed).

**Results**

**Respondent Characteristics**

Of 413 public high schools in Massachusetts, 100 (24.2%) completed the survey,

representing 13 of 14 counties (Nantucket County had no respondents). Compared to

statewide values, respondent schools had higher median enrollment (662 vs. 542 students)

and lower Title I representation (33.0% vs. 45.5%).

The median school enrollment was 662 students (IQR: 433–1,200) with a median staff size of

106 (IQR: 75–200). Thirty-three schools (33.0%) were designated as Title I. Missing and

excluded data ranged from 0% to 20.0% across variables (Table 1).

**Table 1. Missing and Exclusion Data Summary**

| Variable | Missing/Excluded, n (%) |
|---|---|
| Title I status | 0 (0.0) |
| Enrollment | 0 (0.0) |
| Staff count | 0 (0.0) |
| AEDs | 0 (0.0) |
| Manikins | 4 (4.0) |
| Teaches CPR | 0 (0.0) |
| Students trained | 20 (20.0) |
| Staff trained | 11 (11.0)☐ |

☐Includes 3 schools excluded due to implausible values (staff trained > 1,000 per 1,000

staff).

**CPR Training and Equipment**

Seventy-two schools (72.0%) reported teaching CPR in some capacity. The median number

of students trained was 138.2 per 1,000 enrolled (IQR: 44.9–265.0), while the median number

of staff trained was 238.1 per 1,000 staff (IQR: 150.0–375.0). Among 80 schools with non-

missing student training data, 8 schools (10.0%) reported 100% of students trained, and an

additional 4 schools (5.0%) reached 70–99% of students trained. The schools with missing

data (20.0%) reported not knowing how many students were trained. The median manikin

availability was 8.9 per 1,000 students (IQR: 1.9–20.0). Notably, of 96 schools with non-

missing manikin data, 22 (22.9%) reported zero manikins.

Across all schools, the median AED availability was 7.9 devices per 1,000 students (IQR:

4.6–11.9). County-level variation was substantial, ranging from 3.5 per 1,000 in Suffolk

County to 33.3 per 1,000 in Dukes County; however, these estimates were based on small

numbers of schools per county (range: 1–14).

The rate of students trained in CPR was significantly lower in Title I schools compared to

non-Title I schools (median: 50.3 vs. 199.8 per 1,000; $U = 466.5$, adjusted $p = 0.025$), with a

medium effect size ($r = 0.37$; Figure 1). No significant differences were observed for AED

availability, manikin availability, or staff training rates after adjustment for multiple

comparisons. Table 2 presents bivariable comparisons between Title I and non-Title I

schools.

**Table 2. CPR Preparedness by Title I Status**

| Measure | Title I | Non-Title I | Adjusted p-value |
| --- | --- | --- | --- |

| Equipment, median (IQR) | | | |
|---|---|---|---|
| AEDs per 1,000 students | 9.2 (5.1–12.0) | 6.9 (4.4–11.8) | 0.78 |
| Manikins per 1,000 students | 5.4 (0.9–14.8)▢ | 11.1 (3.2–21.8)▢ | 0.27 |
| **Training, median (IQR)** | | | |
| Students trained per 1,000 | 50.3 (12.5–173.6)▢ | 199.8 (84.8–272.7)▢ | **0.025** |
| Staff trained per 1,000 staff | 243.2 (106.2–333.3)▢ | 238.1 (153.8–472.2)▢ | 0.78 |
| **Programs, % (n)** | | | |
| Teaches CPR | 66.7 (22/33) | 74.6 (50/67) | 0.55 |

▢n = 32. ▢n = 64. ▢n = 29. ▢n = 51. ▢n = 30. ▢n = 59.

**Note:** p-values are Holm-Bonferroni adjusted. Continuous variables compared using Mann-

Whitney U tests. Bold indicates statistical significance at $\alpha = 0.05$. Effect size for students

trained: r = 0.37 (medium).

**Exploratory Analyses**

In sensitivity analyses using median imputation for missing student training data within Title

I strata (20 schools with missing values), the disparity in student CPR training between Title I

and non–Title I schools persisted (Mann–Whitney U test, $p < 0.001$). As a sensitivity

analysis, we also fit an ordinary least squares model using log-transformed AEDs per 1,000

students as the outcome. Results were consistent with the negative binomial model: the

direction and statistical significance of all predictors were unchanged.

After excluding counties with fewer than three responding schools, geographic variation was

observed across counties for AED availability (Kruskal-Wallis H = 23.29, df = 11, p = 0.016,

$\varepsilon^2 = 0.24$, indicating a large effect) and student CPR training rates (H = 20.84, df = 10, p =

0.022, $\varepsilon^2 = 0.27$), with large effect sizes. County-level median student training rates ranged

from 19.0 per 1,000 (Hampden County, n = 9) to 712.9 per 1,000 (Middlesex County, n =

10). Teaching CPR ranged from 33.3% (Hampshire and Suffolk counties, n = 6 each) to

100.0% (Barnstable County, n = 3). Given the small sample sizes per county (1–14 schools),

county-specific estimates should be interpreted with caution.

The Poisson model showed overdispersion (deviance/df = 2.42), supporting the use of

negative binomial regression with dispersion parameter $\alpha$ = 0.706. In the negative binomial

regression model with AED count as the outcome and log(enrollment) as an offset, Title I

status was not significantly associated with AED count (IRR = 1.06, 95% CI: 0.78–1.43, p =

0.725).

In multivariable logistic regression (n = 100; 72 events, 28 non-events), neither Title I status

(OR = 0.67,  95% CI: 0.24–1.78, p = 0.427) nor enrollment (OR = 1.05 per 100 students,

95% CI: 0.98–1.19, p = 0.292) were significantly associated with CPR teaching. As expected

for structural predictors, model discrimination was modest (AUC = 0.61) and adequate

calibration (Brier score = 0.196). With 72 events and 2 predictors, the events per variable

ratio was 36, exceeding the recommended threshold of 10.

**Discussion**

In this cross-sectional survey of 100 Massachusetts public high schools (24.2% response

rate), we identified three main patterns in CPR and AED infrastructure. First, although most

responding schools reported offering CPR instruction (72.0%), only a small proportion

achieved universal student training (10.0%), highlighting a substantial gap between offering

instruction and achieving full coverage. Because high school education reaches nearly all

adolescents, incomplete training coverage at graduation may translate into gaps in future

community bystander CPR capacity. Second, student training rates differed markedly by

socioeconomic context: Title I schools had a roughly four-fold lower median student training

rate than non–Title I schools, despite reporting similar levels of AED availability, manikin

supply, staff training, and CPR course offerings. Third, exploratory analyses suggested

geographic variation in training capacity, though these findings should be interpreted

cautiously given small county-level sample sizes.

Although we did not assess implementation processes, the gap between offering CPR

instruction and achieving universal coverage suggests variation in how schools integrate CPR

training into curricula. Equipment access may further constrain implementation. Nearly one

quarter of responding schools reported no manikins, and although median manikin counts did

not differ by Title I status, lower-resource schools may face greater difficulty acquiring,

maintaining, or replacing equipment. Prior studies have documented reliance on external

organizations to support training in resource-limited schools,[10,16] which may partly explain

disparities observed in other settings.

The observed socioeconomic differences have important public health implications. School-

level differences in CPR training may reflect or contribute to broader community inequities

in cardiac arrest preparedness. Survival from OHCA depends heavily on early bystander

CPR, and survival decreases by approximately 7-10% for every minute without

intervention.[3,4] Lower-income and rural communities consistently demonstrate lower

bystander CPR rates and poorer cardiac arrest outcomes.[14] Limited access to CPR instruction

for students in these communities may reinforce community-level disparities by reducing the

number of trained responders. School-based CPR training represents a cost-effective public

health intervention and a potential mechanism for improving equitable access to lifesaving

skills.[10,17]

These findings also have several implications for policy and program development. At the

federal level, the HEARTS Act of 2024 (H.R.6829) would provide funding for CPR

infrastructure but remains unfunded;[18] our data support targeted funding to rural and Title I

districts to address resource inequities. Massachusetts is currently evaluating CPR in Schools

legislation, and these findings may help guide implementation strategies.[19–25] Potential state-

level approaches include centralized purchasing programs or equipment loan libraries to

address equipment shortages. At the school level, several scalable models exist for expanding

CPR education. Student-led programs such as American Heart Association HEART Clubs

provide a low-cost mechanism for increasing student engagement in CPR training.[26] The

American Heart Association's recent $4,500 school-based grant program may also help offset

equipment and implementation costs.[27] Programs such as Project ADAM (Automated

Defibrillators in Adam's Memory) and HEARTSafe Communities offer frameworks for

improving school cardiac preparedness through standardized AED access, emergency

planning, and regular drills.[28,29] Harvard CrimsonEMS's CPR Ambassador Program is also

an example of how collegiate EMS agencies can further support high school CPR advocacy

through training, equipment access, and mentorship.[30] Our study did not evaluate the

effectiveness of these programs; we reference them as examples of existing initiatives that

could be considered in efforts to strengthen school-based CPR and AED infrastructure.

Several limitations should be considered when interpreting these findings. The response rate

was modest (24.2%) but is comparable to other unsolicited surveys of educational and public

sector institutions, where participation depends on administrative availability and perceived

relevance rather than direct incentive. Nonresponse bias is possible; schools without

established CPR programs may have been less likely to participate, potentially leading to

overestimation of overall training coverage. However, the primary findings of this study

relate to differences in coverage and access rather than precise statewide prevalence, and such

bias would be expected to attenuate rather than create the observed disparities. All data were

self-reported by a single respondent per school, introducing possible recall error, incomplete

knowledge of school-wide practices, and social desirability bias. We did not independently

verify AED functionality, equipment condition, or training quality, and the cross-sectional

design precludes causal inference or assessment of temporal trends. These limitations are

common in prior studies of CPR in schools.[5,31] In addition, ambiguity in survey wording

limited our ability to determine the true prevalence of Cardiac Emergency Response Plans

(CERPs) statewide. Future studies should include explicit definitions distinguishing CERPs

and Massachusetts-specific emergency response plans (MBHERPs) to improve measurement

accuracy.

Despite these limitations, the study provides statewide evidence that CPR instruction

availability does not necessarily translate into universal training and that meaningful

inequities exist in student access to lifesaving education. Targeted resource allocation,

structured implementation support, and integration of CPR training into graduation

requirements may improve coverage and reduce disparities in community cardiac arrest

preparedness

**Conclusion**

This statewide survey demonstrates that the presence of CPR instruction in schools does not

necessarily translate into universal student training. Although most responding Massachusetts

public high schools reported offering CPR education, few achieved full student coverage, and

substantial disparities existed between Title I and non-Title I schools. These findings suggest

that school-based CPR training may represent an underrecognized contributor to inequities in

community cardiac arrest preparedness.

Targeted implementation support, equipment access, and integration of CPR training into

graduation requirements may help improve coverage and reduce disparities. The rapid, low-

cost audit approach used in this study provides a practical method for states and organizations

to evaluate school preparedness, monitor progress over time, and guide resource allocation.

Broader adoption of similar assessments may support more equitable access to lifesaving

training and strengthen community response to OHCA.

**Abbreviations and Acronyms**

AED          =          automated external defibrillator

CPR          =          cardiopulmonary resuscitation

OHCA        =          out-of-hospital cardiac arrest

CERP        =          cardiac emergency response plan

MBHERP     =          medical and behavioral health emergency response plan

**Declarations**

**Ethics approval and consent to participate**

This project was reviewed by the Harvard University Committee on the Use of Human

Subjects and determined to be exempt human subjects research in accordance with federal

regulations.

**Consent for publication**

Not applicable

**Availability of data and materials**

The datasets used and/or analysed during the current study are available from the

corresponding author on reasonable request.

**Competing interests**

The authors declare the following potential conflicts of interest: MY, NLS, and IIC are

affiliated with the American Heart Association and CrimsonEMS; both organizations are

referenced in the manuscript. The CrimsonEMS and the American Heart Association CPR

Ambassador Program assisted with survey dissemination and is referenced in the Discussion.

These affiliations did not influence data collection, analysis, or interpretation of findings.

WAP declares no competing interests.

**Funding**

No financial conflicts exist. We did not receive any payment for any aspect of the submitted

work.

**Authors' contributions**

NLS and MY contributed to the conceptualization of the study and development of the research aims. NLS, MY, and IIC collaboratively designed the study methodology. Data collection and investigation were carried out by MY, IIC, and NLS, and all three authors participated in data curation and project administration. NLS and MY conducted the formal analysis of the data. NLS developed the study software and prepared data visualizations. MY, NLS, and IIC each drafted sections of the manuscript and contributed to the original draft preparation. WAP contributed to interpretation of the data and provided substantive critical revision of the manuscript for important intellectual content. All authors contributed to critical revision and editing of the manuscript and approved the final version.

**Acknowledgements**

We would like to thank Dr. Stylianos Maheras and Tristan Milarch for their review and comments on our manuscript.

**References**

1. 2025 Heart Disease and Stroke Statistics: A Report of US and Global Data From the American Heart Association. https://doi.org/10.1161/CIR.0000000000001303.

2. Okubo M, Chan HK, Callaway CW, Mann NC, Wang HE. Characteristics of paediatric out-of-hospital cardiac arrest in the United States. Resuscitation. 2020;153:227–33. https://doi.org/10.1016/j.resuscitation.2020.04.023.

3. Larsen MP, Eisenberg MS, Cummins RO, Hallstrom AP. Predicting survival from out-of-hospital cardiac arrest: a graphic model. Ann Emerg Med. 1993;22:1652–8. https://doi.org/10.1016/s0196-0644(05)81302-2.

4. Hara M, Hayashi K, Hikoso S, Sakata Y, Kitamura T. Different Impacts of Time From Collapse to First Cardiopulmonary Resuscitation on Outcomes After Witnessed Out-of-Hospital Cardiac Arrest in Adults. Circ Cardiovasc Qual Outcomes. 2015;8:277–84. https://doi.org/10.1161/CIRCOUTCOMES.115.001864.

5. Swor R, Grace H, McGovern H, Weiner M, Walton E. Cardiac arrests in schools: Assessing use of automated external defibrillators (AED) on school campuses. Resuscitation. 2013;84:426–9. https://doi.org/10.1016/j.resuscitation.2012.09.014.

6. Lotfi K, White L, Rea T, Cobb L, Copass M, Yin L, et al. Cardiac arrest in schools. Circulation. 2007;116:1374–9. https://doi.org/10.1161/CIRCULATIONAHA.107.698282.

7. Vetter VL, Griffis H, Dalldorf KF, Naim MY, Rossano J, Vellano K, et al. Impact of State Laws: CPR Education in High Schools. J Am Coll Cardiol. 2022;79:2140–3. https://doi.org/10.1016/j.jacc.2022.03.359.

8. CPR in Schools Legislation Map. cpr.heart.org. https://cpr.heart.org/en/training-

programs/community-programs/cpr-in-schools/cpr-in-schools-legislation-map. Accessed 29

Nov 2025.

9. Schroeder DC, Semeraro F, Greif R, Bray J, Morley P, Parr M, et al. KIDS SAVE LIVES:

Basic Life Support Education for Schoolchildren: A Narrative Review and Scientific

Statement From the International Liaison Committee on Resuscitation. Circulation.

2023;147:1854–68. https://doi.org/10.1161/CIR.0000000000001128.

10. Cave DM, Aufderheide TP, Beeson J, Ellison A, Gregory A, Hazinski MF, et al.

Importance and Implementation of Training in Cardiopulmonary Resuscitation and

Automated External Defibrillation in Schools. Circulation. 2011;123:691–706.

https://doi.org/10.1161/CIR.0b013e31820b5328.

11. Blewer AL, Okubo M. Disparities within pediatric out-of-hospital cardiac arrest: A call to

action. Resuscitation. 2023;192. https://doi.org/10.1016/j.resuscitation.2023.109968.

12. Anderson ML, Cox M, Al-Khatib SM, Nichol G, Thomas KL, Chan PS, et al. Rates of

Cardiopulmonary Resuscitation Training in the United States. JAMA Intern Med.

2014;174:201. https://doi.org/10.1001/jamainternmed.2013.11320.

13. Abdulhay NM, Totolos K, McGovern S, Hewitt N, Bhardwaj A, Buckler DG, et al.

Socioeconomic disparities in layperson CPR training within a large U.S. city. Resuscitation.

2019;141:13–8. https://doi.org/10.1016/j.resuscitation.2019.05.038.

14. Sasson C, Magid DJ, Chan P, Root ED, McNally BF, Kellermann AL, et al. Association

of neighborhood characteristics with bystander-initiated CPR. N Engl J Med.

2012;367:1607–15. https://doi.org/10.1056/NEJMoa1110700.

15. Profiles Search. https://profiles.doe.mass.edu/search/search.aspx?leftNavId=11238.

Accessed 29 Nov 2025.

16. Sadjadi M, Brülle R, Onbasilar U, Booke H, Strauß C, von Groote T, et al.

Implementation of school-based CPR training - A systematic review and mixed-methods

meta-analysis. Resusc Plus. 2025;23:100955. https://doi.org/10.1016/j.resplu.2025.100955.

17. Swor R, Compton S. Estimating cost-effectiveness of mass cardiopulmonary resuscitation

training strategies to improve survival from cardiac arrest in private locations. Prehosp Emerg

Care. 2004;8:420–3. https://doi.org/10.1016/j.prehos.2004.06.012.

18. Rep. Pallone F. Text - H.R.6829 - 118th Congress (2023-2024): HEARTS Act of 2024.

2024. https://www.congress.gov/bill/118th-congress/house-bill/6829/text. Accessed 29 Nov

2025.

19. Bill H.556. https://malegislature.gov/Bills/194/H556. Accessed 7 Dec 2025.

20. Bill H.572. https://malegislature.gov/Bills/194/H572. Accessed 7 Dec 2025.

21. Bill H.750. https://malegislature.gov/Bills/194/H750. Accessed 7 Dec 2025.

22. Bill H.2552. https://malegislature.gov/Bills/194/H2552. Accessed 7 Dec 2025.

23. Bill H.4107. https://malegislature.gov/Bills/194/H4107. Accessed 7 Dec 2025.

24. Bill S.1631. https://malegislature.gov/Bills/194/S1631. Accessed 7 Dec 2025.

25. Bill S.2863. https://malegislature.gov/Bills/194/S2863. Accessed 16 Feb 2026.

26. Start a Club. www.heart.org. https://www.heart.org/en/volunteer/high-school-and-

college-students/start-a-club. Accessed 29 Nov 2025.

27. High School Heart Club Grants. www.heart.org. https://www.heart.org/en/volunteer/high-

school-and-college-students/start-a-club/club-resources/high-school-heart-club-grants.

Accessed 29 Nov 2025.

28. Heart Safe Programs. https://projectadam.com/Heartsafe. Accessed 29 Nov 2025.

29. HEARTSAFE COMMUNITY – Citizen CPR. https://citizencpr.org/heartsafe-

community/. Accessed 29 Nov 2025.

30. Advocacy – CrimsonEMS. https://crimsonems.org/advocacy/. Accessed 29 Nov 2025.

31. Brown LE, Lynes C, Carroll T, Halperin H. CPR Instruction in U.S. High Schools. JACC.

2017;70:2688–95. https://doi.org/10.1016/j.jacc.2017.09.1101.

488
489    *Figure 1. Student CPR training rates by Title I status. Box plots show median (horizontal*
490    *line), interquartile range (box), and 1.5x IQR (whiskers); outliers shown as circles.* Title I
491    schools (n=29) trained significantly fewer students than non-Title I schools (n=51; median:
492    50.3 vs. 199.8 per 1,000; p = 0.025, Holm-Bonferroni adjusted).
493
494
495
496

497
498
499 *Figure 2. CPR preparedness resources by Title I status. Bars represent medians for*
500 *continuous variables (top four panels) and percentages for categorical variables (bottom*
501 *panel).* Sample sizes shown below each bar. Three schools with implausible staff training
502 values were excluded from staff analysis. *p = 0.025 (Holm-Bonferroni adjusted).