Population Attributable Mortality Associated with Respiratory Viruses in Ontario

David N. Fisman, MD MPH[1], Alicia A. Grima MPH PhD(c) [1], Natalie J. Wilson[1] MPH PhD (c), Ashleigh R. Tuite PhD MPH[1,2], and Clara Eunyoung Lee[1] MD PhD

[1]Dalla Lana School of Public Health, University of Toronto, Toronto ON, Canada
[2] Centre for Immunization Surveillance and Programs, Public Health Agency of Canada, Ottawa ON, Canada

Address reprint requests and correspondence to:

David Fisman, MD MPH FRCP(C)

Professor, Dalla Lana School of Public Health

University of Toronto

Room 686, 155 College Street, Toronto, Ontario, M5T 3M7

Email: david.fisman@utoronto.ca

Competing Interest Statement

DNF has served on advisory boards related to influenza and SARS-CoV-2 vaccines for Seqirus and Pfizer vaccines. ART was employed by the Public Health Agency of Canada when the research was conducted. The work does not represent the views of the Public Health Agency of Canada. Other authors: no competing interests.

Word count: 2983
Abstract word count: 259

Abstract

Background: Respiratory viruses are major contributors to population mortality, but cause-of-death coding undercounts their impact. Ecological regression models linking viral circulation to mortality fluctuations can address this limitation.

Aim: To estimate the population attributable fraction (PAF) of mortality associated with influenza A and B, respiratory syncytial virus (RSV), and SARS-CoV-2 in Ontario, Canada (1993–2024).

Methods: We analysed monthly all-cause mortality data with laboratory surveillance indicators for influenza A, B, RSV, and SARS-CoV-2. Negative binomial models with secular trends, Fourier seasonal terms, and population offsets were stratified into pre-pandemic (1993-February 2020) and pandemic (March 2020-March 2024) periods. PAFs were derived from counterfactual predictions setting viral coefficients to zero. Sensitivity analyses excluded seasonal terms; Wald tests compared coefficients across model specifications.

Results: Pre-pandemic, influenza A accounted for 1.8% (95% CI 1.4–2.3%) of mortality; influenza B showed no detectable impact. RSV demonstrated inverse associations in seasonally adjusted models but positive associations (PAF 1.9%, 95% CI 1.3–2.4%) without seasonal adjustment. During 2020–2024, amid elevated baseline mortality (IRR 1.050, P=0.027), SARS-CoV-2 dominated, accounting for 6.5% (95% CI 4.5–8.4%) of deaths, 3.6-times the pre-pandemic influenza A burden, despite widespread vaccination and antiviral availability. Model-estimated SARS-CoV-2 deaths (18,052) matched reported COVID-19 deaths (18,603). Meta-analyses showed substantial heterogeneity for influenza A (I^2=93.7%) and RSV (I^2=88.5%) across periods and modeling approaches, but minimal heterogeneity for SARS-CoV-2 (I^2=2.5%).

Conclusion: SARS-CoV-2 demonstrated 3–4-fold higher mortality burden than seasonal influenza A despite available countermeasures. Estimates for influenza A and RSV were sensitive to seasonal adjustment, highlighting the importance of modelling choices when quantifying virus-attributable mortality.

Introduction

Respiratory viral illnesses impose a substantial burden on individuals and the health care system, often resulting in significant morbidity and mortality. Many of these viruses are now vaccine preventable or can have their severity attenuated through vaccination. This holds true in a Canadian context, where two of the leading causes of death are respiratory illnesses; influenza, which prior to the COVID-19 pandemic, caused approximately 3,500 deaths annually ([1](#)), and severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), which as of 2023 had caused a reported 55,282 deaths in Canada ([2](#), [3](#)). Other respiratory viruses, such as respiratory syncytial virus (RSV), can also cause substantial morbidity and mortality, particularly in vulnerable ages groups (e.g., children and older adults) ([4-6](#)).

To prioritize vaccination and other prevention programs against respiratory viruses, it is important to understand the potential for prevention of morbidity and mortality at the population level. One such method of doing so is by analysing the population-attributable fraction (PAF) for a given respiratory virus. PAF quantifies the proportion of cases that could be prevented if the exposure, (in this case, the viral infection), were eliminated, offering a valuable metric for public health planning ([7](#)). PAF can be calculated as the difference between the observed number of cases (O) and the expected number of cases if the exposure was removed (E), divided by the observed number of cases (PAF = (O-E)/O) ([7](#)). "Shortcut formulae" allow PAF to be estimated if the relative risk among the exposed, and the prevalence of exposure, are known ([7](#), [8](#)).

When mortality is the outcome of interest, researchers often rely on cause-specific mortality data to estimate the impact of respiratory viruses. However, these data are subject to well-known limitations, including misclassification and inaccuracies in death certificate coding ([9](#)). As an alternative, regression-based methods offer an indirect means of estimating the PAF

([10](#)). These approaches do not require explicit cause-of-death attribution and are agnostic to the magnitude of the exposure effect. By modeling the statistical association between viral circulation and mortality, often using smoothing techniques to detrend and smooth data, such models can estimate the fraction of mortality attributable to a given virus, based on the observed changes in outcomes corresponding to fluctuations in exposure.

In this study, we sought to quantify the population-level mortality burden attributable to key respiratory viruses in Ontario, Canada, using regression-based approaches. Specifically, we estimated the PAF for influenza A and B, and respiratory syncytial virus (RSV) over the period from 1993 to 2024, and for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) from 2020 to 2024. By applying statistical models that account for temporal variation in viral activity and mortality, our aim was to provide robust, population-based estimates of virus-attributable mortality to inform future vaccination strategies and respiratory virus prevention policies. We also evaluated how modelling choices, specifically the inclusion or exclusion of seasonal adjustment term, affect estimates of virus-attributable mortality.

Methods

Data Sources

We calculated percent positivity for influenza A and B, and RSV in Ontario using laboratory-based surveillance data from FluWatch, Canada's national surveillance system for influenza and other respiratory viruses, available from 1993 onward ([11]). For SARS-CoV-2, we used test-adjusted case counts from Ontario's Case and Contact Management System (CCM) ([12]) from March 2020 to August 2022 (using methodology described in Fisman et al. ([13]) and Bosco et al. ([14])) and percent positivity from FluWatch from September 2022 to August 2024. Mortality data were obtained from Statistics Canada (1991-1993) and the Ontario Deaths Registry (1994-2024) and ([15], [16]). Population denominators were obtained from Statistics Canada ([17]). Individual-level CCM data are not publicly available; all other data sources are public, and aggregated monthly data used in this analysis are provided in a public repository (see Data Availability).

Analysis

All analyses were conducted using Stata version 18. To estimate the number of deaths attributable to seasonal respiratory viruses and SARS-CoV-2, we fit regression models to monthly all-cause mortality data from Ontario, incorporating indicators of viral circulation. We modeled death counts using both Poisson and negative binomial regression, including Fast Fourier Transforms (FFT; sine and cosine terms) to account for seasonal periodicity, and linear and quadratic terms to model secular trends. Population denominators were included as a log-offset to estimate mortality rates.

Poisson model fit was assessed using Pearson and deviance goodness-of-fit tests (estat gof in Stata). Evidence of overdispersion (p < 0.001 for both tests) supported the use of negative

binomial regression for all further analysis. Given that SARS-CoV-2 did not circulate widely before March 2020, analyses for influenza A, influenza B, and RSV were stratified into pre-pandemic (prior to March 2020) and pandemic (March 2020 onwards) periods. To account for potential reporting lags in mortality data, the analysis was restricted to the period ending in March 2024.

As a sensitivity analysis, we re-estimated all models excluding Fourier terms to evaluate the robustness of viral effect estimates to the inclusion of seasonal periodicity. Seasonal adjustment terms may compete with viral predictors for explanatory power, particularly when viruses exhibit strong seasonal patterns. The reduced models were otherwise identical in structure, retaining trend terms, population offsets, and viral covariates.

To formally assess whether removal of Fourier transforms significantly altered viral effect estimates, we compared regression coefficients between primary and sensitivity models using Wald tests. Seemingly unrelated estimation (suest command in Stata) was used to jointly estimate both models and obtain the variance-covariance matrix needed for hypothesis testing. For each viral covariate, we tested the null hypothesis that coefficients were equal across models. P-values < 0.05 were considered evidence of significant modification by seasonal adjustment.

For both primary and sensitivity models, we estimated the fraction of all-cause mortality attributable to each virus by generating model-based predictions under two scenarios: one with observed viral circulation, and a counterfactual in which the coefficients for each virus (or all viruses) were set to zero. PAF was then calculated as the difference between predicted deaths under observed and counterfactual scenarios, divided by the predicted deaths under observed conditions. Confidence intervals for each PAF estimate were obtained using the punaf command,

specifying virus-specific terms to be set to zero in the counterfactual. We validated model-based estimates by comparing them to manually derived estimates from predicted mortality totals.

A secondary sensitivity analysis was conducted to evaluate the impact of using raw SARS-CoV-2 case counts instead of test-adjusted counts. Model fit was compared using the Akaike and Bayesian Information Criteria (AIC/BIC), with the model using test-adjusted counts demonstrating slightly better fit (AIC = 5145.2 vs. 5162.1). Adjusted counts were therefore retained for the primary analysis.

Meta-analysis

We conducted random-effects meta-analyses to generate pooled estimates of the population attributable fraction (PAF) for each viral pathogen individually, and for all respiratory viruses combined, stratified by time-period and model specification. For each virus or virus grouping, and for each combination of modeling approach (with or without Fourier terms) and time-period (pre-pandemic vs. intra-pandemic), we extracted PAF estimates and 95% confidence intervals from counterfactual models using the punaf command in Stata. Meta-analyses were performed using the DerSimonian and Laird method with inverse-variance weighting, as implemented via the metan command in Stata. We assessed heterogeneity using Cochran's Q test and the I² statistic. Stratum-specific estimates were visualized using forest plots.

Significant heterogeneity was observed for estimates associated with all respiratory viruses combined, as well as for influenza A and respiratory syncytial virus (RSV). To explore sources of heterogeneity, we conducted subgroup meta-analyses stratified by inclusion of Fourier terms and by pandemic period, holding the other factor constant. P-values for between-group heterogeneity were calculated for each comparison.

Ethics and Data Availability

All data used in this study were aggregate-level, with viral positivity data and monthly mortality being publicly available. All data and analysis code are available at https://github.com/fismanda/respiratory-virus-mortality-ontario (archived at https://zenodo.org/records/17887720). Approval for use of aggregate level, test-adjusted SARS-CoV-2 case counts was granted by the University of Toronto Research Ethics Board (Protocol # 41690).

RESULTS

Model Fit

Analyses were stratified into pre-pandemic (January 1993–February 2020) and pandemic (March 2020–March 2024) periods. Negative binomial regression demonstrated superior fit to Poisson models (p < 0.001 for overdispersion), and the pandemic period itself was associated with elevated all-cause mortality (IRR 1.050, 95% CI 1.005-1.084, P = 0.027). Mortality and virological surveillance data are shown in **Figure 1**.

Primary Model Results

Separate negative binomial models estimated mortality attributable to respiratory viruses in pre-pandemic and pandemic periods (**Table 1**). In pre-pandemic models (Table 1), only influenza A predicted mortality (IRR 1.0037, 95% CI 1.0028-1.0046). Influenza B did not predict mortality, and RSV showed a non-significant inverse association (IRR 0.9992, 95% CI 0.9979-1.0005, P=0.238). During the pandemic period (March 2020–March 2024), only SARS-CoV-2 activity predicted mortality. Both SARS-CoV-2 percent positivity (IRR 1.0073, 95% CI 1.0039-1.0106) and adjusted case counts (IRR 1.0029 per 10,000 cases, 95% CI 1.0022-1.0037) predicted mortality. Influenza viruses and RSV did not show significant associations with mortality in the pandemic-period models.

Sensitivity Analysis---Models Without Seasonal Adjustment

To evaluate whether seasonal adjustment terms competed with viral predictors for explanatory power, we re-estimated all models excluding Fourier transforms (Table 2). Formal comparison of coefficients using Wald tests revealed substantial differences for viruses with strong seasonal patterns. In the pre-pandemic period, removal of Fourier transforms significantly altered estimates for influenza A and RSV. The influenza A incidence rate ratio increased from

1.0037 to 1.0060 (Wald P<0.001). Most notably, RSV shifted from a counterintuitive non-significant inverse association (IRR 0.9992) to a positive association (IRR 1.0033, 95% CI 1.0023–1.0042, P<0.001), with this reversal being highly significant (Wald P<0.001). Influenza B estimates were unchanged (Wald P=0.497). During the pandemic period, the early SARS-CoV-2 indicator (adjusted cases) increased from 1.0029 to 1.0035 (Wald P=0.042). RSV again demonstrated a significant difference, with the IRR increasing from 1.0032 (P=0.133) to 1.0073 (95% CI 1.0030–1.0116, P=0.001; Wald P=0.006). Other viral predictors did not differ significantly between models.

In a sensitivity analysis using unadjusted SARS-CoV-2 case counts in place of test-adjusted counts, model fit worsened (AIC = 5162.1 vs. 5145.2), supporting the use of test-adjusted data for estimating attributable mortality.

Population Attributable fraction

Using the primary models with seasonal adjustment (Table 1), we calculated the population attributable fraction for each virus. In the pre-pandemic period, influenza A accounted for 1.8% (95% CI 1.4–2.3%) of all-cause mortality. During the pandemic period, SARS-CoV-2 accounted for 6.5% (95% CI 4.5–8.4%) of all-cause mortality. The total number of SARS-CoV-2–attributable deaths predicted by our model (18,052) closely matched the number of COVID-19 deaths reported by Public Health Ontario over the same interval (18,603), providing external validation of our approach. Combined respiratory viruses accounted for 1.5% of mortality pre-pandemic and 7.6% during the pandemic. **Figure 2a** illustrates the temporal patterns of virus-attributable mortality, demonstrating that pre-pandemic combined viral mortality closely tracked influenza A, while pandemic-era combined viral mortality was dominated by SARS-CoV-2.

In models without seasonal adjustment (**Table 2**), attributable fractions were substantially higher. Pre-pandemic, influenza A accounted for 2.9% (95% CI 2.5–3.3%) and RSV for 1.9% (95% CI 1.3–2.4%) of all-cause mortality; combined respiratory viruses accounted for 4.9% (95% CI 4.3–5.4%). The contrast between RSV estimates with and without seasonal adjustment is shown in **Figure 2b**. During the pandemic, SARS-CoV-2 accounted for 8.0% (95% CI 5.7–10.3%) and RSV for 1.8% (95% CI 0.7–2.8%) of mortality; combined respiratory viruses accounted for 10.6% (95% CI 8.0–13.1%).

Meta-Analysis of Population Attributable fraction

Random-effects meta-analyses were conducted to generate pooled estimates stratified by modeling approach (with or without Fourier transforms) and time period (pre-pandemic vs. pandemic) (**Figure 3**). For all respiratory viruses combined, pooled PAF ranged from 2% (95% CI 0.0-0.03) with seasonal adjustment in the pre-pandemic period to 11% (95% CI 0.08-0.13) without seasonal adjustment during the pandemic, with substantial heterogeneity (I^2=95.0%, P<0.001). Influenza A demonstrated high heterogeneity across strata (I^2=93.7%, P<0.001). Pooled estimates ranged from essentially zero during the pandemic with seasonal adjustment (0%, 95% CI -0.00-0.01) to 3% (95% CI 0.02-0.03) in the pre-pandemic period without seasonal adjustment. The pooled estimate across all strata was 2% (95% CI 0.00–0.03).

Influenza B showed no evidence of mortality association in any stratum, with pooled PAF of 0% (95% CI -0.00–0.00) and no heterogeneity (I^2=0.0%, P=0.932). RSV demonstrated substantial heterogeneity (I^2=88.5%, P<0.001). Without seasonal adjustment, RSV accounted for approximately 2% of mortality in both pre-pandemic (PAF 0.02, 95% CI 0.01-0.02) and pandemic (PAF 0.02, 95% CI 0.01-0.03) periods. With seasonal adjustment, estimates approached zero or were slightly negative (pre-pandemic: -0.00, 95% CI -0.01-0.00). The pooled

estimate across all strata was 1% (95% CI -0.00-0.02). SARS-CoV-2 showed minimal

heterogeneity (I^2=2.5%, P=0.311) despite different modeling approaches. The pooled PAF was

7% (95% CI 0.06–0.09), ranging from 6% with seasonal adjustment to 8% without.

Discussion

In this study, we used more than three decades of mortality and virological surveillance data from Ontario to quantify the proportion of all-cause mortality attributable to major respiratory viruses. Prior to the emergence of SARS-CoV-2, influenza A was the only virus with a consistent positive association with all-cause mortality after adjusting for seasonal and secular trends, with a PAF of 1.8% (95% CI 1.4–2.3%), consistent with prior studies ([18], [19]). During the pandemic period (March 2020–March 2024), SARS-CoV-2 fundamentally transformed the landscape of respiratory virus mortality, with a PAF of 6.5% (95% CI 4.5–8.4%) despite the availability of vaccines, antivirals, and non-pharmaceutical interventions. This represents a mortality burden 3-4 times higher than pre-pandemic influenza A, a virus that itself caused thousands of deaths annually, with model-predicted SARS-CoV-2-attributable deaths (18,052) closely matching reported COVID-19 deaths (18,603), supporting the validity of our approach. In contrast, influenza A, influenza B, and RSV did not show significant associations with mortality during the pandemic period, consistent with the well-documented suppression of seasonal respiratory virus circulation due to SARS-CoV-2 dominance and the implementation of non-pharmaceutical interventions (NPIs) ([20], [21]).

A key contribution of our study is the formal comparison of models with and without Fourier-based seasonal adjustment. Seasonal terms are commonly used in time-series mortality models to account for periodic fluctuations ([10]), but because many respiratory viruses also exhibit strong seasonality, these terms may compete with viral predictors for explanatory variance. Similar patterns have been observed in excess mortality studies during the COVID-19 pandemic, where models adjusting for seasonal pathogen circulation produced different estimates than those not accounting for these factors ([22]). Our findings illustrate this tension:

removal of Fourier terms significantly increased estimated effects for pathogens with strong seasonal signatures, particularly influenza A (PAF increased from 1.8% to 2.9%, Wald P < 0.001) and RSV (shifted from PAF –0.45% to PAF 1.9%, Wald P < 0.001).

Meta-analyses revealed substantial heterogeneity for influenza A (I^2=93.7%) and RSV (I^2=88.5%) across modeling approaches, while SARS-CoV-2 showed minimal heterogeneity (I^2=2.5%), indicating a signal robust to modeling choices. Models including Fourier terms provide conservative estimates reflecting mortality variation uniquely attributable to viral indicators after accounting for shared seasonal patterns, while models without such adjustment isolate virus-associated mortality more directly but may conflate viral effects with other seasonal phenomena. The absence of a detectable RSV mortality signal in models with seasonal adjustment initially suggested a statistical artifact driven by viral interference, a well-described phenomenon where staggered epidemic peaks and innate immune responses create negative correlations between viruses ([23]), but the emergence of positive RSV associations in models without Fourier terms suggests the negative coefficients primarily reflect competition between seasonal and viral covariates rather than true protective effects.

Influenza B showed little evidence of measurable mortality impact, aligning with prior studies showing it causes milder illness with a smaller population-wide mortality footprint than influenza A ([24]). The magnitude of SARS-CoV-2's mortality impact cannot be overstated: during March 2020–March 2024, SARS-CoV-2 demonstrated 3.6-fold higher attributable mortality than pre-pandemic influenza A (6.5% vs 1.8%), despite facing countermeasures unavailable for historical influenza seasons including non-pharmaceutical interventions ([25], [26]), multiple vaccine formulations ([27]), antivirals ([28]), and substantial partial immunity from prior infection ([29]). The close concordance between model-based and reported COVID-19 deaths

likely reflects the unique features of SARS-CoV-2 surveillance: unlike influenza and RSV, which are under-attributed in routine cause-of-death coding ([10], [18]), COVID-19 deaths were subject to intensive clinical, laboratory, and public health scrutiny during this period.

Strengths of this analysis include the use of more than 30 years of mortality data, integration of laboratory surveillance indicators, stratification by pandemic period, and formal evaluation of modeling assumptions through sensitivity analyses and Wald tests. Limitations include the use of virological surveillance indicators as imperfect proxies for true infection incidence, the ecological design which cannot establish individual-level causation and assumes viral effects are independent and additive (potentially missing interactions such as interference or co-infections ([30])), and the use of all-cause mortality aggregates which mask age-specific patterns and likely underestimate mortality attributable to age-targeted pathogens such as RSV. Additionally, mortality data for recent months are subject to reporting delays, though we mitigated this by truncating analyses at March 2024.

Despite these limitations, our findings have clear public health relevance. During 2020–2024, SARS-CoV-2 emerged as an unparalleled respiratory virus threat, responsible for a mortality burden 3–4 times higher than pre-pandemic influenza A, a pathogen that itself has historically caused an estimated 3,500 Canadian deaths annually ([1]). Influenza A remains an important contributor warranting continued seasonal vaccination efforts, while RSV's consistent ~2% burden in models without seasonal adjustment and its concentration in vulnerable age groups highlights opportunities for emerging prevention strategies including vaccines for older adults and monoclonal antibodies for infants. The extraordinary mortality impact of SARS-CoV-2, persisting even with widespread vaccination and antiviral availability, demonstrates that as of 2024, this virus represented a fundamentally different magnitude of threat than seasonal

respiratory viruses and demands sustained prioritization of prevention strategies. More broadly, this study illustrates the value of long-term respiratory virus surveillance and the importance of modeling choices when quantifying virus-attributable mortality, emphasizing the need for continued investment in surveillance and analytic methods to monitor shifts in viral circulation, quantify preventable mortality burden, and guide immunization strategies.

Tables

Table 1. Mortality Incidence Rate Ratios for Respiratory Viruses Derived from Negative Binomial Models

Period	Virus	Mortality Incidence Rate Ratio (IRR)*	IRR 95% Confidence Interval	Population Attributable Fraction (PAF)	PAF 95% CI	P-value for Association †
Pre-pandemic (1993 to February 2020)	Influenza A	1.0037	1.0028–1.0046	1.821	1.389 to 2.251	<0.001
	Influenza B	1.0009	0.9992–1.0026	0.153	-0.135-0.441	0.297
	RSV	0.9992	0.9979–1.0005	-0.453	-1.210-0.299	0.238
	Combined Respiratory Viruses	---	---	1.531	0.455-2.295	0.005

Pandemic (March 2020 to March 2024)	SARS-CoV-2 (2020-2022)	1.0029	1.0022–1.0037	‡6.477	4.482-8.429	<0.001
	SARS-CoV-2 (2022-2024)	1.0073	1.0039–1.0106			<0.001
	Influenza A	1.0025	0.9985–1.0065			0.219
	Influenza B	1.0007	0.9789–1.0230			0.95
	RSV	1.0032	0.9990–1.0073			0.133
	Combined Respiratory Viruses	---	---	7.612	5.145-10.015	<0.001

NOTE: *All incidence rate ratios are per percent increase in test positivity except for SARS-CoV-2 (2020-2022), which is incidence rate ratio per 10,000 increase in adjusted cases.

† P-values for multivariable model coefficients and PAF from PUNAF are identical.

‡ SARS-CoV-2 PAF estimated by setting both early (adjusted cases) and later (percent positivity) model terms to zero.

Table 2. Mortality Incidence Rate Ratios for Respiratory Viruses Derived from Negative Binomial Models without Fast Fourier Transforms

Period	Virus	Mortality Incidence Rate Ratio (IRR)*	IRR 95% Confidence Interval	Population Attributable Fraction (PAR) (%)	PAR% 95% CI	P-value for Association †	P-value for Difference from Primary Model with FFT§
Pre-pandemic (1993 to February 2020)	Influenza A	1.0060	1.0051-1.0068	2.889	2.486-3.290	<0.001	<0.001
	Influenza B	1.0013	0.9996-1.0029	0.213	-0.069-0.495	0.139	0.497
	RSV	1.0033	1.0023-1.0042	1.865	1.331-2.396	<0.001	<0.001

	Combined Respiratory Viruses	---	---	4.873	4.337-5.406	<0.001	
Pandemic (March 2020 to March 2024)	SARS-CoV-2 (2020-2022)	1.0035	1.0026-1.0044	‡8.0304	5.733-10.271	<0.001	0.042
	SARS-CoV-2 (2022-2024)	1.0095	1.0055-1.0135			<0.001	0.077
	Influenza A	1.0043	0.9994-1.0092	0.745	-0.106-1.590	0.084	0.134
	Influenza B	1.0107	0.9862-1.0360	0.356	-0.464-1.170	0.393	0.163

RSV	1.0073	1.0030-1.0116	1.770	0.719-2.809	0.001	0.006
Combined Respiratory Viruses	---	---	10.591	8.022-13.089	<0.001	

NOTE: *All incidence rate ratios are per percent increase in test positivity except for SARS-CoV-2 (2020-2022), which is incidence rate ratio per 10,000 increase in adjusted cases. FFT, Fast Fourier Transform.

† P-values for multivariable model coefficients and PAF from PUNAF are identical.

‡ SARS-CoV-2 PAF estimated by setting both early (adjusted cases) and later (percent positivity) model terms to zero.

§ P-value from Wald test comparing coefficient to corresponding estimate in primary model with Fourier transforms (Table 1)

Figure Legends

Figure 1. Mortality and virological surveillance data for Ontario, 1993-2024.

(A) Monthly deaths per 100,000 from Ontario Deaths Registry (red) and Statistics Canada (teal), demonstrating close concordance between data sources (mean difference 0.3%). (B) Monthly percent positivity for influenza A (red) and influenza B (teal) from FluWatch surveillance, showing regular seasonal epidemics and marked suppression during 2020-2021. (C) Monthly RSV percent positivity from FluWatch surveillance, demonstrating seasonal patterns and pandemic-era disruption. (D) SARS-CoV-2 adjusted case counts per 10,000 population (red, left axis) and SARS-2 positivity percentage (teal, right axis) from March 2020 onward, showing multiple pandemic waves. Vertical dashed lines indicate emergence of the SARS-CoV-2 pandemic in Ontario in March 2020. Data truncated at March 2024 to account for reporting delays.

Figure 2. Model-estimated virus-attributable mortality over time.

(A) Monthly death rates per 100,000 population attributable to respiratory viruses from primary models with Fourier seasonal adjustment. Blue line shows all respiratory viruses combined, brown line shows influenza A, and green line shows SARS-CoV-2. Pre-pandemic, the combined viral mortality (blue) nearly overlaps with influenza A (brown), demonstrating that influenza A accounted for essentially all virus-attributable mortality. During the pandemic, combined viral mortality (blue) tracks closely with SARS-CoV-2 (green), illustrating SARS-CoV-2's dominance. (B) RSV-attributable death rates comparing models with Fourier adjustment (blue)

versus without (black). With seasonal adjustment, RSV shows frequently negative values pre-pandemic (PAF-0.45%) and substantially attenuated positive associations during the pandemic. Without seasonal adjustment, RSV demonstrates consistently positive mortality associations of similar magnitude (~1.8-1.9%) in both periods. Both panels exclude a period from 2009-2010 for which data were not available. Vertical dashed lines indicate emergence of the SARS-CoV-2 pandemic in Ontario in March 2020.

Figure 3. Meta-analysis of population attributable fraction (PAF) for respiratory viruses stratified by modeling approach and time period.

Forest plot showing pooled estimates from random-effects meta-analyses for all respiratory viruses combined, influenza A, influenza B, respiratory syncytial virus (RSV), and SARS-CoV-2. Each virus section includes stratum-specific estimates based on four combinations of modeling approach (with or without Fast Fourier Transform seasonal adjustment) and time

period (pre-pandemic: 1993–February 2020; pandemic: March 2020–March 2024). Diamonds represent point estimates with horizontal lines indicating 95% confidence intervals. Diamond size is proportional to inverse-variance weights. Pooled estimates are shown as open diamonds. I² statistics quantify heterogeneity; P-values test for significant heterogeneity across strata. SARS-CoV-2 estimates include only pandemic period strata as this virus did not circulate before March 2020. FFT, Fast Fourier Transform; PAF, population attributable fraction.

References

References

1.   Public Health Agency of Canada. Flu (influenza): For health professionals Government of Canada2024 [updated 2024-10-07. Available from: https://www.canada.ca/en/public-health/services/diseases/flu-influenza/health-professionals.html.

2.   World Health Organization. WHO Coronavirus (COVID-19) Dashboard [Available from: https://data.who.int/dashboards/covid19/deaths?n=c&m49=124.

3.   Statistics Canada. Deaths, 2022. The Daily2023.

4.   Abrams EM, Doyon-Plourde P, Davis P, Brousseau N, Irwin A, Siu W, et al. Burden of disease of respiratory syncytial virus in infants, young children and pregnant women and people. Can Commun Dis Rep. 2024;50(1-2):1-15.

5.   ElSherif M, Andrew MK, Ye L, Ambrose A, Boivin G, Bowie W, et al. Leveraging Influenza Virus Surveillance From 2012 to 2015 to Characterize the Burden of Respiratory Syncytial Virus Disease in Canadian Adults ≥50 Years of Age Hospitalized With Acute Respiratory Illness. Open Forum Infect Dis. 2023;10(7):ofad315.

6.   Tuite AR, Simmons AE, Rudd M, Cernat A, Gebretekle GB, Yeung MW, et al. Respiratory syncytial virus vaccination strategies for older Canadian adults: a cost–utility analysis. Canadian Medical Association Journal. 2024;196(29):E989-E1005.

7.   Mansournia MA, Altman DG. Population attributable fraction. BMJ. 2018;360:k757.

8.   Lin CK, Chen ST. Estimation and application of population attributable fraction in ecological studies. Environ Health. 2019;18(1):52.

9.   Mieno MN, Tanaka N, Arai T, Kawahara T, Kuchiba A, Ishikawa S, et al. Accuracy of

Death Certificates and Assessment of Factors for Misclassification of Underlying Cause of

Death. J Epidemiol. 2016;26(4):191-8.

10.   Zhou H, Thompson WW, Viboud CG, Ringholz CM, Cheng PY, Steiner C, et al.

Hospitalizations associated with influenza and respiratory syncytial virus in the United States,

1993-2008. Clin Infect Dis. 2012;54(10):1427-36.

11.   Public Health Agency of Canada. Overview of influenza monitoring in Canada

Government of Canada2024 [Available from: https://www.canada.ca/en/public-

health/services/diseases/flu-influenza/influenza-surveillance/about-fluwatch.html.

12.   Simcoe Muskoka District Health Unit. Case and Contact Management System (CCM)

HealthSTATS SIMCOE MUSKOKA [Available from:

https://www.simcoemuskokahealth.org/Health-

Stats/HealthStatsHome/Resources/DataSources/CCM.

13.   Fisman DN, Greer AL, Brankston G, Hillmer M, O'Brien SF, Drews SJ, et al. COVID-19

Case Age Distribution: Correction for Differential Testing by Age. Ann Intern Med. 2021.

14.   Bosco S, Peng A, Tuite AR, Simmons A, Fisman DN. Impact of adjustment for

differential testing by age and sex on apparent epidemiology of SARS-CoV- 2 infection in

Ontario, Canada. BMC Infect Dis. 2025;25(1):589.

15.   Statistics Canada. Statistics Canada. Table 13-10-0708-01: Deaths, by month [Internet].

Government of Canada. Available from:

https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=1310070801. Last accessed July 3, 2025.

2025.

16.   Government of Ontario. Vital events data by month [Internet]. Ontario Data Catalogue; Available from: https://data.ontario.ca/en/dataset/vital-events-data-by-month/resource/97622ce6-c06a-4970-afe5-be540c748f24. Last accessed July 3, 2025. 2025.

17.   Statistics Canada. Table 17-10-0009-01: Population estimates, quarterly [Internet]. Government of Canada; Available from: https://doi.org/10.25318/1710000901-eng. Last accessed July 3, 2025. 2025.

18.   Reichert TA, Simonsen L, Sharma A, Pardo SA, Fedson DS, Miller MA. Influenza and the winter increase in mortality in the United States, 1959-1999. Am J Epidemiol. 2004;160(5):492-502.

19.   Goldstein E, Viboud C, Charu V, Lipsitch M. Improving the estimation of influenza-related mortality over a seasonal baseline. Epidemiology. 2012;23(6):829-38.

20.   Song S, Li Q, Shen L, Sun M, Yang Z, Wang N, et al. From Outbreak to Near Disappearance: How Did Non-pharmaceutical Interventions Against COVID-19 Affect the Transmission of Influenza Virus? Front Public Health. 2022;10:863522.

21.   Avolio M, Venturini S, De Rosa R, Crapis M, Basaglia G. Epidemiology of respiratory virus before and during COVID-19 pandemic. Infez Med. 2022;30(1):104-8.

22.   Barnard S, Chiavenna C, Fox S, Charlett A, Waller Z, Andrews N, et al. Methods for modelling excess mortality across England during the COVID-19 pandemic. Stat Methods Med Res. 2022;31(9):1790-802.

23.   Li K, Thindwa D, Weinberger DM, Pitzer VE. The Role of Viral Interference in Shaping RSV Epidemics Following the 2009 H1N1 Influenza Pandemic. Influenza Other Respir Viruses. 2025;19(4):e70111.

24.   Tran D, Vaudry W, Moore D, Bettinger JA, Halperin SA, Scheifele DW, et al. Hospitalization for Influenza A Versus B. Pediatrics. 2016;138(3).

25.   Mendez-Brito A, El Bcheraoui C, Pozo-Martin F. Systematic review of empirical studies comparing the effectiveness of non-pharmaceutical interventions against COVID-19. J Infect. 2021;83(3):281-93.

26.   Peng A, Bosco S, Simmons AE, Tuite AR, Fisman DN. Impact of community mask mandates on SARS-CoV-2 transmission in Ontario after adjustment for differential testing by age and sex. PNAS Nexus. 2024;3(2):pgae065.

27.   Fisman DN, Simmons AE, Tuite AR. Case-cohort design as an efficient approach to evaluating COVID-19 vaccine effectiveness, waning, heterologous immune effect and optimal dosing interval. Vaccine. 2024;42(25):126134.

28.   Faust JS, Kumar A, Shah J, Khadke S, Dani SS, Ganatra S, et al. Oral Nirmatrelvir and Ritonavir for Coronavirus Disease 2019 in Vaccinated, Nonhospitalized Adults Aged 18-50 Years. Clin Infect Dis. 2023;77(9):1257-64.

29.   Goldberg Y, Mandel M, Bar-On YM, Bodenheimer O, Freedman LS, Ash N, et al. Protection and Waning of Natural and Hybrid Immunity to SARS-CoV-2. N Engl J Med. 2022;386(23):2201-12.

30.   Simmons AE, Berry I, Buchan SA, Tuite AR, Fisman DN. Association Between Seasonal Respiratory Virus Activity and Invasive Pneumococcal Disease in Central Ontario, Canada. medRxiv. 2024:2024.09.03.24312990.