1 **Title**: Synchrony between brain age and internalising and externalising symptoms across
2 adolescence
3

4 **Authors**: Dani Beck[1]*, Chloe Carrick[2], Eira R. Aksnes[1,3], Niamh MacSweeney[1,3], Lars T.
5 Westlye[4,5,6], Delia Fuhrmann[2], Christian K. Tamnes[1,3]
6
7
8
[1] PROMENTA Research Center, Department of Psychology, University of Oslo, Oslo,
Norway
[2] Department of Psychology, Institute of Psychiatry, Psychology, and Neuroscience, King's
College London, London, UK
[3] Division of Mental Health and Substance Abuse, Diakonhjemmet Hospital, Oslo, Norway
[4] Department of Psychology, University of Oslo, Oslo, Norway
[5] Center for Precision Psychiatry, Division of Mental Health and Addiction, Oslo University Hospital,
Oslo, Norway
[6] KG Jebsen Centre for Neurodevelopmental Disorders, University of Oslo, Oslo, Norway
24 * Corresponding author: Dani Beck (dani.beck@psykologi.uio.no)

Abstract

Background: Adolescence is a period of rapid neurobiological and behavioural change, yet it remains unclear how deviations from normative brain maturation relate to the development of internalising and externalising symptoms.

Methods: Using data from the Adolescent Brain Cognitive Development (ABCD) Study, we combined brain age prediction with bivariate latent growth curve (BLGC) models to test whether deviations in brain maturation – indexed by the brain age gap (BAG) – relate to mental health development across late childhood and adolescence. Brain age was estimated using T1-weighted, diffusion (dMRI), resting-state fMRI, and multimodal MRI data across four waves (ages ~8.3–17.5). Internalising and externalising symptoms were assessed across ten waves with the self-report Brief Problem Monitor (BPM).

Results: Across T1, dMRI, and multimodal models, deviations from age-expected brain maturation and internalising and externalising symptoms showed coordinated nonlinear development across adolescence. Adolescents whose brains increasingly diverged from age-expected maturation over time also showed increasing symptom trajectories. These associations were small to moderate in magnitude and were most consistent for internalising symptoms in females ($r = .15–.23$), whereas externalising symptoms showed broader but less selective nonlinear associations across modalities in both males and females ($r = .15–.23$). Intercept-level associations were weaker and modality-specific ($r = .06–.11$). Formal tests provided no evidence for robust sex differences in these associations after correction for multiple comparisons.

Conclusion: These results indicate that adolescent development of mental health problems is more strongly linked to nonlinear changes in how individuals diverge from age-expected brain trajectories, rather than to fixed differences in brain age. Shifts in brain maturational tempo may therefore be a key feature underlying vulnerability to psychopathology in youth.

1. Introduction

Internalising and externalising symptoms in childhood and adolescence predict later mental health problems[1], and have been linked to a wide range of neurobiological alterations in youth[2]. Several neuroimaging studies have reported attenuated or delayed cortical maturation in youth with elevated symptoms, including reduced thinning or slower surface-area development[2–9]. Conversely, other work finds accelerated cortical change, including increased thinning or volume loss[3,10,11]. Evidence from subcortical[3,12] and white matter development[13,14] is similarly mixed, ranging from attenuated to accelerated maturational trajectories, and with some studies reporting no robust associations[15]. Functional neuroimaging findings show altered responsivity in affective and reward circuits accompanying elevated symptoms and, in some cases, preceding later symptom progression[16–21]. Adolescence is also a period of rising sex differences, with higher internalising symptom burden in females and differences in the timing and tempo of normative brain maturation[22–24]. Together, these findings suggest that deviations from normative brain maturation trajectories – whether accelerated or delayed – may be centrally implicated in the emergence of youth psychopathology[25].

From a developmental perspective, both accelerated and delayed patterns of brain maturation may confer risk through altered developmental timing. In particular, risk may emerge when neural systems develop out of synchrony with each other or with environmental and socio-emotional demands, potentially increasing vulnerability during sensitive periods of development[26]. Neurodevelopment is highly heterochronous throughout childhood and adolescence, with regional and network-specific trajectories unfolding at different rates and with different sensitivity to environmental and biological influences[27,28]. As a result, deviations between chronological age and neurobiological maturity are not inherently pathological but may reflect individual differences in the timing and coordination of developmental processes. A "more mature" brain at a given chronological age may therefore index earlier maturation of specific systems, altered pruning or myelination schedules, or shifts in functional integration. Such changes may interact with emotional and behavioural regulation during development.

Several theoretical frameworks suggest that such maturational deviations could be relevant for psychopathology. The stress-acceleration model proposes that early adversity or heightened environmental demands may hasten aspects of neurobiological development[29], potentially conferring short-term adaptive advantages but increasing vulnerability to

internalising or externalising problems when regulatory systems mature out of synchrony.

Relatedly, developmental mismatch accounts emphasise that earlier maturation of affective or

motivational systems relative to cognitive control networks may bias behaviour toward

heightened emotional reactivity or impulsivity[30]. From such perspectives, deviation from age-

expected brain development may capture individual differences in developmental timing that

intersect with known pathways to internalising and externalising symptoms across

adolescence.

One approach to quantify maturational deviations involves brain age prediction

models, which estimate an individual's age from multivariate magnetic resonance imaging

(MRI) features[31,32]. The brain age gap (BAG) – predicted minus chronological age – provides

an interpretable proxy for individual differences in neurodevelopmental timing, with positive

BAG values indicating an older-appearing brain relative to age and negative values indicating

a younger-appearing brain. In adults, BAG has been linked to multiple contextual factors and

outcomes, including socioeconomic status[33], cardiometabolic risk and lifestyle factors[34–40],

cognition[41], and mental health and brain disorders[42–45]. In youth, deviations from age-

expected brain maturation have also been associated with several mental health profiles.

Negative BAG values have been linked to higher anxiety and depression symptoms, attention

difficulties, autism spectrum traits, and elevated internalising and externalising symptom

scores[46–51]. Positive BAG values, in contrast, have been associated with greater depressive

and obsessive–compulsive symptoms, higher general psychopathology, and increased

psychosis risk in clinically vulnerable youth[52–55]. Longitudinal evidence, although limited,

suggests that BAG trajectories may carry prognostic significance for later mental health

outcomes[56,57].

Taken together, discrepant findings suggest that BAG–psychopathology associations

in youth may be developmentally heterogeneous, potentially depending on when brain

maturation is assessed, which MRI modalities are indexed, and whether analyses capture

static deviation or change over time. Notably, associations have been reported at both positive

and negative BAG values for some symptom domains, potentially indicating that deviation

from age-expected brain maturation – rather than acceleration or delay in a single direction –

may be relevant for psychopathology risk in youth. Because cross-sectional BAG estimates

conflate stable between-person differences with time-varying deviations, they cannot

determine whether symptoms relate to persistent offsets, coordinated change, or divergence

across development[58]. Addressing these gaps requires longitudinal, multimodal designs and

modelling approaches that can characterise nonlinear, person-specific developmental

dynamics and clarify whether brain–behaviour coupling reflects persistent offsets,

coordinated change, or divergence across development.

To address these challenges, we leverage the Adolescent Brain Cognitive

Development (ABCD) Study (release 6.0) to build T1-weighted, diffusion MRI (dMRI), and

resting-state functional (rs-fMRI) brain age models, as well as a multimodal model

combining features from all three modalities. Focusing on internalising and externalising

symptoms as broad, transdiagnostic dimensions with well-characterised developmental

trajectories, we used the cohort's four MRI waves and ten Brief Problem Monitor (BPM)

waves to examine how developmental trajectories in BAG relate to trajectories of self-

reported internalising and externalising symptoms. Using bivariate latent growth curve

(BLGC) models, we examined (i) individual differences in BAG at the midpoint of the

observation window, as well as linear and quadratic rates of BAG change across late

childhood and adolescence and (ii) the degree to which these developmental components co-

vary with the corresponding growth components of internalising and externalising symptoms.

We hypothesised that trajectories of BAG and internalising and externalising

symptoms would be developmentally coupled across adolescence, such that individual

differences in change over time in BAG would be associated with corresponding differences

in symptom trajectories. Given mixed prior evidence linking both accelerated and delayed

patterns of brain maturation to youth psychopathology, and the relative scarcity of

longitudinal BAG studies, we did not specify a directional hypothesis regarding whether

older- or younger-appearing brains would be associated with more adverse symptom

trajectories. We further tested whether the magnitude and form of BAG–symptom coupling

differed by sex, motivated by established sex differences in the prevalence and development

of internalising and externalising symptoms and in normative brain maturation across

adolescence[24,59]. By combining multimodal brain age prediction with longitudinal growth-

curve modelling, this study addresses a key gap in understanding how the tempo of brain

maturation co-evolves with internalising and externalising symptom trajectories across

adolescence.

2. Methodology

2.1. Sample and ethical approvals

The Adolescent Brain Cognitive Development (ABCD) Study ®[60] is an ongoing longitudinal

study comprising ~11,800 children and adolescents. Participants were recruited from 21 sites

across the United States via outreach to caregivers through local schools, followed by

telephone screening to determine eligibility. Participants were excluded using the ABCD

Study exclusion criteria listed elsewhere[61]. Data used in the present study were drawn from

the ABCD Study release 6.0, containing data from baseline up until the six-year visit (nbdc-

datahub.org/abcd-release-6-0). All ABCD Study data is stored in the NIH Brain Development

Cohorts (NBDC) Data Hub, which is available for authorised users with approved Data Use

Certification (DUC) (lead investigator: Westlye). The 6.0 release has been assigned the DOI

10.82525/jy7n-g441. The Institutional Review Board (IRB) at the University of California,

San Diego, approved all aspects of the ABCD Study[62]. Parents or guardians provided written

consent, while the child provided written assent. The current study was conducted in line with

the Declaration of Helsinki and was approved by the Norwegian Regional Committee for

Medical and Health Research Ethics (REK 2019/943).

2.2. Demographic information and data quality assurance

Our sample consisted of participants drawn from a hold-out (test) sets of each brain age

prediction model from baseline, two-year, four-year, and six-year follow-up waves. T1

included 14,838 observations (male = 7,907 [53.3%], female = 6,931 [46.7%], unique N =

5866), dMRI 12,975 (male = 6,853 [52.8%], female = 6,122 [47.2%], unique N = 5112), rs-

fMRI 12,990 (male = 6,931 [53.4%], female = 6,059 [46.6%], unique N = 5562), and

multimodal 11,770 (male = 6139 [52.2%], female = 5,631 [47.8%], unique N = 4934). The

overall mean age was ~12.3 years (SD ≈ 2.3; range ≈ 8.3–17.7). These MRI observations

were then linked to all available mental health BPM assessments for the same individuals,

with up to ten data waves. Demographic information for each brain MRI modality-specific

test sample can be found in SI Table 1. Comparable distributions for the brain age prediction

training sets can be found SI Table 2, while the full analytic sample is illustrated in Figure 1.

Study sample age distribution

Figure 1. **Age distributions across MRI and BPM waves for the analytic sample.** Density plots
and boxplots show participant age distributions across the ten assessment waves. At Wave 1, two
distributions are shown because MRI Wave 1 (ses-00A) and BPM Wave 1 (ses-00M) occurred at
different visits. Waves 2–10 correspond to session identifier: ses-01A, ses-01M, ses-02A, ses-02M,
ses-03A, ses-03M, ses-04A, ses-05A, and ses-06A, respectively. Beige densities/boxplots represent
BPM-only waves, whereas blue indicates waves in which participants provided both BPM and MRI
data at the same ABCD session (except wave 1). Boxplots summarise the median and interquartile
range, and density plots show the full distributional shape.

2.3. MRI acquisition and processing

Neuroimaging data were acquired at 21 different sites (using 34 scanners) and processed by

the ABCD Study team. A 3-T Siemens Prisma, General Electric 750, or Philips scanner was

used for data acquisition. Protocols used for data acquisition and processing are described in

detail elsewhere[60,63] and described in SI Section 1. Three modalities of brain structural and

functional measures were used in the present study: structural grey matter measures (T1),

white matter microstructural measures from diffusion tensor imaging (DTI), and resting-state

functional connectivity measures (rs-fMRI)[63]. Cortical surface reconstruction and subcortical

segmentation was performed with FreeSurfer v7.1.1[64,65]. White matter microstructural

measures were generated using AtlasTrack, a probabilistic atlas-based method for automated

segmentation of white matter fibre tracts[66]. Measures of functional connectivity were

computed using a seed-based, correlational approach[67], where average time courses were

calculated for cortical surface-based ROIs using a functionally-defined parcellation based on

resting-state functional connectivity patterns[68] and subcortical ROIs[65].

For T1, we extracted tabulated total and regional measures of cortical surface area,

thickness, volume, sulcal depth, and intensity grey/white matter contrast, as well as

subcortical volume (397 measures). For DTI, full and inner (multi) shell tissue properties

including fractional anisotropy (FA) and mean (MD), longitudinal (or axial, AD), and

transverse (or radial, RD) diffusivity were extracted for total and regional features (576

measures). For rs-fMRI, functional connectivity within and between parcellations from the

Gordon network, including subcortical data,[68] were extracted (416 measures). MRI data

underwent quality assurance (QA) per standard procedures outlined elsewhere[63]. Briefly,

participants with excessive head motion or poor data quality were excluded from the curated

data release, in additional to removal of data using the recommendations for data cleaning

provided by the ABCD Study team. Following QA procedures, harmonisation of multi-

scanner effects was carried out using *longComBat*[69] (see SI Section 2 and SI Figures 1-3).

2.4. Brain age prediction

Brain age prediction was carried out using the eXtreme Gradient Boosting (XGBoost)

regression model[70]. Parameters were tuned using five-fold cross-validation (stratified by age)

with two-repeats to average out partition noise. The models were fitted using the best

estimators and optimised models were applied to the (hold-out) test sample. R^2, RMSE, and

MAE were calculated to evaluate prediction accuracy in the test set. For each brain modality

(T1, DTI, rs-fMRI, and multimodal; combining features from all three modalities), 50% of

the data was used as the hold-out test sample and 50% was used for model training and

validation, with all observations from a given individual assigned to the same split.

Consistent with recent ABCD brain age work[57,71,72], confounding effects from

complex family-related factors were minimised using a group shuffle split with family ID as

the group indicator to ensure that siblings were not split across training and test sets. The

brain age model used for the study is publicly available: https://github.com/dani-beck/brain

age-prediction. A detailed overview of T1, DTI, rs-fMRI, and multimodal training and test

samples, including demographic information, are provided in SI Section 3 and SI Tables 1

and 2. Complete lists of all the extracted measures used for brain age prediction per modality

are provided in SI Tables 3-5. Feature importance scores for each model are provided in SI

Figure 4. To adjust for commonly observed age-bias (overestimated predictions for younger

participants and underestimated predictions for older participants)[73], we applied a statistical

correction previously described[74] that involves fitting a linear regression of predicted age on

chronological age and using the residuals. This is visualised in SI Figure 5. The difference

between an individual's predicted brain age and their chronological age (BAG) was

calculated (BAG = predicted age - chronological age) for each of the models, providing T1,

DTI, rs-fMRI, and multimodal-based BAG values for all participants.

2.5. Brief Problem Monitor measures

Youth internalising and externalising problems were measured using the self-report version of

the Brief Problem Monitor (BPM)[75]. The BPM is a brief 19-item instrument derived from the

longer Child Behaviour Checklist (CBCL) for ages 6-18[76], Teacher's Report Form, and Youth

Self-Report[75], and is designed for repeated monitoring of emotional and behavioural

functioning. Prior work indicates that the BPM has good internal consistency (Cronbach's α =

.78–.91 across scales) and strong convergent validity with the full CBCL (total score r = .95;

subscale correlations r = .86–.97), supporting its reliability and validity as a brief assessment

tool in community and clinical samples[77]. Each item is rated as *0 = not true, 1 = somewhat*

true, or 2 = very true over the past week, and sum scores form subscales for internalising and

externalising (and attention) problems. In the present study, ten measurement waves were

available: 6 months after baseline, followed by semi-annual collection until the final two

waves which were annual. For information pertaining to quality control and missing data, see

SI Section 4 and SI Figure 6.

2.6. Statistical analysis

All analyses were carried out using R version 4.3.2[78]. Longitudinal associations between

brain-age gap (BAG) trajectories and internalising and externalising symptom development

were examined using bivariate latent growth curve (BLGC) models estimated in *lavaan*

(version 0.6-20)[79,80]. Latent growth curve models capture development by treating its model

parameters (i.e., the intercept, slope, and where applicable quadratic) as latent variables, and

using factor loadings to indicate the association of each parameter with the observed

timepoints[81,82].

For each imaging modality (T1, dMRI, rs-fMRI, multimodal), we compared linear,

latent-basis, and quadratic growth specifications and selected the most appropriate form

based on model fit and convergence. Quadratic models provided a parsimonious and stable

representation of nonlinear change for all modalities except rs-fMRI, for which change was

adequately captured by a linear slope due to convergence instability in nonlinear

specifications (see SI Section 5 for full model selection procedure). Developmental coupling

was evaluated via cross-process covariances between corresponding BAG and symptom

growth factors (intercept–intercept, slope–slope, and quadratic–quadratic where supported),

indexing associations at the centred time point and coordinated change across the observation

window. Although the multivariate growth model estimates a broader covariance structure

among growth factors, interpretation focused on these corresponding BAG–symptom

covariances as primary indices of developmental synchrony.

All BLGC models were estimated as multigroup models by sex, using robust

maximum likelihood (MLR) with full-information maximum likelihood (FIML) for missing

data. Sex was handled in two stages. First, to determine whether males and females could be

modelled with a shared versus sex-specific latent covariance structure, we compared freely

estimated multigroup models to models in which all latent covariances were constrained

equal across groups. Scaled Satorra–Bentler χ² difference tests were used to evaluate these

omnibus constraints, with significant results indicating that at least one latent covariance

differed by sex; this step was used to guide model specification rather than to draw

conclusions about specific BAG–symptom associations. Second, we tested sex moderation of

the BAG–symptom couplings using targeted likelihood-ratio tests that constrained each

cross-process coupling to be equal across sex one at a time, with multiple-comparison control

using the Benjamini–Hochberg[83] false discovery rate (FDR) procedure.

Model fit was assessed using the χ^2 test, the Root Mean Square Error of

Approximation (RMSEA), the Comparative Fit Index (CFI), and the Standardised Root Mean

Square Residual (SRMR). Models were defined as having good fit if CFI > .97, SRMR < .05,

and RMSEA <.05. Acceptable fit was defined as CFI = .95-.97, SRMR = .05-.10, and

RMSEA = .05-.08[84]. We characterised standardised parameter estimates above 0.10 as small

effects, between 0.20 and 0.30 as moderate, and over 0.30 as large effect sizes[85].

3. Results

3.1. Brain age prediction model performance

Model performance metrics for each brain age prediction model are provided in Table 1. The

correlation between predicted and chronological age for each model is illustrated in Figure 2.

SI Figures 7-10 show raw, corrected, age-binned, and per-wave distributions of corrected

BAG per brain age model. SI Figure 11 shows within-person change and longitudinal

stability of corrected BAG using intercept intraclass correlations (ICC), per modality. SI

Figure 12 illustrates associations between raw and corrected BAG values for all modalities.

Table 1. Model performance metrics (r, r^2, *RMSE, MAE*) for each brain age prediction model in the hold-out (test) sample.

	T1	dMRI	rs-fMRI	Multimodal
Pearson's r	0.83 [0.82, 0.83]	0.86 [0.86, 0.87]	0.69 [0.68, 0.70]	0.88 [0.87, 0.88]
r^2	0.69	0.75	0.47	0.78
RMSE	1.28	1.15	1.67	1.09
MAE	1.00	0.90	1.34	0.85

Note: RMSE = root mean squared error, MAE = mean absolute error.

Figure 2. Predicted age as a function of chronological age. Scatter plots show the correlation (Pearson's r) between predicted age and chronological age for each modality (T1, dMRI, rs-fMRI, multimodal MRI) across four waves (session).

3.2. Bivariate latent growth curve modelling framework and model fit

Across imaging modalities, the quadratic bivariate latent growth curve (BLGC) models provided acceptable-to-good fit (CFI = .95–.98). Full model fit indices are provided in SI Table 6. Multigroup tests indicated that latent covariances differed by sex for all modality-outcome models except dMRI–externalising, for which equality constraints were supported (SI Table 7). We therefore retained sex-stratified models for all other analyses and treated the dMRI–externalising model as sex-invariant.

All BAG–symptom associations reported below reflect standardised latent covariances from these final models. We focus interpretation on the corresponding cross-process covariances (intercept–intercept, slope–slope, and quadratic–quadratic, where applicable) as primary parameters of interest. Intercept–intercept covariances reflect associations between BAG and symptom levels at the centred time point. Linear slope–slope covariances index coordinated linear change over time, such that individuals showing steeper increases in BAG also show steeper increases in symptoms. Quadratic slope–slope covariances index coordinated nonlinear change across the modelled window, indicating that

individuals exhibiting greater acceleration (i.e., curvature) in BAG trajectories across

adolescence also exhibit greater acceleration in symptom trajectories over the same period.

The full results of the final eight multigroup BLGC models, including all the model

parameters, are available in SI Table 8. The results of the parameters of interest are

summarised in SI Table 9 and visualised in Figures 3 and 4. For symptom trajectories over

time at low versus high BAG and coupling scatter plots between BAG and symptom growth

components, see SI Figure 13 and 14.

3.2.1. Longitudinal coupling between brain age gap and internalising symptoms

All significant BAG–internalising couplings were positive in direction, and associations

appeared primarily in the quadratic domain. In the T1 model, the quadratic component of

BAG was positively coupled with the quadratic component of internalising in females ($r =$

$.15$, $p = .010$), while the corresponding estimate in males was not statistically significant. The

dMRI model showed a similar pattern, with a significant quadratic BAG–internalising

coupling in females ($r = .21$, $p < .01$) but not in males. In the multimodal model, the quadratic

component of BAG was again significantly coupled with the quadratic component of

internalising in females ($r = .23$, $p < .001$), whereas the male estimate was not statistically

significant. Together, these results suggest that greater acceleration in BAG trajectories (i.e.,

increasing deviation from age-expected maturation) tended to co-occur with greater

acceleration in internalising symptoms in females over adolescence.

A small intercept–intercept association was observed for males in the multimodal

internalising model ($r = .06$, $p = .04$), with no corresponding effect in females. For rs-fMRI,

where BAG was modelled linearly, a single slope coupling was observed in females ($r = .20$,

$p = .02$), indicating that steeper increases in rs-fMRI BAG were modestly associated with

steeper increases in internalising symptoms; no other BAG–internalising couplings were

significant in the rs-fMRI model. Importantly, although several estimates were significant in

one sex but not the other, targeted coupling-specific likelihood-ratio tests did not indicate

reliable sex differences in these BAG–internalising couplings (Section 3.2.3; SI Table 10).

3.2.2. Longitudinal coupling between brain age gap and externalising symptoms

Several modalities showed significant BAG–externalising associations at the level of

symptom and BAG intercepts. Intercept–intercept couplings between BAG and externalising

symptoms were observed in the multimodal model for both females ($r = .06$, $p = .03$) and

males ($r = .07$, $p = .02$). In the T1 model, an intercept–intercept association was observed in

females ($r = .08$, $p < .01$) but not in males, and in the rs-fMRI model an intercept–intercept

association was likewise observed in females ($r = .09$, $p = .010$) but not males. In the

multimodal externalising model, the linear slope coupling between BAG and externalising

was significant for both females ($r = .10$, $p = .02$) and males ($r = .11$, $p = .03$), indicating that

greater increases in externalising symptoms were associated with greater increases in BAG.

Associations between the quadratic components of BAG and externalising symptom

trajectories were stronger in terms of effect sizes. In the T1 model, the quadratic component

of BAG was coupled with the quadratic component of externalising for both females ($r = .32$,

$p < .001$) and males ($r = .24$, $p < .01$). In the dMRI model, equality-constrained latent

covariances were supported; the BAG–externalising quadratic coupling was therefore

estimated as a single pooled effect across sex ($r = .15$, $p < .01$). In the multimodal model, the

quadratic BAG–externalising coupling was significant in females ($r = .24$, $p < .01$), but not

males. Finally, the rs-fMRI linear model showed no significant BAG–externalising linear

slope couplings. Overall, externalising symptoms showed consistent positive coupling with

BAG trajectories across modalities, indicating that faster or increasingly nonlinear deviations

from age-expected brain maturation tended to co-occur with faster increases in externalising

symptoms over time. As with internalising, targeted coupling-specific tests did not provide

evidence for robust sex differences in BAG–externalising couplings (Section 3.2.3; SI Table

10).

Figure 3. Standardised correlations between BAG and symptom growth factors across imaging modalities and developmental processes. Points represent sex-stratified coupling estimates for all models (female = gold; male = burgundy), except for the dMRI–externalising model, where a single pooled estimate is shown (grey). For T1, dMRI, and multimodal modalities, BAG and symptoms were modelled with centred quadratic growth factors; for rs-fMRI, only linear BAG models converged, so no quadratic coupling is shown. Columns correspond to associations among intercepts (centred levels), linear slopes (rates of change), and quadratic components (curvature). Asterisks indicate statistical significance of the within-sex BAG–symptom covariance (*$p < .05$, **$p < .01$, ***$p < .001$).

Figure 4. **Model-implied symptom trajectories conditional on brain age gap.** The figure visualises the BLGC coupling effects by plotting model-implied internalising and externalising symptom trajectories at low (−1 SD; solid lines) versus high (+1 SD; dotted lines) values of the corresponding BAG growth component. Panels are ordered to mirror the sequence of results presented in Figure 3. Trajectories are shown separately for females (gold) and males (burgundy) where sex-stratified models were estimated; for dMRI externalising, trajectories reflect pooled-sex estimates (grey), consistent with the pooled effect reported in Figure 3. For each modality × outcome coupling combination, the plotted trajectories represent expected symptom development obtained by conditioning the relevant symptom growth factor (intercept, linear slope, or quadratic component) on

417 the corresponding BAG factor, while holding all remaining latent growth factors fixed at their

418 estimated means. Because symptom trajectories are parameterised with quadratic growth functions in

419 the final models, conditional trajectories may appear curved even when visualising intercept–intercept

420 or slope–slope couplings.

422 *3.2.3. Sex differences in brain age gap–symptom coupling*

423 We conducted tests to assess whether specific cross-process couplings between brain age and

424 psychopathology differed by sex. For each imaging modality and outcome, we tested equality

425 constraints on individual latent covariances linking corresponding growth factors. At the

426 nominal level, two coupling-specific sex differences were observed, both involving quadratic

427 component associations with internalising symptoms. Specifically, the quadratic brain age–

428 internalising coupling differed by sex in the T1-weighted model ($\chi^2 = 4.42$, $p = .036$) and in

429 the diffusion MRI model ($\chi^2 = 4.41$, $p = .036$). However, neither of these effects survived

430 correction for multiple testing (p FDR = .285). No other coupling-specific sex differences

431 were observed. Thus, coupling-specific tests provided no robust evidence that the BAG–

432 symptom cross-process couplings differed by sex. The full set of sex-difference tests is

433 reported in SI Table 10.

435 **4. Discussion**

436 Adolescence is a dynamic period in which neurodevelopment and emerging psychopathology

437 unfold in tandem[23,27,86,87], making it a critical window for studying how these processes co-

438 develop. In this longitudinal study spanning late childhood through mid-adolescence, we

439 examined how developmental deviations in brain age align with changes in internalising and

440 externalising symptoms. To our knowledge, this is the first study to apply multimodal brain

441 age modelling within a longitudinal framework to characterise coupled trajectories of brain

442 maturation and development of internalising and externalising symptoms.

443   Using growth curve models, we observed that changes in the pace of brain age

444 maturation were systematically linked to changes in symptom trajectories over time. The

445 most consistent associations emerged for nonlinear components, indicating that accelerations

446 in deviation from age-expected brain maturation were associated with accelerating

447 internalising and externalising symptoms during adolescence. This suggests that nonlinear

448 changes in how individuals diverge from normative brain maturation trajectories, rather than

449 static brain age deviation, may play a key role in linking brain maturation to emerging

450 psychopathology. For internalising symptoms, nonlinear BAG–symptom associations were

most consistently detected in females across T1-weighted, diffusion, and multimodal models,
whereas corresponding male estimates were smaller and not statistically significant. For
externalising symptoms, nonlinear associations were evident in T1 models for both sexes, in
diffusion MRI as a sex-invariant pooled effect, and in multimodal models for females.
Together, these results highlight synchrony in developmental change as a central feature of
brain–psychopathology coupling across adolescence.

Beyond nonlinear effects, a subset of models showed associations between brain age
deviation and symptom levels at the centred time point of the observation window. These
associations were observed for externalising symptoms in multimodal models for both sexes
and in T1-weighted and rs-fMRI models for females, and for internalising symptoms in
multimodal models for males. Linear associations between rates of brain age change and
symptom change were less common but present in specific modality–domain combinations,
indicating that steeper increases in BAG (i.e., increasing deviation from age-expected
maturation) were sometimes accompanied by steeper symptom increases. In contrast, rs-
fMRI–based brain age showed limited evidence of coupling with symptom trajectories. This
modality did not support nonlinear brain age factors and yielded relatively few associations
overall, consistent with prior reports of relatively lower reliability in functional connectivity–
based brain age estimates in youth samples[88,89]. Our findings partially support our hypothesis
that developmental changes in BAG would be coupled with changes in internalising and
externalising symptom trajectories. Importantly, coupling-specific tests did not provide robust
evidence that these cross-process associations differed by sex after correction for multiple
comparisons. As such, sex-stratified estimates should not be interpreted as evidence of sex-
specific coupling.

Our findings are broadly in line with structural and multimodal MRI studies reporting
that more positive brain age gaps – reflecting older-appearing brains – are associated with
greater psychopathology in adolescents[53,54,57]. At the same time, they contrast with reports
that relatively younger-appearing brains are linked to higher symptom burden[46,48,51]. These
discrepancies indicate that BAG–psychopathology relationships in youth may be
heterogeneous across modalities, samples, and analytic approaches. However, most prior
studies are cross-sectional or rely on only two time points, limiting their ability to disentangle
developmental tempo from stable between-person differences. By modelling intercepts,
slopes, and curvature, the present study shows that nonlinear changes in deviation from age-
expected brain maturation, rather than static age deviation or constant rates of change, were
most consistently aligned with increasing symptoms.

Accelerated patterns of brain maturation during adolescence may reflect

developmental mismatch processes in which affective and socioemotional systems mature

more rapidly than regulatory and cognitive control systems[90,91]. Adolescence is characterised

by heightened neuroplasticity and sensitivity to environmental inputs, such that this

maturational asynchrony may increase vulnerability to internalising symptoms when

emotional reactivity outpaces regulatory capacity[92]. In line with the stress-acceleration and

pubertal timing frameworks, faster biological maturation – particularly in females – has been

associated with elevated risk for depressive and internalising symptoms, potentially reflecting

heightened responsivity to psychosocial demands during sensitive developmental windows[93–

96]. Within this framework, the present findings suggest that accelerating deviations from age-

expected brain maturation trajectories may capture an intensified developmental tempo that

aligns with increasing internalising symptom trajectories, particularly in modalities capturing

structural and multimodal maturation, although this is speculative.

The more consistent detectability of acceleration-linked associations in females is

directionally compatible with well-documented sex differences in developmental timing and

risk for internalising psychopathology. Females typically enter puberty earlier and show

earlier developmental inflection points in cortical and white-matter maturation[94,97], coinciding

with the marked rise in internalising symptoms during early–mid adolescence[22], and broader

sex disparities in internalising symptom prevalence and persistence across adolescence[24,98,99].

However, targeted coupling-specific tests did not provide robust evidence of sex moderation

after correction for multiple comparisons, indicating that apparent differences in within-sex

significance should not be interpreted as strong evidence that coupling strength differs

between females and males. One possibility is that differences in precision (e.g., variability,

measurement noise, missingness, or power) contribute to more consistent detection of these

associations in females in the present data. Future work incorporating pubertal, hormonal,

and sex-related measures, alongside adequately powered moderation tests, will be needed to

clarify whether and how sex-linked developmental timing and biological or socio-cultural

factors shape the coupling between brain maturation tempo and evolving emotional

symptoms. Together, these results position nonlinear deviations in brain maturation tempo as

a developmentally meaningful correlate of adolescent psychopathology, while providing

limited evidence for robust sex-specific coupling.

4.1. Strengths and limitations

A major strength of this study is the combination of four waves of multimodal MRI with ten

waves of psychopathology assessment in a large, diverse, community-based cohort. This

design enabled trajectory-level analyses that have not been possible in prior brain age

research in youth, allowing us to examine how brain maturation and mental health symptoms

unfold together across development. The use of bivariate latent growth curve models

permitted decomposition of brain and symptom development into intercept, slope, and

quadratic components, enabling explicit tests of synchrony in development. Our systematic

model-selection process is another key strength. By evaluating linear, interval-based, latent-

basis, and centred-quadratic specifications, we ensured that our models balanced empirical fit

with parameter stability. This allowed us to identify modalities where nonlinear BAG could

be reliably estimated (T1, dMRI, multimodal) and to simplify the rs-fMRI BAG model where

quadratic terms were not supported.

Some limitations should also be acknowledged. First, at the time of analysis, release

6.0 of the ABCD Study only had partial data available from the fourth imaging time point,

limiting the sample size of participants that had four available scans. Second, rs-fMRI BAG

showed low temporal reliability, potentially preventing estimation of curvature and

precluding conclusions about functional maturational dynamics. Third, BAG is a statistical

index of multivariate deviation rather than a direct biological marker, and its interpretation

should remain cautious. Fourth, although the BPM provided dense longitudinal symptom

data, it is a brief self-report instrument that does not capture finer-grained or multi-informant

clinical variation. Fifth, although FIML allowed us to retain participants with incomplete

MRI or symptom data, patterns of missingness and selective attrition may still have

introduced bias, highlighting the need for future work incorporating explicit missing-data

diagnostics or sensitivity analyses. Sixth, because the cohort is drawn from a large U.S.

sample, generalisability to other cultural or socio-environmental contexts may be constrained.

Finally, the BLGC models capture coordinated change but cannot determine directionality or

causal pathways, underscoring the need for future causal designs.

4.2. Conclusion

Adolescence is marked by rapid and uneven change across neurobiological, emotional, and

behavioural domains, and the present findings indicate that it is the *tempo* of brain maturation

that most meaningfully aligns with internalising and externalising symptoms. We show that

accelerations in divergence from normative brain maturation track with parallel accelerations

in psychopathology across adolescence. The results support a broader developmental view in

which vulnerability arises not from absolute levels of neurobiological features, but from asynchronies in their timing and pace of change, offering a powerful framework for understanding why adolescence is a sensitive period for emerging mental health problems.

5. Acknowledgements

Data used in the preparation of this article were obtained from the Adolescent Brain Cognitive Development™ (ABCD) Study, held in the NIH Brain Development Cohorts Data Sharing Platform. This is a multisite, longitudinal study designed to recruit more than 10,000 children aged ~8–10 and follow them over 10 years into early adulthood. The ABCD Study® is supported by the National Institutes of Health and additional federal partners under award numbers: U01DA041048, U01DA050989, U01DA051016, U01DA041022, U01DA051018, U01DA051037, U01DA050987, U01DA041174, U01DA041106, U01DA041117, U01DA041028, U01DA041134, U01DA050988, U01DA051039, U01DA041156, U01DA041025, U01DA041120, U01DA051038, U01DA041148, U01DA041093, U01DA041089, U24DA041123, U24DA041147. A full list of supporters is available at Federal Partners - ABCD Study. ABCD Consortium investigators designed and implemented the study and/or provided data but did not necessarily participate in the analysis or writing of this report. This manuscript reflects the views of the authors and may not reflect the opinions or views of the NIH or ABCD Consortium investigators.

Data was handled inside Service for Sensitive Data (TSD), a platform owned by the University of Oslo, operated, and developed by the TSD service group at the University of Oslo IT-Department (USIT). Computations were also performed using resources provided by UNINETT Sigma2 – the National Infrastructure for High Performance Computing and Data Storage in Norway (NS9666S).

6. Funding

This research is supported by the Research Council of Norway (#288083, #300767, #323951), and the South-Eastern Norway Regional Health Authority (#2021070, #2023012, #500189).

7. Conflict of interest statement

The authors declare that they have no conflict of interest.

8. CRediT author contribution statement

Beck, Dani: Writing – Original draft, review & editing, Project administration, Visualisation,
Methodology, Formal analysis, Data curation, Conceptualisation. **Carrick, Chloe:** Writing –
review & editing, Formal analysis, Conceptualisation. **Aksnes, Eira R.:** Writing – review &
editing, Conceptualisation. **MacSweeney, Niamh.:** Writing – review & editing. **Westlye,**
Lars T.: Writing – review & editing, Funding acquisition. **Tamnes, Christian K.:** Writing –
review & editing, Project administration, Funding acquisition, Conceptualisation.
592
9. References

1. Lewinsohn, P. M., Solomon, A., Seeley, J. R. & Zeiss, A. Clinical implications of
'subthreshold' depressive symptoms. *J. Abnorm. Psychol.* **109**, 345–351 (2000).
2. Whittle, S., Vijayakumar, N., Simmons, J. G. & Allen, N. B. Internalizing and
Externalizing Symptoms Are Associated With Different Trajectories of Cortical
Development During Late Childhood. *J. Am. Acad. Child Adolesc. Psychiatry* **59**,
177–185 (2020).
3. Bos, M. G. N., Peters, S., van de Kamp, F. C., Crone, E. A. & Tamnes, C. K.
Emerging depression in adolescence coincides with accelerated frontal cortical
thinning. *J. Child Psychol. Psychiatry* **59**, 994–1002 (2018).
4. Ducharme, S. *et al.* Anxious/Depressed Symptoms are Linked to Right
Ventromedial Prefrontal Cortical Thickness Maturation in Healthy Children and
Young Adults. *Cereb. Cortex* **24**, 2941–2950 (2014).
5. Newman, E. *et al.* Anxiety is related to indices of cortical maturation in typically
developing children and adolescents. *Brain Struct. Funct.* **221**, 3013–3025
(2016).
6. Papmeyer, M. *et al.* Cortical Thickness in Individuals at High Familial Risk of
Mood Disorders as They Develop Major Depressive Disorder. *Biol. Psychiatry* **78**,
58–66 (2015).

7. Schmaal, L. *et al.* Brain Structural Signatures of Adolescent Depressive Symptom

Trajectories: A Longitudinal Magnetic Resonance Imaging Study. *J. Am. Acad.*

Child Adolesc. Psychiatry **56**, 593-601.e9 (2017).

8. Shaw, P. *et al.* Attention-deficit/hyperactivity disorder is characterized by a delay

in cortical maturation. *Proc. Natl. Acad. Sci.* **104**, 19649–19654 (2007).

9. Perlstein, S. *et al.* Unique Versus Shared Neural Correlates of Externalizing

Psychopathology in Late-Childhood. *J. Psychopathol. Clin. Sci.* **133**, 477–488

(2024).

10. Arango, C. *et al.* Progressive Brain Changes in Children and Adolescents With

First-Episode Psychosis. *Arch. Gen. Psychiatry* **69**, 16–26 (2012).

11. Cannon, T. D. *et al.* Progressive Reduction in Cortical Thickness as Psychosis

Develops: A Multisite Longitudinal Neuroimaging Study of Youth at Elevated

Clinical Risk. *Biol. Psychiatry* **77**, 147–157 (2015).

12. Whittle, S. *et al.* Structural Brain Development and Depression Onset During

Adolescence: A Prospective Longitudinal Study. *Am. J. Psychiatry*

https://doi.org/10.1176/appi.ajp.2013.13070920 (2014)

doi:10.1176/appi.ajp.2013.13070920.

13. Albaugh, M. D. *et al.* Anxious/depressed symptoms are related to microstructural

maturation of white matter in typically developing youths. *Dev. Psychopathol.* **29**,

751–758 (2017).

14. Muetzel, R. L. *et al.* Tracking Brain Development and Dimensional Psychiatric

Symptoms in Children: A Longitudinal Population-Based Neuroimaging Study.

Am. J. Psychiatry https://doi.org/10.1176/appi.ajp.2017.16070813 (2018)

doi:10.1176/appi.ajp.2017.16070813.

15. Dall'Aglio, L., Xu, B., Tiemeier, H. & Muetzel, R. L. Longitudinal Associations

Between White Matter Microstructure and Psychiatric Symptoms in Youth. *J. Am.*

Acad. Child Adolesc. Psychiatry **62**, 1326–1339 (2023).

16. Ho, T. C., Shah, R., Mishra, J., May, A. C. & Tapert, S. F. Multi-level predictors of

depression symptoms in the Adolescent Brain Cognitive Development (ABCD)

study. *J. Child Psychol. Psychiatry* **63**, 1523–1533 (2022).

17. Nelson, B. D., Perlman, G., Klein, D. N., Kotov, R. & Hajcak, G. Blunted Neural

Response to Rewards as a Prospective Predictor of the Development of

Depression in Adolescent Girls. *Am. J. Psychiatry* **173**, 1223–1230 (2016).

18. Stringaris, A. *et al.* The Brain's Response to Reward Anticipation and Depression

in Adolescence: Dimensionality, Specificity, and Longitudinal Predictions in a

Community-Based Sample. *Am. J. Psychiatry* **172**, 1215–1223 (2015).

19. Toenders, Y. J. *et al.* Neuroimaging predictors of onset and course of depression

in childhood and adolescence: A systematic review of longitudinal studies. *Dev.*

Cogn. Neurosci. **39**, 100700 (2019).

20. Thijssen, S., Collins, P. F., Weiss, H. & Luciana, M. The longitudinal association

between externalizing behavior and frontoamygdalar resting-state functional

connectivity in late adolescence and young adulthood. *J. Child Psychol.*

Psychiatry **62**, 857–867 (2021).

21. Nakua, H. *et al.* Investigating cross-sectional and longitudinal relationships

between brain structure and distinct dimensions of externalizing psychopathology

in the ABCD sample. *Neuropsychopharmacology* **50**, 499–506 (2025).

22. Salk, R. H., Hyde, J. S. & Abramson, L. Y. Gender differences in depression in

representative national samples: Meta-analyses of diagnoses and symptoms.

Psychol. Bull. **143**, 783–822 (2017).

23. Keyes, K. M. & Platt, J. M. Annual Research Review: Sex, gender, and

internalizing conditions among adolescents in the 21st century – trends, causes,

consequences. *J. Child Psychol. Psychiatry* **65**, 384–407 (2024).

24. Dalsgaard, S. *et al.* Incidence Rates and Cumulative Incidences of the Full

Spectrum of Diagnosed Mental Disorders in Childhood and Adolescence. *JAMA*

Psychiatry **77**, 155–164 (2020).

25. Kjelkenes, R. *et al.* Deviations from normative brain white and gray matter

structure are associated with psychopathology in youth. *Dev. Cogn. Neurosci.* **58**,

101173 (2022).

26. Andersen, S. L. & Teicher, M. H. Stress, sensitive periods and maturational

events in adolescent depression. *Trends Neurosci.* **31**, 183–191 (2008).

27. Bethlehem, R. a. I. *et al.* Brain charts for the human lifespan. *Nature* **604**, 525–

533 (2022).

28. Giedd, J. N. *et al.* Brain development during childhood and adolescence: a

longitudinal MRI study. *Nat. Neurosci.* **2**, 861–863 (1999).

29. Callaghan, B. L. & Tottenham, N. The Stress Acceleration Hypothesis: effects of

early-life adversity on emotion circuits and behavior. *Curr. Opin. Behav. Sci.* **7**,

76–81 (2016).

30. Casey, B. j., Jones, R. M. & Hare, T. A. The Adolescent Brain. *Ann. N. Y. Acad.*

Sci. **1124**, 111–126 (2008).

31. Cole, J. H. *et al.* Predicting brain age with deep learning from raw imaging data

results in a reliable and heritable biomarker. *NeuroImage* **163**, 115–124 (2017).

32. Franke, K., Ziegler, G., Klöppel, S. & Gaser, C. Estimating the age of healthy

subjects from T1-weighted MRI scans using kernel methods: Exploring the

influence of various parameters. *NeuroImage* **50**, 883–892 (2010).

33. Busby, N. *et al.* Lower socioeconomic status is associated with premature brain

aging. *Neurobiol. Aging* **130**, 135–140 (2023).

34. Beck, D. *et al.* Adipose tissue distribution from body MRI is associated with cross-

sectional and longitudinal brain age in adults. *NeuroImage Clin.* **33**, 102949

(2022).

35. Beck, D. *et al.* Dissecting unique and common variance across body and brain

health indicators using age prediction. *Hum. Brain Mapp.* **45**, e26685 (2024).

36. Beck, D. *et al.* Cardiometabolic risk factors associated with brain age and

accelerate brain ageing. *Hum. Brain Mapp.* **43**, 700–720 (2022).

37. de Lange, A.-M. G. *et al.* Multimodal brain-age prediction and cardiovascular risk:

The Whitehall II MRI sub-study. *NeuroImage* **222**, 117292 (2020).

38. Sanders, F. *et al.* Associations between methylation age and brain age in late

adolescence. 2022.09.08.506972 Preprint at

https://doi.org/10.1101/2022.09.08.506972 (2022).

39. Subramaniapillai, S. *et al.* Sex- and age-specific associations between

cardiometabolic risk and white matter brain age in the UK Biobank cohort. *Hum.*

Brain Mapp. **n/a**, (2022).

40. Leonardsen, E. H. *et al.* Deep neural networks learn general and clinically

relevant representations of the ageing brain. *NeuroImage* **256**, 119210 (2022).

41. Anatürk, M. *et al.* Prediction of brain age and cognitive age: Quantifying brain and

cognitive maintenance in aging. *Hum. Brain Mapp.* **42**, 1626–1640 (2021).

42. Biondo, F. *et al.* Brain-age is associated with progression to dementia in memory

clinic patients. *NeuroImage Clin.* **36**, 103175 (2022).

43. Kaufmann, T. *et al.* Common brain disorders are associated with heritable

patterns of apparent aging of the brain. *Nat. Neurosci.* **22**, 1617–1623 (2019).

44. Rokicki, J. *et al.* Multimodal imaging improves brain age prediction and reveals

distinct abnormalities in patients with psychiatric and neurological disorders.

Hum. Brain Mapp. **42**, 1714–1726 (2021).

45. Tønnesen, S. *et al.* Brain Age Prediction Reveals Aberrant Brain White Matter in

Schizophrenia and Bipolar Disorder: A Multisample Diffusion Tensor Imaging

Study. *Biol. Psychiatry Cogn. Neurosci. Neuroimaging* S2451902220301683

(2020) doi:10.1016/j.bpsc.2020.06.014.

46. Cohen, J. W. *et al.* Relative brain age is associated with socioeconomic status

and anxiety/depression problems in youth. *Dev. Psychol.* **60**, 199–209 (2024).

47. Kurth, F. *et al.* Preliminary evidence for a lower brain age in children with

attention-deficit/hyperactivity disorder. *Front. Psychiatry* **13**, (2022).

48. Luna, A. *et al.* Maturity of gray matter structures and white matter connectomes,

and their relationship with psychiatric symptoms in youth. *Hum. Brain Mapp.* **42**,

4568–4579 (2021).

49. Tunç, B. *et al.* Deviation from normative brain development is associated with

symptom severity in autism spectrum disorder. *Mol. Autism* **10**, 46 (2019).

50. Zhou, Z. *et al.* Differential effects of generalized anxiety and separation anxiety

on brain structural development during adolescence. *J. Affect. Disord.* **339**, 478–

485 (2023).

51. Lund, M. J. *et al.* Brain age prediction using fMRI network coupling in youths and

associations with psychiatric symptoms. *NeuroImage Clin.* **33**, 102921 (2022).

52. Chung, Y. *et al.* Use of Machine Learning to Determine Deviance in

Neuroanatomical Maturity Associated With Future Psychosis in Youths at

Clinically High Risk. *JAMA Psychiatry* **75**, 960–968 (2018).

53. Cropley, V. L. *et al.* Brain-Predicted Age Associates With Psychopathology

Dimensions in Youths. *Biol. Psychiatry Cogn. Neurosci. Neuroimaging* **6**, 410–

419 (2021).

54. Drobinin, V. *et al.* The Developmental Brain Age Is Associated With Adversity,

Depression, and Functional Outcomes Among Adolescents. *Biol. Psychiatry*

Cogn. Neurosci. Neuroimaging **7**, 406–414 (2022).

55. Truelove-Hill, M. *et al.* A Multidimensional Neural Maturation Index Reveals

Reproducible Developmental Patterns in Children and Adolescents. *J. Neurosci.*

40, 1265–1275 (2020).

56. de Nooij, L. *et al.* Longitudinal trajectories of brain age in young individuals at

familial risk of mood disorder from the Scottish Bipolar Family Study. *Wellcome*

Open Res. **4**, 206 (2020).

57. MacSweeney, N. *et al.* Multimodal Brain Age Indicators of Internalizing Problems

in Early Adolescence: A Longitudinal Investigation. *Biol. Psychiatry Cogn.*

Neurosci. Neuroimaging **10**, 475–484 (2025).

58. Whitmore, L. & Beck, D. Current challenges and future directions for brain age

prediction in children and adolescents. *Nat. Commun.* **16**, 7771 (2025).

59. Wierenga, L. M. *et al.* A Key Characteristic of Sex Differences in the Developing

Brain: Greater Variability in Brain Structure of Boys than Girls. *Cereb. Cortex N.*

Y. N 1991 **28**, 2741–2751 (2018).

60. Casey, B. J. *et al.* The Adolescent Brain Cognitive Development (ABCD) study:

Imaging acquisition across 21 sites. *Dev. Cogn. Neurosci.* **32**, 43–54 (2018).

61. Karcher, N. R. *et al.* Assessment of the Prodromal Questionnaire–Brief Child

Version for Measurement of Self-reported Psychoticlike Experiences in

Childhood. *JAMA Psychiatry* **75**, 853–861 (2018).

62. Auchter, A. M. *et al.* A description of the ABCD organizational structure and

communication framework. *Dev. Cogn. Neurosci.* **32**, 8–15 (2018).

63. Hagler, D. J. *et al.* Image processing and analysis methods for the Adolescent

Brain Cognitive Development Study. *NeuroImage* **202**, 116091 (2019).

64. Dale, A. M., Fischl, B. & Sereno, M. I. Cortical Surface-Based Analysis. 16

(1999).

65. Fischl, B. *et al.* Whole Brain Segmentation. *Neuron* **33**, 341–355 (2002).

66. Hagler Jr., D. J. *et al.* Automated white-matter tractography using a probabilistic

diffusion tensor atlas: Application to temporal lobe epilepsy. *Hum. Brain Mapp.*

30, 1535–1547 (2009).

67. Van Dijk, K. R. A. *et al.* Intrinsic Functional Connectivity As a Tool For Human

Connectomics: Theory, Properties, and Optimization. *J. Neurophysiol.* **103**, 297–

321 (2010).

68. Gordon, E. M. *et al.* Generation and Evaluation of a Cortical Area Parcellation

from Resting-State Correlations. *Cereb. Cortex N. Y. NY* **26**, 288–303 (2016).

69. Beer, J. C. *et al.* Longitudinal ComBat: A method for harmonizing longitudinal

multi-scanner imaging data. *NeuroImage* **220**, 117129 (2020).

70. Chen, T. & Guestrin, C. XGBoost: A Scalable Tree Boosting System. in

Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge

Discovery and Data Mining 785–794 (ACM, San Francisco California USA,

2016). doi:10.1145/2939672.2939785.

71. Beck, D. *et al.* Dimensions of Early-Life Adversity Are Differentially Associated

With Patterns of Delayed and Accelerated Brain Maturation. *Biol. Psychiatry*

https://doi.org/10.1016/j.biopsych.2024.07.019 (2024)

doi:10.1016/j.biopsych.2024.07.019.

72. Dehestani, N., Whittle, S., Vijayakumar, N. & Silk, T. J. Developmental brain

changes during puberty and associations with mental health problems. *Dev.*

Cogn. Neurosci. **60**, 101227 (2023).

73. Liang, H., Zhang, F. & Niu, X. Investigating systematic bias in brain age

estimation with application to post-traumatic stress disorders. *Hum. Brain Mapp.*

hbm.24588 (2019) doi:10.1002/hbm.24588.

74. de Lange & Cole. Commentary: Correction procedures in brain-age prediction.

NeuroImage Clin. **26**, 102229 (2020).

75. Achenbach, T., Mcconaughy, S., Ivanova, M. & Rescorla, L. Manual for the

ASEBA Brief Problem Monitor[TM](BPM). *Burlingt. VT ASEBA* (2011).

76. Bilenberg, N. The Child Behavior Checklist (CBCL) and related material:

standardization and validation in Danish population based and clinically based

samples. *Acta Psychiatr. Scand.* **100**, 2–52 (1999).

77. Piper, B. J., Gray, H. M., Raber, J. & Birkett, M. A. Reliability and validity of Brief

Problem Monitor, an abbreviated form of the Child Behavior Checklist. *Psychiatry*

Clin. Neurosci. **68**, 759–767 (2014).

78. R Core Team. *R: A Language and Environment for Statistical Computing.* (R

Foundation for Statistical Computing, Vienna, Austria, 2022).

79. Rosseel, Y. lavaan: An R Package for Structural Equation Modeling. *J. Stat.*

Softw. **48**, 1–36 (2012).

80. Rosseel, Y. *et al.* lavaan: Latent Variable Analysis. (2025).

81. Duncan, T. E. & Duncan, S. C. The ABC's of LGM: An Introductory Guide to

Latent Variable Growth Curve Modeling. *Soc. Personal. Psychol. Compass* **3**,

979–991 (2009).

82. Grimm, K. J., Ram, N. & Hamagami, F. Nonlinear Growth Curves in

Developmental Research. *Child Dev.* **82**, 1357–1371 (2011).

83. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical

and Powerful Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B Methodol.* **57**,

289–300 (1995).

84. Schermelleh-Engel, K., Moosbrugger, H. & Müller, H. Evaluating the Fit of

Structural Equation Models: Tests of Significance and Descriptive Goodness-of-

Fit Measures.

85. Gignac, G. E. & Szodorai, E. T. Effect size guidelines for individual differences

researchers. *Personal. Individ. Differ.* **102**, 74–78 (2016).

86. Edde, M., Leroux, G., Altena, E. & Chanraud, S. Functional brain connectivity

changes across the human life span: From fetal development to old age. *J.*

Neurosci. Res. **99**, 236–262 (2021).

87. Solmi, M. *et al.* Age at onset of mental disorders worldwide: large-scale meta-

analysis of 192 epidemiological studies. *Mol. Psychiatry* **27**, 281–295 (2022).

88. Cornblath, E. J. *et al.* Temporal sequences of brain activity at rest are constrained

by white matter structure and modulated by cognitive demands. *Commun. Biol.*

3, 261 (2020).

89. Satterthwaite, T. D. *et al.* An improved framework for confound regression and

filtering for control of motion artifact in the preprocessing of resting-state

functional connectivity data. *NeuroImage* **64**, 240–256 (2013).

90. Casey, B. j., Jones, R. M. & Hare, T. A. The Adolescent Brain. *Ann. N. Y. Acad.*

Sci. **1124**, 111–126 (2008).

91. Somerville, L. H., Jones, R. M. & Casey, B. J. A time of change: Behavioral and

neural correlates of adolescent sensitivity to appetitive and aversive

environmental cues. *Brain Cogn.* **72**, 124 (2009).

92. Blakemore, S.-J. & Mills, K. L. Is adolescence a sensitive period for sociocultural

processing? *Annu. Rev. Psychol.* **65**, 187–207 (2014).

93. Mendle, J., Harden, K. P., Brooks-Gunn, J. & Graber, J. A. Development's

Tortoise and Hare: Pubertal Timing, Pubertal Tempo, and Depressive Symptoms

in Boys and Girls. *Dev. Psychol.* **46**, 1341–1353 (2010).

94. Vijayakumar, N., Op de Macks, Z., Shirtcliff, E. A. & Pfeifer, J. H. Puberty and the

human brain: Insights into adolescent development. *Neurosci. Biobehav. Rev.* **92**,

417–436 (2018).

95. Colich, N. L. *et al.* Earlier age at menarche as a transdiagnostic mechanism

linking childhood trauma with multiple forms of psychopathology in adolescent

girls. *Psychol. Med.* **50**, 1090–1098 (2020).

96. Pfeifer, J. H. & Allen, N. B. Puberty Initiates Cascading Relationships Between

Neurodevelopmental, Social, and Internalizing Processes Across Adolescence.

Biol. Psychiatry **89**, 99–108 (2021).

97. Gur, R. E. & Gur, R. C. Sex differences in brain and behavior in adolescence:

Findings from the Philadelphia Neurodevelopmental Cohort. *Neurosci. Biobehav.*

Rev. **70**, 159–170 (2016).

98. Wierenga, L. M. *et al.* Recommendations for a Better Understanding of Sex and

Gender in the Neuroscience of Mental Health. *Biol. Psychiatry Glob. Open Sci.* **4**,

100283 (2024).

99. Wierenga, L. M. & Beck, D. Sex/gender associated mechanisms in the (human)

brain: Sex hormones, sex chromosomes, and social-cultural gender factors. in

Reference Module in Neuroscience and Biobehavioral Psychology (Elsevier,

2024). doi:10.1016/B978-0-12-820480-1.00173-X.