1

Predictive value of Acute Physiology and Chronic Health Evaluation II (APACHE II) scoring in patients admitted to intensive care units

Saurav Jha[1*], Shirjan Gautam[2], Samir Shiwakoti[3]

[1] Department of Internal Medicine, Kathmandu Medical College, Kathmandu, Bagmati Province, Nepal

[2] Department of Internal Medicine, Kathmandu Medical College, Kathmandu, Bagmati Province, Nepal

[3] Department of General Surgery, Kathmandu Medical College, Kathmandu, Bagmati Province, Nepal

*Corresponding author

E-mail: sjha101020@gmail.com

Abstract

Background

Risk stratification tools are essential for guiding care and allocating limited resources in intensive care units (ICUs). Evidence on the performance of the Acute Physiology and Chronic Health Evaluation II (APACHE II) score in Nepal is limited. We evaluated the association of APACHE II measured within 24 hours of ICU admission with mortality, ICU length of stay, and discharge disposition in a tertiary hospital in Nepal.

Methods and findings

We conducted a prospective observational cohort study of consecutive adult patients admitted to three multidisciplinary ICUs at a tertiary hospital in Kathmandu, Nepal. Patients with ICU stay <24 hours and those admitted to coronary or neonatal ICUs were excluded. APACHE II was calculated from the worst values in the first 24 hours. The primary outcome was ICU mortality; secondary outcomes were ICU length of stay and discharge disposition (home, ward, high-care/step-down, or in-hospital death). Discrimination for mortality was assessed using receiver operating characteristic (ROC) analysis. Associations with length of stay were examined using linear regression; discharge disposition was evaluated using multinomial logistic regression.

Among 200 patients (54% male; mean age 54.7 ± 21.0 years), ICU mortality was 23.0% (46/200). Non-survivors were older than survivors (mean difference 13.31 years; 95% CI 6.60–20.02; $p < 0.001$). The mean APACHE II score was 13.19 ± 7.89 overall, higher in non-survivors vs survivors (19.85 ± 7.14 vs 11.21 ± 6.98; $p < 0.001$). APACHE II discriminated ICU mortality well (AUC 0.806); a cutoff ≥ 14.5 yielded 76.1% sensitivity and 71.4% specificity. Mortality increased across APACHE II strata (trend $p < 0.001$). Glasgow Coma Scale scores were lower in non-survivors (10.85 ± 4.43) than survivors (13.29 ± 3.11; $p < 0.001$). Higher APACHE II scores were associated with in-hospital death versus ward discharge,

but not with high-care versus ward (OR 1.063; 95% CI 0.994–1.136; p = 0.074). Limitations include the single-center design, modest sample size, and short study duration, which may limit generalizability.

Conclusions

APACHE II measured within 24 hours of ICU admission demonstrated good discrimination for ICU mortality and a modest association with ICU length of stay in a Nepalese tertiary setting, while being less informative for intermediate discharge dispositions. Together with Glasgow Coma Scale, APACHE II offers a pragmatic, cost-effective approach to early prognostication and resource planning in resource-limited ICUs. Multicenter studies are warranted to validate these findings across Nepal.

Introduction

According to Painter et al, "A critically ill or injured patient is defined as one who has an illness or injury impairing one or more vital organ systems such that there is a high probability of imminent or life-threatening deterioration in the patient's condition." [1] It is important to stratify critically ill patients into risk groups to guide their admission into respective units in the hospital and the treatment they are going to receive.[2] Working in an intensive care unit (ICU) requires making early predictions about patient outcomes within the first 24 hours of admission. This involves estimating the likelihood of mortality by analyzing key variables typically used in diagnosing and treating critically ill patients.[3] Some widely used scoring systems for critically ill ICU patients are the Acute Physiology and Chronic Health Evaluation (APACHE) I-IV, the Mortality Prediction Model (MPM) and the Simplified Acute Physiology Score (SAPS) I-III.[4,5] APACHE scoring system was first introduced in 1981 AD by William Knaus et al and was updated 4 years later to APACHE II.[5–7] The APACHE-II system assigns points for 12 physiologic variables, for age, and for chronic health status, in generating a total point score. While measuring APACHE II, patient's temperature, heart rate, respiratory rate, mean arterial blood pressure, oxygenation, arterial pH, serum potassium, sodium, and creatinine, hematocrit, white blood cell (WBC),

and Glasgow Coma Scale are measured.[8] These 12 variables are used to assign an integer score from 0

to 71 for age, and for chronic health status. [9,10]

The APACHE scoring system was developed in USA and its application may or may not be equally

applicable throughout different countries of the globe, especially in the context of Nepal where the

literature regarding the study of APACHE II for predicting mortality in ICU settings is limited.[11] This

study was conducted to study the correlation between the APACHE II score within 24 hours of admission

into ICU and length of ICU stay, outcome of mortality and discharge disposition.

Methods

We conducted a prospective observational cohort study in Kathmandu Medical College and Teaching

hospital, a tertiary care center in Kathmandu, Nepal. There are a total of four ICUs in the hospital,

including a neonatal ICU. Data were taken from three ICU, and neonatal ICU was excluded. The ICUs

were mixed medical-surgical units providing level III critical care. Patients with missing data required to

calculate APACHE II score were excluded as well. Ethical Approval was obtained from the Institutional

Review Committee of the hospital with reference no. 21072025/26. This research was conducted in

accordance with the Declaration of Helsinki. The requirement for written informed consent was waived

by the institutional review committee as only de-identified patient data were collected. Patients were

admitted to these ICUs based on whether they had existing or impending organ failure, or if they

underwent a recent surgery at our center. The ICUs were handled by dedicated critical care experts,

including critical care residents, consultants from different faculty throughout the hospital, on-call

anesthesia residents, junior residents, medical officers, and trained nurses.

We included patients admitted to the ICU and excluded patients admitted to coronary care units, and

neonatal ICUs. The sample size was calculated to estimate the diagnostic performance of APACHE II

score for predicting ICU mortality. Published metanalyses were used to derive the estimates. A pooled

sensitivity of 88%, specificity of 84%, and an AUC of 0.84 for APACHE II in predicting mortality was

used.[12] An expected ICU mortality prevalence of 29.6% was used.[13] Based on these parameters, the

sample size required to estimate sensitivity with 95% confidence and ±10% precision was 138 patients,

which exceeded the number required for estimating specificity or ROC-AUC. To account for

approximately 10% incomplete data, the target sample size was increased to 152 patients. However, the

study included 200 patients to increase the precision of the analysis and reduce random variability. Bias

was minimized using validated scoring systems, structured data collection, consecutive patients inclusion,

and stratification. Recall and misclassification bias were also limited as all information was extracted

from clinical records. Multiple approaches were used to identify and minimize the potential confounders

within the data set. These approaches included comparing baseline characteristics between survivors and

non-survivors, assessing correlations between key variables such as age and APACHE II score,

examining outcome patterns through stratified APACHE II categories and regression analyses.

A prospective study was carried out in our hospital from 1st August 2025 to 30th November 2025. The

data of the patients admitted into different intensive care units of our hospital were recorded over the

period of 4 months. Data were recorded on a daily basis within 24 hours of admission. Patients whose

ICU stay was less than 24 hours were excluded from the study. The variables included were the age, sex

and ethnicity of the patient, the date of admission, the primary diagnosis, any co-morbidities that the

patient had, their mean arterial pressure, body temperature, arterial pH, heart rate, respiratory rate, serum

sodium, potassium, creatinine, hematocrit, WBC count, GCS score, FiO2, PaO2, PaCO2 and eventually

their calculated APACHE II score. The length of ICU stay and the outcome, i.e. mortality or discharge

disposition was also recorded. The primary outcome was ICU mortality; secondary outcomes were ICU

length of stay and discharge disposition (home, ward, high-care/step-down, or in-hospital death). ICU

mortality was defined as death occurring during the ICU stay prior to discharge or transfer. High-care

units were defined as step down units providing continuous monitoring without invasive organ support.

The data of the aforementioned variables were entered into SPSS version 26 and the results were extrapolated and are

mentioned in this study.

Result

A total of 200 patients were included in this study. 108 (54%) of them were men and 92 (46%) were

females. The mean age of participants was 54.7 ± 21 years (Range 19-98). 46 (23%) of the patients died

while 154 (77%) of patients survived. There was a statistically significant difference in the mean age of

survivors (51.63 ± 20.68 years) as compared to non-survivors (64.93 ± 18.73 years). The mean age

difference was 13.31 years (95% CI: 6.60 to 20.02; $p < 0.001$).

The mean APACHE II score of all patients admitted in the ICU was 13.19 ± 7.89. The mean APACHE II

score among non-survivors was 19.85 ± 7.14 and found to be significantly higher ($p<0.001$) than

survivors with score of 11.21 ± 6.98. A moderate positive correlation between age and APACHE II score

was found which was statistically significant ($r = 0.540$, $p < 0.001$).

The predictive value of APACHE II scoring system was evaluated using area under ROC curve. It was

found that the APACHE II score significantly predicted ICU mortality with an area under curve of 0.806

which indicated good diagnostic accuracy, as shown in Fig 1. The optimal value for cut off score for

predicting mortality was ≥ 14.5 with sensitivity of 76.1% and specificity of 71.4%.

Fig 1. Receiver Operating Characteristic (ROC) Curve for APACHE II Score in Predicting ICU

Mortality.

The mortality rates according to APACHE II Scores of the patients are shown in the Table 1. Chi-square

test showed statistically significant association between APACHE II score groups and mortality (χ^2=

42.723, $p < .001$) pointing out that there was significant difference in the mortality rates across APACHE

II score groups. The linear-by-linear association test was also significant ($\chi^2 = 39.218$, $p < .001$),

suggesting a trend of increasing mortality with higher APACHE II scores.

APACHE II Score	Survivors (n)	Non-Survivors (n)	Total (n)	Mortality Rate (%)
0–9	29	0	29	0.0%
10–14	37	5	42	11.9%
15–19	44	6	50	12.0%
20–24	27	13	40	32.5%
25–29	10	11	21	52.4%
30–34	5	6	11	54.5%
≥35	2	5	7	71.4%
Total	**154**	**46**	**200**	**23.0%**

Table 1. Distribution of ICU Mortality Across APACHE II Score Categories

The mean Glasgow Coma Scale (GCS) score among survivors was 13.29 ± 3.11, while among non-

survivors it was 10.85 ± 4.43 and the difference was statistically significant (p<0.001). Linear regression

showed that APACHE II score was a significant predictor of ICU stay duration (B= 0.153, p<0.001). A

weak positive correlation between APACHE II score and ICU stay was observed (R=0.298). About 9% of

variance in the ICU stay was explained by the model.

A multinomial logistic regression was performed to evaluate whether APACHE II score can predict

discharge disposition to High-care, Ward or Home. The model was statistically significant, $\chi^2(2) = 47.61$,

p < .001. Higher APACHE II scores were significantly associated with in-hospital deaths compared to

patients discharged to the ward. However, there was no statistically significant association between

APACHE II score and discharge disposition to high-care unit vs. Ward. (OR = 1.063, 95% CI :0.994 to

1.136, p = 0.074).

The APACHE II Scores and Mortality rate according to the disease category of the patients is shown in

Table 2. The patients with respiratory causes had the highest mortality rate of 35.5% with APACHE II

score ranging 16.55 ± 7.02 closely followed by cardiovascular causes with mortality of 33.3% and

APACHE II score of 16.00 ± 8.86.

Disease Category	APACHE II Score (Mean ± SD)	Mortality Rate (%)
Neurological Causes	11.03 ± 6.29	18.2%
Respiratory Causes	16.55 ± 7.02	35.5%
Gastrointestinal Causes	10.67 ± 6.98	14.3%
Cardiovascular Causes	16.00 ± 8.86	33.3%
Sepsis/Septic Shock	13.38 ± 5.10	25.0%
Other	11.35 ± 9.42	12.5%
Total	**13.20 ± 7.89**	**23.0%**

157   **Table 2.** APACHE II Score and Mortality rate among different disease categories.

Discussion

The mean age of survivors was significantly lower than non-survivors. However, the positive correlation between APACHE II score and age suggests that aging is associated with increased vulnerability to critical illness. This finding was similar to other studies which imply APACHE II score is the predictor of mortality and not the age.[14,15] In countries like Nepal, where access to health care among elderly may be limited, the impact of age on disease severity and outcome may be even more detrimental.[16]

Our study concluded that APACHE II score significantly predicted ICU mortality which is also supported by many similar studies.[17–19] In our study, the cut off value of ≥14.5 was found to predict ICU mortality with a sensitivity of 76.1% and specificity of 71.4%, demonstrating balanced and good predictive accuracy. A wide range of studies have shown APACHE II score having sensitivity from 51% to 93% and specificity from 49% to 97%.[20] Our findings are parallel to the findings by Yan et al. (2021) in a study among gram-negative sepsis patients with APACHE II cut off of 18.5 with sensitivity of 65.7% and specificity of 72.7%. [21]

In our study, stratified analysis using categorized APACHE II scores showed a clear trend of increasing ICU mortality with higher APACHE II scores. Mortality rose from 0% in lowest score group (0-9) to 71.4% in the highest score group (≥35). These findings are consistent with other researches indicating that APACHE II score is a reliable predictor of mortality in intensive care settings.[22,23.]For instance, a cohort study in Pakistan found that all the patients with APACHE II score ≥21 died, while around 70% of survivors had score of 11-20.[24] This weighs in on a clear stratification of outcomes based on score categories. These results support the idea of integrating APACHE II into ICU triage for risk assessment, guiding clinical decisions, and prioritizing resource allocation in resource-limited settings.

Survivors had a significantly higher mean GCS compared to non-survivors, indicating that lower GCS on admission is associated with higher mortality. This is in alignment with the findings of a meta-analysis of Ethiopian ICUs where patients with GCS below 9 were 4.7 times more likely to die in the ICU. [25] The

Pan-Asia Outcomes Study, evaluating over 16000 patients concluded that both full GCS and motor-only

component of GCS has outstanding predictive accuracy for a 30-day mortality.[26] In resource-limited

Nepalese settings, where advanced diagnostic tools may not be readily available, GCS can be a defining

tool for early risk assessment and management.

The APACHE II score was a statistically significant, though modest, predictor of ICU stay duration.

While it accounts for a small portion of the variance, higher scores were associated with slightly longer

ICU stays. This supports the utility of the APACHE II score not only in predicting mortality but also in

anticipating resource utilization in terms of ICU days. In contrast, a study conducted in Pakistan

demonstrated an inverse, although weak, correlation between APACHE II and ICU stay pointing out that

sicker patients often have shorter stays due to higher mortality. [27] Similarly a Japanese study concluded

that increasing APACHE II scores were significantly associated with prolonged ICU stays.[28] These

widely different results point to the fact that the relationship of APACHE II with length of ICU stay may

be context-dependent. In some cases, higher scores mean greater severity and longer need of ICU care,

while in others they may indicate terminal illness and shorter ICU stays due to increasing mortality. In our

Nepalese ICU context, the positive association likely reflects a system where higher-severity illness

corresponds to longer ICU resource utilization but is still tempered by an overall high mortality.

In this study, highest mortality of 35.5% was seen in patients with respiratory diseases closely followed

by patients with cardiovascular diseases with a 33.3% mortality which is similar to the findings of Evran

et. al with 30.7% and 16.5% mortality in patients with respiratory and cardiovascular diseases

respectively being the first and second highest.[29] Sepsis had a mortality rate of 25% with lower mean

APACHE II score which is in accordance with the global averages of 20-40%.[30]

Limitations

204 The major limitation of this study is that it is a single center study so its outcomes may not be generalized

205 to all the ICUs of Nepal. This study also has a small sample size which may limit its statistical power. Its

206 short study duration may not be able to capture seasonal variations in ICU admissions. Comparison with

207 other severity scores like SOFA or SAPS was not performed.

208 Conclusion

Our study concludes that APACHE II score is a significant predictor of ICU mortality, modest predictor

of length of ICU stay and less predictive of intermediate discharge outcomes. Higher APACHE II scores

lead to higher mortality with a cut-off value of ≥ 14.5 demonstrating good diagnostic accuracy. Moreover,

lower GCS scores were significantly associated with mortality. APACHE II and GCS can be used as a

practical and cost-effective tool for early prognosis assessment and resource allocation in the ICUs of

Nepal where resources and advanced health care is limited. Further multicenter studies with larger sample

size are needed to validate and extend the findings of this research.

217 References

1. Painter JR. Critical care in the surgical global period. *Chest*. 2013;143(3):851–5. Available

from: https://pubmed.ncbi.nlm.nih.gov/23460163/

2. Trujillano J, Badia M, Servi L, March J, Rodriguez-Pozo A. Stratification of the severity of

critically ill patients with classification trees. *BMC Med Res Methodol*. 2009;9(1):1–12.

Available from: https://pubmed.ncbi.nlm.nih.gov/20003229/

3. Su YY, Li X, Li SJ, Luo R, Ding JP, Wang L, et al. Predicting hospital mortality using

APACHE II scores in neurocritically ill patients: a prospective study. *J Neurol*.

2009;256(9):1427–33. Available from: https://pubmed.ncbi.nlm.nih.gov/19390767/

4. Keuning BE, Kaufmann T, Wiersema R, Granholm A, Pettilä V, Møller MH, et al. Mortality

prediction models in the adult critically ill: a scoping review. *Acta Anaesthesiol Scand*.

2020;64(4):424–42. Available from: https://pubmed.ncbi.nlm.nih.gov/31828760/

5.   Zimmerman JE, Kramer AA, McNair DS, Malila FM. Acute Physiology and Chronic Health

Evaluation (APACHE) IV: hospital mortality assessment for today's critically ill patients.

Crit Care Med. 2006;34(5):1297–310. Available from:

https://pubmed.ncbi.nlm.nih.gov/16540951/

6.   Knaus WA. APACHE 1978–2001: the development of a quality assurance system based on

prognosis: milestones and personal reflections. *Arch Surg*. 2002;137(1):37–41. Available

from: https://pubmed.ncbi.nlm.nih.gov/11772212/

7.   Ko M, Shim M, Lee SM, Kim Y, Yoon S. Performance of APACHE IV in medical intensive

care unit patients: comparisons with APACHE II, SAPS 3, and MPM0 III. *Acute Crit Care*.

2018;33(4):216–21. Available from: https://pubmed.ncbi.nlm.nih.gov/31723888/

8.   Asmarawati TP, Suryantoro SD, Rosyid AN, Marfiani E, Windradi C, Mahdi BA, et al.

Predictive value of SOFA, qSOFA, APACHE II, and NEWS-2 scores to estimate mortality

of COVID-19 patients. *Indian J Crit Care Med*. 2022;26(4):464–71. Available from:

https://pubmed.ncbi.nlm.nih.gov/35656039/

9.   Godinjak A, Iglica A, Rama A, Tančica I, Jusufović S, Ajanović A, et al. Predictive value of

SAPS II and APACHE II scoring systems for patient outcome in a medical intensive care

unit. *Acta Med Acad*. 2016;45(2):97–103. Available from:

https://pubmed.ncbi.nlm.nih.gov/28000485/

10.   Marsh HM, Krishan I, Naessens JM, Strickland RA, Gracey DR, Campion ME, et al.

Assessment of prediction of mortality using the APACHE II scoring system in intensive care

units. *Mayo Clin Proc*. 1990;65(12):1549–57. Available from:

https://pubmed.ncbi.nlm.nih.gov/2123955/

11. Rowan KM, Kerr JH, Major E, McPherson K, Short A, Vessey MP. Intensive Care Society's

APACHE II study in Britain and Ireland–II: outcome comparisons of intensive care units

after adjustment for case mix by the American APACHE II method. *BMJ*. 1991;307:307–12.

Available from: https://pubmed.ncbi.nlm.nih.gov/8241909/

12. Huang J, Xuan D, Li X, Ma L, Zhou Y, Zou H. The value of APACHE II in predicting

mortality after paraquat poisoning in Chinese and Korean population: A systematic review

and meta-analysis. Medicine (Baltimore). 2017 Jul;96(30):e6838. Available from:

https://pmc.ncbi.nlm.nih.gov/articles/pmid/28746171/

13. Demass TB, Guadie AG, Mengistu TB, Belay ZA, Melese AA, Berneh AA, Mihret LG,

Wagaye FE, Bantie GM. The magnitude of mortality and its predictors among adult patients

admitted to the Intensive care unit in Amhara Regional State, Northwest Ethiopia. Sci Rep.

2023 Jul 25;13(1):12010. Available from:

https://pmc.ncbi.nlm.nih.gov/articles/pmid/37491467/

14. Miniksar ÖH, Özdemir M. Clinical features and outcomes of very elderly patients admitted

to the intensive care unit: a retrospective and observational study. *Indian J Crit Care Med*

[Internet]. 2021;25(6):629–34. Available from: https://pubmed.ncbi.nlm.nih.gov/34316141/

15. Bhandari S, Acharya SP, Shrestha GS, Shrestha PS, Paneru HR. A comparison of clinical

profile and outcomes among different age groups of critically ill elderly patients in a tertiary

level hospital in Nepal: a retrospective study. *Asian J Med Sci* [Internet]. 2023;14(4):126–31.

Available from: https://www.nepjol.info/index.php/AJMS/article/view/50716

16. Adhikari R, Shah R, Khanal V. Older people's challenges accessing health services in

central Nepal: a qualitative study using the domain of access framework. *BMC Health Serv*

Res [Internet]. 2024;24(1):1–10. Available from:

https://pubmed.ncbi.nlm.nih.gov/39443984/

17. Patel P, Gupta S, Patel H, Bashar MA. Assessment of APACHE II score to predict ICU

outcomes of patients with AKI: a single-center experience from Haryana, North India.

Indian J Crit Care Med [Internet]. 2022;26(3):276–82. Available from:

https://pmc.ncbi.nlm.nih.gov/articles/PMC9015919/

18. Gong Y, Ding F, Zhang F, Gu Y. Investigating the predictive capacity of in-hospital

mortality of four severity score systems in critically ill patients with acute kidney injury. *J*

Investig Med [Internet]. 2019;67(8):1103–10. Available from:

https://pmc.ncbi.nlm.nih.gov/articles/PMC6900215/

19. Sungono V, Hariyanto H, Soesilo TEB, Adisasmita AC, Syarif S, Lukito AA, et al. Cohort

study of the APACHE II score and mortality for different types of intensive care unit

patients. *Postgrad Med J* [Internet]. 2022;98(1166):914–8. Available from:

https://pubmed.ncbi.nlm.nih.gov/37063012/

20. Haniffa R, Isaam I, De Silva AP, Dondorp AM, De Keizer NF. Performance of critical care

prognostic scoring systems in low- and middle-income countries: a systematic review. *Crit*

Care [Internet]. 2018;22(1):18. Available from:

https://pmc.ncbi.nlm.nih.gov/articles/PMC5787236/

21. Yan ST, Zhang GQ. Predictive performance of critical illness scores and procalcitonin in

sepsis caused by different gram-stain bacteria. *Clinics* [Internet]. 2021;76:e2610. Available

from: https://pmc.ncbi.nlm.nih.gov/articles/PMC8158675/

22. Edipoglu IS, Dogruel B, Dizi S, Tosun M, Çakar N. The association between APACHE II

scores and age groups for predicting mortality in an intensive care unit: a retrospective study.

Rom J Anaesth Intensive Care [Internet]. 2019;26(1):53–7. Available from:

https://pmc.ncbi.nlm.nih.gov/articles/PMC6502281/

23. Huang Y, Chen J, Zhong S, Yuan J. Role of APACHE II scoring system in prediction of

severity and outcome of acute intracerebral hemorrhage. *Int J Neurosci* [Internet].

2016;126(11):1020–4. Available from:

https://www.tandfonline.com/doi/abs/10.3109/00207454.2015.1099099

24. Mumtaz H, Ejaz MK, Tayyab M, Vohra LI, Sapkota S, Hasan M, et al. APACHE scoring as

an indicator of mortality rate in ICU patients: a cohort study. *Ann Med Surg* [Internet].

2023;85(3):416–22. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC10060092/

25. Endeshaw AS, Tarekegn F, Bayu HT, Ayalew SB, Gete BC. The magnitude of mortality and

its determinants in Ethiopian adult intensive care units: a systematic review and meta-

analysis. *Ann Med Surg* [Internet]. 2022;84:104810. Available from:

https://pmc.ncbi.nlm.nih.gov/articles/PMC9793120/

26. Chien YC, Chiang WC, Chen CH, Sun JT, Jamaluddin SF, Tanaka H, et al. Comparison of

on-scene Glasgow Coma Scale with GCS-motor for prediction of 30-day mortality and

functional outcomes of patients with trauma in Asia. *Eur J Emerg Med* [Internet].

2024;31(3):181–7. Available from: https://pubmed.ncbi.nlm.nih.gov/38100651/

27. APACHE-II score correlation with mortality and length of stay in an intensive care unit.

Indian J Crit Care Med [Internet]. 2011;15(3):159–64. Available from:

https://pubmed.ncbi.nlm.nih.gov/21276376/

28. Takekawa D, Endo H, Hashiba E, Hirota K. Predicting prolonged ICU stay using APACHE

II, APACHE III, and SAPS II scores: a Japanese multicenter retrospective cohort study.

PLoS One [Internet]. 2022;17(6):e0269737. Available from:

https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0269737

29. Evran T, Serin S, Gürses E, Sungurtekin H. Various scoring systems for predicting mortality

in intensive care units. *Niger J Clin Pract* [Internet]. 2016;19(4):530–4. Available from:

https://journals.lww.com/njcp/fulltext/2016/19040/various_scoring_systems_for_predicting_

mortality.19.aspx

30. Rhodes A, Evans LE, Alhazzani W, Levy MM, Antonelli M, Ferrer R, et al. Surviving

Sepsis Campaign: international guidelines for management of sepsis and septic shock: 2016.

Intensive Care Med [Internet]. 2017;43(3):304–77. Available from:

https://pubmed.ncbi.nlm.nih.gov/28101605/