Title:

Proteomic Immune Signatures of Severe HIV-Associated Tuberculosis in Sub-Saharan Africa: A Prospective, Multicenter Analysis from Uganda

Authors and Affiliations:

Jesse E. Ross MD,[1] Alin S. Tomoiaga PhD,[2] Nicholas Owor BA,[3] Xuan Lu MPH,[4] Joseph Shinyale RN,[5] Tonny Kiyingi MBChB,[3] Ignatius Asasira MBChB,[3] Peter James Eliku MS,[3] John Bosco Nsubuga MBChB,[5] Christopher Nsereko MBChB,[5] Irene Nayiga MBChB,[5] Stephen Kyebambe MBChB,[5] Thomas Ochar MBChB,[6] Moses Kiwubeyi DMLT,[6] Rittah Nankwanga RN,[6] Kai Nie MSc,[7] Hui Xie MSc,[7] Sam Miake-Lye MS,[7] Bryan Villagomez BS,[7] Jingjing Qi MS,[7] Steven J Reynolds MD,[8,9,10] Martina Cathy Nakibuuka MHS,[10] John Kayiwa MS,[3] Mercy Haumba BA,[3] Joweria Nakaseegu BS,[3] Xiaoyu Che PhD, [11] Risa Hoffman MD,[12] John A Belperio,[1] Julius J. Lutwama PhD,[3] Seunghee Kim-Schulze PhD,[7] Max R. O'Donnell MD,[4,11,13] Barnabas Bakamutumaho MBChB,[3,14†] Matthew J. Cummings MD[4,11†]

[1]Department of Medicine, Division of Pulmonary, Critical Care, Allergy, and Immunology, David Geffen School of Medicine at University of California, Los Angeles, CA, USA.

[2]Department of Accounting, Business Analytics, Computer Information Systems, and Law, Manhattan College, New York, USA

[3]Department of Arbovirology, Emerging and Re-emerging Infectious Diseases, Uganda Virus Research Institute, Entebbe, Uganda

[4]Department of Medicine, Division of Pulmonary, Allergy, and Critical Care Medicine, Vagelos College of Physicians and Surgeons, Columbia University, New York, NY, USA

[5]Entebbe Regional Referral Hospital, Ministry of Health, Entebbe, Uganda

[6]Tororo General Hospital, Ministry of Health, Tororo, Uganda

[7]Human Immune Monitoring Center, Icahn School of Medicine at Mount Sinai, New York, New York

[8]National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD, USA

[9]Johns Hopkins University School of Medicine, Baltimore, MD, USA

[10]Rakai Health Sciences Program, Kalisizo, Uganda

[11]Center for Infection and Immunity, Mailman School of Public Health, Columbia
University, New York, NY, USA

[12]Department of Medicine, Division of Infectious Diseases, David Geffen School of
Medicine at University of California, Los Angeles, USA.

[13]Department of Epidemiology, Mailman School of Public Health, Columbia University,
New York, NY, USA

[14]Immunizable Diseases Unit, Uganda Virus Research Institute, Entebbe, Uganda

†Co-senior authors

Corresponding Author: Jesse E Ross MD, MS, 200 UCLA Medical Plaza Ste 365B.
Los Angeles CA, 90095-8344

Funding: This work was supported by the National Heart Lung and Blood Institute
(5T32HL105323-08 to J.E.R), the National Institute of Allergy and Infectious Diseases
(K23AI16334 and R01AI184997 to M.J.C and the Division of Intramural Research
[S.J.R]), and the National Center for Advancing Translational Sciences (UL1TR001873
to Columbia University, sub-award to M.R.O), National Institutes of Health. Additional
support was provided by the Burroughs Wellcome Fund /American Society of Tropical
Medicine and Hygiene (M.J.C), the Potts Memorial Foundation (J.E.R), and the Stony
Wold-Herbert Fund, Inc. (J.E.R). This research was support in part by the Intramural
Research Program of the NIH. The contributions of the NIH author(s) were made as part
of their official duties as NIH federal employees, are in compliance with agency policy
requirements, and are considered Works of the United States Government. However, the
findings and conclusions presented in this paper are those of the author(s) and do not
necessarily reflect the views of the NIH or the U.S. Department of Health and Human
Services.

Key Words: HIV, Tuberculosis, Africa, Proteomics, Sepsis

Abstract

Objective: Severe tuberculosis (TB) is a major cause of critical illness and death in people living with HIV (PLWH) worldwide. Despite this, the immunopathology of severe HIV-associated TB (HIV/TB) is poorly understood. We aimed to identify an immunopathologic signature of severe HIV/TB in sub-Saharan Africa.

Design and Setting: We analyzed proteomic data from two prospective observational cohorts of adults hospitalized with severe undifferentiated infection in Uganda: an urban discovery cohort (Entebbe, N=241) and a rural validation cohort (Tororo, N=253).

Patients: Adults (age ≥18 years) hospitalized with severe febrile illness

Interventions: None

Measurements and Main Results: Across both cohorts, severe HIV-associated TB was common, affecting 18% of participants in the discovery cohort and 21% in the validation cohort. Overall mortality was significant (30-day mortality of 22% in the discovery cohort & 60-day mortality of 26% in the validation cohort). Participants were stratified into three HIV/TB phenotypes: HIV-negative without TB, PLWH without TB, and PLWH with microbiologically diagnosed TB. We applied ordinal random forest models in the discovery cohort to identify proteins strongly predictive of progressive HIV/TB phenotype. In both cohorts, PLWH with microbiologically diagnosed TB were at highest risk of critical illness and death (30-day mortality of 42% in the discovery cohort & 60-day mortality of 52% in the validation cohort). An eight-protein signature reliably distinguished this phenotype, reflecting mediators of macrophage/dendritic cell activation (LAMP3), NK- and T-cell stimulation and cytotoxicity (CD70, CRTAM), B-cell activation (IGLC2), protease-mediated tissue injury (PRSS2), dysregulated coagulation (SERPINA5), extracellular matrix remodeling (EFEMP1), and GH/IGF axis dysregulation (IGFBP3).

Conclusions: We identified an immunologic signature of severe HIV-associated TB defined by mediators of macrophage/dendritic cell and cytotoxic lymphocyte activation, extracellular matrix remodeling, and dysregulated coagulation. These findings offer new insight into HIV/TB pathobiology and highlight potential targets for host-directed therapies in this high-risk population.

Key Points

Question: What host-response patterns characterize severe HIV-associated tuberculosis among adults hospitalized with severe febrile illness in sub-Saharan Africa?

Findings: In two prospective cohorts of adults hospitalized with severe febrile illness in Uganda, severe HIV-associated tuberculosis accounted for 18-21% of cases and was associated with higher rates of physiological instability and mortality. An eight-protein host-response signature reproducibly distinguished this high-risk phenotype, reflecting immune activation, tissue injury, extracellular matrix remodeling, and dysregulated coagulation.

Meaning: Severe HIV-associated tuberculosis is associated with a distinct, high-risk clinical phenotype characterized by reproducible host-response patterns that may inform risk stratification and host-directed therapeutic strategies.

Introduction:

Severe tuberculosis (TB) is the leading cause of hospitalization and death among people living with HIV (PLWH) globally, with the highest TB burden in low- and middle-income countries (1–4). In sub-Saharan Africa (SSA), where HIV and TB are co-epidemic, MTB is the predominant cause of critical illness and death among hospitalized PLWH (5–6). In this population, the frequency of disseminated MTB infection can exceed 40%, with short-term mortality approaching 50% (6–9).

Despite this substantial burden, the immunopathology of severe HIV-associated tuberculosis (HIV/TB) remains poorly understood. Few studies have characterized multidimensional host responses in PLWH with severe TB, limiting insight into the mechanisms driving immune dysfunction in this critically ill population and, consequently, limiting the ability to target specific treatment mechanisms that may improve outcomes (1, 10 – 17). Available data suggests that MTB infection may be linked to activation of both pro- and anti-inflammatory innate immune pathways, dysregulated T-cell activation and exhaustion, endothelial dysfunction, and extracellular matrix remodeling (1, 10–17), features that partially overlap with those observed in both HIV and critically ill patients with non-TB bacterial infections (18–20). However, most studies to date have relied on narrow biomarker panels, single-center designs, and lacked comparator groups such as HIV-uninfected and TB-negative controls, which reduces immunologic resolution, hinders attribution of TB-specific responses versus HIV or other critical illness, and limits generalizability across settings (1, 10 – 17). Concurrently, emerging evidence from other severe infections underscores substantial heterogeneity in host immune responses, shaped not only by the pathogen but also by host-specific factors such as comorbidities and immune status (19–21). In PLWH, chronic immune activation, T-cell exhaustion, and endothelial dysfunction further influence host-pathogen interactions, potentially giving rise to distinct immunopathologic signatures of critical illness (23–28). This biological complexity underscores the need for higher-resolution immune profiling in severe HIV/TB.

To address these gaps, we conducted high-throughput serum proteomic profiling in two prospective hospital-based cohorts in Uganda to identify a reproducible immune

signature of severe HIV-associated TB. We hypothesized that hospitalized PLWH with severe TB would exhibit a reproducible serum proteomic signature reflecting exaggerated innate immune activation and T-cell exhaustion.

Methods

Study Settings, Outcomes, and Participants

We analyzed data from two prospective observational cohorts of adults hospitalized with severe febrile illness at two public hospitals in Uganda (**Figure S1**). The discovery cohort (RESERVE-U-1-EBB) was enrolled at Entebbe Regional Referral Hospital (ERRH), an urban referral hospital in central Uganda, between April 2017 and August 2019. The validation cohort (RESERVE-U-2-TOR) was enrolled at Tororo General Hospital (TGH), a rural district hospital in eastern Uganda, between November 2021 and August 2023. Enrollment criteria were uniform across the RESERVE-U cohorts. Non-pregnant adults (age ≥18 years) hospitalized with severe febrile illness (i.e., severe undifferentiated infection) were enrolled based on a reported history of fever or axillary temperature ≥ 37.5°C and illness severe enough to warrant admission. The primary clinical outcomes for RESERVE-U-1-EBB and RESERVE-U-2-TOR were 30- and 60-day vital status, respectively. Details of these cohorts have been published and are included in the supplement along with further details of study sites, procedures, and outcomes (18).

Study Procedures

Enrolled participants underwent standardized clinical assessments, including rapid testing for HIV by a Ministry of Health-recommended algorithm (29) and pathogens implicated as etiologies of severe infection in SSA, as informed by World Health Organization (WHO) guidelines (28). For PLWH, MTB testing included Xpert MTB/RIF Ultra (Cepheid, Sunnyvale, CA, USA) on sputum samples and Determine TB-LAM (Abbott/Alere, Lake Bluff, IL, USA) on urine samples, where specimens were obtainable. TB testing of patients without HIV was at the discretion of treating clinicians.

Participants from both cohorts were individually categorized into one of three
ordinal groups reflecting HIV/TB co-infection phenotypes: (1) HIV-negative and TB-
negative (HIV-/TB-), (2) HIV-positive and TB-negative (HIV+/TB-), (3) HIV-positive
and TB-positive (HIV+/TB+). Microbiological diagnosis of TB was based on a positive
sputum Xpert MTB/RIF Ultra or urine TB-LAM result. HIV-uninfected participants were
classified at TB-negative if sputum Xpert MTB/RIF Ultra was negative and there was no
clinical suspicion of TB. PLWH were classified as TB-negative if both sputum Xpert
MTB/RIF Ultra and urine TB-LAM results were negative and there was no clinical
suspicion of TB. In both cases, no clinical suspicion of TB was defined as the treating
clinicians' decision not to initiate empiric anti-TB therapy. Serum samples were collected
at the time of study enrollment and stored at –80°C until the time of analysis.
174

Olink Proteomics
Proteomic profiling was performed on cryopreserved serum samples using the
Olink Target Immunooncology and Cardiometabolic panels (Olink Proteomics AB,
Uppsala, Sweden), which together quantify concentrations of 184 soluble proteins (18).
These panels were utilized to capture multiple biological domains implicated in the host
response to severe infection (i.e., innate and adaptive immune activation, endothelial
dysfunction, and dysregulated coagulation), including severe TB. For each panel, proteins
were quantified via the Olink Proximity Extension Assay (PEA) platform, and relative
protein abundance was expressed in $\log_2$ normalized protein expression (NPX) units.
Proteins were excluded from downstream analyses if <20% of NPX values were above
each panel's estimated limit of detection (18).
186

Biomarker Selection Using Ordinal Forest Models
To identify soluble proteins associated with HIV/TB phenotypes, we applied an
ordinal random forest model to the RESERVE-U-1-EBB discovery cohort, using the
three-level HIV/TB phenotype classification as the ordinal outcome. To mitigate class
imbalance and optimize feature selection across all HIV/TB phenotypes, we performed
class-balanced upsampling through duplicating existing minority class samples, thereby
equalizing group representation during model training. Upsampling was restricted to the

training data within each resampling split, and the independent validation set was never
upsampled. Random forest model parameters included mtry = 5, nsets = 500, ntreeperdiv
= 100, and nbest = 20. The predictive strength of each protein was quantified using
permutation-based variable importance. The top ten proteins were selected from the
importance rankings to provide a parsimonious set of features that balances model
interpretability with sufficient depth for downstream analyses.

Multivariable Analysis of Soluble Proteins and HIV/TB Phenotypes

To quantify associations between soluble proteins and HIV/TB phenotypes, we
analyzed the top 10 proteins identified in the RESERVE-U-1-EBB discovery cohort
using separate multivariable linear regression models. Each model included NPX-
quantified protein expression as the continuous outcome and HIV/TB phenotype as an
ordinal predictor. Models were adjusted for age, sex, illness duration prior to enrollment,
and malaria status. To avoid collider bias, we did not adjust for physiological severity,
which may be influenced by both HIV/TB infection and host responses. We reported
regression coefficients, 95% confidence intervals, and p-values for linear trend,
correcting the latter for multiple testing using the Benjamini-Hochberg false discovery
rate (FDR) method. Proteins with FDR-adjusted p-values <0.05 in the RESERVE-U-1-
EBB discovery cohort were carried forward for testing in the RESERVE-U-2-TOR
validation cohort using identical regression models.

To explore whether proteomic associations with HIV/TB phenotypes were
independent of HIV-related immunosuppression and/or viremia, we conducted a
sensitivity analysis restricted to PLWH in the RESERVE-U-2-TOR cohort. We fit
multivariable linear regression models for each of the top ten proteins, with NPX-
quantified protein expression as the continuous outcome and HIV/TB phenotype as the
ordinal predictor, adjusted for CD4 count and HIV-1 viral load. CD4 count was modeled
as a continuous variable while HIV-1 viral load was modeled as a binary indicator:
suppressed versus unsuppressed. Details of CD4 count and HIV-1 viral load
quantification methods have been described and are in the supplement (**Table S1**) (18).
Age, sex, illness duration, and malaria status were not included in this model to prevent

overfitting and underpowering of this HIV-restricted exploratory analysis. Similarly,
given the smaller sample size and exploratory intent, FDR correction was not applied and
unadjusted p-values are reported. CD4 count and HIV-1 viral load data were unavailable
from RESERVE-U-1-EBB participants.

All analyses were conducted in R version 4.3.2 using the tidyverse, ordinalForest,
caret, stats, broom, and patchwork packages.

Ethics Statement
Each enrolled participant or their surrogate provided written informed consent.
Study protocols were approved by ethics committees at Columbia University
(AAAR1450), Uganda Virus Research Institute (GC/127/17/02-06/582), and Uganda
National Council for Science and Technology (HS2308).

Results
Participant Characteristics

A total of 494 participants were analyzed: 241 in the RESERVE-U-1-EBB
(discovery) cohort and 253 in the RESERVE-U-2-TOR (validation) cohort.
Characteristics of analyzed patients stratified by HIV/TB phenotype are presented in
Tables 1 and 2. Approximately half of participants were female. The median age was 32
years (IQR 26 – 40) in Entebbe and 48 years (IQR 33 – 61) in Tororo. The proportion of
PLWH was similar in Entebbe and Tororo (51% versus 50% respectively), though the
proportion of PLWH receiving anti-retroviral therapy was higher in Tororo (83% versus
68%). In PLWH enrolled in RESERVE-U-2-TOR, the median CD4 count was 350
cells/µL (IQR 300-600) and 71% had suppressed HIV-1 viral loads (<150 copies/mL).
The frequency of microbiologically diagnosed TB was similar across sites: 21% in
Tororo and 18% in Entebbe. Among those with microbiologically diagnosed TB,
disseminated disease as indicated by positive urine TB-LAM testing was present in 40/47
(85%) of PLWH in Entebbe and 51/53 (96%) of PLWH in Tororo. Quick Sequential
Organ Failure Assessment (qSOFA) score greater than or equal to 1, previously shown to

predict infection-related mortality in low- and middle-income countries, was present in
87% and 88% of participants in the discovery and validation cohorts, respectively (30-
32). Mortality at 30- and 60-days was 22% in RESERVE-U-1-EBB and 26% in
RESERVE-U-2-TOR, respectively. Most deaths occurred after hospital discharge. In
both cohorts, PLWH with microbiologically diagnosed TB were at highest risk of
physiological instability and death.

Biomarker Results

After excluding 11 proteins due to high proportions of expression values below
each Olink panel's estimated limits of detection, we applied an ordinal random forest
model in the RESERVE-U-1-EBB discovery cohort to 173 soluble proteins. The top ten
most important proteins for prediction of HIV/TB phenotype reflected macrophage and
dendritic cell activation (LAMP3), NK and T-cell stimulation and cytotoxic function
(CD70, CRTAM, IL-18), B cell activation and immunoglobulin production (IGLC2),
protease-mediated tissue injury (PRSS2), dysregulated coagulation (SERPINA5),
leukocyte adhesion and immune cell trafficking (ICAM3), extracellular matrix
remodeling (EFEMP1), and growth hormone (GH) / insulin-like growth factor (IGF) axis
dysregulation (IGFBP-3) (**Figure S2**).

In the RESERVE-U-1-EBB discovery cohort, all ten proteins were significantly
associated with HIV/TB co-infection phenotype in multivariable linear regression
models, including after FDR adjustment (**Figure 1A, Table S2**). Expression levels of
eight proteins increased across the HIV-/TB-, HIV+/TB-, and HIV+/TB+ groups, while
SERPINA5 and IGFBP-3 demonstrated an inverse association, with lower expression in
participants with severe HIV/TB. In the validation cohort (RESERVE-U-2-TOR), eight
of the ten proteins remained significantly associated with HIV/TB co-infection phenotype
(**Figure 1B, Table S3**); associations for IL-18 and ICAM3 did not reach statistical
significance. Visual inspection of boxplots confirmed consistent expression gradients
across the HIV-/TB-, HIV+/TB-, and HIV+/TB+ groups in both cohorts (**Figures 2A and
2B**).

In exploratory linear regression models restricted to PLWH in RESERVE-U-2-
TOR and adjusted for CD4 count and HIV-1 viral load (N = 127), three proteins,
EFEMP1, IGLC2, and SERPINA5, remained significantly associated with HIV/TB co-
infection phenotype (**Figure S3, Table S4**).

Discussion

We identified and validated a proteomic immune signature associated with
clinical phenotypes of HIV/TB in critically ill hospitalized adults in Uganda. To our
knowledge, this is the first multicenter study in SSA to apply high-throughput proteomics
to define and validate an immune signature of severe HIV-associated TB. These results
advance translational understanding of immune dysregulation in a leading cause of
critical illness and death among PLWH in the region. They also provide a foundation for
biologically-driven stratification of severe HIV-associated TB and could inform the
design of biomarker-guided treatment strategies for this high-risk population.

Our findings align with contemporary models of TB pathogenesis that emphasize
the balance between protective and pathological inflammation (33, 34). In the context of
HIV, loss and dysfunction of CD4 T cells and antigen-presenting cells (APCs)
compromise immune containment, permitting bacillary dissemination and multiorgan
injury (33, 34). The immune signature identified in our study supports this framework
with *in vivo* data from the global HIV/TB epicenter: LAMP3 reflects
maturation/activation of macrophages and dendritic cells and heightened antigen
presentation; CD70 and CRTAM reflect cytotoxic lymphocyte stimulation and potential
exhaustion; IGLC2 denotes B-cell activation and heightened humoral response; PRSS2
and EFEMP1 implicate proteolytic tissue injury and extracellular matrix remodeling;
lower SERPINA5 is consistent with unrestrained serine-protease activity and coagulation
dysregulation leading to an immune-thrombotic milieu; and lower IGFBP-3 is associated
with GH/IGF axis dysregulation, a pattern previously reported with AIDS-associated
wasting and increasing severity of sepsis (35, 36). Collectively, these features define a
reproducible HIV/TB signature linking APC activation, cytotoxic and humoral
lymphocyte stimulation, tissue injury, disordered coagulation, and GH/IGF-axis
perturbation.

315   Prior studies from South Africa have highlighted the complexity of immune
dysregulation in PLWH with severe TB, characterized by simultaneous activation and
exhaustion across innate and adaptive pathways (1, 11, 15). Using relatively low-
dimensional cytokine profiling, one study showed that non-survivors exhibited elevated
concentrations of both pro-inflammatory (CSF-3, TNF, IL-6) and anti-inflammatory (IL-
1Ra) cytokines, alongside expansion of pro-inflammatory monocytes but impaired
capacity of innate cells to mount cytokine responses to bacterial antigens (15). Broader
cytokine panels have identified a clinically high-risk profile dominated by innate and
chemotactic mediators (IL-6, IL-1Ra), whereas T-cell–associated mediators were
inversely correlated with mortality (1). More recently, anemia, upregulation of matrix
metalloproteinases (MMPs), and extracellular matrix turnover have also been linked with
disease severity and poor outcomes in severe HIV/TB (11, 13).

327   A complementary study from South Africa in PLWH quantified circulating MTB
burden using blood-based Xpert MTB/RIF Ultra and demonstrated a dose–response
relationship between bacillary load, host-response signatures of inflammation and tissue
damage, and mortality risk (12). Related work from our group in Uganda showed that
higher urine TB-LAM grade, a marker of bacillary burden, was significantly associated
with increased pro-inflammatory innate and T-cell activation and chemotaxis (10).
Building on these findings, our 173-protein analysis provides a higher-resolution view of
HIV/TB pathogenesis, revealing axes of immune dysregulation not captured by narrower
cytokine panels. Additionally, our work demonstrates external validation across two
hospitals with HIV-negative and TB-negative comparators, improving dissection of
HIV/TB-specific biology across the clinical spectrum. Together, these findings point to a
coordinated and multifaceted host response involving both innate and adaptive pathways,
building on prior studies implicating immune activation in TB severity and poor
outcomes among PLWH. In conjunction with pathogen-centric work, this motivates a
combined pathogen-load plus host-response staging framework to inform risk
stratification and potentially therapeutic trial design in severe HIV/TB.

343   Across the clinical spectrum of HIV/TB co-infection, the respective contributions
of HIV and MTB to immune dysregulation, including mechanisms that drive illness

severity, remain poorly defined (34, 37). Prior studies from South Africa have shown that

HIV viremia and CD4 lymphocytopenia are associated with host inflammatory responses,

with IL-17A, which was not included in our Olink panels, emerging as a central

immunological mediator (33, 34, 37). In contrast, a prior study from Malawi showed that

blunted TNF production in monocytes, suggesting innate immunoparalysis, was strongly

predictive of poor outcomes in severe TB while HIV co-infection was not (17). In an

exploratory analysis restricted to PLWH in the RESERVE-U-2-TOR cohort, three

proteins including IGLC2, SERPINA5, and EFEMP1 remained significantly associated

with HIV/TB phenotype after adjustment for CD4 count and HIV-1 viral load. These

persistent associations suggest that although HIV-related immunosuppression and/or

viremia may contribute to immune dysregulation observed in our cohorts, it does not

fully account for the proteomic signature observed in severe HIV/TB. Nonetheless, given

the exploratory nature of this analysis, our results should be interpreted cautiously and

considered hypothesis-generating.

Beyond identifying an immune signature of severe HIV/TB, our findings advance

understanding of TB-specific immune responses, in contrast to those observed in other

severe HIV-associated infections (18–20). Prior analyses of the RESERVE-U cohorts

identified two pathobiologically-driven "Uganda Sepsis Signatures" (USS-1/USS-2),

with USS-2, enriched for severe HIV/TB and higher mortality, defined by myeloid cell

and inflammasome activation, T-cell exhaustion, endothelial barrier dysfunction and

prothrombotic imbalance (18). Our findings build on these insights, suggesting the

presence of a reproducible immunopathological signature specific to severe HIV-

associated TB in the global sepsis epicenter.

Our findings have direct relevance for the development of host-directed

therapeutics in severe HIV/TB. As diagnostic capabilities expand, protein-based

signatures such as ours could enable precise, biomarker-guided treatment strategies and

support the design of immunomodulatory treatment trials (12, 38). Several candidate

therapies are already being evaluated, including in randomized controlled trials. For

example, corticosteroids in NewStrat-TB aim to attenuate hyperinflammatory immune

activation; doxycycline, a known MMP inhibitor, may mitigate matrix injury and

remodeling, as suggested by preclinical and phase 2 data; aspirin targets the
immunothrombotic axis, with signals of safety and potential benefit demonstrated in TB
meningitis (11, 39–42). Conceptually, a parsimonious protein signature could be
operationalized to enrich randomization in ongoing or future host-directed treatment trials
and serve as an early biomarker endpoint. Moving forward, combining longitudinal
proteomic profiling with quantitative pathogen measurements could jointly stage
bacillary burden and host response. In a serial sampling study of disseminated HIV-
associated TB, MTB DNA remained detectable in blood and urine for up to 14 days after
therapy initiation (43), underscoring the sustained disseminated burden early in treatment
and the potential value of pairing dynamic pathogen and host-response data to refine
therapeutic strategies.

Our study has several strengths. Our primary analysis was performed in two
independent prospective cohorts, across distinct geographic and temporal settings,
encompassing nearly 500 hospitalized adults. These cohorts reflect real-world clinical
populations at high risk for HIV/TB co-infection and mortality, enhancing the
translational relevance of our findings. We employed a tiered analytical approach
combining robust machine learning–based feature selection with prespecified
multivariable models and validation. By treating HIV/TB phenotype as an ordinal
variable, our models captured gradations in clinical and microbiological phenotypes,
rather than relying on binary group comparisons. Predictive associations largely held
across cohorts, increasing generalizability in high-burden settings. The Olink platform
enabled relatively high-throughput measurement of soluble proteins across multiple
biological domains. Importantly, several proteins retained significant associations even
after adjustment for CD4 count and viral load, suggesting components of the signature
reflect TB-specific immunopathology. Together, these strengths support the biological
plausibility, scalability, and external validity of the identified immune signature.

Our study also has limitations. Although the two cohorts were geographically and
temporally distinct, we did not conduct external validation outside Uganda. However,
participant characteristics were broadly comparable to those of comparable cohorts in
SSA (44). While we included only microbiologically confirmed TB cases in the

HIV+/TB+ group, undiagnosed or subclinical TB may have been present in the HIV-/TB-
and HIV+/TB- groups, potentially resulting in misclassification and attenuation of
observed differences biasing results towards the null. Although we attempted to adjust for
co-infections where data were available, including malaria, adjustment for cryptococcal
and other bacterial infections was not possible due to limited testing. Blood cultures were
not performed because of resource limitations. Therefore, residual confounding from
unmeasured or incompletely characterized infections is possible, and our signature may
reflect a more complex infectious background not limited to TB and HIV. We also did
not include a direct measure of circulating mycobacterial burden, which could have
strengthened our interpretation of immune activation patterns. Additionally, CD4 count
and HIV-1 viral load data were only available in the RESERVE-U-2-TOR cohort,
limiting adjustment for HIV disease severity in the discovery cohort. CD4 counts were
multiply imputed in a proportion of PLWH in RESERVE-U-2-TOR (18). However, we
believe this approach is justified, as simulation studies have demonstrated that multiple
imputation reduces bias compared to complete case analysis, even at higher levels of
missingness than our CD4 data (45). Proteomic data were collected at a single time
point, which limits our ability to assess dynamic immune responses over the course of
illness or treatment. Finally, our analysis focused on soluble protein expression, which
may not fully reflect tissue-level immune processes, particularly in lung or intra-cellular
compartments central to TB pathogenesis. Moreover, the Olink platform provides relative
(NPX) assessments of protein expression rather than absolute quantification, which may
limit cross-platform comparability.

Conclusions

In conclusion, we identified a reproducible eight-protein immune signature
associated with HIV/TB co-infection among critically ill hospitalized adults in Uganda.
These findings provide new insights into the immunopathology of severe HIV/TB in this
high-risk population and highlight potential candidate biomarkers that could support
future prognostic and therapeutic research.

Author Contributions

JER, BB, MRO'D, and MJC conceived the study and design. JJL, NO, CN, IN, SK, TO,

MK, RN, JK, MH, and JN collected, organized and entered clinical data and blood
samples and contributed to pathogen diagnostics. SKS, KN, HX, SML, BV contributed to
proteomic analyses. JER, MJC, AT, and XL had full access to all data in the study and
contributed to data cleaning, processing, verification, and analysis. JER developed the
first draft of the manuscript. JER, AT, NO, XL, JS, TK, IA, PJE, JBN, CN, IN, SK, TO,
MK, RN, KN, HX, SML, BV, JQ, SJR, MCN, JK, MH, JN, XC, WIL, RH, JAP, JJL,
SKS, MRO, BB, MJC reviewed and edited the final version of the manuscript. JER had
final responsibility for the decision to submit for publication.

Declaration of Interests

MJC reports consulting fees from Vertex Pharmaceuticals and Veracyte unrelated to the
submitted work. The remaining authors declare no conflicts of interest.

Data sharing

Proteomic data are available in Dryad at https://doi.org/10.5061/dryad.b2rbnzsq2.
Analytic code will be deposited in Github prior to publication.

References

1. Schutz C, Barr D, Andrade B, et al. Clinical, microbiologic, and immunologic determinants of mortality in hospitalized patients with HIV-associated tuberculosis: a prospective cohort study. *PLoS Med*. 2019;16(7): e1002840.

2. UNAIDS. Fact sheet—World Tuberculosis Day 2022. Geneva, Switzerland: Joint United Nations Programme on HIV/AIDS; 2022. Available at: https://www.unaids.org.

3. Ford N, Matteelli A, Shubber Z, et al. TB as a cause of hospitalization and in-hospital mortality among people living with HIV worldwide: a systematic review and meta-analysis. *J Int AIDS Soc.* 2016;19(1):20714.

4. World Health Organization. Global tuberculosis report 2022. Geneva, Switzerland: World Health Organization; 2022. Available at: https://www.who.int.

5. Rudd K, Johnson S, Agesa K, et al. Global, regional, and national sepsis incidence and mortality, 1990–2017: analysis for the Global Burden of Disease Study. *Lancet.* 2020;395:200–211.

6. Cummings M, O'Donnell M. Inverting the pyramid: increasing awareness of mycobacterial sepsis in sub-Saharan Africa. *Int J Tuberc Lung Dis.* 2015;19(10):1128–1134.

7. Reddy E, Shaw A, Crump J. Community-acquired bloodstream infections in Africa: a systematic review and meta-analysis. *Lancet Infect Dis.* 2010;10(6):417–32.

8. Barr D, Lewis J, Feasey N, et al. *Mycobacterium tuberculosis* bloodstream infection prevalence, diagnosis, and mortality risk in seriously ill adults with HIV: a systematic review and meta-analysis of individual patient data. *Lancet Infect Dis.* 2020;20(6):742–752.

9. Gupta R, Lucas S, Fielding K, et al. Prevalence of tuberculosis in post-mortem studies of HIV-infected adults and children in resource-limited settings: a systematic review and meta-analysis. *AIDS.* 2015;29(15):1987–2002.

10. Cummings M, Bakamutumaho B, Jain K, et al. Detection of urine lipoarabinomannan is associated with proinflammatory innate immune activation, impaired host defense, and organ dysfunction in adults with severe HIV-associated tuberculosis in Uganda. *J Acquir Immune Defic Syndr.* 2023;93(1):79–85.

11. Walker N, Schutz C, Ward A, et al**.** Elevated plasma matrix metalloproteinases associate with *Mycobacterium tuberculosis* bloodstream infection and mortality in HIV-associated tuberculosis. *J Infect Dis.* 2025;231(1):109–114

12. Boloko L, Schutz C, Sibiya N, et al. Xpert Ultra testing of blood in severe HIV-associated tuberculosis to detect and measure *Mycobacterium tuberculosis* bloodstream infection: a diagnostic and disease biomarker cohort study. *Lancet Microbe.* 2022;3(7):e521–e532.

13. Araújo-Pereira M, Schutz C, Barreto-Duarte B, et al. Interplay between systemic inflammation, anemia, and mycobacterial dissemination and its impact on mortality in TB-associated HIV: a prospective cohort study. *Front Immunol.* 2023;14:1177432.

14. Ravimohan S, Tamuhla N, Steenhoff A, et al. Immunological profiling of tuberculosis-associated immune reconstitution inflammatory syndrome and non-IRIS death in HIV-infected adults with pulmonary tuberculosis starting antiretroviral therapy: a prospective observational cohort study. *Lancet Infect Dis.* 2015;15(4):429–438.

15. Janssen S, Schutz C, Ward A, et al**.** Mortality in severe human immunodeficiency virus–tuberculosis associates with innate immune activation and dysfunction of monocytes. *Clin Infect Dis.* 2017;65(1):73–82.

16. Clark D, Banura P, Bandeen-Roche K, et al. Biomarkers of endothelial activation/dysfunction distinguish subgroups of Ugandan patients with sepsis and differing mortality risks. *JCI Insight.* 2019;5(10):e127623.

17. Waitt C, Banda N, White S, et al. Early deaths during tuberculosis treatment are associated with depressed innate responses, bacterial infection, and tuberculosis progression. *J Infect Dis.* 2011;204(3):358–62.

18. Cummings M, Lutwama J, Owor N, et al. Unsupervised classification of the host response identifies dominant pathobiological signatures of sepsis in sub-Saharan Africa. *Am J Respir Crit Care Med.* 2025;211(3):357–369.

19. Reddy K, Sinha P, O'Kane C, et al. Subphenotypes in critical care: translation into clinical practice. *Lancet Respir Med.* 2020;8(6):631–643.

20. van der Poll T, Shankar-Hari M, Wiersinga W. The immunology of sepsis. *Immunity.* 2021;54(11):2450–2464

21. Singer M, Deutschman C, Seymour C, et al. The third international consensus definitions for sepsis and septic shock (Sepsis-3). *JAMA.* 2016;315:801–810.

22. Schutz C, Meintjes G, Almajid F, et al. Clinical management of tuberculosis and HIV-1 co-infection. *Eur Respir J.* 2010;36(6):1460–81.

23. Kanmogne GD. HIV infection, antiretroviral drugs, and the vascular endothelium. *Cells.* 2024;13(8):672.

24. Mu W, Patankar V, Kitchen S, et al. Examining chronic inflammation, immune metabolism, and T cell dysfunction in HIV infection. *Viruses.* 2024;16(2):219.

25. Walker N, Meintjes G, Wilkinson R. HIV-1 and the immune response to TB. *Future Virol.* 2013;8(1):57–80.

26. Cummings M, Bakamutumaho B, Price A, et al. HIV infection drives pro-inflammatory immunothrombotic pathway activation and organ dysfunction among adults with sepsis in Uganda. *AIDS.* 2023;37:233–245.

27. Hunt P, Lee S, Siedner M. Immunologic biomarkers, morbidity, and mortality in treated HIV infection. *J Infect Dis.* 2016;214(Suppl 2):S44–S50.

28. World Health Organization. IMAI district clinician manual: hospital care adolescents and adults. Geneva, Switzerland: World Health Organization; 2012. Available at: https://www.who.int/publications/i/item/imai-district-clinician-manual-hospital-care-adolescents-and-adults.

29. Ministry of Health. National HIV Testing Services Policy and Implementation Guidelines: Uganda. 4th Edition. October 2016. Available at: https://library.health.go.ug/sites/default/files/resources/UGANDA%20HTS%20POLICY%20AND%20IMPLEMENTATION%20GUIDELINES%202016%20%282%29.pdf

30. Rudd KE, Seymour CW, Aluisio AR, et al; Sepsis Assessment and Identification in Low Resource Settings (SAILORS) Collaboration: Association of the quick Sequential (Sepsis-Related) Organ Failure Assessment (qSOFA) score with excess hospital mortality in adults with suspected infection in low- and middle-income countries. JAMA 2018; 319:2202–2211

31. Bonnewell JP, Rubach MP, Madut DB, et al: Performance assessment of the universal vital assessment score vs other illness severity scores for predicting risk of in-hospital death among adult febrile inpatients in Northern Tanzania, 2016-2019. JAMA Netw Open 2021; 4:e2136398

32. Cummings M, Lutwama J, Owor N, et al. "Host Response Stratification in Malarial and Non-Malarial Sepsis: A Prospective, Multicenter Analysis From Uganda." *Critical Care Medicine.* [[Baltimore, MD]□:], vol. 53, no. 4, 2025, pp. e984–91, https://doi.org/10.1097/CCM.0000000000006591.

33. Scriba T, Maseeme M, Young C, et al. Immunopathology in human tuberculosis. *Sci Immunol.* 2024;9(102):eado5951.

34. Meintjes G, Maartens G. HIV-associated tuberculosis. *N Engl J Med.* 2024;391(4):343–355.

35. Frost R, Fuhrers J, Steigbigels R, et al. Wasting in the acquired immune deficiency syndrome is associated with multiple defects in the serum insulin-like growth factor system. *Clinical Endocrinology.* 1996;44:501-514

36. Papastathi C, Mavrommatis A, Mentzelopoulos S, et al. Insulin-like growth factor I and its
binding protein 3 in sepsis. *Growth Hormone & IGF Research.* 2013;23(4):98-104.
37. Du Bruyn E, Fukutani K, Rockwood N, et al. Inflammatory profile of patients with tuberculosis
with or without HIV-1 co-infection: a prospective cohort study and immunological network
analysis. *Lancet Microbe.* 2021;2(8):e375–e385.
38. Ross J, Cummings M, Bakamutumaho B, et al. Xpert MTB/RIF Ultra enables sensitive detection
of *Mycobacterium tuberculosis* in blood. *Int J Tuberc Lung Dis.* 2024;28(12):594–596.
39. Namale P, Boloko L, Vermeulen M, et al. Testing novel strategies for patients hospitalised with
HIV-associated disseminated tuberculosis (NewStrat-TB): protocol for a randomised controlled
trial. *Trials.* 2024;25(1):311.
40. Miow Q, Vallejo A, Wang Y, et al. Doxycycline host-directed therapy in human pulmonary
tuberculosis. *J Clin Invest.* 2021;131(15):e141895.
41. Walker N, Clark S, Oni T, et al. Doxycycline and HIV infection suppress tuberculosis-induced
matrix metalloproteinases. *Am J Respir Crit Care Med.* 2012;185(9):989–997.
42. Rohilla R, Shafiq N, Malhotra S. Efficacy and safety of aspirin as an adjunctive therapy in
tubercular meningitis: a systematic review and meta-analysis. *EClinicalMedicine.*
2021;34:100819.
43. Boloko L, Vermeulen M, Sossen B, et al. Blood and urine early treatment response biomarkers in
HIV-associated disseminated tuberculosis. *S Afr J HIV Med.* 2025;26(1):1664.
44. Lewis JM, Feasey NA, Rylance J. Aetiology and outcomes of sepsis in adults in sub-Saharan
Africa: a systematic review and meta-analysis. *Crit Care.* 2019;23:212.
45. Madley-Dowd P, Hughes R, Tilling K, et al. The proportion of missing data should not be used to
guide decisions on multiple imputation. *J Clin Epidemiol.* 2019;110:63–73.
582
583

Table 1: Participant characteristics stratified by HIV/TB phenotype in RESERVE-U-1-EBB cohort.

Characteristics [a]	All participants N = 241	HIV - / TB - N = 117	HIV + / TB - N = 77	HIV + / TB + N=47
Demographics				
Female sex	149/241 (62%)	71/117 (61%)	52/77 (68%)	26/47 (55%)
Age, years	32 (26, 40)	28 (23, 38)	37 (30, 43)	32 (27, 40)
Mid-upper arm circumference, cm	--	--	--	--
Illness duration prior to enrollment, days	4 (2, 6)	3 (2, 5)	4 (3, 7)	6 (3, 7)
Clinical signs and physiological severity scores				
KPS at hospital admission	--	--	--	--
Whole-blood lactate, mmol/L	--	--	--	--
qSOFA score ≥ 1[b]	209/241 (87%)	97/117 (83%)	67/77 (87%	45/47 (96%)
qSOFA score ≥ 2[b]	102/241 (42%)	30/117 (26%)	39/77 (51%)	33/47 (70%)
Modified Early Warning Score	3 (2, 4)	3 (2, 3)	3 (2, 4)	5 (3, 6)
Universal Vital Assessment score	2 (1, 4)	1 (0, 2)	3 (2, 5)	3 (3, 7)
HIV Characteristics				
PLWH	124/241 (51%)	NA	77/77 (100%)	47/47 (100%)
WHO HIV clinical stage 3 or 4	97/124 (78%)	NA	50/77 (65%)	47/47 (100%)
Receiving ART before hospitalization	73/124 (59%)	NA	45/77 (58%)	28/47 (60%)
CD4 cell count (cells/mm^3)	--	--	--	--
HIV-1 viral suppression	--	--	--	--
Pathogen diagnostics				
Malaria RDT positive	54/237 (23%)	30/116 (26%)	20/76 (26%)	4/45 (8.9%)
Urine TB-LAM positive if PLWH	40/99 (40%)	NA	NA	40/47 (85%)
Cryptococcal ag. positive if PLWH	9/29 (31%)	NA	6/13 (46%)	3/16 (19%)

Influenza PCR positive	14/219 (6.4%)	8/106 (7.5%)	5/71 (7.0%)	1/42 (2.4%)
Outcomes				
Death in hospital	14/241 (5.8%)	4/117 (3.4%)	4/77 (5.2%)	6/47 (13%)
Death at 30 days	49/219 (22%)	10/106 (9.4%)	20/68 (29%)	19/45 (42%)

Abbreviations:, KPS: Karnofsky Performance Status, qSOFA: quick Sepsis-related Organ Failure
Assessment, PLWH: person living with HIV; WHO: World Health Organization, ART: anti-retroviral
therapy, RDT: rapid diagnostic test, TB: tuberculosis, LAM: lipoarabinomannan, PCR: polymerase chain
reaction
Legend: [a]Dichotomous and categorical variables summarized as counts (proportions), continuous variables
summarized as medians (interquartile ranges), n if fewer patients with available data than the total number
of patients in the cohort; [b]Systolic blood pressure ≤100mmHg, respiratory rate ≥22 breaths/min, and altered
mental status, latter defined using

Table 2: Participant characteristics stratified by HIV/TB phenotype in RESERVE-
U-2-TOR cohort.

Characteristics [a]	All participants N = 253	HIV - / TB - N = 126	HIV + / TB - N = 73	HIV + / TB + N=54
Demographics				
Female sex	158/253 (62%)	84/126 (67%)	39/73 (53%)	35/54 (65.0%)
Age, years	48 (33, 61)	48 (25, 70)	47 (40, 56)	48 (40, 56)
Mid-upper arm circumference, cm	24 (22, 26)	24 (23, 26)	24 (23, 26)	22 (20, 24)
Illness duration prior to enrollment, days	6 (4, 10)	6 (4, 9)	7 (4, 10)	9 (4, 14)
Clinical signs, lactate, and physiological severity scores				
KPS at hospital admission	60 (20, 70)	70 (20, 80)	60 (40, 70)	50 (20, 70)
Whole-blood lactate, mmol/L	2.4 (1.8, 3.0)	2.4 (1.8, 3.0)	2.4 (1.7, 2.9)	2.4 (2.0, 3.0)
qSOFA score ≥ 1 [b]	222/253 (88%)	110/126 (87%)	65/73 (89%)	47/54 (87%)
qSOFA score ≥ 2 [b]	100/253 (40%)	46/126 (37%)	29/73 (40%)	25/54 (46%)
Modified Early Warning Score	3 (2, 5)	3 (2, 5)	4 (2, 5)	3 (2, 5)
Universal Vital Assessment score	2 (1, 5)	1 (0, 4)	3 (2, 6)	4 (2, 6)
HIV Characteristics				
PLWH	127/253 (50%)	NA	73/73 (100%)	54/54 (100%)
WHO HIV clinical stage 3 or 4	93/127 (73%)	NA	51/73 (70%)	42/54 (78%)
Receiving ART before hospitalization	105/127 (83%)	NA	57/73 (78%)	48/54 (89%)
CD4 count, cells/mm^3	350 (300, 600)	NA	400 (300, 600)	344 (200, 500)
HIV-1 viral suppression	85/120 (71%)	NA	49/68 (72%)	36/52 (69%)
Pathogen diagnostics				
Malaria RDT positive	74/253 (29%)	41/126 (33%)	18/73 (25%)	15/54 (28%)
Urine TB-LAM positive	51/122 (42%)	NA	0/69 (0%)	51/53 (96%)
Cryptococcal ag. positive if PLWH	10/123 (8.1%)	NA	6/70 (8.6%)	4/53 (7.5%)
Influenza PCR positive	14/246 (5.7%)	9/123 (7.3%)	5/72 (6.9%)	0/51 (0%)
SARS-CoV-2 PCR positive	13/246 (5.3%)	8/123 (6.5%)	1/72 (1.4%)	4/51 (7.8%)
Outcomes				

Death in hospital	5/253 (2.0%)	1/126 (0.8%)	1/73 (1.4%)	3/54 (5.6%)
Death at 30 days	51/252 (20%)	11/126 (8.7%)	15/72 (21%)	25/54 (46%)
Death at 60 days	65/252 (26%)	18/126 (14%)	19/72 (26%)	28/54 (52%)

Abbreviations:, KPS: Karnofsky Performance Status, qSOFA: quick Sepsis-related Organ Failure
Assessment, PLWH: person living with HIV; WHO: World Health Organization, ART: anti-retroviral
therapy, RDT: rapid diagnostic test, TB: tuberculosis, LAM: lipoarabinomannan, PCR: polymerase chain
reaction
Legend: [a]Dichotomous and categorical variables summarized as counts (proportions), continuous variables
summarized as medians (interquartile ranges), n if fewer patients with available data than the total number
of patients in the cohort; [b]Systolic blood pressure ≤100mmHg, respiratory rate ≥22 breaths/min, and altered
mental status, latter defined using

Figure 1A. Forest plot showing adjusted regression coefficients and 95% confidence
intervals (CI) for the association between HIV/TB phenotype and protein expression in
the RESERVE-U-1-EBB discovery cohort. Coefficients and 95% CIs generated from
linear regression models including NPX-quantified protein expression as the continuous
outcome and HIV/TB phenotype as an ordinal predictor, adjusted for age, sex, illness
duration prior to enrollment, and malaria status (N = 241).

Figure 1B. Forest plot showing adjusted regression coefficients and 95% confidence

Figure 1B. Forest plot showing adjusted regression coefficients and 95% confidence
intervals (CI) for the association between progressive HIV/TB phenotype and protein
expression in the RESERVE-U-2-TOR validation cohort. Coefficients and 95% CIs
generated from linear regression models including NPX-quantified protein expression as
the continuous outcome and HIV/TB phenotype as an ordinal predictor, adjusted for age,
sex, illness duration prior to enrollment, and malaria status (N = 253).

Protein expression across HIV/TB phenotype (RESERVE-U-1-EBB)

669
670   **Figure 2A**. Boxplots showing distribution of top 10 most predictive proteins across
671 HIV/TB phenotypes in RESERVE-U-1-EBB discovery cohort (N = 241).

Protein expression across HIV/TB phenotype (RESERVE-U-2-TOR)

Figure 2B. Boxplots showing distribution of top 10 most predictive proteins across HIV/TB phenotypes in RESERVE-U-2-TOR validation cohort (N = 253).