Late start of eating is linked to hyperinsulinemia, insulin resistance, and increased body fat in prediabetes

Beeke Peters[1,2] §, Noelle Fröhlich[1,3] §, Jürgen Machann[2,4,5], Ulrike Dambeck[6], Caroline Honsek[6], Christiana Gerbracht[6], Anna Sachno[2,7], Margrit Kemper[2,6], Stefan Kabisch[2,6,7], Andreas Fritsche[2,8,9], Knut Mai[2,7,10,11,12], Andreas F.H. Pfeiffer[2,6,7,11], Olga Pivovarova-Ramich[1,2,6,7] *

[1]*Department of Molecular Metabolism and Precision Nutrition, German Institute of Human Nutrition Potsdam-Rehbruecke, Nuthetal, Germany;*

[2]*German Center for Diabetes Research (DZD), München-Neuherberg, Germany;*

[3]*University of Potsdam, Potsdam, Germany;*

[4]*Institute for Diabetes Research and Metabolic Diseases, Helmholtz Center Munich, Eberhard-Karls University of Tübingen, Tübingen, Germany*

[5]*Department of Radiology, Section on Experimental Radiology, Eberhard-Karls University of Tübingen, Tübingen, Germany*

[6]*Department of Clinical Nutrition, German Institute of Human Nutrition Potsdam-Rehbruecke, Nuthetal, Germany;*

[7]*Department of Endocrinology and Metabolism, European Reference Network on Rare Endocrine Diseases (ENDO-ERN), Charité – Universitätsmedizin Berlin, Corporate Member of Freie Universität Berlin and Humboldt-Universität zu Berlin, 10117 Berlin, Berlin, Germany.*

[8]*Internal Medicine IV, Department of Diabetology, Endocrinology and Nephrology, University of Tübingen, Tübingen, Germany.*

[9]*Institute for Diabetes Research and Metabolic Diseases of the Helmholtz Center Munich at the University of Tübingen, Tübingen, Germany;*

[10]*Department of Human Nutrition, German Institute of Human Nutrition Potsdam-Rehbrücke, Nuthetal, Germany*

[11]*NutriAct-Competence Cluster Nutrition Research Berlin-Potsdam 10117, Germany.*

[12]*Berlin Center for Rare Diseases, Charité-Universitätsmedizin Berlin, Corporate Member of Freie Universität Berlin and Humboldt-Universität zu Berlin, 10117 Berlin, Germany.*

§ B.P. and N.F. contributed equally to this manuscript.

Short title: Eating timing and glucose metabolism in prediabetes

Corresponding author: Olga Pivovarova-Ramich, German Institute of Human Nutrition Potsdam-Rehbruecke, Arthur-Scheunert-Allee 114-116, 14558 Nuthetal, Germany, Tel.: +49 33200882749, Email: olga.ramich@dife.de

Trial registration: NCT01947595, NCT02609243

Keywords: glucose metabolism, insulin sensitivity, hyperinsulinemia, diabetes, eating timing, circadian clocks

Main text word count: 4525; references: 67; tables: 4; figures: 2; supplemental tables: 5; supplemental figures: 1.

Abstract

Background: Prediabetes is associated with an increased risk of progression to type 2 diabetes. While dietary interventions in prediabetes traditionally focus primarily on energy and macronutrient intake, the role of eating timing has recently been highlighted. This study aimed to examine the relationships between eating timing pattern and glycaemic parameters in prediabetes.

Methods: 297 individuals with prediabetes, i.e. impaired fasting glucose or impaired glucose tolerance (age 59.5 y, BMI 31.3 kg/m^2), underwent detailed metabolic phenotyping including an oral glucose tolerance test and magnetic resonance imaging/spectroscopy. Parameters of temporal eating pattern (eating timing, daily calorie distribution, and meal number) were extracted from four-day food records. Eating start (ES) was defined as the start of the first caloric event of the day.

Results: Among eating timing parameters, ES showed the most associations with glycaemic traits. Each later hour of ES was associated with a 1.08 mU/l higher fasting insulin (95% CI, 0.40 to 1.77 mU/l), a 16.86 mU/l higher 120-min insulin (95% CI, 5.62 to 28.10 mU/l), a 1147 mU/l*min higher AUC insulin (95% CI, 390 to 1904 mU/l*min), a 0.32 units higher HOMA-IR (95% CI, 0.14 to 0.50), and a -0.18 units lower Matsuda insulin sensitivity index (95% CI, -0.32 to -0.04) after adjustment for age, sex, and daily energy intake. Glycemic associations remained after additional adjustment for food composition as well as body fat percentage, which was increased in individuals with later ES (0.81% per ES hour; 95% CI, 0.27 to 1.34%). No associations of eating timing with visceral or liver fat were found.

Conclusion: Late ES is associated with estimates of hyperinsulinemia and lower insulin sensitivity in prediabetes, independent of energy intake, food composition, and body fat. Earlier ES might relate to improved glucose metabolism and lower the risk of progression to diabetes.

Funding: This work was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation, RA 3340/4-1 to OP-R, project number 530918029). The DZD is funded

by the German Federal Ministry for Education and Research (BMBF, funding code 01GI0925).

KM was supported by the Competence Cluster Nutrition Research Berlin-Potsdam, funded by

the Federal Ministry of Education and Research (BMBF, funding code 01EA1408) and by the

Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – TRR 412/1 –

project number 535081457.

Abstract word count: 293

Abbreviations

85	AUC	Area under the curve
86	BMI	Body mass index
87	CM	Caloric midpoint
88	ELISA	Enzyme linked immunosorbent assay
89	ES	Eating start
90	FM	Start of the first meal
91	HbA1c	Haemoglobin A1c
92	HDL	High-density lipoprotein
93	HIC	Hepatic insulin clearance
94	HOMA-IR	Homeostatic Model Assessment for Insulin Resistance
95	IFG	Impaired fasting glucose
96	IGT	Impaired glucose tolerance
97	ISI	Insulin sensitivity index
98	LDL	Low-density lipoprotein
99	MASLD	Metabolic dysfunction-associated steatotic liver disease
100	MRI	Magnetic resonance imaging
101	MRS	Magnetic resonance spectroscopy
102	OGTT	Oral glucose tolerance test
103	SD	Standard deviation
104	T2D	Type 2 diabetes
105	TRE	Time-restricted eating

1. Introduction

The pandemic of diabetes mellitus affects millions of people worldwide. Type 2 diabetes (T2D),

the most common form, is preceded by prediabetes, characterized by increasing blood glucose

levels which does not yet achieve the diagnostic criteria for diabetes (1). Dietary interventions,

aiming to impede the progression of prediabetes to diabetes, have traditionally focused on

energy and macronutrient intake. Recent research has highlighted the importance of eating

timing in managing of blood glucose levels and potentially in the prevention of T2D (2; 3). This

fact is based on the diurnal variation of metabolic functions resulting in different metabolic

responses to the same food consumed at different times of the day (4; 5).

Eating timing has emerged as a significant factor in glucose metabolism and insulin

sensitivity (2; 6). Cross-sectional studies have shown that early eating patterns aligning with

individual circadian rhythms are associated with lower body weight and metabolic health (3; 6-

8). In interventional studies, eating earlier in the biological day resulted in greater reductions

in body weight, BMI, and waist circumference, and improved glycaemic control. For instance,

early time-restricted eating (TRE), limiting food intake to a 6-10 hour window early in the day,

improved glucose tolerance and insulin sensitivity in several studies (9-12), although other

research did not confirm this finding (13; 14) and numerous cofounders (longer fasting, fewer

meals, lower calorie intake, and possibly different meal composition) has to be considered. In

contrast, late and night eating showed an association with increased risk of obesity,

cardiovascular diseases, and related traits such as higher BMI and body fat (6; 15-19).

Timing of specific eating events, particularly breakfast, appears to play a crucial role in

glycaemic control. Most previous studies suggest that skipping breakfast has been linked to a

higher risk of developing T2D as partly mediated by BMI (20), while the earlier breakfast timing

was related to the lower T2D incidence (21). Nevertheless, metabolic risks of dinner skipping

were also described (22). This suggests that not only timing of meals but also daily pattern of

calorie distribution may support managing blood glucose levels in diabetes (23; 24).

Additionally, the number of eating events and the duration of overnight fasting have been

investigated for their potential impact on glucose regulation and diabetes risk (3; 25; 26).

Nevertheless, it remains insufficiently studied how temporal eating pattern and its specific

parameters, such as timing, frequency, and diurnal calorie distribution, contribute to the

parameters of glucose metabolism in prediabetes and generally to diabetes risk. Therefore,

this study aimed to examine relationships between eating timing and parameters of glucose

metabolism in individuals with prediabetes.

2. Methods

2.1. Study participants and study protocol

This cross-sectional analysis included baseline data of 297 individuals with prediabetes

obtained in two clinical trials, the Prediabetes Lifestyle Intervention Study (PLIS,

NCT01947595, n=174) and the Diabetes Nutrition Algorithm - Prediabetes (DiNA-P,

NCT02609243, n=123). Complete details of these trials have been described elsewhere (27-

29). Both trials were conducted in accordance with the principles of the Declaration of Helsinki

and Good Clinical Practice guidelines. The study protocol was approved by the ethics

committee of the Charité-Universitätsmedizin Berlin. All patients provided written informed

consent before enrollment.

Data included in this study were collected at the German Institute of Human Nutrition

Potsdam-Rehbruecke and at the Charité-Universitätsmedizin Berlin between 2013 and 2019.

Inclusion criteria in both trials were age between 18 and 75 years, BMI < 45 kg/m², and

diagnosis of impaired fasting glucose (IFG; fasting blood glucose 100-126 mg/dl) and/or

impaired glucose tolerance (IGT; blood glucose at 120 min of oral glucose tolerance test

[OGTT] 140-200 mg/dl) according to the criteria of the American Diabetes Association (1).

Exclusion criteria included overt diabetes mellitus or antidiabetic medication, BMI > 45 kg/m²,

current pregnancy or breastfeeding, serious disease (e.g. symptomatic coronary heart

disease, serious symptomatic malignant disease, severe liver or kidney disease, systemic

infection, severe mental illness), drug abuse, treatment with systemic corticosteroids.

Claustrophobia, metal implants and excessive body circumference were deemed as

contraindication of magnetic-resonance imaging, only. Participants underwent a detailed

metabolic phenotyping including measurement of anthropometric parameters, body

composition, liver fat content, and an OGTT.

Study subjects completed handwritten food records, noting the start and end of each

eating event, the amount and kind of food consumed, for four days including three working

days and one free day to reflect their dietary habits. They did not receive any dietary

recommendations or restrictions before completing the food records.

2.2.   Anthropometric measurements

For the assessment of BMI, body weight was measured using a digital scale and body height

was measured using a stadiometer. Waist and hip circumferences were measured using a

metric tape. Fat mass was analysed using bioelectrical impedance analyser Nutriguard MS

(Data Input GmbH, Pöcking, Germany). Total and visceral adipose tissue (VAT) compartments

of the trunk were assessed by magnetic resonance imaging (MRI). Liver fat content was

determined by localized proton magnetic resonance spectroscopy (^{1}H-MRS) applying a single-

voxel stimulated echo acquisition mode sequence in the posterior hepatic segment 7 as

described (27).

2.3.   OGTT, blood sample analyses, and assessment of glycaemic indices

A standardized 2-hour OGTT was conducted after an overnight fast (>10 h since last food

intake) starting between 8:00 and 9:00. After a fasting blood sampling, participants ingested

standardised glucose solution (75 g glucose in 300 ml water; Accu-Chek Dextro O.G.T.,

Roche), and further blood samples were drawn from the forearm vein 30, 60, 90, and 120 min

after glucose intake. Routine serum parameters were determined using an automated analyzer

(ABX Pentra 4000; ABX, Montpellier, France). Glucose concentrations were analysed locally

by certified laboratories using the glucose oxidase method. Serum insulin and C-peptide

concentrations were determined by immunoassays with the ADVIA Centaur XP Immunoassay

System (Siemens Healthineers, Erlangen, Germany) or with a Tecan infinite M200 microplate

reader (Tecan, Maennedorf, Switzerland).

191 Glucose and insulin values in OGTT were used to calculate indices of insulin sensitivity

192 (homeostasis model assessment of insulin resistance [HOMA-IR] and Matsuda index of insulin

193 sensitivity [ISI Matsuda]), as well as insulin secretion (insulinogenic and disposition indices).

194 HOMA-IR was calculated as ($Gluc_0$ (mmol/l) x Ins_0 (mU/l)) / 22.5. The insulinogenic index was

195 calculated as Ins_{30-0} (mU/l) /$Gluc_{30-0}$ (mg/dl) (30). ISI Matsuda and the disposition index, which

196 reflects the relation between β-cell secretion capacity and insulin sensitivity, were assessed

197 as described (31; 32).

199 *2.4. Analysis of eating timing pattern*

200 The handwritten food records were digitalized using the PRODI software version 4.5

201 (NutriScience GmbH, Hofstetten-Flüh, Switzerland) by an experienced nutritionist. The digital

202 versions were used to assess time, energy intake, and food composition (carbohydrates,

203 protein, fat, and fiber) of each eating event for each of the four consecutive days (three working

204 days and one weekend day) for all participants. Incomplete dietary records were excluded from

205 the analysis. The temporal eating pattern was characterised by the assessment of (i) eating

206 timing itself (i.e., timing of the first and last eating event, eating window duration), (ii) calorie

207 distribution across the day (i.e., time of caloric midpoint and calories in the first and last meal),

208 and (iii) eating frequency (i.e., number of eating events per day) as described in details

209 previously (6). In brief, an eating event was defined by the following rules: (1) at least 50 kcal,

210 and (2) a time gap to another food or meal of at least 15 minutes (33; 34). Start and end of

211 eating were defined as the start of the first caloric event of the day and the end of the last

212 caloric event of the day, respectively. Eating window duration was calculated as the time

213 difference from the start of the first eating event to the end of the last eating event, and the

214 caloric midpoint (CM) was assessed as the time when the 50% of total calorie intake was

215 reached (33).

216 To assess the timing and energy intake in the first and last meals, eating events were

217 classified as a meal or as a snack, based on the time of the day, nutrient density, and regularity

218 as described (6). In brief, an eating event was defined as a "meal" if it had > 15.0 % of daily

energy intake (35), was consumed at the main meal times between 08:00-10:00, 12:00 –

14:00, and 18:00 – 20:00 (typical breakfast, lunch, and dinner times in Germany) (36), and

was regular.

2.5. Statistical Analysis

Statistical analyses were carried out using SPSS 28.0 (IBM SPSS, Chicago, IL). Data

were expressed as mean (SD). Associations between eating timing parameters and glycaemic

and anthropometric traits were examined using multivariable linear regression models to

estimate a standardized regression coefficient (β) and 95% confidence interval (CI). Primary

models were adjusted for potential confounders, including age, sex, and daily energy intake.

Some models were additionally adjusted for macronutrient composition and body fat. Further,

for glycaemic and anthropometric traits, an unstandardized regression coefficient B was shown

to estimate their alterations per each 1-unit change in original predictor scale (e.g. per hour for

the eating time). Group comparison was conducted by Student's t test or by Mann-Whitney U

test. Statistical significance was defined as $p < 0.05$.

3. Results

3.1. Clinical characteristics of participants

Eating pattern was assessed in 297 individuals (age: 59.5 (8.9) years, BMI: 31.3 (6.1) kg/m²,

106 males and 191 females) who completed food records and provided full metabolic data.

Body fat content was 34.9 (8.1) %. Visceral fat accounted for 24.4 (9.2) % of total fat, and liver

fat content was 9.12 (8.00) %. Average fasting glucose was 106.7 (9.5) mg/dL, and HbA1c of

5.7 (0.4) %. Anthropometric measures and parameters of glucose metabolism are shown in

Table 1.

3.2. Eating timing pattern of participants

Daily distribution of eating events is shown in **Figure S1.** Temporal eating pattern of study

participants, including eating timing, calorie distribution across the day, and eating frequency,

is shown in **Table 2**. Participants consumed 4.07 (0.92) meals per day, with an eating start

(ES) at 08:08 (01:24) and an eating end at 20:06 (02:13), resulting in an eating window duration

of 12:09 (01:49) hours (**Table 2**). Energy intake was 27.7 (10.1) % of daily calories in the first

and 33.8 (11.3) % in the last meal, while the CM was estimated at 15:19 (02:04) (**Table 2**).

Eating timing parameters were tightly interrelated. In particular, ES was linked to the

daily calorie distribution, showing positive associations with caloric midpoint (CM, β = 0.332, p

< 0.001) and the percentage of energy in the first (β = 0.178, p = 0.002) and last meal (β =

0.167, p = 0.004). ES was negatively associated with eating duration (β = -0.582, p < 0.001)

and number of meals per day (β = -0.182, p = 0.002). This indicates that individuals who started

eating later had shorter eating windows and consumed fewer meals per day, but the individual

meals, at least the first and last ones, tended to be larger (**Table S1**). Further, CM was

positively associated with calorie intake during the last meal, total daily calorie intake, and the

duration of the daily eating window, and negatively associated with calorie intake during the

first meal. This suggests that individuals with a late CM tended to have a longer eating window

and to consume more evening and daily calories (**Table S1**).

3.3.   Association of eating start and glucose metabolism parameters

In an association analysis between eating timing characteristics and glycaemic traits, most

associations were found for ES. In non-adjusted models, ES showed positive associations with

fasting and 120-min insulin, AUC insulin, HOMA-IR, and a negative association with ISI

Matsuda (**Figure 1A**). To account for potential confounders, the model was adjusted for age,

sex, and daily energy intake, but these associations remained significant (**Figure 1B**). In

adjusted models, each later hour of ES was associated with a 1.08 mU/l higher fasting insulin

(95% CI, 0.40 to 1.77 mU/l), a 16.86 mU/l higher 120-min insulin (95% CI, 5.62 to 28.10 mU/l),

a 1147 mU/l*min higher AUC insulin (95% CI, 390 to 1904 mU/l*min), a 0.32 units higher

HOMA-IR (95% CI, 0.14 to 0.50), and a -0.18 units lower ISI Matsuda (95% CI, -0.32 to -0.04)

(**Table 3**). This suggests that individuals with a late ES tend to have higher insulin levels and

lower insulin sensitivity, independently of the energy intake. Additional adjustment for the daily

intake of carbohydrates, fats, and proteins did not affect these relationships (**Table S2**).

ES was also positively associated with percentage of body fat (0.81% per ES hour;

95% CI, 0.27 to 1.34%) and waist circumference (1.4 cm per ES hour; 95% CI, 0.2 to 2.6 cm),

but not with BMI or liver fat (**Figure 1B**). To prove whether ES associations with insulin levels

and sensitivity are mediated by the excessive body fat, we additionally adjusted the models for

this parameter, but all associations remained significant (**Table 3**).

3.4. ***Associations of other eating timing parameters with glycaemic and***

anthropometric traits

Non-adjusted analysis of other eating timing parameters revealed several associations with

glycaemic traits, most of which disappeared in adjusted models (**Figure 1, Table S3)**. Only,

CM remained positively associated with fasting glucose (0.62 mg/dl per ES hour; 95% CI, 0.10

to 1.15 mg/dl) after adjusting for age, sex, and daily energy intake. Further, calory intake in the

last meal remained positively associated with waist circumference, while daily eating window

was negatively associated with total body fat (**Table S3**). No associations with glycaemic or

anthropometric traits were found for the time of eating end and the daily number of meals

(**Figure 1**).

3.5. ***Comparison of the glycaemic and anthropometric parameters in individuals***

with early vs late start of eating

Based on the observed associations between ES and glycaemic parameters, all participants

were categorized into groups with early ES (before 08:09, which was the average ES in the

study cohort) and with late ES (after 08:09). The groups did not differ in BMI, waist

circumference, visceral and liver fat, and blood lipids, but individuals with late ES showed 2.7

(0.9) % higher body fat (p=0.004) and were on average 1.9 (1.0) years older (p=0.018) (**Table**

3, Table S4).

Participants with late ES ended to eat at a similar time of day as individuals with early

ES, resulting in an average 1 h 42 min shorter eating window (p<0.001) and a 42 min later CM

(p=0.003) (**Table 4**). Daily food composition and calorie intake did not differ between groups,

whereas percentage of energy within the first meal was slightly larger in individuals with late

ES (p=0.041) (**Table S5**). In accordance with the association analysis, in the OGTT,

participants with late ES showed higher insulin (p=0.029) and C-peptide (p=0.034) levels at

120 min of test, and a tendency towards higher fasting insulin (p=0.088), HOMA-IR (p=0.070),

and lower ISI Matsuda (p=0.096) compared to the early ES group, whereas the glucose values

were similar in both groups (**Figure 2, Table 4**).

4. Discussion

As the prevalence of diabetes and prediabetes continues to rise globally, understanding the

relationship between eating timing and glucose metabolism becomes increasingly important,

potentially offering an additional non-pharmacological approach to prevent or delay the

development of T2D. In this study, we demonstrated associations of ES timing with

hyperinsulinemia and insulin resistance, which represent key players in the progression of

prediabetes towards T2D (37) and typically develop 10-15 years before diabetes diagnosis

(38). In our study, individuals with prediabetes underwent comprehensive metabolic

phenotyping, including OGTT, MRI/MRS, and documentation of their nutritional habits using

food diaries. Given this comprehensive approach, the highlight of this study is the detailed

characterization of individual temporal eating patterns through assessment of eating timing

itself and the diurnal distribution of energy intake.

Our analysis revealed a number of associations between eating timing characteristics

and glycaemic parameters, with most associations found for ES. We demonstrated that a later

ES is associated with higher fasting and postprandial insulin values and lower insulin

sensitivity, while no relationship was found between ES timing and glucose values.

Remarkably, these associations were very robust, remaining significant after adjusting for

potential cofounders - age, gender, and energy intake. While ES impact on fasting parameters

(insulin, HOMA-IR) was relatively small or moderate, ES-related differences in post-challenge

insulin responses in OGTT (16.86 mU/L higher 120-min insulin and 1147 mU/L*min higher

AUC insulin for every hour later ES) were more pronounced. For individuals with a 3-hour later

ES, this would translate to an increase of 50.58 mU/L 120-min insulin and of 3441 mU/L*min

AUC insulin, representing substantial hyperinsulinemia and possibly a delayed/late insulin

peaks, strongly associated with increased T2D risk (39). Considering the ES associations with

HOMA-IR and ISI Matsuda, our data suggest that individuals with a late ES have more

pronounced insulin resistance, which apparently results in higher compensatory

hyperinsulinemia to maintain glucose control due to still sufficient beta-cell capacities.

Grouping individuals based on early versus late ES supports this hypothesis, as the late ES

group showed higher 120-minute insulin and C-peptide levels, despite comparable glucose

values.

Importantly, in our study, later ES was also associated higher body fat, a factor, which

strongly relates to obesity and might be another cofounding factor contributing to observed

associations between ES and glycemic traits. Specifically, each later hour of ES was

associated with a 0.81-unit higher body fat percentage after adjusting for age, sex, and energy

intake. Hence, individuals with a 3-hour difference in ES timing would exhibit~2.5% difference

in percentage of body fat, which might affect metabolic and cardiovascular risks, especially if

it crosses established thresholds (40). Notably, body fat content better reflects obesity and is

superior to BMI for obesity-related health risks (41). Our finding on body fat aligns with

published data showing an association of late eating with increased risk of obesity and higher

body fat (7; 42), although we did not observe the previously reported relationship with BMI (6;

8; 15). To investigate whether ES associations with insulin levels and sensitivity are mediated

by the excessive body fat, we additionally adjusted the regression models for this parameter.

Remarkably, ES timing remained associated with glycemic traits, demonstrating that the

effects of ES on glucose metabolism are largely independent of body fat.

Our finding aligns with previously published evidence that timing, calorie intake and

composition of breakfast play an important role in glucose homeostasis and diabetes

management (20; 23; 24; 43). Concerning breakfast timing, large cohort meta-analyses

demonstrate an association between breakfast skipping and increased T2D risk (20; 43). T2D

risk increased with every additional day of breakfast skipping, but the curve reached a plateau

at 4-5 day per week, showing an increased risk of 55%, whereas this effect was only partly

mediated by the weight gain (20). Cross-sectional studies revealed an association of breakfast

skipping with a later chronotype, both of which contribute to poorer glycaemic control, as

indicated by higher HbA1c levels (44). Several observational studies suggest that early ES at

the day confers significant metabolic advantages including lower BMI, blood lipids, and better

glycaemic control (3; 45). For example, data from the National Health and Nutrition

Examination Survey (NHANES) demonstrated that every hour later that eating commenced

was associated with approximately 0.6% higher glucose level and 3% higher HOMA-IR after

adjustment for BMI and other confounders (3). A very recent study has shown that early ES is

associated with lower nocturnal glucose in women with gestational diabetes (46). Notably,

although in our study we did not observe difference in blood glucose, we found robust insulin

changes providing clear evidence that eating timing affects glucose metabolism.

ES timing is tightly associated with calorie distribution over the day. Several studies

demonstrated beneficial effects of implementing a high-calorie breakfast on glycaemic control

in individuals with T2D (23; 24). In contrast, breakfast skipping is frequently accompanied by

late-night dinner consumption, leading to calorie distribution towards the later part of the day.

This misaligns with endogenous metabolic rhythms and can contribute to metabolic

dysfunctions (47), as discussed in detail below. Interestingly, we made a similar observation in

our study, showing a later CM in the late ES group, although the timing of the last eating event

remained unchanged. Growing observational and experimental evidence, summarized in our

and other reviews (5; 48), shows that consuming the majority of calories early in the day

supports lower body weight and better glucose homeostasis compared to late or nighttime

calorie intake, although detailed timing recommendations are still lacking. Notably, in our study,

the calorie intake during the first meal and during the day did not differ in individuals with early

and late ES. Taken together, these data suggest that the timing of ES per se, and not the

calorie intake, is related to glycaemic parameters of study participants.

Interestingly, we also found that individuals with a late ES showed a shorter daily eating

duration as their end of eating window did not differ from participants with an early ES.

However, they showed poorer glycaemic parameters despite the shorter eating window. This

finding contrasts with data from human TRE studies, reducing daily eating duration to less than

10 hours, which demonstrated beneficial effects on glucose homeostasis (49; 50). Thus, our

data suggest that the timing of eating might have a stronger impact on glycaemic parameters

compared to eating duration when daily calorie intake is similar in both groups. Notably, eating

duration in both early and late ES groups was over eleven hours per day, which may also

explain the absence of effects related to eating duration. Finally, similarly to our data, several

cross-sectional or prospective human cohorts have found no link between eating window

duration and body weight or metabolic traits (3; 6; 51), therefore the role of the eating duration

in free-living populations remains debatable.

The next possible cofounding factor, which might affect the observed ES effects, is the

daily macronutrient composition or composition of breakfast meals, which exerts profound

short-term and long-term effects on glycaemic and insulinemic responses. Studies reducing

breakfast carbohydrates and replacing them with proteins or fats demonstrated lower glucose,

decreased glycaemic variability, improved β-cell responsiveness, and HbA1c in subjects with

T2D (24; 52; 53). Based on this data, we examined in our study whether food composition

might contribute to the observed ES associations with insulin levels or insulin sensitivity, but

they remained significant after the additional adjustment for the macronutrient intake. We also

found no differences in the macronutrient composition of the first meal or the daily average

when comparing individuals with early and late ES.

Previous research has largely highlighted the CM as an eating time-related parameter,

demonstrating robust associations with obesity-related traits such as BMI, waist

circumference, and body fat (7; 8; 15). In addition, we recently provided evidence linking CM

and insulin sensitivity in healthy subjects (6), underscoring the essential role of daily calorie

distribution for glucose metabolism. In contrast to published data, our present study observed

a much stronger relationship of ES with glycaemic regulation, specifically with insulin levels

and sensitivity, than CM. This difference likely stems from the different study cohorts, as the

CM associations were primarily observed in cohorts without focus on diabetes (7; 8; 15).

The mechanistic basis for these timing effects likely involves the circadian regulation of

glucose metabolism and insulin sensitivity by endogenous clocks (54). In healthy subjects,

insulin sensitivity in skeletal muscle and adipose tissue is generally higher in the early morning,

facilitating rapid and efficient postprandial glucose uptake (55; 56). Similarly, pancreatic β-cells

show enhanced insulin release in the morning, which effectively suppresses hepatic

gluconeogenesis after breakfast (57). As the day progresses, insulin action gradually declines

due to both reduced peripheral glucose uptake and diminished β-cell responses, resulting in

larger glycaemic excursions in response to the same glycaemic load (57). Aligning food intake

with the body's natural metabolic rhythms – namely, early in the day – putatively leads to

optimal metabolic regulation. Eating late opposes these natural rhythms and has the potential

to cause circadian misalignment and metabolic dysfunction (47; 58). Remarkably, several

studies showed altered circadian rhythms in individuals with T2D, with a flattened profile of

insulin sensitivity and β-cell function, and elevated gluconeogenesis in the morning (59-61).

Whether individuals with prediabetes also have altered metabolic rhythmicity remains

insufficiently investigated (62), but if this is the case, it may help explain why our association

data differ from those observed in metabolically healthy subjects. Nevertheless, our findings in

early and late ES groups support the idea that individuals with prediabetes may also benefit

from early eating, possibly acting as a zeitgeber and, thereby, improving their metabolic

rhythmicity (5; 47).

Our study has several important strengths. The main strength is the detailed metabolic

and dietary habit phenotyping in a large cohort of individuals with prediabetes, which allowed

us to assess the association of eating timing with glycaemic traits – an assessment never

conducted previously. Using of OGTT allowed the analysis of both fasting and postprandial

metabolism, including insulin levels and the calculation of specific indices. A further strength

of our study is the analysis of multiple components of the eating pattern and their potential

relation to glycaemic regulation.

The prominent novelty and strength of our study is the analysis of the association of

eating timing with liver fat content. Liver fat accumulation, described as MASLD, often

accompanies prediabetes and T2D and strongly contributes to metabolic dysfunctions (63).

Liver fat content is regulated by circadian clocks, as shown in animal models (64). Further,

TRE interventions have been shown to ameliorate liver fat content in mice fed with high-fat diet

(65) and in humans, with an effect size comparable to calorie restriction (66). Based on known

associations of eating timing with abdominal obesity (7) and insulin sensitivity (6), we expected

to observe an association of eating timing with liver fat in prediabetes. Surprisingly, no eating

timing parameters showed a relation to liver fat content in our cohort. This data expands our

knowledge in this field and requires further investigation in larger cohorts and diverse

categories of individuals with and without metabolic dysfunctions.

Some limitations have to be also discussed when interpreting the results of this work.

First, food intake and timing data were self-reported using the food diaries. Although this

method is recognized as a gold standard for nutritional data collection, it does not exclude

measurement errors, such as over- and underreporting. Future studies should include

objective monitoring of eating behaviour using wearables or smartphone apps. Secondly, no

data on sleep timing or quality, which are strongly associated with metabolism and can worsen

glycaemic parameters upon the sleep deficit (67; 68), were collected. Moreover, collecting

sleep time data or questionnaires to assess individual chronotypes would help to decipher how

the relation between eating time and chronotype is linked to the diabetes development. Thirdly,

the analysis did not consider the socioeconomic status of participants. Breakfast skipping

associated with poor socioeconomic status could also explain obesity and the impaired

metabolic state. Further, practicing of intermittent fasting in form of breakfast skipping could

also be associated with poor metabolic state in the sense of reverse causality: as an attempt

to overcome prediabetes instead of being the reason for its development. Finally, the selection

bias by cutting of patients with stronger glycemic impairment (overt diabetes) has to be

mentioned as a limitation.

In conclusion, our study showed that a late ES is with estimates of hyperinsulinemia

and lower insulin sensitivity in prediabetes, independent of energy intake, food composition,

and body fat. Our findings suggest that an earlier ES might relate to improved glucose

metabolism and lower the risk of progression from prediabetes to diabetes. This non-

pharmacological approach needs to be investigated in interventional studies.

Acknowledgements

We thank all study participants for their cooperation. We gratefully acknowledge the excellent
assistance of dietitians, study nurses, and lifestyle advisors involved in this study. We also
thank Elwira Gliwska for her contribution to the algorithm development for the analysis of meal
timing.

Data availability

The data that support the findings of this study are available from the corresponding author
upon reasonable request

Funding

This work was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research
Foundation, RA 3340/4-1 to OP-R, project number 530918029). The DZD is funded by the
German Federal Ministry for Education and Research (BMBF, funding code 01GI0925). KM
was supported by the Competence Cluster Nutrition Research Berlin-Potsdam, funded by the
German Federal Ministry of Education and Research (BMBF, funding code 01EA1408) and by
the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – TRR 412/1 –
project number 535081457. The funders had no role in study design, data collection and
analysis, decision to publish, or preparation of the manuscript.

Authors' relationships and activities

S.K. received grants from the German Diabetes Association, the Wilhelm Doerenkamp
Foundation, the Almond Board of California and the California Walnut Commission, and lecture
fees from Sanofi, Boehringer Ingelheim, Berlin Chemie, Lilly Deutschland and Juzo Akademie.
The remaining authors declare that there are no relationships or activities that might bias, or
be perceived to bias, their work.

Contribution statement

References

1. American Diabetes Association Professional Practice C. 2. Diagnosis and Classification of Diabetes: Standards of Care in Diabetes-2025. Diabetes care 2025;48:S27-S49

2. Diaz-Rizzolo DA, Santos Baez LS, Popp CJ, Borhan R, Sordi-Guth A, Manoogian ENC, Panda S, Cheng B, Laferrere B. Late eating is associated with poor glucose tolerance, independent of body weight, fat mass, energy intake and diet composition in prediabetes or early onset type 2 diabetes. Nutrition & diabetes 2024;14:90

3. Ali M, Reutrakul S, Petersen G, Knutson KL. Associations between Timing and Duration of Eating and Glucose Metabolism: A Nationally Representative Study in the U.S. Nutrients 2023;15

4. Asher G, Sassone-Corsi P. Time for food: the intimate interplay between nutrition, metabolism, and the circadian clock. Cell 2015;161:84-92

5. Peters B, Vahlhaus J, Pivovarova-Ramich O. Meal timing and its role in obesity and associated diseases. Frontiers in endocrinology 2024;15:1359772

6. Vahlhaus J, Peters B, Hornemann S, Ost AC, Kruse M, Busjahn A, Pfeiffer AFH, Pivovarova-Ramich O. Later eating timing in relation to an individual internal clock is associated with lower insulin sensitivity and affected by genetic factors. EBioMedicine 2025;116:105737

7. McHill AW, Phillips AJ, Czeisler CA, Keating L, Yee K, Barger LK, Garaulet M, Scheer FA, Klerman EB. Later circadian timing of food intake is associated with increased body fat. Am J Clin Nutr 2017;106:1213-1219

8. Martinez-Lozano N, Tvarijonaviciute A, Rios R, Baron I, Scheer F, Garaulet M. Late Eating Is Associated with Obesity, Inflammatory Markers and Circadian-Related Disturbances in School-Aged Children. Nutrients 2020;12

9. Sutton EF, Beyl R, Early KS, Cefalu WT, Ravussin E, Peterson CM. Early Time-Restricted Feeding Improves Insulin Sensitivity, Blood Pressure, and Oxidative Stress Even without Weight Loss in Men with Prediabetes. Cell metabolism 2018;27:1212-1221 e1213

10. Jones R, Pabla P, Mallinson J, Nixon A, Taylor T, Bennett A, Tsintzas K. Two weeks of early time-restricted feeding (eTRF) improves skeletal muscle insulin and anabolic sensitivity in healthy men. Am J Clin Nutr 2020;112:1015-1028

11. Jamshed H, Beyl RA, Della Manna DL, Yang ES, Ravussin E, Peterson CM. Early Time-Restricted Feeding Improves 24-Hour Glucose Levels and Affects Markers of the Circadian Clock, Aging, and Autophagy in Humans. Nutrients 2019;11

12. Teong XT, Liu K, Vincent AD, Bensalem J, Liu B, Hattersley KJ, Zhao L, Feinle-Bisset C, Sargeant TJ, Wittert GA, Hutchison AT, Heilbronn LK. Intermittent fasting plus early time-restricted eating versus calorie restriction and standard care in adults at risk of type 2 diabetes: a randomized controlled trial. Nature medicine 2023;29:963-972

13. Andriessen C, Fealy CE, Veelen A, van Beek SMM, Roumans KHM, Connell NJ, Mevenkamp J, Moonen-Kornips E, Havekes B, Schrauwen-Hinderling VB, Hoeks J, Schrauwen P. Three weeks of time-restricted eating improves glucose homeostasis in adults with type 2 diabetes but does not improve insulin sensitivity: a randomised crossover trial. Diabetologia 2022;65:1710-1720

14. Peters B, Schwarz J, Schuppelius B, Ottawa A, Koppold DA, Weber D, Steckhan N, Mai K, Grune T, Pfeiffer AFH, Michalsen A, Kramer A, Pivovarova-Ramich O. Intended isocaloric time-restricted eating shifts circadian clocks but does not improve cardiometabolic health in women with overweight. Sci Transl Med 2025;17:eadv6787

15. Crispim CA, Rinaldi AEM, Azeredo CM, Skene DJ, Moreno CRC. Is time of eating associated with BMI and obesity? A population-based study. Eur J Nutr 2024;63:527-537

16. Dashti HS, Gomez-Abellan P, Qian J, Esteban A, Morales E, Scheer F, Garaulet M. Late eating is associated with cardiometabolic risk traits, obesogenic behaviors, and impaired weight loss. Am J Clin Nutr 2021;113:154-161

17. Shim JS, Kim HC. Late eating, blood pressure control, and cardiometabolic risk factors among adults with hypertension: results from the Korea National Health and Nutrition Examination Survey 2010-2018. Epidemiol Health 2021;43:e2021101

18. Nakajima K, Suwa K. Association of hyperglycemia in a general Japanese population with late-
night-dinner eating alone, but not breakfast skipping alone. J Diabetes Metab Disord 2015;14:16
19. Scheer FA, Hilton MF, Mantzoros CS, Shea SA. Adverse metabolic and cardiovascular
consequences of circadian misalignment. Proceedings of the National Academy of Sciences of the
United States of America 2009;106:4453-4458
20. Ballon A, Neuenschwander M, Schlesinger S. Breakfast Skipping Is Associated with Increased Risk
of Type 2 Diabetes among Adults: A Systematic Review and Meta-Analysis of Prospective Cohort
Studies. The Journal of nutrition 2019;149:106-113
21. Palomar-Cros A, Srour B, Andreeva VA, Fezeu LK, Bellicha A, Kesse-Guyot E, Hercberg S,
Romaguera D, Kogevinas M, Touvier M. Associations of meal timing, number of eating occasions and
night-time fasting duration with incidence of type 2 diabetes in the NutriNet-Sante cohort. Int J
Epidemiol 2023;52:1486-1497
22. Yamamoto R, Tomi R, Shinzawa M, Yoshimura R, Ozaki S, Nakanishi K, Ide S, Nagatomo I, Nishida
M, Yamauchi-Takihara K, Kudo T, Moriyama T. Associations of Skipping Breakfast, Lunch, and Dinner
with Weight Gain and Overweight/Obesity in University Students: A Retrospective Cohort Study.
Nutrients 2021;13
23. Jakubowicz D, Wainstein J, Ahren B, Bar-Dayan Y, Landau Z, Rabinovitz HR, Froy O. High-energy
breakfast with low-energy dinner decreases overall daily hyperglycaemia in type 2 diabetic patients:
a randomised clinical trial. Diabetologia 2015;58:912-919
24. Jakubowicz D, Wainstein J, Landau Z, Ahren B, Barnea M, Bar-Dayan Y, Froy O. High-energy
breakfast based on whey protein reduces body weight, postprandial glycemia and HbA(1C) in Type 2
diabetes. J Nutr Biochem 2017;49:1-7
25. Dai J, Nianogo R, Wong ND, Moin T, McClain AC, Alver S, Cordero C, Daviglus ML, Qi Q, Sotres-
Alvarez D, Chen L. Energy Intake and Dietary Glycemic Load in Late Morning and Risk of Type 2
Diabetes: The Hispanic Community Health Study/Study of Latinos-A Multicenter Prospective Cohort
Study. Diabetes care 2024;47:1673-1681
26. Kahleova H, Belinova L, Malinska H, Oliyarnyk O, Trnovska J, Skop V, Kazdova L, Dezortova M,
Hajek M, Tura A, Hill M, Pelikanova T. Eating two larger meals a day (breakfast and lunch) is more
effective than six smaller meals in a reduced-energy regimen for patients with type 2 diabetes: a
randomised crossover study. Diabetologia 2014;57:1552-1560
27. Fritsche A, Wagner R, Heni M, Kantartzis K, Machann J, Schick F, Lehmann R, Peter A, Dannecker
C, Fritsche L, Valenta V, Schick R, Nawroth PP, Kopf S, Pfeiffer AFH, Kabisch S, Dambeck U, Stumvoll
M, Bluher M, Birkenfeld AL, Schwarz P, Hauner H, Clavel J, Seissler J, Lechner A, Mussig K, Weber K,
Laxy M, Bornstein S, Schurmann A, Roden M, de Angelis MH, Stefan N, Haring HU. Different Effects of
Lifestyle Intervention in High- and Low-Risk Prediabetes: Results of the Randomized Controlled
Prediabetes Lifestyle Intervention Study (PLIS). Diabetes 2021;70:2785-2795
28. Wagner R, Heni M, Kantartzis K, Sandforth A, Machann J, Schick F, Peter A, Fritsche L, Szendrodi J,
Pfeiffer AFH, Schurmann A, Bluher M, Hauner H, Seissler J, Bornstein S, Roden M, Stefan N,
Birkenfeld AL, White MF, Haring HU, Fritsche A. Lower Hepatic Fat Is Associated With Improved
Insulin Secretion in a High-Risk Prediabetes Subphenotype During Lifestyle Intervention. Diabetes
2023;72:362-366
29. Kabisch S, Bather S, Dambeck U, Kemper M, Gerbracht C, Honsek C, Sachno A, Pfeiffer AFH. Liver
Fat Scores Moderately Reflect Interventional Changes in Liver Fat Content by a Low-Fat Diet but Not
by a Low-Carb Diet. Nutrients 2018;10
30. Stumvoll M, Mitrakou A, Pimenta W, Jenssen T, Yki-Jarvinen H, Van Haeften T, Renn W, Gerich J.
Use of the oral glucose tolerance test to assess insulin release and insulin sensitivity. Diabetes care
2000;23:295-301
31. Bergman RN, Ader M, Huecking K, Van Citters G. Accurate assessment of beta-cell function: the
hyperbolic correction. Diabetes 2002;51 Suppl 1:S212-220
32. Matsuda M, DeFronzo RA. Insulin sensitivity indices obtained from oral glucose tolerance testing:
comparison with the euglycemic insulin clamp. Diabetes care 1999;22:1462-1470
33. Gill S, Panda S. A Smartphone App Reveals Erratic Diurnal Eating Patterns in Humans that Can Be
Modulated for Health Benefits. Cell metabolism 2015;22:789-798

34. Gibney MJ, Wolever TM. Periodicity of eating and human health: present perspective and future
directions. Br J Nutr 1997;77 Suppl 1:S3-5
35. Johnson GH, Anderson GH. Snacking definitions: impact on interpretation of the literature and
dietary recommendations. Crit Rev Food Sci Nutr 2010;50:848-871
36. Silva CM, Teixeira BS, Wright KP, Jr., Maia YCP, Crispim CA. Time-Related Eating Patterns Are
Associated with the Total Daily Intake of Calories and Macronutrients in Day and Night Shift Workers.
Nutrients 2022;14
37. Ahlqvist E, Storm P, Karajamaki A, Martinell M, Dorkhan M, Carlsson A, Vikman P, Prasad RB, Aly
DM, Almgren P, Wessman Y, Shaat N, Spegel P, Mulder H, Lindholm E, Melander O, Hansson O,
Malmqvist U, Lernmark A, Lahti K, Forsen T, Tuomi T, Rosengren AH, Groop L. Novel subgroups of
adult-onset diabetes and their association with outcomes: a data-driven cluster analysis of six
variables. Lancet Diabetes Endocrinol 2018;6:361-369
38. Freeman AM, Acevedo LA, Pennings N. Insulin Resistance. In *StatPearls* Treasure Island (FL), 2025
39. Kekalainen P, Sarlund H, Pyorala K, Laakso M. Hyperinsulinemia cluster predicts the development
of type 2 diabetes independently of family history of diabetes. Diabetes care 1999;22:86-92
40. Macek P, Biskup M, Terek-Derszniak M, Stachura M, Krol H, Gozdz S, Zak M. Optimal Body Fat
Percentage Cut-Off Values in Predicting the Obesity-Related Cardiovascular Risk Factors: A Cross-
Sectional Cohort Study. Diabetes Metab Syndr Obes 2020;13:1587-1597
41. Salmon-Gomez L, Catalan V, Fruhbeck G, Gomez-Ambrosi J. Relevance of body composition in
phenotyping the obesities. Rev Endocr Metab Disord 2023;24:809-823
42. Thomas EA, Zaman A, Cornier MA, Catenacci VA, Tussey EJ, Grau L, Arbet J, Broussard JL, Rynders
CA. Later Meal and Sleep Timing Predicts Higher Percent Body Fat. Nutrients 2020;13
43. Bi H, Gan Y, Yang C, Chen Y, Tong X, Lu Z. Breakfast skipping and the risk of type 2 diabetes: a
meta-analysis of observational studies. Public Health Nutr 2015;18:3013-3019
44. Reutrakul S, Hood MM, Crowley SJ, Morgan MK, Teodori M, Knutson KL. The relationship
between breakfast skipping, chronotype, and glycemic control in type 2 diabetes. Chronobiology
international 2014;31:64-71
45. Kant AK, Graubard BI. Clock Time of First Eating Episode and Prospective Risk of All-Cause
Mortality in US Adults. The Journal of nutrition 2022;152:217-226
46. Cunningham HA, Ward L, Butler MP, Valent AM. Early meal timing improves nocturnal glucose in
pregnancies complicated by gestational diabetes. Diabetologia 2026;
47. Qian J, Scheer FA. Circadian System and Glucose Metabolism: Implications for Physiology and
Disease. Trends in endocrinology and metabolism: TEM 2016;27:282-293
48. Lopez-Minguez J, Gomez-Abellan P, Garaulet M. Timing of Breakfast, Lunch, and Dinner. Effects
on Obesity and Metabolic Risk. Nutrients 2019;11
49. Ezpeleta M, Cienfuegos S, Lin S, Pavlou V, Gabel K, Tussing-Humphreys L, Varady KA. Time-
restricted eating: Watching the clock to treat obesity. Cell metabolism 2024;36:301-314
50. Schuppelius B, Peters B, Ottawa A, Pivovarova-Ramich O. Time Restricted Eating: A Dietary
Strategy to Prevent and Treat Metabolic Disturbances. Frontiers in endocrinology 2021;12:683140
51. Zhao D, Guallar E, Woolf TB, Martin L, Lehmann H, Coughlin J, Holzhauer K, Goheer AA, McTigue
KM, Lent MR, Hawkins M, Clark JM, Bennett WL. Association of Eating and Sleeping Intervals With
Weight Change Over Time: The Daily24 Cohort. J Am Heart Assoc 2023;12:e026484
52. Chang CR, Francois ME, Little JP. Restricting carbohydrates at breakfast is sufficient to reduce 24-
hour exposure to postprandial hyperglycemia and improve glycemic variability. Am J Clin Nutr
2019;109:1302-1309
53. Bauer LB, Reynolds LJ, Douglas SM, Kearney ML, Hoertel HA, Shafer RS, Thyfault JP, Leidy HJ. A
pilot study examining the effects of consuming a high-protein vs normal-protein breakfast on free-
living glycemic control in overweight/obese 'breakfast skipping' adolescents. Int J Obes (Lond)
2015;39:1421-1424
54. Poggiogalle E, Jamshed H, Peterson CM. Circadian regulation of glucose, lipid, and energy
metabolism in humans. Metabolism: clinical and experimental 2018;84:11-27
55. Carrasco-Benso MP, Rivero-Gutierrez B, Lopez-Minguez J, Anzola A, Diez-Noguera A, Madrid JA,
Lujan JA, Martinez-Augustin O, Scheer FA, Garaulet M. Human adipose tissue expresses intrinsic

circadian rhythm in insulin sensitivity. FASEB journal : official publication of the Federation of
American Societies for Experimental Biology 2016;30:3117-3123

56. van Moorsel D, Hansen J, Havekes B, Scheer FAJL, Jörgensen JA, Hoeks J, Schrauwen-Hinderling
VB, Duez H, Lefebvre P, Schaper NC, Hesselink MKC, Staels B, Schrauwen P. Demonstration of a day-
night rhythm in human skeletal muscle oxidative capacity. Molecular Metabolism 2016;5:635-645

57. Saad A, Dalla Man C, Nandy DK, Levine JA, Bharucha AE, Rizza RA, Basu R, Carter RE, Cobelli C,
Kudva YC, Basu A. Diurnal pattern to insulin secretion and insulin action in healthy individuals.
Diabetes 2012;61:2691-2700

58. McHill AW, Butler MP. Eating Around the Clock: Circadian Rhythms of Eating and Metabolism.
Annu Rev Nutr 2024;44:25-50

59. SHAPIRO ET, POLONSKY KS, COPINSCHI G, BOSSON D, TILLIL H, BLACKMAN J, LEWIS G, CAUTER
EV. Nocturnal Elevation of Glucose Levels during Fasting in Noninsulin-Dependent Diabetes*. The
Journal of Clinical Endocrinology & Metabolism 1991;72:444-454

60. Boden G, Chen X, Urbain JL. Evidence for a Circadian Rhythm of Insulin Sensitivity in Patients With
NIDDM Caused by Cyclic Changes in Hepatic Glucose Production. Diabetes 1996;45:1044-1050

61. Mason IC, Qian J, Adler GK, Scheer F. Impact of circadian disruption on glucose metabolism:
implications for type 2 diabetes. Diabetologia 2020;63:462-472

62. Sonnier T, Rood J, Gimble JM, Peterson CM. Glycemic control is impaired in the evening in
prediabetes through multiple diurnal rhythms. J Diabetes Complications 2014;28:836-843

63. Demir M, Bornstein SR, Mantzoros CS, Perakakis N. Liver fat as risk factor of hepatic and
cardiometabolic diseases. Obesity reviews : an official journal of the International Association for the
Study of Obesity 2023;24:e13612

64. de Assis LVM, Demir M, Oster H. The role of the circadian clock in the development, progression,
and treatment of non-alcoholic fatty liver disease. Acta Physiol (Oxf) 2023;237:e13915

65. Hatori M, Vollmers C, Zarrinpar A, DiTacchio L, Bushong EA, Gill S, Leblanc M, Chaix A, Joens M,
Fitzpatrick JAJ, Ellisman MH, Panda S. Time-Restricted Feeding without Reducing Caloric Intake
Prevents Metabolic Diseases in Mice Fed a High-Fat Diet. Cell metabolism 2012;15:848-860

66. Oh JH, Yoon EL, Park H, Lee S, Jo AJ, Cho S, Kwon E, Nah EH, Lee JH, Park JH, Ahn SB, Jun DW.
Efficacy and safety of time-restricted eating in metabolic dysfunction-associated steatotic liver
disease. Journal of hepatology 2025;

67. Reutrakul S, Van Cauter E. Sleep influences on obesity, insulin resistance, and risk of type 2
diabetes. Metabolism: clinical and experimental 2018;84:56-66

68. Schmid SM, Hallschmid M, Schultes B. The metabolic burden of sleep loss. Lancet Diabetes
Endocrinol 2015;3:52-62

Figure 1. Association of eating timing, glycaemic and anthropometric parameters. Heatmap represents beta-coefficient values calculated using

linear regression models without (A) and with (B) adjustment for sex, age, and daily energy intake. Eating timing parameters were averaged over

four days for each person. Intensity of colour represents the strength of the association (beta-coefficients), whereas red indicates a positive and blue

a negative association.

Figure 2. Glucose and insulin values in OGTT in individuals with early vs late start of eating. Glucose (A), insulin (B), and C-peptide (C) values in OGTT are presented in individuals with early (blue) and late (red) start of eating. Data are shown as mean ± SEM. * P<0.05 in the Student's t test. ES, eating start.

717 **Table 1. Clinical characteristics of study participants**

Parameters	All
Male / female	106 / 191
Age [years]	59.5 (8.9)
Anthropometric parameters	
BMI [kg/m²]	31.3 (6.1)
Waist circumference [cm]	101.8 (14.7)
Total body fat [%]	34.9 (8.1)
Visceral fat [% of total fat]	24.4 (9.1)
Liver fat [%]	9.12 (8.00)
Blood pressure	
Systolic blood pressure [mmHg]	136.2 (15.1)
Diastolic blood pressure [mmHg]	88.2 (10.3)
Lipid parameters	
Triglycerides [mg/dl]	139.3 (76.0)
Total cholesterol [mg/dl]	211.9 (36.7)
HDL cholesterol [mg/dl]	54.1 (15.2)
LDL cholesterol [mg/dl]	137.6 (32.6)
Glycaemic parameters	
Fasting glucose [mg/dl]	106.7 (9.5)
AUC glucose [mg/dl*min]	19839 (3212)
Fasting insulin [mU/l]	14.4 (8.6)
AUC insulin [mU/l*min]	13795 (9242)
HOMA-IR [AU]	3.82 (2.32)
ISI Matsuda [AU]	2.63 (1.65)
Insulinogenic index [AU]	1.23 (1.14)
Disposition index [AU]	2.67 (2.90)
HbA1c [%]	5.7 (0.4)

719 Data are shown as mean (SD).

721 **Table 2. Dietary composition and meal timing pattern of study participants**

Parameters	All
Daily dietary composition	
Daily calorie intake [kcal]	1941 (547)
Carbohydrates [g]	198.9 (65.1)
Carbohydrates [EN%]	42.6 (7.5)
Protein [g]	83.6 (24.7)
Protein [EN%]	18.2 (4.6)
Fat [g]	80.6 (30.5)
Fat [EN%]	36.7 (7.6)
Fibre [g]	24.0 (8.3)
Fibre [EN%]	2.56 (1.02)
Eating timing	
Eating window [hh:mm]	12:09 (01:49)
Start of first eating event [hh:mm]	08:09 (01:24)
End of last eating event [hh:mm]	20:06 (02:13)
Daily calorie distribution	
Caloric midpoint [hh:mm]	15:19 (02:04)
Calories in first meal [%]	27.7 (10.1)
Calories in last meal [%]	33.8 (11.3)
Eating frequency	
Number of eating events	4.07 (0.92)

723 Data are shown as mean (SD).

Table 3. Associations of eating start with insulin levels, insulin sensitivity, and body fat

with adjustments

Parameter[a]	Coefficient B[b]	95% CI	p
Glycaemic parameters			
Fasting insulin [mU/l]			
Non-adjusted	1.06	0.37 to 1.75	**0.003**
+Adjusted for age and sex	1.06	0.38 to 1.74	**0.002**
++Adjusted for daily energy intake	1.08	0.40 to 1.77	**0.002**
+++Adjusted for body fat	0.84	0.18 to 1.51	**0.013**
Insulin 120 min [mU/l]			
Non-adjusted	17.70	6.50 to 28.90	**0.002**
+Adjusted for age and sex	17.11	5.89 to 28.34	**0.003**
++Adjusted for daily energy intake	16.86	5.62 to 28.10	**0.003**
+++Adjusted for body fat	15.52	4.17 to 26.87	**0.008**
AUC insulin [mU/l*min]			
Non-adjusted	1170	419 to 1921	**0.002**
+Adjusted for age and sex	1155	400 to 1910	**0.003**
++Adjusted for daily energy intake	1147	390 to 1904	**0.003**
+++Adjusted for body fat	951	199 to 1704	**0.013**
HOMA-IR [AU]			
Non-adjusted	0.31	0.13 to 0.49	**0.001**
+Adjusted for age and sex	0.31	0.13 to 0.49	**0.001**
++Adjusted for daily energy intake	0.32	0.14 to 0.50	**0.001**
+++Adjusted for body fat	0.26	0.08 to 0.43	**0.005**
ISI Matsuda [AU]			
Non-adjusted	-0.17	-0.31 to -0.04	**0.013**
+Adjusted for age and sex	-0.18	-0.31 to -0.04	**0.011**
++Adjusted for daily energy intake	-0.18	-0.32 to -0.04	**0.029**
+++Adjusted for body fat	-0.12	-0.24 to 0.01	0.079
Anthropometric parameters			
Total body fat [%]			
Non-adjusted	1.08	0.42 to 1.73	**0.001**
+Adjusted for age and sex	0.83	0.30 to 1.36	**0.002**
++Adjusted for daily energy intake	0.81	0.27 to 1.34	**0.003**
Waist circumference [cm]			
Non-adjusted	1.18	-0.01 to 2.38	0.053
+Adjusted for age and sex	1.38	0.23 to 2.54	**0.019**
++Adjusted for daily energy intake	1.40	0.25 to 2.56	**0.018**

[a] Associations between eating start (ES) and glycaemic or anthropometric traits were examined using multivariable linear regression models adjusted for age, sex, and daily energy intake. For glycemic parameters, models were additionally adjusted for a percentage of body fat. Each subsequent line with plus sign (+, ++, +++) indicates that the adjustment is in addition

732   to adjustments in previous line. Only parameters, demonstrated associations in the adjusted
733   models, are shown.
734   [b] An unstandardized regression coefficient B shows parameter alterations per each later hour
735   of ES.
736

737   **Table 4. Characteristics of individuals with early vs late start of eating**

Parameters	Early eating start	Late eating start	P
Male / female	65 / 95	41 / 96	
Age [years]	58.6 (8.6)	60.5 (9.2)	**0.018**
Anthropometric parameters			
BMI [kg/m²]	31.3 (6.3)	31.3 (6.0)	0.994
Waist circumference [cm]	101.6 (14.3)	102.1 (15.2)	0.781
Fat mass [%]	33.6 (8.6)	36.3 (7.3)	**0.004**
Visceral fat [% of total fat]	24.4 (9.4)	24.4 (8.9)	0.743
Liver fat [%]	8.84 (8.03)	9.42 (7.99)	0.466
Glycaemic parameters			
Fasting glucose [mg/dl]	106.2 (9.3)	107.2 (9.6)	0.366
Glucose 120 min [mg/dl]	137.2 (30.9)	141.2 (30.8)	0.270
AUC glucose [mg/dl*min]	19752 (3201)	19942 (3233)	0.612
Fasting insulin [mU/l]	13.8 (8.3)	15.2 (9.0)	0.088
Insulin 120 min [mU/l*min]	120.7 (117.2)	150.6 (157.5)	**0.029**
AUC insulin [mU/l*min]	12877 (7944)	14851 (10469)	0.146
Fasting C-peptide [µg/l] [a]	2.18 (2.60)	2.63 (3.89)	0.282
C-peptide 120 min [µg/l] [a]	9.61 (3.65)	10.84 (4.98)	**0.034**
AUC C-peptide [µg/l*min] [a]	931 (296)	1001 (474)	0.372
HOMA-IR	3.62 (2.20)	4.05 (2.44)	0.070
ISI Matsuda [AU]	2.75 (1.64)	2.49 (1.66)	0.096
Insulinogenic index	1.17 (0.96)	1.31 (1.31)	0.587
Disposition index [AU]	2.64 (2.33)	2.70 (3.45)	0.460
HbA1c [%]	5.7 (0.4)	5.7 (0.3)	0.663
Eating timing			
Eating window [hh:mm]	12:56 (1:34)	11:14 (1:39)	**<0.001**
Start of eating [hh:mm]	7:11 (0:43)	9:15 (1:09)	**<0.001**
End of eating [hh:mm]	20:06 (1:35)	20:06 (2:47)	0.411
Daily calorie distribution			
Daily calorie intake [kcal]	1983 (586)	1891 (496)	0.302
Caloric midpoint [hh:mm]	14:59 (1:58)	15:41 (2:08)	**0.003**
Calories in first meal [%]	26.7 (10.4)	28.7 (9.7)	**0.041**
Calories in last meal [%]	33.1 (11.2)	34.7 (11.4)	0.181
Eating frequency			
Number of eating events	4.21 (0.95)	3.91 (0.87)	**0.006**

738
739   Data are shown as mean (SD). Group comparison was conducted by Student's t test or by
740   Mann-Whitney U test. [a] C-peptide data was available for 239 individuals.
741

Supplementary material

Supplementary Figures

Figure S1. Daily distribution of eating events.

Supplementary Tables

Table S1. Correlation analysis of eating timing parameters and daily calorie distribution

Table S2. Associations of eating start with insulin levels and body fat with adjustments for
macronutrient intake

Table S3. Associations of other eating timing parameters with glycaemic and anthropometric
traits

Table S4. Lipid parameters in individuals with early vs. late start of eating

Table S5. Food composition and calorie intake in individuals with early vs. late start of eating

SUPPLEMENTARY FIGURES

Figure S1. Daily distribution of eating events. Distribution of eating events is shown as

percentage of all eating events in all study participants (n=297). Each number at the time axis

represents an hour starting from 0 (24:00 – 1:00) and ending with 23 (23:00 – 24:00).

763 # SUPPLEMENTARY TABLES

764 **Table S1. Correlation analysis of eating timing parameters and daily calorie distribution**

	Eating window	Start of first eating event	End of last eating event	Caloric midpoint	Calories in first meal	Calories in last meal	Daily calorie intake
Start of first eating event	-0.582**	-	n.s.	n.s.	n.s.	n.s.	n.s.
End of last eating event	0.461**	n.s.	-	n.s.	n.s.	n.s.	n.s.
Caloric midpoint	0.180**	0.332***	n.s.	-	n.s.	n.s.	n.s.
Calories in first meal	-0.347**	0.178**	-0.146*	-0.211**	-	n.s.	n.s.
Calories in last meal	-0.221**	0.167**	-0.152**	0.454**	0.159**	-	n.s.
Daily calorie intake	0.298**	n.s.	n.s.	0.224**	-0.182**	n.s.	-
Number of meals	0.502**	-0.182**	0.243**	n.s.	-0.494**	-0.500**	0.232**

765
766 Correlation analysis was conducted by the Pearson test. Only significant correlation
767 coefficients are shown. * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$.
768

769 **Table S2. Associations of eating start with insulin levels and body fat with adjustments**

770 **for macronutrient intake**

Parameter[a]	Coefficient B[b]	95% CI	P
Glycaemic parameters			
Fasting insulin [mU/l]			
+Adjusted for carb intake	0.97	0.21 to 2.53	**0.007**
+Adjusted for fat intake	0.99	0.29 to 1.70	**0.006**
+Adjusted for protein intake	0.96	0.26 to 1.66	**0.008**
Insulin 120 min [mU/l]			
+Adjusted for carb intake	15.68	4.29 to 27.07	**0.007**
+Adjusted for fat intake	16.23	4.84 to 27.62	**0.005**
+Adjusted for protein intake	15.30	3.91 to 26.69	**0.009**
AUC insulin [mU/l*min]			
+Adjusted for carb intake	1061	298 to 1825	**0.007**
+Adjusted for fat intake	1120	355 to 1884	**0.004**
+Adjusted for protein intake	1057	292 to 1822	**0.007**
HOMA-IR [AU]			
+Adjusted for carb intake	0.28	0.09 to 0.47	**0.004**
+Adjusted for fat intake	0.29	0.10 to 0.48	**0.003**
+Adjusted for protein intake	0.28	0.09 to 0.47	**0.004**
ISI Matsuda [AU]			
+Adjusted for carb intake	-0.18	-0.31 to -0.04	**0.012**
+Adjusted for fat intake	-0.19	-0.32 to -0.05	**0.008**
+Adjusted for protein intake	-0.17	-0.31 to -0.04	**0.012**
Anthropometric parameters			
Total body fat [%]			
+Adjusted for carb intake	0.84	0.30 to 1.37	**0.002**
+Adjusted for fat intake	0.86	0.33 to 1.39	**0.002**
+Adjusted for protein intake	0.81	0.28 to 1.35	**0.003**
Waist circumference [cm]			
+Adjusted for carb intake	1.37	0.21 to 2.53	**0.021**
+Adjusted for fat intake	1.47	0.32 to 2.62	**0.013**
+Adjusted for protein intake	1.38	0.24 to 2.52	**0.017**

771
[a] Associations between eating start (ES) and glycaemic or anthropometric traits were
examined using multivariable linear regression models adjusted for age, sex, and daily energy
intake. Only parameters, demonstrated associations in the adjusted models, are shown. Each
line with plus sign (+) indicates the additional adjustment of this model to daily carb, fat, or
protein intake.
[b] An unstandardized regression coefficient B shows parameter alterations per each later hour
of ES.
779
780

781 **Table S3. Associations of other eating timing parameters with glycaemic and**

782 **anthropometric traits**

Parameter[a]	Coefficient B[b]	95% CI	P
Eating window			
Total body fat [%]			
Non-adjusted	-0.82	-1.33 to -0.31	**0.002**
+Adjusted for age and sex	-0.60	-1.01 to -0.18	**0.005**
++Adjusted for daily energy intake	-0.53	-0.97 to -0.10	**0.016**
Caloric midpoint			
Fasting glucose [mg/dl]			
Non-adjusted	0.65	0.14 to 1.16	**0.013**
+Adjusted for age and sex	0.70	0.19 to 1.22	**0.008**
++Adjusted for daily energy intake	0.62	0.10 to 1.15	**0.020**
+++Adjusted for body fat	0.60	0.07 to 1.12	**0.026**
Glucose 120 min [mg/dl]			
Non-adjusted	-1.88	-3.57 to -0.19	**0.030**
+Adjusted for age and sex	-1.63	-3.33 to -0.08	0.061
++Adjusted for daily energy intake	-1.48	-3.21 to 0.25	0.094
+++Adjusted for body fat	-1.57	-3.30 to 0.17	0.076
HOMA-IR [AU]			
Non-adjusted	0.14	0.02 to 0.27	**0.026**
+Adjusted for age and sex	0.13	-0.001 to 0.25	0.053
++Adjusted for daily energy intake	0.11	-0.02 to 0.24	0.094
+++Adjusted for body fat	0.09	-0.03 to 0.22	0.135
Waist circumference [cm]			
Non-adjusted	0.91	0.10 to 1.72	**0.028**
+Adjusted for age and sex	0.66	-0.13 to 1.45	0.101
++Adjusted for daily energy intake	0.65	-0.15 to 1.46	0.111
Calories in the first meal			
Waist circumference [cm]			
Non-adjusted	0.19	0.02 to 0.36	**0.024**
+Adjusted for age and sex	0.14	-0.02 to 0.30	0.088
++Adjusted for daily energy intake	0.16	-0.01 to 0.33	0.062
Calories in the last meal			
Waist circumference [cm]			
Non-adjusted	0.22	0.08 to 0.37	**0.003**
+Adjusted for age and sex	0.15	0.001 to 0.30	**0.048**
++Adjusted for daily energy intake	0.16	0.01 to 0.31	**0.041**
Visceral fat [% of total fat]			
Non-adjusted	0.001	0.0002 to 0.002	**0.020**
+Adjusted for age and sex	0.0002	-0.0004 to 0.0008	0.458
++Adjusted for daily energy intake	0.0003	-0.0003 to 0.001	0.309

[a] Associations between eating timing parameters and glycaemic or anthropometric traits were
examined using multivariable linear regression models adjusted for age, sex, and daily energy
intake. For glycemic parameters, models were additionally adjusted for a percentage of body
fat. Each subsequent line with plus sign (+, ++, +++) indicates that the adjustment is in addition
to adjustments in previous line. Only parameters, demonstrated associations in the non-
adjusted models, are shown.
[b] An unstandardized regression coefficient B shows parameter alterations per each later hour
of ES.

Table S4. Lipid parameters in individuals with early vs. late start of eating

Parameters	Early eating start	Late eating start	P
Triglycerides [mg/dl]	137.8 (70.9)	140.9 (81.8)	0.812
Total cholesterol [mg/dl]	213.0 (38.8)	210.5 (34.1)	0.553
HDL cholesterol [mg/dl]	53.5 (15.0)	54.8 (15.6)	0.467
LDL cholesterol [mg/dl]	138.8 (34.9)	136.2 (29.9)	0.511

793 Data are shown as mean (SD). N=297.

795 **Table S5. Food composition and calorie intake in individuals with early vs. late**
796 **start of eating**

Parameters	Early eating start	Late eating start	P*
Daily dietary composition			
Calorie intake [kcal]	1983 (586)	1891 (496)	0.302
Carbohydrates [g]	205.3 (66.3)	191.4 (63.0)	0.062
Carbohydrates [EN%]	43.0 (7.5)	42.2 (7.6)	0.375
Protein [g]	85.7 (28.0)	81.2 (20.0)	0.353
Protein [EN%]	18.2 (5.0)	18.1 (4.0)	0.954
Fat [g]	82.0 (32.4)	78.9 (28.2)	0.748
Fat [EN%]	36.2 (7.3)	37.2 (7.8)	0.271
Fibre [g]	25.0 (8.4)	22.7 (8.1)	**0.024**
Fibre [EN%]	2.60 (0.87)	2.52 (1.17)	0.300
Dietary composition of the first meal			
Calorie intake [kcal]	488 (202)	509 (218)	0.524
Calorie intake [% of daily intake]	26.7 (10.4)	28.7 (9.7)	**0.041**
Carbohydrates [g]	57.5 (26.1)	56.9 (27.0)	0.902
Carbohydrates [EN%]	48.8 (12.0)	46.9 (11.6)	0.171
Protein [g]	20.2 (11.8)	20.8 (10.0)	0.416
Protein [EN%]	16.4 (5.6)	16.6 (4.9)	0.981
Fat [g]	18.8 (11.4)	20.7 (12.4)	0.210
Fat [EN%]	31.8 (11.2)	33.6 (12.3)	0.176
Fibres [g]	7.29 (3.88)	7.07 (3.95)	0.527
Fibres [EN%]	3.00 (1.26)	2.82 (1.25)	0.155

798 Data are shown as mean (SD).

800