Late start of eating is linked to lower insulin sensitivity and hyperinsulinemia in prediabetes

Beeke Peters[1,2], Noelle Fröhlich[1,3], Jürgen Machann[2,4,5], Ulrike Dambeck[6], Caroline Honsek[6], Anna Sachno[2,7], Margrit Kemper[2,6], Stefan Kabisch[2,6,7], Andreas Fritsche[2,4,8], Knut Mai[2,7,9,10,11], Andreas F.H. Pfeiffer[2,6,7,10], Olga Pivovarova-Ramich[1,2,6,7] *

[1]*Department of Molecular Metabolism and Precision Nutrition, German Institute of Human Nutrition Potsdam-Rehbruecke, Nuthetal, Germany;*

[2]*German Center for Diabetes Research (DZD), München-Neuherberg, Germany;*

[3]*University of Potsdam, Potsdam, Germany;*

[4]*Institute for Diabetes Research and Metabolic Diseases, Helmholtz Center Munich, University of Tübingen, Tübingen, Germany*

[5]*Department of Radiology, Section on Experimental Radiology, Eberhard-Karls University of Tübingen, Tübingen, Germany*

[6]*Department of Clinical Nutrition, German Institute of Human Nutrition Potsdam-Rehbruecke, Nuthetal, Germany;*

[7]*Department of Endocrinology and Metabolism, European Reference Network on Rare Endocrine Diseases (ENDO-ERN), Charité – Universitätsmedizin Berlin, Corporate Member of Freie Universität Berlin and Humboldt-Universität zu Berlin, 10117 Berlin, Berlin, Germany.*

[8]*Internal Medicine IV, Department of Diabetology, Endocrinology and Nephrology, University of Tübingen, Tübingen, Germany.*

[9]*Department of Human Nutrition, German Institute of Human Nutrition Potsdam-Rehbrücke, Nuthetal, Germany*

[10]*NutriAct-Competence Cluster Nutrition Research Berlin-Potsdam 10117, Germany.*

[11]*Berlin Center for Rare Diseases, Charité-Universitätsmedizin Berlin, Corporate Member of Freie Universität Berlin and Humboldt-Universität zu Berlin, 10117 Berlin, Germany.*

Corresponding author: Olga Pivovarova-Ramich, German Institute of Human Nutrition

Potsdam-Rehbruecke, Arthur-Scheunert-Allee 114-116, 14558 Nuthetal, Germany, Tel.: +49

33200882749, Email: olga.ramich@dife.de

Short title: Eating timing and glucose metabolism

Main text word count: 4571; references: 69; tables: 3; figures: 2; supplemental tables: 6; supplemental figures: 2.

Abstract

Background: Prediabetes is associated with an increased risk of progression to type 2 diabetes. While dietary interventions in prediabetes traditionally primary focus on energy and macronutrient intake, the role of eating timing has recently been highlighted. This study aimed to examine the relationships between components of eating timing patterns and glycaemic parameters in prediabetes.

Methods: In 297 individuals with prediabetes, i.e. impaired fasting glucose or impaired glucose tolerance (age 59.4 ± 9.0 y, BMI 31.3 ± 6.2 kg/m2), glycaemic traits were assessed using the oral glucose tolerance test (OGTT). Parameters of eating timing pattern (eating timing itself, daily calorie distribution, and meal number) were extracted from four-day food records. Eating start (ES) was defined as the start of the first caloric event of the day.

Results: Among eating timing components, the most associations with glycaemic parameters were observed for ES. Individuals with late ES showed higher fasting insulin ($p = 0.006$), AUC insulin ($p = 0.020$), insulinogenic index ($p = 0.049$), HOMA-IR ($p = 0.004$), and lower Matsuda insulin sensitivity index ($p = 0.004$) compared to those with early ES, even after adjustment for age, sex, daily energy intake, and body fat. Participants with late ES had decreased insulin clearance ($p<0.001$) upon comparable glucose and C-peptide levels.

Conclusion: Late ES is associated with estimates of lower insulin sensitivity and hyperinsulinemia in prediabetes, independent of energy intake and body fat. Our findings suggest that an earlier ES is related to improved glucose metabolism which might lower the risk of progression to diabetes.

Funding: This work was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation, RA 3340/4-1 to OP-R, project number 530918029, and TRR 418, project number 541063275 to OP-R). The DZD is funded by the German Federal Ministry for Education and Research (BMBF, funding code 01GI0925). KM was supported by the

Competence Cluster Nutrition Research Berlin-Potsdam, funded by the Federal Ministry of

Education and Research (BMBF, funding code 01EA1408) and by the Deutsche

Forschungsgemeinschaft (DFG, German Research Foundation) – TRR 412/1 – project

number 535081457.

Abstract word count: 251

Keywords: glucose metabolism, insulin sensitivity, hyperinsulinemia, diabetes, eating timing,

circadian clocks

Abbreviations

AUC   Area under the curve

BMI   Body mass index

CM   Caloric midpoint

ELISA   Enzyme linked immunosorbent assay

FM   Start of the first meal

HbA1c   Haemoglobin A1c

HDL   High-density lipoprotein

HIC   Hepatic insulin clearance

HOMA-IR   Homeostatic Model Assessment for Insulin Resistance

IFG   Impaired fasting glucose

IGT   Impaired glucose tolerance

ISI   Insulin sensitivity index

LDL   Low-density lipoprotein

MRI   Magnetic resonance imaging

MRS   Magnetic resonance spectroscopy

OGTT   Oral glucose tolerance test

SD   Standard deviation

T2D   Type 2 diabetes

1. Introduction

The pandemic of diabetes mellitus affects millions of people worldwide. Type 2 diabetes (T2D), the most common form, is preceded by prediabetes, characterized by increasing blood glucose levels which does not yet achieve the diagnostic criteria for diabetes (1). Dietary interventions, aiming to impede the progression of prediabetes to diabetes, have traditionally focused on energy and macronutrient intake. Recent research has highlighted the importance of eating timing in managing blood glucose levels and potentially in the prevention of T2D (2; 3). This fact is based on the diurnal variation of metabolic functions resulting in different metabolic responses to the same food consumed at different times of the day (4; 5).

Eating timing has emerged as a significant factor in glucose metabolism and insulin sensitivity (2; 6). Cross-sectional studies have shown that early eating patterns aligning with individual circadian rhythms are associated with lower body weight and metabolic health (3; 6-8). In interventional studies, eating earlier in the biological day resulted in greater reductions in body weight, BMI, and waist circumference, and improved glycaemic control. For instance, early time-restricted eating (TRE), limiting food intake to a 6-10 hour window early in the day, improved glucose tolerance and insulin sensitivity in several studies (9-12), although other research did not confirm this finding (13; 14) and numerous cofounders (longer fasting, fewer meals, lower calorie intake, and possibly different meal composition) has to be considered. In contrast, late and night eating showed an association with increased risk of obesity, cardiovascular diseases, and related traits such as higher BMI and body fat (6; 15-19).

Timing of specific eating events, particularly breakfast, appears to play a crucial role in glycaemic control. Most previous studies suggest that skipping breakfast has been linked to a higher risk of developing T2D as partly mediated by BMI (20), while the earlier breakfast timing was related to the lower T2D incidence (21). Nevertheless, metabolic risks of dinner skipping were also described (22). This suggests that not only timing of meals but also daily pattern of calorie distribution may support managing blood glucose levels in diabetes (23; 24). Additionally, the number of eating events and the duration of overnight fasting have been investigated for their potential impact on glucose regulation and diabetes risk (3; 25; 26).

Nevertheless, it remains insufficiently studied how eating timing and its specific parameters,

such as timing, frequency, and diurnal calorie distribution, contribute to the parameters of

glucose metabolism in prediabetes and generally to diabetes risk. Therefore, this study aimed

to examine relationships between components of the eating timing pattern and parameters of

glucose metabolism in individuals with prediabetes.

2. Methods

2.1. Study participants and study protocol

This cross-sectional analysis included baseline data of 297 individuals with prediabetes

obtained in two clinical trials, the Prediabetes Lifestyle Intervention Study (PLIS,

NCT01947595, n=174) and the Diabetes Nutrition Algorithm - Prediabetes (DiNA-P,

NCT02609243, n=123). Complete details of both trials have been described elsewhere (27-

29). Both trials were conducted in accordance with the principles of the Declaration of Helsinki

and Good Clinical Practice guidelines. The study protocol was approved by the ethics

committee of the Charité-Universitätsmedizin Berlin. All patients provided written informed

consent before enrollment.

Data included in this study were collected at the German Institute of Human Nutrition

Potsdam-Rehbruecke and at the Charité-Universitätsmedizin Berlin between 2012 and 2016.

Inclusion criteria in both trials were age between 18 and 75 years, BMI < 45 kg/m², and

diagnosis of impaired fasting glucose (IFG; fasting blood glucose 100-126 mg/dl) and/or

impaired glucose tolerance (IGT; blood glucose at 120 min of oral glucose tolerance test

[OGTT] 140-200 mg/dl) according to the criteria of the American Diabetes Association (1).

Exclusion criteria included overt diabetes mellitus or antidiabetic medication, BMI > 45 kg/m²,

current pregnancy or breastfeeding, serious disease (e.g. symptomatic coronary heart

disease, serious symptomatic malignant disease, severe liver or kidney disease, systemic

infection, severe mental illness), drug abuse, treatment with systemic corticosteroids.

Claustrophobia, metal implants and excessive body circumference were deemed as

contraindication of magnetic-resonance imaging, only. Participants underwent a detailed

metabolic phenotyping including measurement of anthropometric parameters, body

composition, liver fat content, and an OGTT.

Study subjects completed handwritten food records, noting the start and end of each

eating event, the amount and kind of food consumed, for four days including three working

days and one free day to reflect their dietary habits. They did not receive any dietary

recommendations or restrictions before completing the food records.

2.2.   Anthropometric measurements

For the assessment of BMI, body weight was measured using a digital scale and body height

was measured using a stadiometer. Waist and hip circumferences were measured using a

metric tape. Fat mass was analysed using bioelectrical impedance analyser Nutriguard MS

(Data Input GmbH, Pöcking, Germany). Total and visceral adipose tissue (VAT) compartments

of the trunk were assessed by magnetic resonance imaging (MRI). Liver fat content was

determined by localized proton magnetic resonance spectroscopy (^{1}H-MRS) applying a single-

voxel stimulated echo acquisition mode sequence in the posterior hepatic segment 7 as

described (27).

2.3.   OGTT, blood sample analyses, and assessment of glycaemic indices

A standardized 2-hour OGTT was conducted after an overnight fast (>10 h since last food

intake) starting between 8:00 and 9:00. After a fasting blood sampling, participants ingested

standardised glucose solution (75 g glucose in 300 ml water; Accu-Chek Dextro O.G.T.,

Roche), and further blood samples were drawn from the forearm vein 30, 60, 90, and 120 min

after glucose intake. Routine serum parameters were determined using an automated analyzer

(ABX Pentra 4000; ABX, Montpellier, France). Glucose concentrations were analysed locally

by certified laboratories using the glucose oxidase method. Serum insulin and C-peptide

concentrations were analysed by ELISA (Mercodia, Uppsala, Sweden).

Glucose and insulin values in OGTT were used to calculate indices of insulin sensitivity

(homeostasis model assessment of insulin resistance [HOMA-IR] and Matsuda index of insulin

sensitivity [ISI Matsuda]), insulin secretion (insulinogenic and disposition indices), and hepatic

insulin clearance (HIC). HOMA-IR was calculated as ($Gluc_0$ (mmol/l) x Ins_0 (µU/ml)) / 22.5. The

insulinogenic index was calculated as Ins_{30-0} (µU/ml) /$Gluc_{30-0}$ (mg/dl) (30). ISI Matsuda and

the disposition index, which reflects the relation between β-cell secretion capacity and insulin

sensitivity, were assessed as described (31; 32). The hepatic insulin clearance (HIC) was

determined as the ratio of the incremental areas under the OGTT curve: $iAUC_{C\text{-peptide }0-120min}$

(pmol/l*min) / $iAUC_{insulin\ 0-120min}$ (pmol/l*min) (33).

2.4.   Analysis of eating timing pattern

The handwritten food records were digitalized using the PRODI software version 4.5

(NutriScience GmbH, Hofstetten-Flüh, Switzerland) by an experienced nutritionist. The digital

versions were used to assess time, energy intake, and macronutrient composition

(carbohydrates, protein, fat, and fiber) of each eating event for each of the four consecutive

days for all participants. Incomplete dietary records were excluded from the analysis. The

eating timing pattern was characterised by the assessment of (i) eating timing itself (i.e., timing

of the first and last eating event, eating window duration), (ii) calorie distribution across the day

(i.e., time of caloric midpoint and calories in the first and last meal), and (iii) eating frequency

(i.e., number of eating events per day) as described in details previously (6). In brief, an eating

event was defined by the following rules: (1) at least 50 kcal, and (2) a time gap to another

food or meal of at least 15 minutes (34; 35). Start and end of eating were defined as the start

of the first caloric event of the day and the end of the last caloric event of the day, respectively.

Eating window duration was calculated as the time difference from the start of the first eating

event to the end of the last eating event, and the caloric midpoint (CM) was assessed as the

time when the 50% of total calorie intake was reached (34).

To assess the timing and energy intake in the first and last meals, eating events were

classified as a meal or as a snack, based on the time of the day, nutrient density, and regularity

as described (6). In brief, an eating event was defined as a "meal" if it had > 15.0 % of daily

energy intake (36), was consumed at the main meal times between 08:00-10:00, 12:00 –

14:00, and 18:00 – 20:00 (typical breakfast, lunch, and dinner times in Germany) (37), and

was regular.

2.5. Statistical Analysis

Statistical analyses were carried out using SPSS 25.0 (IBM SPSS, Chicago, IL). Data

were expressed as mean (SD). Group comparison was conducted by Student's t test for

continuous variables with normal distribution or by Mann-Whitney U test if the normality

assumption was violated. Associations between eating timing parameters itself as well as

between eating timing and glycaemic and anthropometric parameters were examined using

linear regression models and adjusted for potential confounders. Model 1 included no covariate

adjustment. Model 2 was adjusted for sex and age, and model 3 for sex, age, and average

daily energy intake. Comparison between groups with early and late ES was conducted using

a one-way ANOVA with adjustment for sex, age, average daily energy intake, and total body

fat. Statistical significance was defined as $p < 0.05$.

3. Results

3.1. Clinical characteristics of participants

Eating pattern was assessed in 297 individuals (age: 59.0 (9.0) years, BMI: 31.3 (6.2) kg/m²,

106 males and 191 females) who completed food records and provided full metabolic data.

Body fat content was 34.9 (8.1) %. Visceral fat accounted for 24.4 (9.2) % of total fat, and liver

fat content was 9.12 (8.00) %. Average fasting glucose was 106.7 (9.5) mg/dL and HbA1c of

5.67 (0.36) %. Anthropometric measures and parameters of glucose metabolism are shown in

Table 1.

3.2. Eating timing pattern of participants

Daily distribution of eating events is shown in **Figure S1.** Eating timing pattern of study

participants, including the eating timing itself, calorie distribution across the day, and eating

frequency, is shown in **Table 2**. Participants consumed 4.07 (0.92) meals per day, with an

eating start (ES) at 08:08 (01:24) and eating end at 20:06 (02:13), resulting in an eating window

duration of 12:09 (01:49) hours (**Table 2**). Energy intake was 27.7 (10.1) % of daily calories in

the first and 33.8 (11.3) % in the last meal, while the CM was estimated at 15:19 (02:04) (**Table**

2).

Eating timing parameters were tightly interrelated. In particular, ES was linked to the

daily calorie distribution, showing positive associations with CM (β = 0.332, p < 0.001) and

percentage of energy in the first (β = 0.178, p = 0.002) and last meal (β = 0.167, p = 0.004).

ES was negatively associated with eating duration (β = -0.582, p < 0.001) and number of meals

per day (β = -0.182, p = 0.002). This indicates that individuals who started eating later showed

shorter eating window and consumed fewer meals per day, but the individual meals, at least

the first and last ones, tended to be larger (**Table S1**). Further, the caloric midpoint (CM) was

positively associated with calorie intake during the last meal (β = 0.454, p < 0.001), with total

daily calorie intake (β = 0.224, p < 0.001) as well as with duration of the daily eating window

(β = 0.180, p = 0.002), and negatively associated with calorie intake during the first meal (β =

-0.211, p < 0.001). This suggests that individuals with a late CM tended to have a longer eating

window and to consume more evening and daily calories (**Table S1**).

3.3. *Association of eating timing and glucose metabolism parameters*

The association analysis revealed several interrelations between eating timing

characteristics and parameters of glucose metabolism, with most associations found for ES

and CM. ES showed positive associations with fasting insulin (β = 0.221, p = 1.7 x 10^{-4}), AUC

insulin (β = 0.208, p = 4.1 x 10^{-4}), HOMA-IR (β = 0.236, p = 6.2 x 10^{-5}), and insulinogenic index

(β = 0.186, p = 0.002), and a negative association with the insulin sensitivity index ISI Matsuda

(β = -0.234, p = 6.9 x 10^{-5}) as calculated in the non-adjusted linear regression model (model

1) (**Figure 1a**). To account for potential confounders, the model was adjusted for sex and age

(model 2), and additionally for daily energy intake (model 3). After adjustment for

abovementioned confounders, the associations remained significant (fasting insulin: β = 0.221,

p = 1.4 x 10^{-4}, AUC insulin: β = 0.204, p = 4.8 x 10^{-4}, HOMA-IR: β = 0.238, p = 4.4 x 10^{-5},

insulinogenic index: β = 0.187, p = 0.001; ISI Matsuda: β = -0.235, p = 6.1 x 10^{-5}) **(Figure 1b,**

Figure S2, Table S2). ES was also positively associated with fat mass and waist

circumference – even after adjustment for sex, age, and energy intake (β = 0.143, p = 0.002

and β = 0.126, p = 0.025, respectively), but not with BMI **(Figure 1b, Table S2)**.

CM showed fewer interrelations with glycaemic parameters, which were, however,

similar to those of ES. For the CM, positive associations with fasting glucose, fasting insulin,

and HOMA-IR were found, which remained significant after adjustment to potential

confounders (fasting glucose: β = 0.155, p = 0.009, fasting insulin: β = 0.125, p = 0.038, HOMA-

IR: β = 0.143, p = 0.017 in model 3) (**Figure 1, Table S3**). CM also showed an association with

waist circumference, which disappeared in adjusted models (**Table S3**).

Positive associations of energy intake during the first meal with waist circumference

and of energy intake during the last meal with visceral fat and waist circumference were

observed in non-adjusted models, but after adjustment for confounders, only the last relation

remained significant (β = 0.172, p = 0.022) (**Figure 1b, Table S4**). No associations with

glycaemic or anthropometric traits were found for the end of the last eating event and daily

number of meals (**Figure 2**).

3.4.   Comparison of the glycaemic parameters in individuals with early vs late

start of eating

Based on the observed associations between eating time and glycaemic parameters,

we categorized all participants into groups with early ES (before 08:09, which was the average

ES in the study cohort) and with late ES (after 08:09). The groups did not differ in sex, BMI,

waist circumference, visceral and liver fat, and blood lipids, but individuals with late ES were

on average two years older (p=0.018) and showed a higher percentage of body fat (p=0.004)

(**Table 3**).

Participants with late ES ended to eat at a similar time as subjects with early ES,

resulting in an average 1h 42 min shorter eating window (p<0.001) and a later CM (p=0.003)

(**Table 3**). Macronutrient composition and calorie intake per day did not differ between groups,

whereas percentage of calories within the first meal was slightly larger in individuals with late

ES (p=0.041) (**Table S5**). In the OGTT, glucose and C-peptide values did not differ between

groups at any sampling point (**Figure 2a, c**). Insulin values were greater at 0, 30, 90, and 120

min (p<0.05) in individuals with late ES compared to the early ES group (**Figure 2b**). In a line

with the association analysis, participants with late ES showed higher insulin levels (AUC

insulin: p = 0.020, insulinogenic index: p = 0.049) and lower insulin sensitivity (HOMA-IR: p =

0.004; ISI Matsuda: p = 0.004) compared to individuals with early ES (**Table 3**). These

differences remained significant after adjustment for sex, age, energy intake, and additionally

for body fat (**Table S6**). To explain the different insulin values upon comparable C-peptide

levels, we assessed a hepatic insulin clearance (HIC) and found that it was markedly lower in

participants with late ES (p<0.001).

4. Discussion

As the prevalence of diabetes and prediabetes continues to rise globally, understanding the

relationship between eating timing and glucose metabolism becomes increasingly important,

potentially offering an additional diet-based non-pharmacological approach to prevent or delay

the development of T2D. In this study, we demonstrated robust associations of eating timing

parameters with hyperinsulinemia and insulin resistance, which represent key players in the

progression of prediabetes towards T2D (38) and typically develop 10-15 years before

diabetes diagnosis (39). In our study, individuals with prediabetes underwent comprehensive

metabolic phenotyping, including OGTT, and documentation of their nutritional habits using

food diaries over several days. Given this comprehensive approach, the highlight of this study

is the detailed characterization of individual eating timing patterns through assessment of

eating timing itself and the diurnal distribution of energy intake.

Our analysis revealed a number of associations between eating timing characteristics

and parameters of glucose metabolism, with most associations found for ES. We

demonstrated that a late ES is associated with higher fasting and postprandial insulin values

and lower insulin sensitivity, as assessed in OGTT, whereas glucose values were not related

to ES. Remarkably, these associations were very robust, remaining significant after adjustment

for age, gender, and energy intake. Thus, our data suggest that individuals with a late ES have

more pronounced insulin resistance, which apparently results in higher compensatory

hyperinsulinemia to maintain glucose control due to still sufficient beta-cell capacities.

Our finding aligns with previously published evidence that timing, calorie intake and

composition of breakfast play an important role in glucose homeostasis and diabetes

management (20; 23; 24; 40). Concerning breakfast timing, large cohort meta-analyses

demonstrate an association between breakfast skipping and increased T2D risk (20; 40). T2D

risk increased with every additional day of breakfast skipping, but the curve reached a plateau

at 4-5 day per week, showing an increased risk of 55%, whereas this effect was only partly

mediated by the weight gain (20). Cross-sectional studies revealed an association of breakfast

skipping with a later chronotype, both of which contribute to poorer glycaemic control, as

indicated by higher HbA1c levels (41). Several observational studies suggest that early ES at

the day confers significant metabolic advantages including lower BMI, blood lipids, and better

glycaemic control (3; 42). For example, data from the National Health and Nutrition

Examination Survey (NHANES) demonstrated that every hour later that eating commenced

was associated with approximately 0.6% higher glucose level and 3% higher HOMA-IR after

adjustment for BMI and other confounders (3). Notably, although in our study we did not

observe difference in blood glucose, we found marked insulin changes providing clear

evidence that eating timing affects metabolism.

ES timing is tightly associated with calorie distribution over the day. Several studies

demonstrated beneficial effects of implementing a high-calorie breakfast on glycaemic control

in individuals with T2D (23; 24). In contrast, breakfast skipping is frequently accompanied by

late-night dinner consumption, leading to calorie distribution towards the later part of the day.

This misaligns with endogenous metabolic rhythms and can contribute to metabolic

dysfunctions (43), as discussed in detail below. Interestingly, we made a similar observation in

our study, showing a later CM in the late ES group, although the timing of the last eating event

remained unchanged. Growing observational and experimental evidence, summarized in our

and other reviews (5; 44), shows that consuming the majority of calories early in the day

supports lower body weight and better glucose homeostasis compared to late or nighttime

calorie intake, although detailed timing recommendations are still lacking. Notably, in our study,

the calorie intake during the first meal (as well as total daily calorie intake) did not differ in

individuals with early and late ES. Taken together, these data suggest that the timing of ES

per se, and not the calorie intake or food composition, is related to glycaemic parameters of

study participants.

Interestingly, we also found that individuals with a late ES showed a shorter daily eating

duration as their end of eating window did not differ from participants with an early ES.

However, they showed poorer glycaemic parameters despite the shorter eating window. This

finding contrasts with data from human TRE studies, reducing daily eating duration to less than

10 hours, which demonstrated beneficial effects on glucose homeostasis (45; 46). Thus, our

data suggest that the timing of eating might have a stronger impact on glycaemic parameters

compared to eating duration when daily calorie intake is similar in both groups. Notably, eating

duration in both early and late ES groups was over eleven hours per day, which may also

explain the absence of effects related to eating duration. Finally, similarly to our data, several

cross-sectional or prospective human cohorts have found no link between eating window

duration and body weight or metabolic traits (3; 6; 47), therefore the role of the eating duration

in free-living populations remains debatable.

The next possible cofounding factor, which might affect the observed ES effects, is the

macronutrient composition of breakfast meals, which exerts profound short-term and long-term

effects on glycaemic and insulinemic responses. Studies reducing breakfast carbohydrates

and replacing them with proteins or fats demonstrated lower glucose, decreased glycaemic

variability, improved β-cell responsiveness, and HbA1c in subjects with T2D (24; 48; 49).

Based on this data, we examined in our study whether food composition might contribute to

the observed associations with insulin secretion or insulin sensitivity, and we found no

differences in the macronutrient composition of the first meal or the daily average when

comparing individuals with early and late ES.

Previous research has largely highlighted the CM as an eating time-related parameter,

demonstrating robust associations with obesity-related traits such as BMI, waist

circumference, and body fat (7; 8; 15). In addition, we recently provided evidence linking CM

and insulin sensitivity in healthy subjects (6), underscoring the essential role of daily calorie

distribution for glucose metabolism. In contrast to published data, our present study observed

a much stronger relationship of ES with glycaemic regulation, specifically with insulin levels

and sensitivity, than CM. This difference likely stems from the different study cohorts, as the

CM associations were primarily observed in cohorts without focus on diabetes (7; 8; 15).

The mechanistic basis for these timing effects likely involves the circadian regulation of

glucose metabolism and insulin sensitivity by endogenous clocks (50). In healthy subjects,

insulin sensitivity in skeletal muscle and adipose tissue is generally higher in the early morning,

facilitating rapid and efficient postprandial glucose uptake (51; 52). Similarly, pancreatic β-cells

show enhanced insulin release in the morning, which effectively suppresses hepatic

gluconeogenesis after breakfast.(53) As the day progresses, insulin action gradually declines

due to both reduced peripheral glucose uptake and diminished β-cell responses, resulting in

larger glycaemic excursions in response to the same glycaemic load.(53) Aligning food intake

with the body's natural metabolic rhythms – namely, early in the day – putatively leads to

optimal metabolic regulation. Eating late opposes these natural rhythms and has the potential

to cause circadian misalignment and metabolic dysfunction (43; 54). Remarkably, circadian

rhythms are often altered in individuals with T2D, showing a flattened profile of insulin

sensitivity and β-cell function, and elevated gluconeogenesis in the morning (55-57). Whether

individuals with prediabetes also have altered metabolic rhythmicity remains insufficiently

investigated (58), but if this is the case, it may help explain why our association data differ from

those observed in metabolically healthy subjects. Nevertheless, our findings in early and late

ES groups support the idea that individuals with prediabetes may also benefit from early eating,

possibly acting as a zeitgeber and, thereby, improving their metabolic rhythmicity (5; 43).

Our data on hormone secretion in OGTT provide another important insight into the

mechanisms underlying the observed ES associations, which was not described previously.

Increased insulin values were accompanied by unchanged C-peptide levels in participants with

a late ES, suggesting that insulin clearance, rather than insulin secretion, was affected, as

confirmed by the calculation of the HIC index. Insulin clearance refers to the uptake and

degradation of insulin in insulin-sensitive tissues, which contributes to the regulation of

circulating insulin after it is secreted from pancreatic β-cells (59). The liver is the primary site

for insulin clearance, accounting for the degradation up to 80% of endogenously secreted

insulin, so HIC strongly correlates with whole-body insulin clearance. HIC is decreased in

individuals with insulin resistance, metabolic dysfunction-associated steatotic liver disease

(MASLD), and metabolic syndrome (60; 61). Decreased HIC contributes to hyperinsulinemia,

which, according to one hypothesis, may be a key factor in T2D development rather than

simply a consequence of the disease(59; 62). Insulin clearance exhibits robust circadian

variation, being lower in the morning than in the evening (63), which may contribute to better

glucose tolerance in the morning in healthy subjects. Taken together, our finding on HIC

suggests, that a late ES might dysregulate the rhythm of insulin degradation, contributing to

more pronounced hyperinsulinemia and, in this way, increasing the risk of diabetes

progression.

The prominent novelty of our study is the analysis of the association of eating timing

with liver fat content. Liver fat accumulation, described as MASLD, often accompanies

prediabetes and T2D and strongly contributes to metabolic dysfunctions (64). Liver fat content

is regulated by circadian clocks, as shown in animal models (65). Further, TRE interventions

have been shown to ameliorate liver fat content in mice fed with high-fat diet (66) and in

humans, with an effect size comparable to calorie restriction (67). Based on known

associations of eating timing with abdominal obesity (7) and insulin sensitivity (6), we expected

to observe an association of eating timing with liver fat in prediabetes. Surprisingly, no eating

timing parameters showed a relation to liver fat content in our cohort. This data expands our

knowledge in this field and requires further investigation in larger cohorts and diverse

categories of individuals with and without metabolic dysfunctions.

Our study has several important strengths. The main strength is the detailed metabolic

and dietary habit phenotyping in a large cohort of individuals with prediabetes, which allowed

us to assess the association of eating timing with glycaemic traits – an assessment never

conducted previously. Using of OGTT allowed the analysis of both fasting and postprandial

metabolism, including hormone secretion and the calculation of specific indices. A further

strength of our study is the analysis of multiple components of the eating pattern and their

potential relation to glycaemic regulation.

Some limitations have to be discussed when interpreting the results of this work. First,

food intake and timing data were self-reported using the food diaries. Although this method is

recognized as a gold standard for nutritional data collection, it does not exclude measurement

errors, such as over- and underreporting. Future studies should include objective monitoring

of eating behaviour using wearables or smartphone apps. Secondly, no data on sleep timing

or quality, which are strongly associated with metabolism and can, for example, worsen

metabolic parameters upon the sleep deficit (68), were collected. Moreover, collecting sleep

time data or questionnaires to assess individual chronotypes would help to decipher how the

relation between eating time and chronotype is linked to the diabetes development. Thirdly,

the analysis did not consider the socioeconomic status of participants. Breakfast skipping

associated with poor socioeconomic status could also explain obesity and the impaired

metabolic state. Further, practicing of intermittent fasting in form of breakfast skipping could

also be associated with poor metabolic state in the sense of reverse causality: as an attempt

to overcome prediabetes instead of being the reason for its development. Finally, the selection

bias by cutting of patients with stronger glycemic impairment (overt diabetes) has to be

mentioned as a limitation.

In conclusion, our study showed that a late ES is associated with lower insulin

sensitivity and hyperinsulinemia in prediabetes, independent of energy intake or other

confounding factors. Our findings suggest that an earlier ES might improve glucose

metabolism and lower the risk of progression from prediabetes to diabetes. This non-

pharmacological approach needs to be investigated in interventional studies.

Contributors

All authors read and approved the final version of the manuscript.

O.P.-R. conceived the study. J.M., U.D., C.H., A.S., and M.K. collected the data. B.P. and N.F. analyzed the data. B.P. and O.P.-R. wrote the manuscript. N.F., B.P., S.K., A.F., K.M., A.F.H.P., and O.P.-R. contributed to interpretation of the data and edited the manuscript. All other authors contributed to data acquisition and approved the final version of the manuscript. O.P.-R. attested that all listed authors meet authorship criteria and no others meeting the criteria have been omitted. O.P.-R. affirmed that the manuscript is an honest, accurate, and transparent account of the study being reported and that no important aspects of the study have been omitted. O.P.-R. is the guarantor of this work and, as such, had full access to all data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Data Sharing Statement

The datasets generated during and/or analyzed during the current study are not publicly available but are available from the corresponding author on reasonable request.

Declaration of Interests

Acknowledgements

We thank all study participants for their cooperation. We gratefully acknowledge the excellent assistance of Christiana Gerbracht and other dietitians, study nurses, and lifestyle advisors involved in this study. We also thank Elwira Gliwska for her contribution to the algorithm development for the analysis of eating timing. This work was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation, RA 3340/4-1 to OP-R, project

486 number 530918029). The DZD is funded by the German Federal Ministry for Education and

487 Research (BMBF, funding code 01GI0925). KM was supported by the Competence Cluster

488 Nutrition Research Berlin-Potsdam, funded by the Federal Ministry of Education and Research

489 (BMBF, funding code 01EA1408) and by the Deutsche Forschungsgemeinschaft (DFG,

490 German Research Foundation) – TRR 412/1 – project number 535081457.

References

1. American Diabetes Association Professional Practice C. 2. Diagnosis and Classification of Diabetes: Standards of Care in Diabetes-2025. Diabetes care 2025;48:S27-S49

2. Diaz-Rizzolo DA, Santos Baez LS, Popp CJ, Borhan R, Sordi-Guth A, Manoogian ENC, Panda S, Cheng B, Laferrere B. Late eating is associated with poor glucose tolerance, independent of body weight, fat mass, energy intake and diet composition in prediabetes or early onset type 2 diabetes. Nutrition & diabetes 2024;14:90

3. Ali M, Reutrakul S, Petersen G, Knutson KL. Associations between Timing and Duration of Eating and Glucose Metabolism: A Nationally Representative Study in the U.S. Nutrients 2023;15

4. Asher G, Sassone-Corsi P. Time for food: the intimate interplay between nutrition, metabolism, and the circadian clock. Cell 2015;161:84-92

5. Peters B, Vahlhaus J, Pivovarova-Ramich O. Meal timing and its role in obesity and associated diseases. Frontiers in endocrinology 2024;15:1359772

6. Vahlhaus J, Peters B, Hornemann S, Ost AC, Kruse M, Busjahn A, Pfeiffer AFH, Pivovarova-Ramich O. Later eating timing in relation to an individual internal clock is associated with lower insulin sensitivity and affected by genetic factors. EBioMedicine 2025;116:105737

7. McHill AW, Phillips AJ, Czeisler CA, Keating L, Yee K, Barger LK, Garaulet M, Scheer FA, Klerman EB. Later circadian timing of food intake is associated with increased body fat. Am J Clin Nutr 2017;106:1213-1219

8. Martinez-Lozano N, Tvarijonaviciute A, Rios R, Baron I, Scheer F, Garaulet M. Late Eating Is Associated with Obesity, Inflammatory Markers and Circadian-Related Disturbances in School-Aged Children. Nutrients 2020;12

9. Sutton EF, Beyl R, Early KS, Cefalu WT, Ravussin E, Peterson CM. Early Time-Restricted Feeding Improves Insulin Sensitivity, Blood Pressure, and Oxidative Stress Even without Weight Loss in Men with Prediabetes. Cell metabolism 2018;27:1212-1221 e1213

10. Jones R, Pabla P, Mallinson J, Nixon A, Taylor T, Bennett A, Tsintzas K. Two weeks of early time-restricted feeding (eTRF) improves skeletal muscle insulin and anabolic sensitivity in healthy men. Am J Clin Nutr 2020;112:1015-1028

11. Jamshed H, Beyl RA, Della Manna DL, Yang ES, Ravussin E, Peterson CM. Early Time-Restricted Feeding Improves 24-Hour Glucose Levels and Affects Markers of the Circadian Clock, Aging, and Autophagy in Humans. Nutrients 2019;11

12. Teong XT, Liu K, Vincent AD, Bensalem J, Liu B, Hattersley KJ, Zhao L, Feinle-Bisset C, Sargeant TJ, Wittert GA, Hutchison AT, Heilbronn LK. Intermittent fasting plus early time-restricted eating versus calorie restriction and standard care in adults at risk of type 2 diabetes: a randomized controlled trial. Nature medicine 2023;29:963-972

13. Andriessen C, Fealy CE, Veelen A, van Beek SMM, Roumans KHM, Connell NJ, Mevenkamp J, Moonen-Kornips E, Havekes B, Schrauwen-Hinderling VB, Hoeks J, Schrauwen P. Three weeks of time-restricted eating improves glucose homeostasis in adults with type 2 diabetes but does not improve insulin sensitivity: a randomised crossover trial. Diabetologia 2022;65:1710-1720

14. Peters B, Schwarz J, Schuppelius B, Ottawa A, Koppold DA, Weber D, Steckhan N, Mai K, Grune T, Pfeiffer AFH, Michalsen A, Kramer A, Pivovarova-Ramich O. Intended isocaloric time-restricted eating shifts circadian clocks but does not improve cardiometabolic health in women with overweight. Sci Transl Med 2025;17:eadv6787

15. Crispim CA, Rinaldi AEM, Azeredo CM, Skene DJ, Moreno CRC. Is time of eating associated with BMI and obesity? A population-based study. Eur J Nutr 2024;63:527-537

16. Dashti HS, Gomez-Abellan P, Qian J, Esteban A, Morales E, Scheer F, Garaulet M. Late eating is associated with cardiometabolic risk traits, obesogenic behaviors, and impaired weight loss. Am J Clin Nutr 2021;113:154-161

17. Shim JS, Kim HC. Late eating, blood pressure control, and cardiometabolic risk factors among adults with hypertension: results from the Korea National Health and Nutrition Examination Survey 2010-2018. Epidemiol Health 2021;43:e2021101

18. Nakajima K, Suwa K. Association of hyperglycemia in a general Japanese population with late-
night-dinner eating alone, but not breakfast skipping alone. J Diabetes Metab Disord 2015;14:16
19. Scheer FA, Hilton MF, Mantzoros CS, Shea SA. Adverse metabolic and cardiovascular
consequences of circadian misalignment. Proceedings of the National Academy of Sciences of the
United States of America 2009;106:4453-4458
20. Ballon A, Neuenschwander M, Schlesinger S. Breakfast Skipping Is Associated with Increased Risk
of Type 2 Diabetes among Adults: A Systematic Review and Meta-Analysis of Prospective Cohort
Studies. The Journal of nutrition 2019;149:106-113
21. Palomar-Cros A, Srour B, Andreeva VA, Fezeu LK, Bellicha A, Kesse-Guyot E, Hercberg S,
Romaguera D, Kogevinas M, Touvier M. Associations of meal timing, number of eating occasions and
night-time fasting duration with incidence of type 2 diabetes in the NutriNet-Sante cohort. Int J
Epidemiol 2023;52:1486-1497
22. Yamamoto R, Tomi R, Shinzawa M, Yoshimura R, Ozaki S, Nakanishi K, Ide S, Nagatomo I, Nishida
M, Yamauchi-Takihara K, Kudo T, Moriyama T. Associations of Skipping Breakfast, Lunch, and Dinner
with Weight Gain and Overweight/Obesity in University Students: A Retrospective Cohort Study.
Nutrients 2021;13
23. Jakubowicz D, Wainstein J, Ahren B, Bar-Dayan Y, Landau Z, Rabinovitz HR, Froy O. High-energy
breakfast with low-energy dinner decreases overall daily hyperglycaemia in type 2 diabetic patients:
a randomised clinical trial. Diabetologia 2015;58:912-919
24. Jakubowicz D, Wainstein J, Landau Z, Ahren B, Barnea M, Bar-Dayan Y, Froy O. High-energy
breakfast based on whey protein reduces body weight, postprandial glycemia and HbA(1C) in Type 2
diabetes. J Nutr Biochem 2017;49:1-7
25. Dai J, Nianogo R, Wong ND, Moin T, McClain AC, Alver S, Cordero C, Daviglus ML, Qi Q, Sotres-
Alvarez D, Chen L. Energy Intake and Dietary Glycemic Load in Late Morning and Risk of Type 2
Diabetes: The Hispanic Community Health Study/Study of Latinos-A Multicenter Prospective Cohort
Study. Diabetes care 2024;47:1673-1681
26. Kahleova H, Belinova L, Malinska H, Oliyarnyk O, Trnovska J, Skop V, Kazdova L, Dezortova M,
Hajek M, Tura A, Hill M, Pelikanova T. Eating two larger meals a day (breakfast and lunch) is more
effective than six smaller meals in a reduced-energy regimen for patients with type 2 diabetes: a
randomised crossover study. Diabetologia 2014;57:1552-1560
27. Fritsche A, Wagner R, Heni M, Kantartzis K, Machann J, Schick F, Lehmann R, Peter A, Dannecker
C, Fritsche L, Valenta V, Schick R, Nawroth PP, Kopf S, Pfeiffer AFH, Kabisch S, Dambeck U, Stumvoll
M, Bluher M, Birkenfeld AL, Schwarz P, Hauner H, Clavel J, Seissler J, Lechner A, Mussig K, Weber K,
Laxy M, Bornstein S, Schurmann A, Roden M, de Angelis MH, Stefan N, Haring HU. Different Effects of
Lifestyle Intervention in High- and Low-Risk Prediabetes: Results of the Randomized Controlled
Prediabetes Lifestyle Intervention Study (PLIS). Diabetes 2021;70:2785-2795
28. Wagner R, Heni M, Kantartzis K, Sandforth A, Machann J, Schick F, Peter A, Fritsche L, Szendrodi J,
Pfeiffer AFH, Schurmann A, Bluher M, Hauner H, Seissler J, Bornstein S, Roden M, Stefan N,
Birkenfeld AL, White MF, Haring HU, Fritsche A. Lower Hepatic Fat Is Associated With Improved
Insulin Secretion in a High-Risk Prediabetes Subphenotype During Lifestyle Intervention. Diabetes
2023;72:362-366
29. Kabisch S, Bather S, Dambeck U, Kemper M, Gerbracht C, Honsek C, Sachno A, Pfeiffer AFH. Liver
Fat Scores Moderately Reflect Interventional Changes in Liver Fat Content by a Low-Fat Diet but Not
by a Low-Carb Diet. Nutrients 2018;10
30. Stumvoll M, Mitrakou A, Pimenta W, Jenssen T, Yki-Jarvinen H, Van Haeften T, Renn W, Gerich J.
Use of the oral glucose tolerance test to assess insulin release and insulin sensitivity. Diabetes care
2000;23:295-301
31. Bergman RN, Ader M, Huecking K, Van Citters G. Accurate assessment of beta-cell function: the
hyperbolic correction. Diabetes 2002;51 Suppl 1:S212-220
32. Matsuda M, DeFronzo RA. Insulin sensitivity indices obtained from oral glucose tolerance testing:
comparison with the euglycemic insulin clamp. Diabetes care 1999;22:1462-1470

33. Pivovarova O, Isken F, Bernigau W, Bobbert T, Spranger J, Weickert MO, Pfeiffer AF, Rudovich N.
Decreased hepatic insulin clearance is closely related to metabolic syndrome components. Exp Clin
Endocr Diab 2013;121
34. Gill S, Panda S. A Smartphone App Reveals Erratic Diurnal Eating Patterns in Humans that Can Be
Modulated for Health Benefits. Cell metabolism 2015;22:789-798
35. Gibney MJ, Wolever TM. Periodicity of eating and human health: present perspective and future
directions. Br J Nutr 1997;77 Suppl 1:S3-5
36. Johnson GH, Anderson GH. Snacking definitions: impact on interpretation of the literature and
dietary recommendations. Crit Rev Food Sci Nutr 2010;50:848-871
37. Silva CM, Teixeira BS, Wright KP, Jr., Maia YCP, Crispim CA. Time-Related Eating Patterns Are
Associated with the Total Daily Intake of Calories and Macronutrients in Day and Night Shift Workers.
Nutrients 2022;14
38. Ahlqvist E, Storm P, Karajamaki A, Martinell M, Dorkhan M, Carlsson A, Vikman P, Prasad RB, Aly
DM, Almgren P, Wessman Y, Shaat N, Spegel P, Mulder H, Lindholm E, Melander O, Hansson O,
Malmqvist U, Lernmark A, Lahti K, Forsen T, Tuomi T, Rosengren AH, Groop L. Novel subgroups of
adult-onset diabetes and their association with outcomes: a data-driven cluster analysis of six
variables. Lancet Diabetes Endocrinol 2018;6:361-369
39. Freeman AM, Acevedo LA, Pennings N. Insulin Resistance. In *StatPearls* Treasure Island (FL), 2025
40. Bi H, Gan Y, Yang C, Chen Y, Tong X, Lu Z. Breakfast skipping and the risk of type 2 diabetes: a
meta-analysis of observational studies. Public Health Nutr 2015;18:3013-3019
41. Reutrakul S, Hood MM, Crowley SJ, Morgan MK, Teodori M, Knutson KL. The relationship
between breakfast skipping, chronotype, and glycemic control in type 2 diabetes. Chronobiology
international 2014;31:64-71
42. Kant AK, Graubard BI. Clock Time of First Eating Episode and Prospective Risk of All-Cause
Mortality in US Adults. The Journal of nutrition 2022;152:217-226
43. Qian J, Scheer FA. Circadian System and Glucose Metabolism: Implications for Physiology and
Disease. Trends in endocrinology and metabolism: TEM 2016;27:282-293
44. Lopez-Minguez J, Gomez-Abellan P, Garaulet M. Timing of Breakfast, Lunch, and Dinner. Effects
on Obesity and Metabolic Risk. Nutrients 2019;11
45. Ezpeleta M, Cienfuegos S, Lin S, Pavlou V, Gabel K, Tussing-Humphreys L, Varady KA. Time-
restricted eating: Watching the clock to treat obesity. Cell metabolism 2024;36:301-314
46. Schuppelius B, Peters B, Ottawa A, Pivovarova-Ramich O. Time Restricted Eating: A Dietary
Strategy to Prevent and Treat Metabolic Disturbances. Frontiers in endocrinology 2021;12:683140
47. Zhao D, Guallar E, Woolf TB, Martin L, Lehmann H, Coughlin J, Holzhauer K, Goheer AA, McTigue
KM, Lent MR, Hawkins M, Clark JM, Bennett WL. Association of Eating and Sleeping Intervals With
Weight Change Over Time: The Daily24 Cohort. J Am Heart Assoc 2023;12:e026484
48. Chang CR, Francois ME, Little JP. Restricting carbohydrates at breakfast is sufficient to reduce 24-
hour exposure to postprandial hyperglycemia and improve glycemic variability. Am J Clin Nutr
2019;109:1302-1309
49. Bauer LB, Reynolds LJ, Douglas SM, Kearney ML, Hoertel HA, Shafer RS, Thyfault JP, Leidy HJ. A
pilot study examining the effects of consuming a high-protein vs normal-protein breakfast on free-
living glycemic control in overweight/obese 'breakfast skipping' adolescents. Int J Obes (Lond)
2015;39:1421-1424
50. Poggiogalle E, Jamshed H, Peterson CM. Circadian regulation of glucose, lipid, and energy
metabolism in humans. Metabolism: clinical and experimental 2018;84:11-27
51. Carrasco-Benso MP, Rivero-Gutierrez B, Lopez-Minguez J, Anzola A, Diez-Noguera A, Madrid JA,
Lujan JA, Martinez-Augustin O, Scheer FA, Garaulet M. Human adipose tissue expresses intrinsic
circadian rhythm in insulin sensitivity. FASEB journal : official publication of the Federation of
American Societies for Experimental Biology 2016;30:3117-3123
52. van Moorsel D, Hansen J, Havekes B, Scheer FAJL, Jörgensen JA, Hoeks J, Schrauwen-Hinderling
VB, Duez H, Lefebvre P, Schaper NC, Hesselink MKC, Staels B, Schrauwen P. Demonstration of a day-
night rhythm in human skeletal muscle oxidative capacity. Molecular Metabolism 2016;5:635-645

53. Saad A, Dalla Man C, Nandy DK, Levine JA, Bharucha AE, Rizza RA, Basu R, Carter RE, Cobelli C,
Kudva YC, Basu A. Diurnal pattern to insulin secretion and insulin action in healthy individuals.
Diabetes 2012;61:2691-2700
54. McHill AW, Butler MP. Eating Around the Clock: Circadian Rhythms of Eating and Metabolism.
Annu Rev Nutr 2024;44:25-50
55. SHAPIRO ET, POLONSKY KS, COPINSCHI G, BOSSON D, TILLIL H, BLACKMAN J, LEWIS G, CAUTER
EV. Nocturnal Elevation of Glucose Levels during Fasting in Noninsulin-Dependent Diabetes*. The
Journal of Clinical Endocrinology & Metabolism 1991;72:444-454
56. Boden G, Chen X, Urbain JL. Evidence for a Circadian Rhythm of Insulin Sensitivity in Patients With
NIDDM Caused by Cyclic Changes in Hepatic Glucose Production. Diabetes 1996;45:1044-1050
57. Mason IC, Qian J, Adler GK, Scheer F. Impact of circadian disruption on glucose metabolism:
implications for type 2 diabetes. Diabetologia 2020;63:462-472
58. Sonnier T, Rood J, Gimble JM, Peterson CM. Glycemic control is impaired in the evening in
prediabetes through multiple diurnal rhythms. J Diabetes Complications 2014;28:836-843
59. Bergman RN, Piccinini F, Kabir M, Kolka CM, Ader M. Hypothesis: Role of Reduced Hepatic Insulin
Clearance in the Pathogenesis of Type 2 Diabetes. Diabetes 2019;68:1709-1716
60. Pivovarova O, Bernigau W, Bobbert T, Isken F, Mohlig M, Spranger J, Weickert MO, Osterhoff M,
Pfeiffer AF, Rudovich N. Hepatic insulin clearance is closely related to metabolic syndrome
components. Diabetes care 2013;36:3779-3785
61. Kotronen A, Juurinen L, Tiikkainen M, Vehkavaara S, Yki-Jarvinen H. Increased liver fat, impaired
insulin clearance, and hepatic and adipose tissue insulin resistance in type 2 diabetes.
Gastroenterology 2008;135:122-130
62. Najjar SM, Caprio S, Gastaldelli A. Insulin Clearance in Health and Disease. Annual review of
physiology 2023;85:363-381
63. Boden G, Ruiz J, Urbain JL, Chen X. Evidence for a circadian rhythm of insulin secretion. The
American journal of physiology 1996;271:E246-252
64. Demir M, Bornstein SR, Mantzoros CS, Perakakis N. Liver fat as risk factor of hepatic and
cardiometabolic diseases. Obesity reviews : an official journal of the International Association for the
Study of Obesity 2023;24:e13612
65. de Assis LVM, Demir M, Oster H. The role of the circadian clock in the development, progression,
and treatment of non-alcoholic fatty liver disease. Acta Physiol (Oxf) 2023;237:e13915
66. Hatori M, Vollmers C, Zarrinpar A, DiTacchio L, Bushong EA, Gill S, Leblanc M, Chaix A, Joens M,
Fitzpatrick JAJ, Ellisman MH, Panda S. Time-Restricted Feeding without Reducing Caloric Intake
Prevents Metabolic Diseases in Mice Fed a High-Fat Diet. Cell metabolism 2012;15:848-860
67. Oh JH, Yoon EL, Park H, Lee S, Jo AJ, Cho S, Kwon E, Nah EH, Lee JH, Park JH, Ahn SB, Jun DW.
Efficacy and safety of time-restricted eating in metabolic dysfunction-associated steatotic liver
disease. Journal of hepatology 2025;
68. Reutrakul S, Van Cauter E. Sleep influences on obesity, insulin resistance, and risk of type 2
diabetes. Metabolism: clinical and experimental 2018;84:56-66

Figure 1. Association of eating timing, glycaemic and anthropometric parameters. Heatmap represents beta-coefficient values calculated using linear regression models without (a) and with (b) adjustment for sex, age, and daily energy intake (n=297). Eating timing parameters were averaged over four days for each person. Intensity of colour represents the strength of the association (beta-coefficients), whereas red indicates a positive and blue a negative association.

693 **Figure 2. Glucose and insulin values in OGTT in individuals with early vs late start of eating.** Glucose (A), insulin (B), and C-peptide (C) values

694 in OGTT are presented in individuals with early (blue) and late (red) start of eating (n=297). Data are shown as mean ± SEM. ** P<0.01, * P<0.05

695 by the unpaired T-test. ES, eating start.

697
698

699 **Table 1. Clinical characteristics of study participants**

Parameters	All
Male / female	106 / 191
Age [years]	59.5 (8.8)
Anthropometric parameters	
BMI [kg/m²]	31.3 (6.1)
Waist circumference [cm]	101.8 (14.7)
Total fat mass [%]	34.9 (8.1)
Visceral fat mass [% of total fat]	24.4 (9.1)
Liver fat [%]	9.12 (8.00)
Blood pressure	
Systolic blood pressure [mmHg]	136.2 (15.1)
Diastolic blood pressure [mmHg]	88.2 (10.3)
Lipid parameters	
Triglycerides [mg/dl]	139.3 (76.0)
Total cholesterol [mg/dl]	211.9 (36.7)
HDL cholesterol [mg/dl]	54.1 (15.2)
LDL cholesterol [mg/dl]	137.6 (32.6)
Glycaemic parameters	
Fasting glucose [mg/dl]	106.7 (9.5)
AUC glucose [mg/dl*min]	19839 (3212)
Fasting insulin [mU/l]	11.0 (7.6)
AUC insulin [mU/l*min]	10269 (7908)
HOMA-IR	2.90 (2.03)
ISI Matsuda [AU]	3.67 (2.42)
Insulinogenic index	0.89 (0.80)
Disposition index [AU]	2.67 (2.96)
HbA1c [%]	5.67 (0.36)

701 Data are shown as mean (SD).

Table 2. Dietary composition and meal timing pattern of study participants

Parameters	All
Daily dietary composition	
Daily calorie intake [kcal]	1941 (547)
Carbohydrates [g]	198.9 (65.1)
Carbohydrates [EN%]	42.6 (7.5)
Protein [g]	83.6 (24.7)
Protein [EN%]	18.2 (4.6)
Fat [g]	80.6 (30.5)
Fat [EN%]	36.7 (7.6)
Fibre [g]	24.0 (8.3)
Fibre [EN%]	2.56 (1.02)
Eating timing	
Eating window [hh:mm]	12:09 (01:49)
Start of first eating event [hh:mm]	08:09 (01:24)
End of last eating event [hh:mm]	20:06 (02:13)
Daily calorie distribution	
Caloric midpoint [hh:mm]	15:19 (02:04)
Calories in first meal [%]	27.7 (10.1)
Calories in last meal [%]	33.8 (11.3)
Eating frequency	
Number of eating events	4.07 (0.92)

Data are shown as mean (SD).

Table 3. Characteristics of individuals with early vs late start of eating

Parameters	Early eating start	Late eating start	P
Male / female	65 / 95	41 / 96	
Age [years]	58.6 (8.6)	60.5 (9.2)	**0.018**
Anthropometric parameters			
BMI [kg/m²]	31.3 (6.3)	31.3 (6.0)	0.994
Waist circumference [cm]	101.6 (14.3)	102.1 (15.2)	0.781
Fat mass [%]	33.6 (8.6)	36.3 (7.3)	**0.004**
Visceral fat [% of total fat]	24.4 (9.4)	24.4 (8.9)	0.743
Liver fat [%]	8.84 (8.03)	9.42 (7.99)	0.465
Lipid parameters			
Triglycerides [mg/dl]	137.8 (70.9)	140.9 (81.8)	0.812
Total cholesterol [mg/dl]	213.0 (38.8)	210.5 (34.1)	0.553
HDL cholesterol [mg/dl]	53.5 (15.0)	54.8 (15.6)	0.467
LDL cholesterol [mg/dl]	138.8 (34.9)	136.2 (29.9)	0.511
Glycaemic parameters			
Fasting glucose [mg/dl]	106.2 (9.3)	107.2 (9.6)	0.366
AUC glucose [mg/dl*min]	19752 (3201)	19942 (3233)	0.612
Fasting insulin [mU/l]	10.0 (6.3)	12.2 (8.8)	**0.006**
AUC insulin [mU/l*min]	9213 (6067)	11493 (9490)	**0.020**
Fasting C-peptide [µg/l]	1.52 (0.89)	1.68 (1.16)	0.211
AUC C-peptide [µg/l*min]	763.1 (315.3)	758.3 (427.9)	0.647
HOMA-IR	2.60 (1.60)	3.25 (2.39)	**0.004**
ISI Matsuda [AU]	4.03 (2.67)	3.24 (2.03)	**0.004**
Insulinogenic index	0.79 (0.65)	1.01 (0.93)	**0.049**
Disposition index [AU]	2.61 (2.43)	2.75 (3.48)	0.715
Hepatic insulin clearance [l/min]	4.91 (2.20)	4.05 (2.36)	**<0.001**
HbA1c [%]	5.7 (0.4)	5.7 (0.3)	0.063
Eating timing			
Eating window [hh:mm]	12:56 (1:34)	11:14 (1:39)	**<0.001**
Start of eating [hh:mm]	7:11 (0:43)	9:15 (1:09)	**<0.001**
End of eating [hh:mm]	20:06 (1:35)	20:06 (2:47)	0.411
Daily calorie distribution			
Daily calorie intake [kcal]	1983 (586)	1891 (496)	0.302
Caloric midpoint [hh:mm]	14:59 (1:58)	15:41 (2:08)	**0.003**
Calories in first meal [%]	26.7 (10.4)	28.7 (9.7)	**0.041**
Calories in last meal [%]	33.1 (11.2)	34.7 (11.4)	0.181
Eating frequency			
Number of eating events	4.21 (0.95)	3.91 (0.87)	**0.006**

Data are shown as mean (SD).

Late start of eating is linked to lower insulin sensitivity and hyperinsulinemia in prediabetes

Beeke Peters, Noelle Fröhlich, Jürgen Machann, Ulrike Dambeck, Caroline Honsek, Anna Sachno, Margrit Kemper, Stefan Kabisch, Andreas Fritsche, Knut Mai, Andreas F.H. Pfeiffer, Olga Pivovarova-Ramich

Supplementary material

Supplementary Figures

Supplementary Tables

SUPPLEMENTARY FIGURES

Figure S1. Daily distribution of eating events. Distribution of eating events is shown as percentage of all eating events in all study participants (n=297). Each number at the time axis represents an hour starting from 0 (24:00 – 1:00) and ending with 23 (23:00 – 24:00).

741 **Figure S2. Associations of eating start with insulin levels and insulin sensitivity.**

742 Associations with (A) fasting insulin, (B) AUC insulin, (C) insulinogenic index (IGI), (D) HOMA-

743 IR, and (E) ISI Matsuda are shown. Beta-coefficient and p-values calculated using linear

744 regression models adjusted for sex, age, and average daily energy intake are represented.

Table S1. Correlation analysis of eating timing parameters and daily calorie distribution

	Eating window	Start of first eating event	End of last eating event	Caloric midpoint	Calories in first meal	Calories in last meal	Daily calorie intake
Start of first eating event	-0.582**	-	n.s.	n.s.	n.s.	n.s.	n.s.
End of last eating event	0.461**	n.s.	-	n.s.	n.s.	n.s.	n.s.
Caloric midpoint	0.180**	0.332**	n.s.	-	n.s.	n.s.	n.s.
Calories in first meal	-0.347**	0.178**	-0.146*	-0.211**	-	n.s.	n.s.
Calories in last meal	-0.221**	0.167**	-0.152**	0.454**	0.159**	-	n.s.
Daily calorie intake	0.298**	n.s.	n.s.	0.224**	-0.182**	n.s.	-
Number of meals	0.502**	-0.182**	0.243**	n.s.	-0.494**	-0.500**	0.232**

Only significant correlation coefficients are shown. * if $p < 0.05$, ** if $p < 0.01$, and *** if $p < 0.001$.

Table S2. Associations of start of the first eating event with glycaemic and anthropometric traits

	Model 1		Model 2		Model 3	
	β	p	β	p	β	p
Glycaemic traits						
Fasting glucose	0.096	0.099	0.100	0.086	0.106	0.067
AUC glucose	-0.010	0.866	-0.001	0.984	0.001	0.984
Fasting insulin	0.221	**1.7×10^{-4}**	0.217	**1.8×10^{-4}**	0.221	**1.4×10^{-4}**
AUC insulin	0.208	**4.1×10^{-4}**	0.205	**4.5×10^{-4}**	0.204	**4.8×10^{-4}**
HOMA-IR	0.236	**6.2×10^{-5}**	0.233	**6.3×10^{-5}**	0.238	**4.4×10^{-4}**
ISI Matsuda	-0.234	**6.9×10^{-5}**	-0.233	**7.0×10^{-5}**	-0.235	**6.1×10^{-5}**
Insulinogenic index	0.186	**0.002**	0.187	**0.001**	0.187	**0.001**
Disposition index	-0.002	0.969	0.000	1.000	-0.002	0.969
Anthropometric traits						
BMI	0.104	0.073	0.100	0.082	0.099	0.086
Waist circumference	0.103	0.077	0.125	**0.026**	0.126	**0.025**
Fat mass	0.189	**0.001**	0.147	**0.002**	0.143	**0.002**
Visceral fat	0.023	0.716	0.074	0.068	0.077	0.056
Liver fat	0.086	0.166	0.089	0.152	0.094	0.130

Linear regression analyses were performed to estimate the strength of association between variables. Model 1 is without any covariate adjustment. model 2 is adjusted for sex and age, and model 3 for sex, age, and daily energy intake. β are the standardized coefficients.

Table S3. Associations of caloric midpoint with glycaemic and anthropometric traits

	Model 1		Model 2		Model 3	
	β	p	β	p	β	p
Glycaemic traits						
Fasting glucose	0.161	**0.005**	0.172	**0.003**	0.155	**0.009**
AUC glucose	-0.027	0.646	-0.023	0.691	-0.033	0.580
Fasting insulin	0.142	**0.016**	0.135	**0.022**	0.125	**0.038**
AUC insulin	0.033	0.576	0.024	0.687	0.029	0.631
HOMA-IR	0.163	**0.006**	0.156	**0.008**	0.143	**0.017**
ISI Matsuda	-0.098	0.099	-0.090	0.131	-0.086	0.160
Insulinogenic index	0.062	0.301	0.040	0.500	0.044	0.458
Disposition index	-0.019	0.754	-0.036	0.550	-0.027	0.663
Anthropometric traits						
BMI	0.095	0.102	0.084	0.148	0.092	0.122
Waist circumference	0.126	**0.030**	0.092	0.105	0.092	0.112
Fat mass	-0.004	0.949	0.062	0.192	0.079	0.104
Visceral fat	0.115	0.064	0.035	0.398	0.027	0.511
Liver fat	0.005	0.933	-0.001	0.989	-0.016	0.800

Linear regression analyses were performed to estimate the strength of association between variables. Model 1 is without any covariate adjustment. model 2 is adjusted for sex and age, and model 3 for sex, age, and daily energy intake. β are the standardized coefficients.

Table S4. Associations of other eating timing parameters with glycaemic and anthropometric traits

	Model 1		Model 2		Model 3	
	β	p	β	p	β	p
Calories in the first meal						
Waist circumference	0.169	**0.004**	0.086	0.145	0.099	0.138
Calories in the last meal						
Waist circumference	0.224	**1.0×10^{-4}**	0.129	**0.039**	0.172	**0.022**
Visceral fat	0.322	**9.9×10^{-8}**	0.084	0.065	0.067	0.210

Linear regression analyses were performed to estimate the strength of association between variables. Model 1 is without any covariate adjustment, model 2 is adjusted for sex and age, and model 3 for sex, age, and daily energy intake. Only parameters showing significant associations in model 1 are shown. β are the standardized coefficients.

Table S5. Food composition and calorie intake in individuals with early vs. late start of eating

Parameters	Early eating start	Late eating start	P*
Daily dietary composition			
Calorie intake [kcal]	1983 (586)	1891 (496)	0.302
Carbohydrates [g]	205.3 (66.3)	191.4 (63.0)	0.062
Carbohydrates [EN%]	43.0 (7.5)	42.2 (7.6)	0.375
Protein [g]	85.7 (28.0)	81.2 (20.0)	0.353
Protein [EN%]	18.2 (5.0)	18.1 (4.0)	0.954
Fat [g]	82.0 (32.4)	78.9 (28.2)	0.748
Fat [EN%]	36.2 (7.3)	37.2 (7.8)	0.271
Fibre [g]	25.0 (8.4)	22.7 (8.1)	**0.024**
Fibre [EN%]	2.60 (0.87)	2.52 (1.17)	0.300
Dietary composition of the first meal			
Calorie intake [kcal]	488 (202)	509 (218)	0.524
Calorie intake [% of daily intake]	26.7 (10.4)	28.7 (9.7)	**0.041**
Carbohydrates [g]	57.5 (26.1)	56.9 (27.0)	0.902
Carbohydrates [EN%]	48.8 (12.0)	46.9 (11.6)	0.171
Protein [g]	20.2 (11.8)	20.8 (10.0)	0.416
Protein [EN%]	16.4 (5.6)	16.6 (4.9)	0.981
Fat [g]	18.8 (11.4)	20.7 (12.4)	0.210
Fat [EN%]	31.8 (11.2)	33.6 (12.3)	0.176
Fibres [g]	7.29 (3.88)	7.07 (3.95)	0.527
Fibres [EN%]	3.00 (1.26)	2.82 (1.25)	0.155

Data are shown as mean (SD).

Table S6. Effects early vs. late start of eating on previously significant glycaemic parameters after adjusting for potential confounders

Adjusting covariates	Sex Age	Sex Age Energy intake	Sex Age Energy intake Body fat
Fasting insulin [mU/l]	0.002	0.002	0.006
AUC insulin [mU/l*min]	0.010	0.010	0.022
HOMA-IR	0.001	0.001	0.003
ISI Matsuda [AU]	0.002	0.001	0.004
Insulinogenic index	0.011	0.012	0.019

P-values calculated using the one-way ANOVA with adjustment for sex, age, average daily energy intake, and total body fat are shown.