Implementation and first results of the Czech nationwide prostate cancer

screening pilot programme

Marcela Koudelková[a,b], Kateřina Hejcmanová[a,b], Marek Babjuk[c], Roman Zachoval[d],

Jiří Ferda[e], Ola Bratt[f,g], Renata Chloupková[a,b], Ondřej Ngo[a,b], Karel Hejduk[a,b], Vlastimil

Válek[h], Ladislav Dušek[a,b,i], Ondřej Májek[a,b,i]

[a]National Screening Centre, Institute of Health Information and Statistics of the Czech Republic, Prague,

Czech Republic

[b]Institute of Biostatistics and Analyses, Faculty of Medicine, Masaryk University, Brno, Czech Republic

[c]Department of Urology, 2nd Faculty of Medicine, Charles University, Motol University Hospital, Prague,

Czech Republic

[d]Department of Urology, 3rd Faculty of Medicine, Charles University and Faculty Thomayer Hospital,

Prague, Czech Republic

[e]Department of Imaging Methods Faculty of Medicine in Pilsen

[f]Department of Urology, Institute of Clinical Sciences, Sahlgrenska Academy, University of Gothenburg,

Gothenburg, Sweden

[g]Department of Urology, Sahlgrenska University Hospital, Gothenburg, Sweden

[h]Department of Radiology and Nuclear Medicine, Masaryk University, Brno, Czech Republic

[i]RECETOX, Faculty of Science, Masaryk University, Kotlarska 2, Brno, Czech Republic

Corresponding author

Ondřej Májek, PhD

Institute of Biostatistics and Analyses, Faculty of Medicine, Masaryk University

Kamenice 126/3, 625 00 Brno, Czech Republic

+420 778 543 692

majek@iba.muni.cz

Keywords

Cancer screening, Early detection, Population, Prostate cancer, Prostate-specific

antigen, Screening

Abstract

Background

In 2022, the European Union recommended piloting organised prostate cancer screening. Subsequently, the Czech Republic launched a nationwide pilot programme fully reimbursed by public health insurance in January 2024 to reduce the high incidence of late-stage prostate cancer and the widespread unorganised prostate-specific antigen (PSA) testing.

Objective

To describe the strategy, methodology, and first results of the Czech nationwide prostate cancer screening pilot programme.

Methods

We analysed data from the first 12 months of the programme, which targets men aged 50–69 years without a previous prostate cancer diagnosis. General practitioners and urologists offer men an initial PSA test. Those with PSA ≥3.0 µg/l are referred to certified urologists, based on urology assessment selected for magnetic resonance imaging (MRI) and, if indicated, targeted biopsy. Reported key indicators are PSA testing coverage and overall use of pre-biopsy MRI.

Key Findings and Limitations

Screening was offered to 150,498 men; PSA results were available for 146,109 (97.1%), 8.8% of whom had ≥3.0 µg/l. Within 6 months, 67.2% with PSA ≥3.0 µg/l

underwent urological evaluation. From 2023 to 2024, the 2-year PSA testing rate rose

from 47.4% to 53.3%, and pre-biopsy MRI use from 28.0% to 38.3% (10.3 percentage-

point difference, 95% CI 8.3–12.2). Data on diagnostic outcomes are not yet analysed.

Conclusions and Clinical Implications

These results demonstrate the feasibility of transforming widespread unorganised PSA

testing to an organised screening programme with improved adherence to the

recommended diagnostic pathway. Further evaluation to assess outcomes and efforts

to reduce opportunistic testing will follow.

Advancing Practice

What does this study add?

This is the first population-level evaluation of a nationwide organised prostate cancer

screening programme in Central Europe. The findings demonstrate the feasibility of

integrating a structured screening programme with PSA testing in routine primary care

into a national health system. The programme improved adherence to guideline-based

diagnostic pathways.

Patient Summary

This report presents the initial results of a new nationwide, organised prostate cancer

screening programme in the Czech Republic. The programme contributed to increased

screening coverage in the target population (50-69 years) and improved the use of MRI

before biopsy, which is now the recommended practice.

Introduction

Prostate cancer is one of the most common malignancies among men in the European Union, and the incidence is steadily increasing [1]. Late diagnosis is a persistent challenge, with approximately one-third of cases detected in stage III or IV in the Czech Republic [2], when therapeutic options are limited and the prognosis is poor [2]. In contrast, early detection significantly increases the chance of curative treatment and long-term survival [3] [4]. Approximately 8,000 new prostate cancer cases are diagnosed annually in the Czech Republic, nearly three times more than two decades ago [2]. This rise is attributable not only to the ageing population but also to widespread use of opportunistic prostate-specific antigen (PSA) testing. International randomized trials have demonstrated that organised PSA-based screening can reduce prostate cancer mortality [4]. However, it also carries risks of overdiagnosis and overtreatment, although less so than unorganised (opportunistic) testing [4] [5]. Screening may thus lead to clinically unjustified diagnostic interventions and put an unnecessary burden on healthcare services [6]. In response to current challenges and the potential benefits of organised screening using modern prostate cancer diagnostics, including magnetic resonance imaging (MRI) and targeted biopsy, the EU in 2022 recommended that member states assess the feasibility and effectiveness of organised screening programmes [8]. This initiative was also endorsed by the 2022 International Prostaforum declaration, which supported the launch of pilot programmes for early detection across Europe [9].

Based on these international and national frameworks, a nationwide prostate cancer

screening pilot programme was launched in the Czech Republic in January 2024 [10].

The primary objective of the programme is to assess the feasibility of organised

prostate cancer screening using PSA as a simple and cost-effective biomarker to guide

diagnostics and reduce advanced-stage disease. A further aim is to validate the

structured early detection algorithm incorporating PSA-based risk stratification and

MRI-guided biopsy within the Czech healthcare system [11].

We here describe the strategy and methods employed in the Czech national prostate

cancer screening programme and discuss its initial outcomes, to provide insights for

future screening.

Patients and Methods

Target population

106 The nationwide prostate cancer screening pilot targets men aged 50–69 years with no

107 prior prostate cancer or clinical suspicion. Men over 70 are not actively enrolled but

108 may be offered the same scheme based on individual health, at the physician's

109 discretion.

Enrolment and diagnostic pathway

The screening algorithm was developed based on international evidence-based

recommendations and the consensus of relevant professional societies [12–14], while

incorporating local specifics of the Czech healthcare system. A detailed schematic

representation of the algorithm is provided in Figure 1 and the Supplementary material.

Enrolment is undertaken by general practitioners (GPs) at routine preventive check-

ups, recommended biennially in this age group. Urologists additionally enrol patients

in the context of managing other urological conditions. After providing information and

obtaining oral consent, a PSA test is sampled. Management is stratified by PSA: <1.0

µg/l, re-test in 4 years; 1.0–2.99 µg/l, in 2 years; ≥3.0 µg/l, referral to a Czech Urological

Society–certified urologist for further assessment. If indicated (Fig. 1 and Supplement),

men undergo biparametric MRI at a Ministry of Health–certified centre. MRI is

interpreted using PI-RADS (Prostate Imaging Reporting and Data System): PI-RADS

1–2, repeat evaluation in 1 year; PI-RADS 3, repeat multiparametric MRI with contrast

(peripheral zone lesion) or PSA re-test in 6 months (transitional zone lesion); PI-RADS

4–5, image fusion biopsy at a certified centre. Men with prostate cancer are referred to

a multidisciplinary team, preferably at a certified onco-urological centre.

Data sources and statistical methods

Patient-level data from the National Health Information System (NHIS) were used for

monitoring and evaluation. The primary data source was the National Registry of

Reimbursed Health Services (NRRHS), which collects data on reimbursed care from

health insurance companies in the Czech Republic. Demographic data from the Czech

Statistical Office (CZSO) were used to determine the target population.

The primary objective of the present analysis was the identification of men who were

approached to enter the programme and their subsequent PSA values. As national

registries do not currently allow the collection of quantitative values, it was only

possible to obtain information on whether a man had a PSA value <1 µg/l, 1–2.99 µg/l,

or ≥3 µg/l. All these results were also analysed by 5-year age groups. The involvement

of GPs (in connection with preventive check-ups) and urologists was also monitored.

The data on enrolment and baseline PSA testing were assessed for the first year of

the organised programme, from January through December 2024. Men who were

identified as having PSA ≥3 µg/l were analysed to see if they had a follow-up

appointment with a urologist within 6 months. Due to data availability, this indicator was

only assessed in men with a PSA result in January-June 2024.

The second objective was to analyse how the organised screening programme affects

opportunistic PSA testing. Coverage of the target population was defined as the

proportion of men aged 50–69 years with a PSA test in the past two years, divided by

the number of men in the target population. Men who died before the year for which

the indicator was evaluated were not included in the coverage. The age of men was

determined for the year the coverage was evaluated, i.e., all men who turned 50 years

of age in 2024 were categorised as aged 50 years.

The third objective was to assess diagnostic guidelines adherence before and after

initiation of the screening programme. For men aged 50–69 years who underwent PSA

testing (by GPs or urologists) in the first half of 2023 and the first half of 2024 (within

or outside the screening programme), it was monitored whether they subsequently had

a prostate biopsy and, if so, whether it was preceded by a prostate MRI, or the MRI

was performed afterwards. Two-sided 95% confidence intervals (CI) were calculated

for the presented proportions and their difference.

Results

Recruitment and baseline PSA testing

In the first twelve months of the Czech national pilot prostate cancer screening programme, a total of 150,498 men aged 50–69 years were offered to participate. Most men (97.1%) were enrolled in the screening programme by GPs, mainly at a routine preventive health check-up. In 2024, 3,291 GPs and 89 urologists enrolled men in the programme.

PSA results were available for 146,109 (diagram of participation flow in Supplementary material), whereas 1,501 men chose not to participate. For 2,888 men, information about their willingness to enter the programme was not available. Willingness to participate was calculated only for men for whom PSA values or non-participation was known (99.0%).

The largest 5-year age group was 50–54 years (33.5%). Over half of the men (56.5%, n = 82,613) had PSA <1.0 µg/l. The proportion of men with PSA <1.0 µg/l decreased with age, from 66.5% in men aged 50–54 to 44.4% in those aged 65–69 years. A PSA value ≥3.0 µg/l was identified in 8.8% (n = 12,874). This proportion increased with age, from 4.0% in the 50–54 age group to 15.7% in the 65–69 age group (Table 1).

Of the 6,595 men with a PSA result ≥3.0 µg/l between January and June 2024, 4,431 (67.2%) underwent a urological examination within six months of the PSA result. This proportion declined slightly with age, from 69.9% in the youngest 5-year age group to 64.8% in the oldest (Supplementary material: Table 1).

Transition from opportunistic to organised screening: Results on a population

level

During 2024, 53.3% of the target population of men aged 50–69 years without

a prostate cancer diagnosis had a registered PSA test in the past 2 years, 5.9

percentage points more than in 2023 (47.4%; Figure 2).

PSA testing coverage consistently increased across all age groups from 2023, the year

before the organised screening programme, to its first year, 2024. The greatest

increase was observed in the youngest age group (50–54 years), with coverage rising

from 37.7% in 2023 to 44.6% in 2024 (Figure 3).

We assessed adherence to the screening diagnostic algorithm the first 6 months of the

programme and compared with the same 6-month period before the implementation of

the programme. The proportion of men undergoing prostate MRI before biopsy

increased from 28.0% to 38.3%, a difference of 10.3 percentage points (95% CI 8.3–

12.2 percentage points), while the rate of post-biopsy MRI decreased by 0.5

percentage points (Table 2).

Discussion

This report presents the first results of an organised nationwide prostate cancer screening pilot programme in the Czech Republic, targeting men aged 50–69 years.

Uptake of prostate cancer screening

Overall, 99.0% of the invited men decided to participate. This high participation rate is likely attributable to the fact that recruitment was conducted directly by GPs and urologists when men were physically attending healthcare and could have a PSA test with minimal extra effort. This personalized approach contrasts with many population-based screening programmes and trials that rely on centrally mailed invitation letters. It has been shown that men tend to have a higher level of trust in medical recommendations than in invitation letters [15].

Examples of population-based screening models that rely on mailed invitation letters are the Swedish Gothenburg-1 trial, where participation in the first rounds reached 59% among men aged 50–69 years [15], the Swedish regional organised prostate cancer testing (OPT) programmes with 35% participation among men aged 50 years [16], and the Finnish ProScreen trial with 51% participation among men aged 50–63 years [17].

The Czech approach builds on recommended regular preventive check-ups in primary care, enabling GPs to invite men for prostate screening during fully reimbursed visits, where PSA testing can be added with minimal extra burden. Urologist involvement in screening was limited due to payer contracts not automatically established by the end of 2024, delaying patient enrolment in the screening programme.

PSA results

216 The observed proportion of 8.8% of men aged 50–69 years with PSA ≥3 µg/l was in
217 the expected range, and so was the increase of this proportion with age [18]. In
218 comparison, 6.7% of men aged 50–60 in the Gothenburg-2 trial [19], 12% of men aged
219 50–74 in the STHLM3-MRI trial [14], and 10.7% of men aged 50–74 in the Lithuanian
220 population-based program had a PSA of ≥3 µg/l [20].

Urology assessment

The proportion of men with PSA of ≥3 µg/l who undergo the recommended subsequent
urological assessment was lower in the first six months of the Czech programme (67%)
than in the four Swedish regions with a similar diagnostic pathway for OPT (83%–99%)
[21]. The Czech proportion may be underestimated, as only a 6-month period was
analysed and waiting times for appointments may sometimes be longer. One barrier to
urological assessment is that it should be carried out by a certified urologist (lists are
available on an official website [11]), who may be located some distance away from
the participant's residence. Another factor is that the Czech programme has no
dedicated coordinators who actively communicate with participants and facilitate
appointments. In contrast, Swedish OPT has regional offices that does this [21].
Consequently, continued participation in the Czech programme largely depends on
individual motivation.

Prostate MRI

The use of pre-biopsy MRI and targeted biopsy to reduce unnecessary systematic
biopsy and overdiagnosis is crucial in modern prostate cancer screening [22]. After

programme initiation in the Czech Republic, the overall proportion of men undergoing

MRI before biopsy increased from 28.0% in the first half of 2023 to 38.3% in 2024,

indicating improved alignment of diagnostic pathways.

MRI capacity is a known bottleneck when implementing screening [23]. In the Czech

programme, an initial urology assessment is used to estimate risk and select men for

MRI, in line with European Association of Urology recommendations and the PRAISE-

U project [3] [13]. The impact of this pathway on MRI utilisation and diagnostic

outcomes will be analysed once sufficient data are available. Standardised protocols

are essential for ensuring high-quality MRI scanning and reading, and subsequent

targeted biopsy sampling [23]. The Czech programme applies mandatory MRI

protocols published by the Ministry of Health (Bulletin 15/2023).

Change of coverage and practice

The observed increase in 2-year PSA testing coverage from 47.4% in 2023 to 53.3%

in 2024 among men aged 50–69 years is likely an effect of the introduction of the

organized screening programme. Although non-organized PSA testing still is practiced

in the Czech Republic, the implementation of a systematic screening pathway has

enhanced awareness and accessibility, especially through the engagement of GPs.

The nearly 6 percentage point increase in coverage within a single year suggests

successful early adoption of the programme.

In Lithuania, a primary care–based prostate cancer early detection programme was

initiated in 2006 [20]. Sweden started implementation of regional, population-based

organised testing (OPT) within public healthcare in 2020 [21]. This Swedish approach

substantially increased the proportion of men aged 50 years undergoing PSA testing,

from 14% to 35% [24]. Notably, unlike the Czech and Lithuanian programmes, the

Swedish OPT relies on centralised invitations mailed to entire birth cohorts, without the

involvement of general practitioners.

The planning of the Czech prostate cancer screening programme was facilitated by

networking with the EU-funded PRAISE-U project [6]. PRAISE-U coordinates pilot

projects in Poland, Ireland, Spain, Lithuania, and Estonia that are structured similarly

to the Czech programme. The Italian Society of Urology has also proposed a structured

model for PSA screening with defined roles for GPs and specialists [25]. Collaborative

development of algorithms and shared performance standards are essential for

effective prostate cancer screening in Europe.

Strengths and limitations

The strengths of this report are the programme's nationwide organised design, its full

coverage by public health insurance, and its integration into routine care with

enrolment at GPs' preventive health check-ups. The programme focuses on clearly

defined age groups, enhancing comparability and targeted evaluation.

Limitations include current lack of a national clinical register, restricting follow-up data

such as biopsy results and urological assessments. Only 67% of men with elevated

PSA were seen by a urologist within six months, which is suboptimal. A national

register is under development. While initially observed uptake is promising, further

efforts are needed to optimise informing the men, increase participation in follow-up

steps, and assess its impact on reducing opportunistic PSA testing.

Conclusions

In 2024, a nationwide prostate cancer screening programme was successfully

launched in the Czech Republic. Early results show increased PSA testing coverage

in men aged 50–69 years, particularly in the 50–54 age group, and greater use of pre-

biopsy MRI in line with the diagnostic algorithm. These findings indicate that organised

screening promotes both higher participation and more risk-adapted diagnostic

practice. Recruitment through routine preventive check-ups in primary care, fully

covered by public insurance, has proven feasible and establishes the foundation for

future implementation of the routine programme. Further evaluation will further

elaborate on the compliance of participants and providers with the recommended

screening pathway.

Data access: Hejcmanová, K. (2026). Czech nationwide prostate cancer screening

pilot programme (DATA) [Data set]. Zenodo. https://doi.org/10.5281/zenodo.18151231

Funding/Support and role of the sponsor: The study was supported by Programme

JAC – project SALVAGE (CZ.02.01.01/00/22_008/0004644), financed by MEYS – Co-

funded by the European Union.

The preparatory work for the nationwide pilot programme was supported by the project

„Complex information background for improving the quality of cancer screening

programmes   within   the   National   Screening   Centre"

(CZ.31.8.0/0.0/0.0/23_075/0008430) funded by the European Union from the

Recovery and Resilience Facility through the National Recovery Plan of the Czech

Republic.

Acknowledgements: We would like to acknowledge close collaboration with the

PRAISE-U project, which received funding from the EU4Health programme under

grant agreement 101101217, co-funded by the European Union. Notably, we are

grateful to Hendrik Van Poppel, Sarah Collen, and Monique Roobol-Bouts for the

support of the Prague Prostaforum Declaration and preparatory works of the Czech

programme. We also sincerely thank Eva Nevrtalová for her valuable consultations

during the development of this manuscript and Markéta Vranová for her close

collaboration in the joint coordination and implementation of the national programme

in the Czech Republic.

References

[1] Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, et al.
Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality
Worldwide for 36 Cancers in 185 Countries. CA Cancer J Clin 2021;71:209–49.
https://doi.org/10.3322/caac.21660.

[2] Krejčí D, Mužík J, Šnábl I, Gregor J, Komenda M, Dušek L. Portal of Cancer
Epidemiology in the Czech Republic [online] n.d. https://www.svod.cz.

[3] Van Poppel H, Hogenhout R, Albers P, van den Bergh RCN, Barentsz JO, Roobol
MJ. Early Detection of Prostate Cancer in 2020 and Beyond: Facts and
Recommendations for the European Union and the European Commission. Eur
Urol 2021;79:327–9. https://doi.org/10.1016/j.eururo.2020.12.010.

[4] Roobol MJ, de Vos II, Månsson M, Godtman RA, Talala KM, den Hond E, et al.
European Study of Prostate Cancer Screening - 23-Year Follow-up. N Engl J Med
2025;393:1669–80. https://doi.org/10.1056/NEJMoa2503223.

[5] Arnsrud Godtman R, Holmberg E, Lilja H, Stranne J, Hugosson J. Opportunistic
Testing Versus Organized Prostate-specific Antigen Screening: Outcome After
18 Years in the Göteborg Randomized Population-based Prostate Cancer
Screening Trial. Eur Urol 2015;68:354–60.
https://doi.org/10.1016/j.eururo.2014.12.006.

[6] Chandran A, van Harten M, Singh D, Vilaseca J, Patasius A, Tupikowski K, et al.
Risk-stratified Approach to Implementing Population-based Prostate Cancer

Screening in Five Pilot Sites in the European Union: A Protocol for the PRAISE-U Project. Eur Urol Open Sci 2024;70:8–17. https://doi.org/10.1016/j.euros.2024.09.003.

[7] Vynckier P, Annemans L, Raes S, Amrouch C, Lindgren P, Májek O, et al. Systematic Review on the Cost Effectiveness of Prostate Cancer Screening in Europe. Eur Urol 2024;86:400–8. https://doi.org/10.1016/j.eururo.2024.04.036.

[8] Beyer K, Leenen R, Venderbos LDF, Helleman J, Denijs F, Bramer W, et al. Health Policy for Prostate Cancer Early Detection in the European Union and the Impact of Opportunistic Screening: PRAISE-U Consortium. J Pers Med 2024;14:84. https://doi.org/10.3390/jpm14010084.

[9] Májek O, Babjuk M, Roobol MJ, Bratt O, Van Poppel H, Zachoval R, et al. How to follow the new EU Council recommendation and improve prostate cancer early detection: the Prostaforum 2022 declaration. Eur Urol Open Sci 2023;53:106–8. https://doi.org/10.1016/j.euros.2023.05.011.

[10] Leenen RCA, Venderbos LDF, Helleman J, Gómez Rivas J, Vynckier P, Annemans L, et al. Prostate Cancer Early Detection in the European Union and UK. Eur Urol 2025;87:326–39. https://doi.org/10.1016/j.eururo.2024.07.019.

[11] Koudelková M, Zachoval R, Ferda J, Babjuk M, Matoušková M, Herber O, et al. Czech nationwide prostate cancer screening pilot programme. Prostascreening.cz 2024. https://www.prostascreening.cz/ (accessed August 6, 2025).

[12] Kohestani K, Chilov M, Carlsson SV. Prostate cancer screening-when to start

and how to screen? Transl Androl Urol 2018;7:34–45.

https://doi.org/10.21037/tau.2017.12.25.

[13] Van Poppel H, Hogenhout R, Albers P, van den Bergh RCN, Barentsz JO, Roobol

MJ. A European Model for an Organised Risk-stratified Early Detection

Programme for Prostate Cancer. Eur Urol Oncol 2021;4:731–9.

https://doi.org/10.1016/j.euo.2021.06.006.

[14] Eklund M, Jäderling F, Discacciati A, Bergman M, Annerstedt M, Aly M, et al.

MRI-Targeted or Standard Biopsy in Prostate Cancer Screening. N Engl J Med

2021;385:908–20. https://doi.org/10.1056/nejmoa2100852.

[15] Svensson L, Bratt O, Hugosson J, Stinesen K. Prostate Cancer Screening

Decisions: Which Aspects Do Men Value Most? An Interview Study With Men

Invited to a Population-Based Program. Am J Mens Health

2025;19:15579883251344563. https://doi.org/10.1177/15579883251344563.

[16] Bratt O, Godtman RA, Jiborn T, Wallström J, Akre O, Carlsson S, et al.

Population-based Organised Prostate Cancer Testing: Results from the First

Invitation of 50-year-old Men. Eur Urol 2024;85:207–14.

https://doi.org/10.1016/j.eururo.2023.11.013.

[17] Auvinen A, Tammela TLJ, Mirtti T, Lilja H, Tolonen T, Kenttämies A, et al.

Prostate Cancer Screening With PSA, Kallikrein Panel, and MRI: The ProScreen

Randomized Trial. JAMA 2024;331:1452–9.

https://doi.org/10.1001/jama.2024.3841.

[18] Deantoni EP. Age-Specific Reference Ranges for PSA in the Detection of
Prostate Cancer. Oncology 1997;11:475–85.

[19] Hugosson J, Månsson M, Wallström J, Axcrona U, Carlsson SV, Egevad L, et al.
Prostate Cancer Screening with PSA and MRI Followed by Targeted Biopsy Only.
N Engl J Med 2022;387:2126–37. https://doi.org/10.1056/nejmoa2209454.

[20] Patasius A, Krilaviciute A, Smailyte G. Prostate cancer screening with psa: Ten
years' experience of population based early prostate cancer detection
programme in Lithuania. J Clin Med 2020;9:1–9.
https://doi.org/10.3390/jcm9123826.

[21] Bratt O, Butt ST, Carlsson C, Jelf-Eneqvist L, Gunnarsson O, Ihre A, et al.
Swedish regional population-based organised prostate cancer testing: why, what
and how? Scand J Urol 2025;60:97–104. https://doi.org/10.2340/sju.v60.43809.

[22] Schoots IG, Ahmed HU, Albers P, Asbach P, van den Bergh RCN, Godtman RA,
et al. Magnetic Resonance Imaging–based Biopsy Strategies in Prostate Cancer
Screening: A Systematic Review. Eur Urol 2025;88:247–60.
https://doi.org/10.1016/j.eururo.2025.05.038.

[23] Singh D, Chandran A, Panebianco V, Vilaseca J, Patasius A, Tupikowski K, et al.
MRI capacity assessment for prostate cancer screening in five sites of Europe.
Eur J Radiol 2025;190:112235. https://doi.org/10.1016/j.ejrad.2025.112235.

[24] Järbur E, Holmberg E, Björk-Eriksson T, Bratt O, Arnsrud Godtman R.
Associations between socioeconomic factors and PSA testing in a population-
based organised testing programme and routine healthcare: a register-based

study   of   50-year-old   men.   BMJ   Oncol   2024;3:e000400.

https://doi.org/10.1136/bmjonc-2024-000400.

[25] Ficarra V, Italian Society of Urology (SIU) panel, Bartoletti R, Borghesi M, Caffo

O, De Nunzio C, et al. Organized prostate cancer screening program: a proposal

from the Italian Society of Urology (SIU). Minerva Urol Nephrol 2024;76.

https://doi.org/10.23736/s2724-6051.24.06117-2.

Figures

Figure 1. Screening algorithm: MRI = Magnetic Resonance Imaging; PI-RADS =

Prostate Imaging Reporting and Data System; PSA = Prostate-Specific Antigen; PSAD

= PSA Density; PZ = the peripheral zone; TZ = the transition zone; MDT =

Multidisciplinary Team; *Includes a series of clinical examinations defined by the

programme algorithm (Supplement).

Figure 2. Coverage of the target population by the PSA test in a two-year interval in

men aged 50–69 years. Data source: National Registry of Reimbursed Health

Services; PSA = prostate-specific antigen

Figure 3. Target population coverage by age for a PSA test in the past two years, in

2023 and 2024. Data source: National Registry of Reimbursed Health Services; PSA

= prostate-specific antigen

Tables

Table 1: Age distribution of men approached for the screening programme and

prostate-specific antigen (PSA) results by 5-year age groups. Data source: National

Registry of Reimbursed Health Services, January-December 2024

Age (years)	50–54	55–59	60–64	65–69	Total
Screening offer, N (%)	50,405 (33.5)	38,103 (25.3)	33,124 (22.0)	28,866 (19.2)	150,498 (100.0)
PSA screening results available*, N	48,988	37,033	32,176	27,912	146,109
PSA <1.0 µg/l, N (%)	32,599 (66.5)	21,480 (58.0)	16,137 (50.2)	12,397 (44.4)	82,613 (56.5)
PSA 1.0–2.9 µg/l, N (%)	14,453 (29.5)	12,798 (34.6)	12,234 (38.0)	11,137 (39.9)	50,622 (34.6)
PSA ≥ 3 µg/l, N (%)	1,936 (4.0)	2,755 (7.4)	3,805 (11.8)	4,378 (15.7)	12,874 (8.8)

Screening results with known PSA values are presented. For less than 2% of men,

the result is unknown due to incorrect reporting of data to healthcare payers, and

another 1% chose not to be screened (Supplementary material: Figure 1).

Table 2: Comparison of the proportion of men with magnetic resonance imaging (MRI)

before and after prostate biopsy, the first half of 2023 vs. the first half of 2024. Data

source: National Registry of Reimbursed Health Services

	January-June 2023	January-June 2024
Men with a PSA result, N	227,640	274,887
PSA sampled in primary care, N	119,107	164,606
PSA sampled in urology, N	108,533	110,281
Men with a prostate biopsy*, N (%)	4,309 (1.9)	4,670 (1.7)
PSA sampled in primary care, N (%)	1,113 (0.9)	1,504 (0.9)
PSA sampled in urology, N (%)	3,196 (2.9)	3,166 (2.9)
Men with MRI before** prostate biopsy, N (%)	1,208 (28.0)	1,789 (38.3)
Difference between periods, N (percent points; 95% CI)	10.3 (8.3–12.2)	
PSA sampled in primary care, N (%)	217 (19.5)	464 (30.9)
PSA sampled in urology, N (%)	991 (31.0)	1,325 (41.9)
Men with MRI after*** prostate biopsy, N (%)	360 (8.4)	371 (7.9)
Difference between periods, N (percent points; 95% CI)	0.5 (-1.5–0.7)	
PSA sampled in primary care, N (%)	147 (13.2)	177 (11.8)
PSA sampled in urology, N (%)	213 (6.7)	194 (6.1)

431   ** Biopsy taken within 180 days after PSA test.*

432   *** MRI done between PSA test and prostate biopsy.*

433   **** MRI done within 90 days after prostate biopsy.*

Supplementary information

Detailed description of the programme's algorithm

Enrolment into the programme is facilitated through general practitioners (routine
preventive health check-ups, which are recommended every two years for all
individuals) and through urologists, who invite men already under follow-up for other
urological conditions. In both settings, physicians provide education about importance
of PSA testing, prostate cancer prevention and inform them about next steps of the
programme. In accordance with Czech legislation (Act No. 372/2011), informed
consent is needed but not necessarily in written form.

Information on participation is reported by physicians to health insurance companies
and subsequently included in the National Registry of Reimbursed Health Services
(NRRHS).

If the informed consent is given, a blood sample is collected to determine the serum
PSA value.

Follow-up algorithm is based on PSA levels:

PSA < 1.0 µg/l: Repeat PSA screening in 4 years.

PSA 1.0–2.99 µg/l: Repeat PSA screening in 2 years.

PSA ≥ 3.0 µg/l: Considered a positive screening result – the man is referred for
specialised urological evaluation by a certified urologist.

Certified urologist assessment (PSA ≥ 3.0 µg/l or suspicious digital rectal examination):

Certification is granted by the Czech Urological Society based on predefined criteria.

In men with PSA ≥ 3.0 µg/l, a certified urologist carries out examinations according to

the programme algorithm:

repeat PSA test,

digital rectal examination,

ultrasonographic examination of the kidneys, urinary bladder,

transrectal ultrasound (TRUS) for prostate volume measurement,

calculation of PSA density (PSAD),

estimation of PSA velocity (PSAV) if more than one PSA test is available.

Based on the findings (PSAD ≥ 0.15 or PSAV cut-off 0.75 ng/ml/year), the urologist

may refer for magnetic resonance imaging (MRI), either biparametric (bpMRI, contrast-

free) or multiparametric (mpMRI, with intravenous contrast), at an MRI unit certified by

the Ministry of Health.

In the case of PSA >1 µg/l and a suspicious digital rectal examination, the urologist

can initiate either an MRI or a prostate biopsy.

MRI Evaluation within the programme (PI-RADS — the Prostate Imaging Reporting

and Data System):

PI-RADS 1 and 2 (negative): Further management follows the approved algorithm.

The man is returned to certified urologist for, urological examination with a repeat

MRI scheduled in one year.

PI-RADS 3 (equivocal) in the peripheral zone (PZ): Repeat MRI with intravenous

gadolinium-based contrast enhancement (mpMRI) within two weeks. If then PI-

RADS 2 findings, repeat PSA testing and urological examination after one year. If

PI-RADS 4, targeted biopsy with MRI/ultrasound image fusion.

PI-RADS 3 (equivocal) in the transitional zone (TZ): Repeat PSA test and urological

examinations (TRUS, PSAD, PSAV) after six months; additional mpMRI or prostate

biopsy (high suspicion) if considered indicated.

PI-RADS 4 and 5 (positive): MRI/ultrasound fusion biopsy at a certified biopsy

centre. Certification is granted by the Ministry of Health based on established

criteria.

Men diagnosed with prostate cancer are referred to a multidisciplinary team for further

management. If the biopsy is negative, follow-up MRI is scheduled after one year.

Supplementary Table 1: Proportion of men with PSA ≥ 3 µg/l and a follow-up

examination by a urologist within 6 months by 5-year age group. Data source: National

Registry of Reimbursed Health Services January-June 2024

Age (years)	50–54	55–59	60–64	65–69	Total
Men with PSA ≥ 3 µg/l, N	1,000	1,412	1,945	2,238	6,595
Men with PSA ≥ 3 µg/l examined by urologist, N (%)	699 (69.9)	964 (68.3)	1,317 (67.7)	1,451 (64.8)	4,431 (67.2)

491   Supplementary Figure 1: Participation rate, PSA results in first 12 months, and follow-

492   up examination by a urologist within 6 months. Data source: National Registry of

493   Reimbursed Health Services January-June 2024.

Take-home Message

Organised prostate cancer screening is feasible at the national level. Early results show increasing population coverage and better alignment with evidence-based diagnostic practices. Active involvement of GPs and urologists is crucial for improving screening uptake and adherence to guidelines.