What Do Clinicians Edit in Ambient AI-Drafted Clinical Documentation? A Qualitative Content Analysis

Yawen Guo MISM[1], Di Hu MS[1], Ziqi Yang MS[1], Seungjun Kim MS[1], Brian Tran PhD[1], Jamie Lee[1], Sitha Vallabhaneni MS[1], Rachael Zehrung MS[1], Sairam Sutari MS[2], Steven Tam MD[3], Emilie Chow MD[3], Danielle Perret MD[4], Deepti Pandita MD[3] , Kai Zheng PhD[1]

1. Department of Informatics, University of California, Irvine, Irvine, CA, USA

2. Institute for Clinical and Translational Science, University of California, Irvine, Irvine, CA, USA

3. Department of Medicine, University of California, Irvine, Irvine, CA, USA

4. Department of Physical Medicine & Rehabilitation, University of California, Irvine, Irvine, CA, USA

Corresponding Author: Yawen Guo

Mailing Address: 6136 Donald Bren Hall, Irvine, CA 92697, United States

E-mail: yaweng5@uci.edu

Telephone: (412) 641-0453

Keywords: ambient listening; artificial intelligence; documentation; electronic health records[E05.318.308.940.968.625.500]; qualitative coding; thematic analysis

Objective

Ambient artificial intelligence (AI) documentation is increasingly used to draft notes from clinical conversations, but what clinicians change and how they revise these drafts in routine care

are not well described. This qualitative content analysis characterizes real-world edits to AI drafts and identifies practical targets for tool improvement and implementation.

Materials and Methods

Eight coders analyzed 200 pairs of ambient AI drafts and clinician-final notes at a sentence level, spanning 33 specialties. We developed an inductive coding framework with 11 codes across three categories: clinical content, terminology and wording, and language and style. We then applied thematic analysis to synthesize patterns across the coded edits. Interrater reliability was assessed using Cohen's kappa.

Results

Clinical content modifications were most common, led by procedure, test, and lab order details (39.9%), followed by symptom refinements (30.3%), medication modifications (27.3%), and diagnosis refinements (25.9%). Terminology and wording edits (11.6%) and language and style edits (7.2%) were less frequent. Our thematic analysis identified five recurring themes in how clinicians revised AI drafts, including correcting factual errors, refining generic drafts into specialty-appropriate documentation, calibrating diagnostic certainty to the available evidence, converting transcript-like narrative into professional documentation, and reorganizing or condensing content to improve clarity.

Conclusion and Discussion

Clinicians routinely revise ambient AI drafts to improve accuracy and clinical specificity. Future implementation and model development should emphasize specialty customization and support personalized documentation practices, alongside clinician education that promotes a consistent review routine and appropriate use in daily workflows.

Introduction

Ambient artificial intelligence (AI) systems are increasingly integrated into clinical workflows to support documentation by generating note drafts. [1,2] Early studies suggest they may reduce documentation burden, improve efficiency, and enhance note quality. [3,4] However, despite growing adoption, the real-world reliability and clinical appropriateness of AI-drafted documentation remain underexplored.

Prior evaluations have largely emphasized clinician and patient perspectives through surveys and interviews, focusing on usability, trust, and time savings rather than the content of the generated text. [5–9] A few studies compared AI-generated notes with reference text using automated performance and text-similarity metrics including the F1 score, overlap-based measures such as Recall-Oriented Understudy for Gisting Evaluation (ROUGE) and embedding-based scores. However, these approaches typically produce aggregate, note-level outcomes without explicating what content was changed. [10–12] Expert-based note quality assessments, such as structured reviewer ratings and instruments like the Physician Documentation Quality Instrument, provide an overall quality assessment but are often conducted on simulated or curated data, offering limited insight into the specific edits clinicians make or the reasons behind them. [13]

Limited work has compared real-world ambient AI drafts and final notes committed to electronic health record (EHR) systems to directly characterize editing behavior. Understanding these edits is essential because they show where AI drafts fall short and can reveal systematic gaps with implications for clinical safety, billing, and communication. [14] They can also inform model improvement and implementation support to better align ambient AI with clinical documentation practice.

To address this gap, we conducted a content analysis of ambient AI-generated note sections from 200 paired ambient AI drafts and clinician-final versions, including approximately 1,800 word-level edit operations (insertions, deletions, and replacements) across 33 specialties and 78 clinicians in an ambient AI pilot program. Using qualitative coding and thematic analysis, we examined the types of textual changes clinicians made, the content areas most frequently modified, and the broader themes that describe how clinicians refine AI-generated drafts into finalized documentation. This study offers a fine-grained empirical characterization of real-world clinician editing in ambient AI documentation, with the goal of informing more reliable and more workflow-aligned documentation support systems.

Method

This study employed qualitative coding and thematic analysis to characterize clinician edits to AI-generated clinical note drafts at the content level. The research was conducted as part of a quality improvement initiative at the University of California, Irvine Medical Center (UCI Health), involving clinicians across ambulatory primary and specialty care clinics. They utilized two commercially available ambient AI systems, Nuance DAX and Abridge, which were integrated into the Epic electronic health record (EHR) via mobile and web interfaces. [15,16] The study protocol was approved by the University of California, Irvine Institutional Review Board (IRB #7234). All data processing occurred within HIPAA-compliant secure computing environments, including UCI Health's Protected Virtual Computing Environment (PVCE) and Amazon Web Services (AWS) infrastructure.

Data Source and Sampling Strategy. Ambient AI drafts clinical content by sections, covering history of present illness (HPI), assessment and plan (A&P), physical exam, and results, which

clinicians can selectively insert into EHR notes. The overall corpus of this study consisted of all AI-generated draft note sections and their corresponding clinician-finalized versions recorded between September 26, 2024, and August 5, 2025, from the University of California, Irvine Medical Center. To build an annotation dataset for content analysis representative of the full corpus, we took 200 stratified note samples that contained clinician edits from 33 clinician specialties. Sampling quotas were aligned with the overall corpus distribution, ensuring the inclusion of at least two notes per specialty or one in cases where only a single note was available.

Edit Detection Using the Myers Diff Algorithm. We detected editing activity in the sample by quantifying insertions, deletions, and replacements between each ambient AI draft and the clinician-finalized note section. For each draft–final pair, we applied a word-level Myers diff algorithm to align tokens and identify edit operations, recording the span and sequence of each change. [17] We also generated side-by-side HTML views with highlighted edits, allowing for direct comparison of the AI output against clinician revisions. Figure 1 provides a synthetic example for illustration only and does not contain patient data.

Figure 1. Synthetic Example of Side-by-Side Visualization of Clinician Edits to an AI-Generated Clinical Note

Sentence-Level Unit of Analysis for Content Analysis. We defined an edit unit as the smallest contiguous set of sentences that fully captured each diff-detected edit span, including adjacent or overlapping edits, to preserve clinical context and meaning. To align draft and final text at the sentence level, we used a dynamic approach that allowed flexible mappings, including one-to-one, one-to-many (split), many-to-one (merge), and many-to-many alignments. Each edit unit was labeled by its alignment pattern and quantified by the number of insertions, deletions, and replacements it contained, capturing both within-sentence phrasing changes and larger structural revisions.

Coding Framework and Procedure. An inductive coding framework was developed to characterize the types and functions of clinician edits. [18] Two researchers (YG, DH) independently reviewed a subset of edit units to develop an initial codebook. The codebook was then iteratively refined through consensus meetings and multiple rounds of coding with eight coders, all with experiences in health informatics research and qualitative analysis. The full coding team then conducted an initial calibration round in which all eight coders applied the draft codebook to the same set of edit units; discrepancies were resolved through adjudication meetings and email discussion, and the codebook was iteratively refined. Interrater reliability was assessed using Cohen's κ across major code categories. [19] We finalized the codebook after achieving acceptable interrater agreement across coders (Cohen's kappa). The final refined framework consisted of three top-level code categories capturing distinct categories of editing behavior (entity-level clinical content changes, terminology and wording adjustments, or

language and style modifications), along with 11 subcodes representing more granular

phenomena. [20,21]

Using the finalized codebook, coders applied labels to each sentence-level edit unit. We treated

each code as a binary label, present or absent, for each sentence-level edit unit, allowing for

multiple codes per unit. Across eight coders, we computed Cohen's kappa for each code for each

annotator pair and macro-averaged kappas across codes with equal weight; overall reliability was

the mean of the 28 pairwise macro-averaged kappa values. All coding was conducted in

structured spreadsheets using the finalized codebook. To maintain independent coding, each

coder worked in a separate spreadsheet. For reference when needed, coders could consult

color-coded HTML diff visualizations of the paired draft and final text to view each edit unit

alongside the exact textual changes within the full note section.

Table 1. Qualitative coding schema

Primary Code Category	Code	Example(s)	Description
Entity-level: Clinical Content	Medication (E–Med)	Before: "Keppra one pump BID"; After: "Keppra 500 mg BID"	Edits that add, remove, or modify medication information, including medication name, dose, frequency, route, formulation, timing, or adherence details.
	Symptom (E–Sym)	Before: "Cramping"; After: "Cramping at night"	Edits that add, remove, or modify symptom mentions, including presence/absence, severity, location, timing, duration, triggers, or associated features.
	Diagnosis (E–Dx)	Before: "epilepsy"; After: "refractory focal epilepsy"	Edits that add, remove, or modify diagnoses/conditions or diagnostic labels, including specificity (type/subtype), qualifiers, or

			diagnostic framing.
	Procedure / test / lab order (E–Test)	Before: "EEG not done"; After: *(deleted)*	Edits that add, remove, or modify diagnostic tests, labs, imaging, procedures, referrals, or orders (including whether performed vs planned).
	Demographics / ID (E–ID)	Before: "[WRONG-SPELLED NAME]"; After: "[CORRECTED NAME]"	Edits that add, remove, or modify patient identifiers or demographic details (e.g., age, name, pronouns, PHI placeholders)
	Social context (E–Soc)	Before: *(none)*; After: "lives with spouse"	Edits that add, remove, or modify social history/context (e.g., tobacco/alcohol/substance use, living situation, education, occupation, support system, marital status, habits/behaviors) [22]
	Past medical history (E-Hx)	Before: (none); After: "Previously saw nephrologist at [LOCATION]"	Edits that add, remove, or modify past medical history, including comorbidities, prior events, prior specialty care, surgeries, and relevant historical conditions.
Terminology and Wording	Lay–jargon substitution (T–Lay)	Before: "pain"; After: "myalgia"	Edits that substitute clinical jargon with lay language (or vice versa) while preserving the underlying meaning.
	Abbreviation Expansion/Contraction (T–Abbr)	Before: "right arm"; After: "R arm"	Edits that expand, introduce, or standardize abbreviations and acronyms.

Language Style	Empathy / rapport / politeness (S–Emp)	Before: *(none)*; After: "It was a pleasure to see Ms. [NAME]"	Edits that change interpersonal tone (rapport, empathy, politeness, patient-centered phrasing) without changing core clinical content.
	Certainty / hedging (S–Cert)	Before: "due to"; After: "likely due to"	Edits that increase or decrease expressed certainty, probability, attribution, or diagnostic commitment (e.g., hedging, qualifiers, speculative language).

Thematic Analysis. After all edit units were coded using the finalized codebook, we conducted a thematic synthesis to identify higher-level patterns in clinician revision behavior. Two lead researchers (YG, DH) independently reviewed code distributions, representative edit units, and analytic memos to examine how frequently co-occurring codes reflected shared underlying revision behavioral patterns. Codes were iteratively grouped based on conceptual similarity and the clinical intent they represented (e.g., factual correction, clinical precision, documentation standardization). Through this process, we aggregated code-level edits into five higher-level themes that capture recurring clinician revision patterns in AI-drafted notes. Themes were refined through iterative team discussion and disagreements in theme boundaries were resolved through consensus. The resulting themes represent interpretive syntheses of multiple code types and are intended to characterize how clinicians systematically revise AI-drafted documentation.

Results

Dataset Characteristics

A total of 200 note pairs containing ambient AI drafted sections and finalized versions were included in the annotated sample, representing 78 clinicians, 200 patients, and 33 clinical specialties (Table 2). Clinicians could generate more than one ambient AI section draft within the same note (for example, only A&P, or both HPI and A&P). We coded section-level draft–final pairs, grouped by note ID for tracking. The complete sampled notes for qualitative annotation included a history of present illness section (n = 149) and an assessment and plan section (n = 135), while fewer included physical examination (n = 18) or results (n = 12). Across all edit units, clinicians made 1,804 token-level modifications, including 556 insertions, 393 deletions, and 855 replacements. The sample covered 33 specialties; family practice contributed the most notes (n = 54), followed by rheumatology (n = 22) and internal medicine (n = 20), with the remaining notes distributed across a broad set of lower-volume specialties (Table 2).

Table 2. Characteristics for paired ambient AI drafts and clinician-finalized notes

Category	Subcategory	Value
Overall (sample-level)	Study period	2024-09-26 to 2025-08-05
	# of notes	200
	# of clinicians	78
	# of patients	200
AI- drafted Note Section Pairs	# of History of Present Illness (HPI) pairs	149
	# of Assessment & Plan (A&P) pairs	135
	# of Physical Exam pairs	18
	# of Results pairs	12
Token-level editing activity	# of Insertions	556
	# of Deletions	393
	# of Replacements	855
Specialty Distribution (n = # of notes, % of 200 notes)	Family Practice (n=54, 27.0%), Rheumatology (n=22, 11.0%), Internal Medicine (n=20, 10.0%), Orthopaedic Surgery (n=15, 7.5%), Geriatric Medicine (n=15, 7.5%), Hand Surgery (n=10, 5.0%), Neurology (n=7, 3.5%), Nurse Practitioner (n=6, 3.0%), Gastroenterology (n=3, 1.5%), Ob/Gyn (n=3, 1.5%), Physician Assistant (n=3, 1.5%), Psychiatry (n=2, 1.0%), Urology (n=2, 1.0%), Physical Medicine & Rehab (n=2, 1.0%), Otolaryngology (n=2, 1.0%), Vascular Neurology (n=2, 1.0%), Optometry (n=2, 1.0%), Ophthalmology (n=2, 1.0%), Cardiology (n=2, 1.0%), Neurosurgery (n=2, 1.0%), General Surgery (n=2, 1.0%), Colon/Rectal Surgery (n=2, 1.0%), Integrative Medicine (n=2, 1.0%), Infectious Diseases (n=2, 1.0%), Hospitalist (n=2, 1.0%), Hematology/Oncology (n=2, 1.0%), Head & Neck Surgery (n=2, 1.0%), Endocrinology (n=2, 1.0%), Emergency Medicine (n=2, 1.0%), Dermatology (n=2, 1.0%), Orthopedics (n=1, 0.5%), Vascular Surgery (n=1, 0.5%).	

Editing activity varied by note section (Table 3). HPI and A&P contained the highest editing

volume, averaging approximately six edits per section (HPI: 6.07; A&P: 6.16), driven primarily

by insertions and replacements. Physical Exam and Results sections exhibited substantially fewer modifications, averaging fewer than three and two edits per section, respectively.

Table 3. Average token-level edit counts by note section

Note Section	Avg # of Inserts	Avg # of Deletes	Avg # of Replaces	Avg # of Total Edits
A&P (Assessment & Plan)	2.01	1.22	2.93	6.16
HPI (History of Present Illness)	1.68	1.46	2.93	6.07
Physical Exam	1.56	0.28	1.11	2.94
Results	0.5	0.5	0.25	1.25

Distribution of Qualitative Edit Codes

We coded 714 sentence-level edit units extracted from 200 AI draft–final note pairs. After an initial training phase and two rounds of independent coding with consensus discussions, interrater reliability across eight coders reached a moderate level of agreement (overall Cohen's $\kappa = 0.568$). Because a single edit unit could receive multiple codes, results are reported as the frequency of edit units containing each code; percentages therefore do not sum to 100 percent.

The most common edit types involved clinical content. Procedures, tests, and lab orders edits (E–Test) appeared in 285 edit units (39.9%). A common pattern was clarifying order-specific metadata, including the test type, the intended location or service, and the timing or status of the order when that detail affected interpretability. This was followed by symptom changes (E–Sym) in 216 (30.3%). For symptom edits, clinicians commonly refined the symptom description to improve clinical specificity. This included adjusting the symptom label, and specifying severity

or pattern when the draft was vague. Clinicians also removed symptom descriptions that were not part of the patient's current presentation or that appeared to be carried over from prior context without supporting evidence from the encounter. Medication modifications (E–Med) appeared in 195 (27.3%), and edits most often made the regimen actionable by adding or correcting dose, frequency, route, or administration details. Clinicians also standardized medication nomenclature, such as correcting the drug name, aligning the term with common clinical usage, or clarifying which formulation was intended. A smaller but recurring pattern was medication reconciliation, where clinicians removed a medication that was not relevant to the current plan or clarified whether the patient was currently taking it. Diagnosis refinements (E–Dx) appeared in 185 (25.9%). Clinicians frequently revise the diagnostic framing by narrowing a broad label to a more specific condition, aligning the diagnosis term with the symptom description, or clarifying whether the diagnosis was established versus under evaluation. Edits involving past medical history (18.9%) and social/context information (17.8%) were also frequent, whereas terminology/wording edits such as jargon↔lay (7.1%) and abbreviation changes (4.5%) and language/style edits such as certainty/hedging (4.1%) and empathy/rapport (3.1%) were less common.

Thematic Synthesis of Clinician Edits

Representative before/after examples are shown in Table 4.

Theme 1- Clinicians correct factual errors in AI drafts.

These edits corrected misspelled names and pronouns introduced during transcription and fixed inaccurate time-stamp information. They also refined event descriptions by updating prior medical history details, such as changing a hospitalization to a more accurate emergency department visit, and by correcting anatomic and severity information (e.g., laterality, joint

location, stenosis severity). Clinicians further corrected medication and test/procedure details, including drug names and dosing frequency, sedation type, and imaging modality. These edits improved the accuracy of the clinical note in aspects, including medical history, demographics, and treatments.

Theme 2- Clinicians refine generic drafts into specialty-appropriate documentation.

Clinicians added clinical details commonly required for specialty documentation, such as symptom onset and trajectory, medication regimen specifics (dose, schedule, and whether use was as needed), and test or treatment context that shaped the plan. In neurology/epilepsy, edits emphasized temporal progression and neuroanatomical localization, reframing brief summaries into problem-oriented assessments that track symptom course, differentiate neurologic problems, and document medication intervals or titration. For example, a neurology progress note was expanded by adding detailed trauma history, symptom characterization across neurologic problems, and rewriting the assessment into a specialized, multi-layered plan. In procedure-oriented specialties such as hand surgery and orthopaedics, edits replaced general descriptions with structured, objective findings and procedure- or imaging-specific terminology, making decisions explicit through digit-level exam metrics or dated radiology findings. A hand surgery note was refined by replacing general symptom and exam language with digit-specific, quantified findings (e.g., finger-level localization, monofilament scores) and by stating clear procedural intent, such as proceeding with right index and ring trigger finger release. In cardiology, clinicians expanded symptom documentation beyond brief general phrasing by specifying exertional context and exercise tolerance, linking dyspnea, chest symptoms, or palpitations to what the patient can do and under what conditions, so the note reflected cardiovascular-style functional assessment rather than a generic symptom checklist.

Theme 3- Clinicians revise overly certain language to match the available evidence with clinical wording.

Across notes, clinicians often replaced definitive causal or diagnostic wording with more appropriately qualified language when the documentation did not support a firm conclusion, such as changing "due to" to "likely due to" and revising "no retinal or optic nerve disease" to "no obvious retinal or optic nerve disease." They also clarified when a condition was being evaluated rather than established, for example, by reframing plans aiming to "definitively rule out or confirm AFib" as monitoring in the setting of "no confirmed diagnosis of AFib." Similarly, diagnostic labels were adjusted to match test evidence, such as shifting a stated dermatomyositis diagnosis to a suspected drug eruption after a negative myositis panel.

Theme 4- Clinicians normalize conversational narratives into standardized chart language.

Clinicians frequently edited transcript-like language to align the note with clinician-facing documentation standards by removing or reframing patient-attributed causal explanations and speculative risk phrasing, and by replacing them with objective, clinically interpretable statements. For example, subjective clauses such as "which she believes are related to her thyroid condition" or "she recalls an instance when…" were removed or rewritten as verifiable clinical information, such as the current levothyroxine dose or the patient's expressed concern about thyroid level fluctuations. More broadly, conversational and vague phrasing was rewritten into conventional chart language and medical terminology, such as changing "she's doing okay" to "patient reports stable condition" and replacing lay descriptions like "a disintegrating disc" with "a bulging disc".

Edits also standardized abbreviations and shorthand, for example, using "MCV" and inserting routine clinical abbreviations such as "EGD", "abnl", and "CIN II". Clinicians increased

technical specificity by naming the exact device, procedure, or structure, such as revising "partial amputation" to "ray amputation," and specifying the ECU tendon. Medication references were similarly clarified by replacing generic wording with the specific drug name, such as "used the medication" to "used Qsymia" and "Lexapro" to "escitalopram," with accompanying refinements to dosing or decision language.

Theme 5-Clinicians reorganize, relocate, and condense AI drafts.

Clinicians frequently restructure AI-generated drafts to make the note easier to navigate and use, reorganizing content to improve scanability and clarity of clinical plans while trimming material that did not match the visit. Dense narrative was often converted into structured formats, for example, with long paragraphs rewritten as bullet lists, problem narratives turned into numbered plans, and free text reorganized under problem-oriented headings. Clinicians also relocated information to more appropriate sections, such as moving social or surgical history into labeled past history fields and separating imaging descriptions into a dedicated radiology-style section with dated, objective findings. Edits also reflected the process of relevance triage where duplicative or boilerplate text was consolidated, and content that appeared misaligned with the clinical context was removed, including passages on specialty-irrelevant medication management.

Table 4. Representative examples of clinician edits to ambient AI–drafted notes by five qualitative themes.

Theme 1- Clinicians correct factual errors in AI drafts		
Edit type	**AI draft text** [a]	**Clinician-final text** [a]

Demographic correction	"Seen by [PROVIDER_NAME_A]."	"Seen by [PROVIDER_NAME_B].
Demographic correction	"The patient is [AGE_A] years old."	"The patient is [AGE_B] years old."
Temporal correction	"...in March"	"...in next March"
Tense correction	"The patient has been taking..."	"The patient had been taking..."
Pronoun correction	"...he..."	"...she..."
Clinical event type correction	"a recent hospitalization"	"a recent ER visit on 12/15/24"
Medication detail correction	"has been using fluconazole 200 mg weekly"	"has been using fluconazole 200 mg a few times per week"
Medication name correction	"Zepal"	"albuterol"
Laterality correction	"Cerumen impaction… more on the left"	"Cerumen impaction… more on the right"
Anatomic site correction	"MCP joint"	"DIP joint"
Procedure/test label correction	"colonoscopy"	"sigmoidoscopy"
Test detail correction	"MRI of brain"	"MRA of head and neck"
Device name correction	"a cardio device"	"a Kardia device"
Followup detail correction	"return in 6 week"	"return in 2–3 Month"

Theme 2- Clinicians refine generic drafts into specialty-appropriate documentation

Edit type	AI draft text	Clinician-final text
Diagnosis specificity	"headache"	"possibly migraine with aura"

Diagnosis specificity	"hypoglycemia and hyperglycemia"	"hyperglycemia and rare hypoglycemia"
Diagnosis specificity	"Psoriatic Arthritis…"	"Rheumatoid arthritis, double seropositive… high titer anti-CCP"
Symptom specificity	"noise sensitivity"	"light and sound sensitivity"
Symptom specificity	"pain in back and neck"	"neck pain without acute changes"
Symptom specificity	"Achilles"	"Calf pain"
Temporal course specification	"headaches"	"persistent daily headaches since November 2024"
Medication regimen specification	"Vitamin B2"	"Vitamin B2 400 mg per day"
Medication regimen specification	"Prescribe meloxicam once daily…"	"Restart meloxicam 15 mg daily with dinner."

Theme 3- Clinicians revise overly certain language to match the available evidence with clinical wording

Edit type	AI draft text	Clinician-final text
Certainty calibrated to evidence	"due to"	"likely due to"
Certainty calibrated to evidence	"No retinal or optic nerve disease."	"No obvious retinal or optic nerve disease."
Certainty calibrated to evidence	"to definitively rule out or confirm AFib"	"as there is currently no confirmed diagnosis of AFib."

Theme 4- Clinicians convert transcript-like narrative into professional documentation

Edit type	AI draft text	Clinician-final text

Clinical register normalization	"She's doing okay"	"Patient reports stable condition"
Medical terminology standardization	"a disintegrating disc"	"a bulging disc"
Clinician framing / documentation precision	"He is on HIV medications."	"He is compliant with HIV medications."
Abbreviation standardization	"mean corpuscular volume"	"MCV"
Technical specificity increase	"potential removal of the stones"	"potential removal of the dropped gall stones in hepatorenal fossa"
Procedure terminology standardization	"partial amputation"	"ray amputation"
Generic to specific entity	"used the medication"	"used Qsymia"
Medication name normalization	"Lexapro"	"escitalopram"
Patient narrative reframed into clinician documentation	Subjective self-attributions ("which she believes are related…", "she recalls…")	Reframed into objective clinician documentation

Theme 5- Clinicians reorganize or condense content to improve clarity [b]

Edit type	AI draft structure	Clinician-final structure
Format conversion	Long paragraph	Bullet list (e.g., "• Continue healthy diet…")
Format conversion	Problem paragraph (prose)	Numbered list (1–7)
Format conversion	Prose narrative	Structured "#condition:" headings
Section relocation	Social + surgical history embedded mid-paragraph	Moved under labeled "past medical history"

		Deleted and/or reorganized; "psychiatric history" section
Relevance trimming	Large chunks of past history	added/structured

4. Discussion

Clinicians substantially modify AI-generated drafts, and those edits follow repeated patterns that align with the five themes in our qualitative analysis, which point to specific opportunities for improvement in ambient AI documentation tools and their implementation.

Clinicians often correct inaccuracies or hallucinations, especially in discrete details that can change clinical meaning, such as names, medication dosages, and test or procedure attributes. Even when the overall narrative in AI drafts was broadly accurate, these errors are difficult for downstream readers to detect and can propagate if copied forward. This finding suggests that improving factual reliability of ambient AI tools should be a top priority, paired with workflow features that help clinicians quickly verify the fields that most often need to be double-checked. [23] At present, most ambient AI tools do not directly access the patient's chart, so certain errors, such as names or other identifiers, cannot be automatically cross-checked even when the correct information already exists in the EHR. If chart integration becomes feasible, many of these errors could be corrected upstream. In the meantime, for institutions integrating ambient AI into routine practice, training for clinicians should go beyond a generic requirement to "review the draft" and instead teach a consistent verification routine that prioritizes the most error-prone details. [24,25]

Clinicians tailoring AI drafts to be more specialty-specific suggest that a single, generic note structure and style is often insufficient for specialty documentation needs. An actionable response is to support specialty-tuned drafting behavior through templates and prompts that reliably elicit the details that matter for a given discipline, rather than broadly relying on the subjective–objective–assessment–plan (SOAP) structured format. [26–28] For institutions, this implies that ambient AI adoption should be paired with specialty-level configuration and evaluation aligned with each specialty's documentation norms and clinical priorities. [29]

Overly certain language in ambient AI drafts can outpace the available clinical evidence, creating extra work for clinicians and increasing the risk of carrying forward unverified conclusions. Beyond clinical risk, this pattern may raise medicolegal concerns if AI-generated phrasing systematically nudges notes toward stronger diagnostic commitments than the encounter supports. [30] In our data, clinicians spent substantial effort dialing back overly certain AI language. Actionably, AI developers can in response design models that default to appropriately qualified language and conventional diagnostic framing, and that avoid stating conclusions more strongly than the note supports. [31] At the practice level, this theme points to a clear focus for governance and education, reinforcing a consistent review habit around diagnostic uncertainty and ensuring the system's outputs do not nudge documentation toward over-commitment when diagnoses are not yet established.

Clinicians often rewrote transcript-like text into a usable clinical record, shifting conversational phrasing into objective, clinically interpretable documentation. This mismatch may be amplified by how the system structures conversational content into the Subjective section (SOAP). [26] In our data, clinicians frequently condensed and reorganized that section, suggesting potential misalignment between how the draft is assembled and how clinicians typically write and review

notes. Ambient AI developers can strengthen attribution-aware writing that keeps a clear boundary between patient reports and clinician assessment without adding interpretation that the note does not support. Terminology normalization can also help increase precision without causing note bloat. [32–35] For clinicians, this theme is a reminder that review is not only error correction; it is also ensuring the note reads as a professional record that others can quickly interpret and act on.

Recurring patterns of note restructuring and condensing suggest that draft organization is a major determinant of whether ambient notes are usable at the point of care. For example, ambient capture can include social history and social determinants of health (SDOH) information that is clinically relevant because it can shape access to care, home and community support, and adherence. However, in the context of certain encounters, some of these elements may be secondary to the immediate diagnostic and clinical decision making, particularly when captured with high granularity (e.g., transportation needs narratives). [36] Clinicians often shortened or removed these details when they did not inform the assessment or plan. These edits reflect a familiar tradeoff in real-world documentation. As notes get longer, they become harder to review efficiently, and important information can be harder to find. [37,38] An actionable implication is to prioritize tool behaviors that produce more structured drafts, route content into predictable sections, and give clinicians simple ways to control verbosity without losing clinically important context. At the organizational level, this supports setting expectations for note structure and routinely monitoring note length and usability as part of ambient AI governance.

This study has several limitations. Our qualitative analysis drew on 200 AI draft–final note pairs, and some specialties were represented by only a small number of cases, so we may not fully capture specialty-specific nuances. The dataset also comes from a single health system and a

single deployment context, which may limit how well these patterns generalize to other institutions, care settings, or ambient AI products. We analyzed text changes only. Without the source audio, user interface cues, or clinicians' stated intent, we cannot determine why a given edit was made or whether it reflects individual style, local documentation norms, or risk tolerance. Interrater agreement was moderate (Cohen's kappa = 0.568). This is consistent with a multi-coder, multi-label codebook, but it suggests some overlap at code boundaries. [39] Finally, we did not assess the clinical correctness of the finalized notes or downstream impacts such as care outcomes, time burden, or corresponding change of billing codes.

Taken together, our findings translate routine clinician edits into concrete actionable insights for both tool improvement and real-world implementation. The patterns we observed suggest that ambient AI should be evaluated and tuned for specialty needs and clinician-to-clinician variation. Future work should incorporate targeted semi-structured interviews to clarify the rationales and workflow factors behind common edits and to validate our interpretation of these patterns. Future work should also investigate whether specific edit patterns are associated with continuity of care. For example, edits that clarify the problem list or tighten the assessment and plan may make notes more usable for downstream readers. Studies should also examine operational performance, including turnaround time, downstream review effort or follow-up work, and whether term-level changes affect billing or coding outcomes. Finally, future work should evaluate safety-critical signals, especially the correction of factual inaccuracies and the moderation of overly certain language when diagnoses remain uncertain.

5. Conclusion

Our qualitative content analysis of clinician edits to AI drafts provides a comprehensive view of what clinicians changed and where improvement is needed. Clinicians corrected mistakes,

replaced overly generic statements with specialty-appropriate detail, and tempered overly certain language to better match the available evidence using standard clinical wording. They also rewrote conversational, patient-attributed narrative into professional documentation and reorganized drafts into clearer, more structured notes. Ambient AI tools should be designed around how clinicians document in different specialties and should avoid wording that goes beyond the evidence in the encounter. Health systems can use recurring edit patterns to identify target implementation support and training that improve day-to-day documentation efficiency.

ACKNOWLEDGEMENT

This study did not receive any external funding.

COMPETING INTERESTS STATEMENT

The authors have no competing interests to declare.

CONTRIBUTORSHIP STATEMENT

Yawen Guo (YG) led the study, contributing to the study concept and design, data acquisition, data analysis and interpretation, and drafting of the manuscript. Di Hu (DH) contributed to notebook development and qualitative coding. Ziqi Yang (ZY), Brian Tran (BT), Jamie Lee (JL), Sitha Vallabhaneni (SV), Seungjun Kim (SK), and Rachael Zehrung (RZ) contributed to qualitative coding. Sairam Sutari (SS) provided data curation and administrative support. Steven Tam (ST), Emilie Chow (EC), Danielle Perret (DPe), and Deepti Pandita (DPa) provided clinical perspective and contributed to interpretation of findings. Kai Zheng (KZ) provided overall guidance and supervision throughout the project. All authors critically reviewed and revised the manuscript for important intellectual content and approved the final version.

DATA AVAILABILITY STATEMENT

The underlying dataset contains protected health information (PHI) and is not publicly available.

Reference

[1] Hundal J, Jain M, McCollom J. Ambient Artificial Intelligence in Health Care Documentation: A Review of Tools, Integration, and Clinical Implications. AI Precis Oncol 2025.

[2] Kanaparthy NS, Villuendas-Rey Y, Bakare T, et al. Real-World Evidence Synthesis of Digital Scribes Using Ambient Listening and Generative Artificial Intelligence for Clinician Documentation Workflows: Rapid Review. JMIR AI 2025;4:e76743.

[3] Tierney AA, Gayre G, Hoberman B, et al. Ambient Artificial Intelligence Scribes to Alleviate the Burden of Clinical Documentation. NEJM Catal 2024;5. https://doi.org/10.1056/CAT.23.0404.

[4] Tierney AA, Gayre G, Hoberman B, et al. Ambient Artificial Intelligence Scribes: Learnings after 1 Year and over 2.5 Million Uses. NEJM Catal 2025;6. https://doi.org/10.1056/CAT.25.0040.

[5] Guo Y, Hu D, Wang J, et al. Ambient Listening in Clinical Practice: Evaluating EPIC Signal Data Before and After Implementation and Its Impact on Physician Workload 2025. https://doi.org/10.48550/arXiv.2504.13879.

[6] Guo Y, Wang J, Hu D, et al. Evaluating ambient artificial intelligence documentation: effects on work efficiency, documentation burden, and patient-centered care. J Am Med Inform Assoc 2025:ocaf180.

[7] Albrecht M, Shanks D, Shah T, et al. Enhancing clinical documentation with ambient artificial intelligence: a quality improvement survey assessing clinician perspectives on work burden, burnout, and job satisfaction. JAMIA Open 2024;8:ooaf013. https://doi.org/10.1093/jamiaopen/ooaf013.

[8] Nguyen OT, Turner K, Charles D, et al. Implementing Digital Scribes to Reduce Electronic Health Record Documentation Burden Among Cancer Care Clinicians: A Mixed-Methods Pilot Study. JCO Clin Cancer Inform 2023:e2200166. https://doi.org/10.1200/CCI.22.00166.

[9] Haberle T, Cleveland C, Snow GL, et al. The impact of nuance DAX ambient listening AI documentation: a cohort study. J Am Med Inform Assoc JAMIA 2024;31:975–9. https://doi.org/10.1093/jamia/ocae022.

[10] Jorg T, Kämpgen B, Feiler D, et al. Efficient structured reporting in radiology using an intelligent dialogue system based on speech recognition and natural language processing. Insights Imaging 2023;14:47. https://doi.org/10.1186/s13244-023-01392-y.

[11] Yim W, Fu Y, Ben Abacha A, et al. Aci-bench: a novel ambient clinical intelligence dataset for benchmarking automatic visit note generation. Sci Data 2023;10:586.

[12] Sezgin E, Sirrianni JW, Kranz K. Evaluation of a digital scribe: conversation summarization for emergency department consultation calls. Appl Clin Inform 2024;15:600–11.

[13] Siddiqui S, Etingen B, Patrianakos J, et al. Use of the Veterans Health Administration online patient portal among Veterans with spinal cord injuries and disorders. J Spinal Cord Med 2023;46:917–28. https://doi.org/10.1080/10790268.2022.2084967.

[14] Suhail D, Wong ZY, Ubhi J, et al. Current evidence and future directions of metrics used to evaluate ambient clinical documentation: A scoping review. Int J Med Inf 2026;205:106113. https://doi.org/10.1016/j.ijmedinf.2025.106113.

[15] Generative AI for Clinical Conversations | Abridge n.d. https://www.abridge.com/ (accessed May 5, 2025).

[16] Microsoft Dragon Copilot | Microsoft Cloud for Healthcare n.d. https://www.microsoft.com/en-us/health-solutions/clinical-workflow/dragon-copilot (accessed May 5, 2025).

[17] The Myers Difference Algorithm n.d. https://www.nathaniel.ai/myers-diff/ (accessed December 13, 2025).

[18] Fereday J, Muir-Cochrane E. Demonstrating Rigor Using Thematic Analysis: A Hybrid Approach of Inductive and Deductive Coding and Theme Development. Int J Qual Methods 2006;5:80–92. https://doi.org/10.1177/160940690600500107.

[19] McHugh ML. Interrater reliability: the kappa statistic. Biochem Medica 2012;22:276–82.

[20] Colicchio TK, Dissanayake PI, Cimino JJ. The anatomy of clinical documentation: an assessment and classification of narrative note sections format and content. AMIA Annu Symp Proc AMIA Symp 2020;2020:319–28.

[21] Campbell JR, Carpenter P, Sneiderman C, et al. Phase II Evaluation of Clinical Coding Schemes: Completeness, Taxonomy, Mapping, Definitions, and Clarity. J Am Med Inform Assoc 1997;4:238–51. https://doi.org/10.1136/jamia.1997.0040238.

[22] Navarro V. What We Mean by Social Determinants of Health. Int J Health Serv 2009;39:423–41. https://doi.org/10.2190/HS.39.3.a.

[23] Magrabi F, Ammenwerth E, McNair JB, et al. Artificial Intelligence in Clinical Decision Support: Challenges for Evaluating AI and Practical Implications: A Position Paper from the IMIA Technology Assessment & Quality Development in Health Informatics Working Group and the EFMI Working Group for Assessment of Health Information Systems. Yearb Med Inform 2019;28:128–34. https://doi.org/10.1055/s-0039-1677903.

[24] Varghese J. Artificial Intelligence in Medicine: Chances and Challenges for Wide Clinical Adoption. Visc Med 2020;36:443–9. https://doi.org/10.1159/000511930.

[25] Schubert T, Oosterlinck T, Stevens RD, et al. AI education for clinicians. eClinicalMedicine 2025;79:102968. https://doi.org/10.1016/j.eclinm.2024.102968.

[26] Cameron S, Turtle‑Song I. Learning to Write Case Notes Using the SOAP Format. J Couns Dev 2002;80:286–92. https://doi.org/10.1002/j.1556-6678.2002.tb00193.x.

[27] Pearce PF, Ferguson LA, George GS, et al. The essential SOAP note in an EHR age. Nurse Pract 2016;41:29–36. https://doi.org/10.1097/01.NPR.0000476377.35114.d7.

[28] Seo J-H, Kong H-H, Im S-J, et al. A pilot study on the evaluation of medical student documentation: assessment of SOAP notes. Korean J Med Educ 2016;28:237–41. https://doi.org/10.3946/kjme.2016.26.

[29] Acampora G, Cook DJ, Rashidi P, et al. A Survey on Ambient Intelligence in Healthcare. Proc IEEE 2013;101:2470–94. https://doi.org/10.1109/JPROC.2013.2262913.

[30] McCradden MD, Stedman I. Explaining decisions without explainability? Artificial intelligence and medicolegal accountability. Future Healthc J 2024;11:100171. https://doi.org/10.1016/j.fhj.2024.100171.

[31] Hatherley JJ. Limits of trust in medical AI. J Med Ethics 2020;46:478–81. https://doi.org/10.1136/medethics-2019-105935.

[32] Kahn D, Stewart E, Duncan M, et al. A Prescription for Note Bloat: An Effective Progress Note Template. J Hosp Med 2018;13:378–82. https://doi.org/10.12788/jhm.2898.

[33] Liu J, Capurro D, Nguyen A, et al. "Note Bloat" impacts deep learning-based NLP models for clinical prediction tasks. J Biomed Inform 2022;133:104149. https://doi.org/10.1016/j.jbi.2022.104149.

[34] Lowry LZ, Ramaiah M, Prettyman SS, et al. Examining the Copy and Paste Function in the Use of Electronic Health Records. Gaithersburg, MD: National Institute of Standards and Technology; 2017. https://doi.org/10.6028/NIST.IR.8166.

[35] Vawdrey DK, Cauthorn C, Francis D, et al. A Practical Approach for Monitoring the Use of Copy-Paste in Clinical Notes. AMIA Annu Symp Proc AMIA Symp 2021;2021:1178–85.

[36] Wang M, Pantell MS, Gottlieb LM, et al. Documentation and review of social determinants of health data in the EHR: measures and associated insights. J Am Med Inform Assoc 2021;28:2608–16. https://doi.org/10.1093/jamia/ocab194.

[37] Baumann LA, Baker J, Elshaug AG. The impact of electronic health record systems on clinical documentation times: A systematic review. Health Policy 2018;122:827–36. https://doi.org/10.1016/j.healthpol.2018.05.014.

[38] Hultman GM, Marquard JL, Lindemann E, et al. Challenges and Opportunities to Improve the Clinician Experience Reviewing Electronic Progress Notes. Appl Clin Inform 2019;10:446–53. https://doi.org/10.1055/s-0039-1692164.

[39] Cole R. Inter-Rater Reliability Methods in Qualitative Case Study Research. Sociol Methods Res 2024;53:1944–75. https://doi.org/10.1177/00491241231156971.