Title: Nasal Gene Expression in ART-Naive Adults with HIV and Pulmonary Tuberculosis in Uganda

Authors

Nisreen Khambati*[1]

Kattya Lopez Tamara*[2]

Elizabeth Nakabugo[3]

Arthur Van Valkenburg[4]

Jessica K Anderson[4]

Sean Lu[4]

Rinn Song[1]

Suryaram Gummuluru[5]

Andrew J Pollard[1]

Jerrold Ellner[2]

Padmini Salgame[2]

Else Margreet Bijker[1,6]

Lydia Nakiyingi[3]

Daniel O Connor[1]

W Evan Johnson[4]

*shared first author

Affiliations

1. Oxford Vaccine Group, Department of Paediatrics, University of Oxford, Oxford, United
Kingdom

2. Department of Medicine, Centre for Emerging Pathogens, Rutgers-New Jersey Medical
School, Newark, NJ, USA

3. Makerere University College of Health Sciences, Kampala, Uganda

4. Division of Infectious Disease, Center for Data Science, Rutgers New Jersey Medical School,

Newark, NJ, 07103, USA

5. Department of Virology, Immunology & Microbiology, Boston University Chobanian and

Avedisian School of Medicine, Boston MA 02118

6. Department of Paediatrics, Maastricht University Medical Center, MosaKids Children's

Hospital, Maastricht, The Netherlands

Corresponding author

Nisreen Khambati, nisreen.khambati@paediatrics.ox.ac.uk

Department of Paediatrics, Oxford University, Children's Hospital, John Radcliffe

Headington, Oxford, OX3 9DU

Abstract

Background: Diagnosis of pulmonary tuberculosis (TB) in people living with HIV remains difficult.

Since the first pathogen-host interaction in TB occurs in the upper airway, we hypothesized that host

transcriptomic analysis on nasal specimens may identify novel diagnostic biomarkers. We aimed to

demonstrate differences in nasal gene expression between people with HIV and TB disease versus

people with HIV without TB, evaluate the performance of nasal signatures in classifying TB and

compare nasal gene profiles with blood gene profiles from the same cohort.

Methods: We enrolled adults in Uganda with newly diagnosed HIV and symptoms of pulmonary TB

disease We collected nasal cells and blood for RNA sequencing to identify differentially expressed

genes and enriched pathways between people with HIV and TB disease and people with HIV without

TB. Supervised machine-learning of gene expression data was used to predict TB status.

Results: 40 adults living with HIV were enrolled (median age: 34 years, median CD4 count: 182),

including 20 with TB disease and 20 without. We identified 44 nasal differentially expressed genes

and 238 blood differentially expressed genes, with three overlapping genes between sample types.

Models trained using all 44 nasal differentially expressed genes had a cross-validated area under the

curve between 0.87-0.90 for predicting TB disease amongst adults living with HIV. A simplified four-

gene signature (*SPIB, SHISA2, TESPA1* and *CD1B*) met the World Health Organization criteria for a TB

triage test. Among adults with TB, pathways related to the inflammatory response and innate

immune system were downregulated in nasal samples and upregulated in blood.

Conclusion: This proof-of-concept study demonstrated that there were distinct nasal gene expression

patterns associated with TB, not seen in blood. Differences in nasal gene expression in people with

HIV who have TB disease, versus those without TB, highlight their potential as diagnostic biomarkers.

Further validation studies of gene signatures using minimally invasive nasal samples are

recommended in other difficult to diagnose groups.

Keywords

Manuscript

Background

Tuberculosis (TB) is the primary reason for hospitalization and death in people living with HIV

(PLHIV)[1], with the highest TB-HIV coinfection in the African Region[2]. TB-related mortality in PLHIV is

partly due to difficulties in diagnosis. Autopsy studies in PLHIV demonstrated that TB was

undiagnosed in nearly half of cases identified post-mortem[3]. PLHIV are more likely to have

paucibacillary disease and difficulty in expectorating sputum than those without HIV[1], limiting

sputum-based tests such as culture or Xpert MTB/RIF Ultra (Ultra), a cartridge-based polymerase-

chain-reaction (PCR) assay. Recognizing this, the World Health Organization (WHO) has advocated for

development of non-sputum-based biomarker tests for diagnosis of pulmonary TB[4].

Several blood-based mRNA biomarkers have shown promise as triage tests for TB[5]. However, their

potential has been limited by a poorer performance in PLHIV[6,7], particularly in antiretroviral therapy

(ART) naive individuals[8], or when differentiating TB from other respiratory diseases[9]. Lower

specificity in PLHIV could be due to the dominance of interferon-stimulated genes (ISGs) in blood and

HIV-related viraemia or opportunistic infections upregulating interferon-driven pathways[8-10].

Therefore, alternative sample types for biomarker discovery are worth exploring. With the first

pathogen-host interaction in TB occurring in the upper airway, transcriptomic analysis of nasal

specimens may identify host gene profiles and pathways not previously identified in blood.

Overlap in patterns of gene expression across the upper and lower airways in respiratory

inflammatory conditions[11-13] suggests that nasal samples could be proxies for studying lung disease.

Moreover, nasal specimens are also minimally invasive and easy to collect.

A rich transcriptomic response in nasal samples has been demonstrated for biomarker discovery and

understanding pathophysiology for several non-TB respiratory illnesses, including respiratory

syncytial virus[14,15], SARS-CoV-2[16-18], pneumonia[19], asthma[20] and lung cancer[12]. Pandya *et al* analyzed

1,555 nasal samples across ten cohorts and developed a 33-gene nasal signature that could classify

viral acute respiratory infections from healthy or non-viral samples with over 80% specificity and

sensitivity[21]. Amongst SARS-CoV-2 patients, suppressed interferon responses in nasal samples were

linked to more severe disease[17]. In children, nasal gene signatures could accurately predict

respiratory viral infections[14,22], and in some cases, performed better than blood classifiers[14]. Such

studies highlight the value of studying the local nasal response to respiratory infections.

Despite these potential benefits, to our knowledge, nasal gene expression profiling has not been

studied in TB. To address this gap, we sought to determine if nasal gene expression differs in ART-

naive PLHIV and pulmonary TB disease compared with PLHIV without TB disease. We identified

differentially expressed genes (DEGs) and biological pathways from nasal cells and assessed the

performance of nasal signatures in classifying TB disease. As secondary objectives, we compared

nasal-derived DEGs, biological pathways and gene signatures with paired blood samples from the

same cohort and evaluated the diagnostic performance of existing published blood TB signatures in

our nasal dataset.

155 **Methods**

156 **Figure 1** provides an overview of the methods. Computational code for the methodology is available

157 at: https://github.com/nisreenkhambati/uganda_nasal_cells.

Figure 1: Overview of the methods. Adults in Uganda with newly diagnosed HIV and TB symptoms were enrolled and nasal

and blood samples were collected for RNA sequencing. Analysis aimed to determine if nasal gene expression differed in

PLHIV and pulmonary TB disease compared with PLHIV without TB disease. Figure created in BioRender.com, license

obtained by Anderson, J., 2026, https://BioRender.com/3fuig8f)

Ultra= Xpert MTB/RIF Ultra; AUC= area under the curve; PLHIV=people with HIV

Study cohort

This study was conducted at Mulago National Referral Hospital and Infectious Diseases Institute-

affiliated clinics in Kampala, Uganda between 2020 and 2021. Individuals between 18-50 years with

presumptive TB (defined as having symptoms of pulmonary TB including cough, fever, night sweats,

weight loss and hemoptysis), no previous HIV or TB treatment (including isoniazid prophylaxis) and

newly confirmed HIV were consecutively enrolled. The HIV status of those with presumptive TB was

determined as part of study screening, i.e. HIV was diagnosed due to the presentation with TB

symptoms. Those who were current or former smokers, had diabetes, severe malnutrition, a history

of TB, or had used corticosteroids or antibiotics in the past three months were excluded. Informed

consent was obtained from all participants, and the study was approved by institutional review

boards of all partnering institutions (Boston University Medical Campus, Joint Clinical Research

Centre and Uganda National Council for Science and Technology).

Definitions and sample collection

Each adult had one early morning expectorated sputum tested with Ultra (Cepheid, CA, USA). Adults

were classified as having active TB disease based on a positive sputum Ultra. Control subjects were

those with no microbiological evidence of TB.

Study personnel followed a standardized protocol to collect nasal cells and blood. A cytology brush

was used to collect nasal cells by insertion into each nostril and rotating it against the lateral wall for

three seconds. The brush head was placed into a cryovial containing RNA-preserving agent. Blood

was collected into PAXgene RNA tubes. Samples were obtained before TB or HIV treatments were

initiated and frozen at -80°C.

RNA sequencing and quality control

Nasal and blood samples were shipped on dry ice to Rutgers University. Total cellular RNA from nasal

and PAXgene blood samples was extracted using the RNeasy Plus Micro Kit protocol (QIAGEN Inc.,

Hilden, Germany) at Rutgers University according to the manufacturer's instructions. The integrity

of RNA was analyzed using an Agilent TapeStation. To generate the cDNA library, the NEBNext®

rRNA Depletion Kit was used to remove the ribosomal RNA. Illumina-compatible RNA-seq library was

then prepared using the NEB Next Ultra II RNASeq library preparation kit. The cDNA libraries were

purified with AmpureXP beads and quantified using an Agilent TapeStation and a Qubit 4

Fluorometer. Equimolar amounts of barcoded libraries were pooled and sequenced on the

Illumina NovaSeq S2 FC 6000 (Illumina, San Diego, CA), producing 40-50 million paired-end 150 base-

pair reads per sample. All samples were run across two lanes in a single flow cell in the same

sequencing run.

Raw FASTQ data were assessed for quality using FastQC[23]. Trimmomatic was used to remove low

quality bases and adaptors (LEADING:3 TRAILING:3 SLIDINGWINDOW:4:15 MINLEN:36)[24]. Reads were

aligned to the human genome (GRCh38.p14) using the Spliced Transcripts Alignment to a Reference

(STAR) tool[25] and annotated using the GENCODE v44 primary assembly.[26]

Gene count data were exported to R (version 4.3.1) and subset for protein coding genes using the

biomaRt package[27]. Mitochondrial, hemoglobin, ribosomal and RNA genes, as well as low-count

genes, were filtered out. Outliers were identified by examining the Euclidean distances between

pairwise combinations of samples, visualized using ComplexHeatmap[28] and principal component

analysis (PCA). TB status, age, sex, CD4 count, and HIV stage were investigated as potential

expression-associated variables using PCA and corrected if needed (excluding TB status) for

downstream analyses. Differences in principal components for variables separating TB and controls

were statistically tested using the Student's t-test, given normality of data distribution. All PCA and

heatmap analyses were performed on transformed normalized counts (variance stabilizing

transformation).

Differential gene expression, machine learning and ROC analysis

Differential gene expression analysis between PLHIV and TB disease and PLHIV without TB disease

(controls) was performed in the nasal and blood datasets using DESeq2[29]. DEGs were defined as

having an adjusted p-value of <0.05 and log2 fold-change ≥1. Identified DEGs were used to cluster

samples, visualized with ComplexHeatmap[28], using Euclidean distance as a measure of dissimilarity

and complete linkage for between-cluster separation. Gene set enrichment analysis (GSEA) was

performed using fgsea[30] to evaluate whether DEGs from one tissue were enriched in the ranked list

of genes from the other tissue. Genes were ranked based on the test statistic from DESeq2[29].

Models were trained with six machine learning algorithms based on all DEGs using caret[31]: random

forest (RF), support vector machine (SVM) with radial and linear kernels, elastic net, partial least

squares (PLS) discriminant analysis and k-nearest neighbors (KNN), with 10-fold cross-validation

repeated 10 times. These models incorporate both linear and non-linear data relationships and are

commonly used in TB diagnostic transcriptomic research[32-36]. Prior to model training, gene counts of

nasal and blood DEGs underwent variance stabilizing transformation. Predictors were pre-processed

by removing near zero-variance variables and centering and scaling for models sensitive to feature

scaling (elastic net, SVM, KNN and PLS) were also performed. Receiver operating characteristic (ROC)

curves for each model in diagnosing TB were constructed with the MLeval package[37] to estimate the

area under the curve (AUC), sensitivity, and specificity with 95% confidence intervals. Cut-off

thresholds were determined by the maximum Youden index[38].

Models with a reduced gene panel are potentially easier to translate to a diagnostic test. Variable

importance for each nasal model trained using all DEGs was estimated using caret[31]. The top 10

predictive genes for each model were identified and genes common to all six nasal models were

retained to refit a new model using only these genes. The AUC, sensitivity and specificity of this

parsimonious nasal signature in classifying TB was calculated using MLeval[37].

Finally, the INTERFEROME database[39] was used to identify which top DEGs were interferon-

stimulated genes, which we defined as genes that were significantly up or down regulated (>2.0-fold

change, p<0.05) in human samples treated with interferon versus controls.

Pathway analysis and sample deconvolution

Pathway analyses for nasal and blood samples were performed with GSEA using the fgsea package[30],

and over-representation analysis (ORA) using the clusterProfiler package[40]. For fGSEA, ranked gene

lists using the test statistic were compared to Hallmark gene sets and KEGG and Reactome pathways

from the Molecular Signatures Database (MSigDB)[41]. For ORA, gene ontology biological processes[42]

(GO BP) were evaluated in the differentially expressed upregulated and downregulated genes

separately in adults with TB. Pathways with adjusted p-value < 0.05 were considered functionally

enriched.

Nasal and blood gene lists from DESeq2 were also uploaded to the Qiagen's Ingenuity Pathway

Analysis (IPA) tool[43] and filtered by adjusted p-value (false discovery rate) < 0.1 to determine

canonical pathways and upstream regulators in TB. IPA uses a Fisher's exact test to compute p-values

and uses the log2 fold-change to compute z-scores. For visualization, canonical pathways and

upstream regulators with a p-value <0.05 and IPA z-score≥ ±1.5 were shown.

Nasal samples consist of both immune and epithelial cell components. The CIBERSORTX tool and the

validated LM22 gene signature matrix[44,45] were used to examine the immune cell composition in

nasal and blood samples. CIBERSORTX is a deconvolution tool that employs support vector regression

to estimate the relative proportion of immune cell subtypes from mixed bulk tissue transcriptome

profiles[45]. The LM22 matrix consists of 547 genes that can accurately distinguish 22 immune cell

populations, which can further be summarized into 11 major cell types based on shared lineage,

including B cells, CD4 cells, CD8 cells, gamma delta T cells, eosinophils, plasma cells, NK cells,

macrophages, neutrophils, dendritic cells and mast cells[44]. Counts data were first normalized as

transcripts per million (TPM) to match the signature matrix. CIBERSORTX was performed with B-

mode batch correction, quantile normalization disabled, and the number of permutations set to 500.

Results were expressed as fractions of total leukocyte content. Proportions of estimated immune cell

populations were compared by TB status using either Student's t-test or Mann-Whitney U test

depending on normality of data distribution, with analyses corrected for multiple comparisons.

Performance of published blood RNA signatures

The TBSignatureProfiler package[46] was used to apply published blood signatures that diagnose TB

disease to our nasal and blood RNA sequencing datasets. We selected blood signatures that were

shown to distinguish TB disease from latent TB infection, TB from some combination of other

diseases or TB from HIV. Raw counts were first converted into log counts per million (CPM).

Signature performance scores were calculated using gene set variation analysis (GSVA) and single-

sample GSEA (ssGSEA) and compared using two-sample t-tests and bootstrapped AUCs. Only

published blood signatures with two or more genes that coincided with genes in our sequencing data

were used in the analysis.

294   <u>**Results**</u>

Study participants

Forty PLHIV were enrolled, including 20 with pulmonary TB disease and 20 without (controls). The median age was 35 years, 45% were male, 55% had HIV stage 3 and the median CD4 count was 182. PAXgene samples were collected from all, whereas nasal samples were available from 35 (17 with TB and 18 controls). Adults with TB and controls did not differ in age, sex, CD4 count or HIV stage **(Supplementary Table 1).**

Quality control and PCA

67% of nasal and 57% of blood reads uniquely mapped to the human genome **(Supplementary Table 2).** Three nasal samples (S4, S20, S31) demonstrated high average sample-to-sample distances from most other samples and had low reads (<5 million reads) **(Supplementary Figure 1).** These samples were removed, with 15 adults with TB and 17 controls remaining for downstream nasal analyses. No blood samples failed quality control, hence none were excluded.

Using PCA, nasal samples from adults with TB and controls clustered differently on PC4 (p= 0.048) **(Supplementary Figure 2A, 2C).** In blood, differences in PC1 between TB and controls were significant (p = 6.853×10^{-4}) **(Supplementary Figure 2B, 2D).** Males clustered separately from females for PC3 (p= 2.798×10^{-3}) and PC4 (p= 4.943×10^{-7}) in nasal samples and for PC3 (p=5.449×10^{-5}) and PC4 (p=1.303×10^{-5}) for blood, whereas HIV-related variables and age did not separate samples **(Supplementary Figure 3).** After sex-correction, PC3 in nasal samples and PC1 in blood samples demonstrated significant differences between TB and controls (nasal PC3: p = 0.046, blood PC1: p = 2.706×10^{-4}) **(Supplementary Figure 4).**

Differential gene expression and ROC curve analysis

319   44 nasal DEGs and 238 blood DEGs were identified between TB and controls **(Figure 2,**

320   **Supplementary Table 3).** Nine of the top ten blood DEGs were classified as ISGs compared to four of

321   the top nasal DEGs. Hierarchical clustering based on the DEGs showed grouping of samples by TB

322   status, although overlap remained between the two groups **(Supplementary Figure 5).**

Figure 2: Volcano plots and box plots illustrating the top DEGs in PLHIV and TB compared to PLHIV without TB (controls).

Volcano plots show the effect size by the $-\log_{10}P$ value for nasal (A) and blood (B) samples. DEGs are highlighted in red

(upregulated) or blue (downregulated), with gene names shown for the top 5 upregulated and downregulated DEGs.

Boxplots are of normalized counts ($\log_{10}$ scale) for the top ten DEGs, which were all downregulated for nasal samples (C)

and upregulated for blood samples (D) in TB cases. Individual data points are shown as jittered dots.

DEGs=differentially expressed genes

In TB compared with controls, most nasal DEGs were downregulated whereas most blood DEGS were

upregulated, with only three overlapping DEGs **(Figure 3).** Based on fGSEA, nasal DEGs were

significantly negatively enriched in the ranked blood gene list and blood DEGs were also negatively

enriched in the ranked nasal gene list **(Supplementary Figure 6),** suggesting that genes respond

differently to TB disease in the two tissues, with compartment-specific immune responses.

Figure 3: Agreement plot of log2FC for genes in blood samples (x-axis) and nasal samples (y-axis). Colored features are

genes which were differentially expressed with log2 FC ≥1 and an adjusted p-value of <0.05. Dashed x=y line represents

agreement. Dotted lines are shown at +/-1 log2FC. Labelled DEGs are those which are common to both nasal cells and

blood and were downregulated in TB.

DEGs= differentially expressed genes; log2 FC= log2 fold-change.

Models trained using all 44 nasal DEGs had a cross-validated AUC between 0.87-0.90 for predicting

TB status in the nasal dataset, which was similar to models trained with the 238 blood DEGs (AUC:

0.89-0.98) in blood **(Figure 4, Table 1).** The highest AUC for the nasal models was RF. All four adults

with TB that were missed by the nasal RF model were correctly identified as TB by the blood RF

model. Likewise, the two adults with TB missed by the blood RF model were correctly classified by

the nasal RF model.

Figure 4: ROC curves and the respective AUC of different machine learning models in distinguishing TB from controls,

trained using the (A) 44 nasal DEGs and (B) 238 blood DEGs on the respective datasets, with 10-fold cross-validation

repeated 10 times.

Table 1: Diagnostic performance with 95% CI of each model

	NASAL DATA			BLOOD DATA		
	AUC (95%CI)	sensitivity (95%CI)	specificity (95%CI)	AUC (95%CI)	sensitivity (95%CI)	specificity (95%CI)
k-Nearest Neighbors	0.88 (0.75-1.01)	0.80 (0.55-0.93)	0.88 (0.66-0.97)	0.89 (0.78-1)	0.85 (0.64-0.95)	0.80 (0.58-0.92)
Support Vector Machine (radial)	0.89 (0.77-1.01)	0.73 (0.48-0.89)	0.94 (0.73-0.99)	0.93 (0.85-1.01)	0.90 (0.7-0.97)	0.85 (0.64-0.95)
Support Vector Machine (linear)	0.87 (0.74-1)	0.87 (0.62-0.96)	0.82 (0.59-0.94)	0.93 (0.85-1.01)	0.90 (0.7-0.97)	0.85 (0.64-0.95)
Partial Least Squares	0.88 (0.75-1.01)	0.73 (0.48-0.89)	0.94 (0.73-0.99)	0.93 (0.85-1.01)	0.95 (0.76-0.99)	0.80 (0.58-0.92)
Elastic Net	0.89 (0.77-1.01)	0.80 (0.55-0.93	0.88 (0.66-0.97)	0.98 (0.93-1.03)	0.95 (0.76-0.99)	0.95 (0.76-0.99)
Random Forest	0.90 (0.78-1.02)	0.73 (0.48-0.89)	1.00 (0.82-1)	0.91 (0.81-1.01)	0.90 (0.7-0.97)	0.80 (0.58-0.92)

Amongst the top ten nasal DEGs with the highest variable importance for each model, *SPIB, SHISA2,*

TESPA1 and *CD1B* were common to all nasal models (**Supplementary Figure 7).** Models trained using

this parsimonious four-gene signature (NASAL_4) had an AUC between 0.87-0.94 for predicting TB

status in the nasal data, with 5/6 models meeting the WHO target criteria for a TB triage test (>90%

sensitivity and >70% specificity)[4] (**Supplementary Figure 8 and Supplementary Table 4**).

Enrichment pathway analysis

GSEA results are summarized in **Table 2.** The top GSEA enriched nasal pathways were interferon

gamma (IFN-γ), neutrophil degranulation and chemokine signaling, all downregulated in adults with

TB compared with controls. Enriched downregulated pathways related to cytokine signaling and the

inflammatory response, particularly the innate immune system. Upregulated enriched pathways

were associated with cilia function, organelle maintenance and cholesterol metabolism. In contrast,

in blood, upregulated enriched pathways in TB related to immune activity and hem metabolism,

whereas downregulated pathways were associated with transcription and RNA processing. Enriched

pathways in interferon and interleukin signaling, neutrophil degranulation, TNF-alpha signaling,

complement, inflammatory response, phagocytosis, and leishmania infection were downregulated in

nasal samples and upregulated in blood samples amongst adults with TB.

Using ORA, no enriched nasal pathways were identified with an adjusted p-value of <0.05. With an

adjusted p-value of <0.1, upregulated nasal DEGs in TB pertained mainly to metabolic biological

processes and cell division and differentiation whereas inflammatory and immune pathways were

enriched in the downregulated genes **(Supplementary Figure 9A, 9C).** In blood, upregulated DEGs

were enriched in the inflammatory and immune response, whereas biological processes in the

downregulated DEGs included synaptic transmission, odontogenesis and regulation of cell migration

(adjusted p-value <0.05) **(Supplementary Figure 9B, 9D).**

Canonical pathways in IPA also demonstrated suppressed immune activity in nasal samples from

adults with TB, including reduced interferon signaling, phagosome formation, and S100 family and G-

protein-coupled receptor signaling. In contrast, blood showed activation of innate immune pathways,

such as neutrophil degranulation, IFN-γ, toll-like receptor signaling, inflammasomes, pyroptosis and

iNOS signaling **(Supplementary Figure 10A, 10B)**. This opposing pattern of inflammation and

interferon response was supported by upstream regulator analysis which predicted inhibition of type

I and type III interferon expression (IFNA2, IFNL1), interferon regulatory factors (IRF1), JAK- STAT

components and pathogen-recognition receptors (TLR9) in the nose and activation of inflammatory

regulators (IFN-γ, CSF2, IL5, TNF, IL15, TLR2, SELPG) in blood **(Supplementary Figure 10C, 10D)**.

Overall, GSEA, ORA and IPA suggested there was reduced nasal mucosal inflammation and increased

systemic inflammation in TB disease.

Table 2 should appear here. Table 2 is larger than one A4 page in length and as per BMC Infectious Diseases Submission Guidelines is found at the end of
the document text file.

Immune cell composition

Enrichment analysis suggested that the immune response, particularly the innate immune system,

was downregulated in the nasal samples and upregulated in blood in adults with TB. Heterogeneity

in immune cell composition was explored using CIBERSORTX and LM22. There were no statistically

significant differences in the proportions of immune cell subtypes in nasal samples based on TB

status. However, for blood, the proportion of neutrophils was higher in TB (adjusted p-value =

0.0376), whereas natural killer cells were lower in TB (adjusted p-value = 0.0376). Other types of

immune cells in blood showed no significant differences **(Supplementary Figure 11).**

TB Signature Profiler

In the RNA sequencing dataset, only 6/59 published TB diagnostic blood signatures distinguished TB

from controls with an AUC ≥0.7 and p-value <0.05 in the nasal dataset (Maertzdorf_100, Lee_5,

Sloot_2, Natarajan_7, Jacobensen_3 and Gjoen_10[46]) **(Figure 5A, 5C).** In contrast, most published

blood signatures distinguished TB from controls in our blood dataset (44/59 and 41/59 had an AUC

≥0.7 and p-value <0.05 using ssGSEA and GSVA respectively) **(Figure 5B, 5D).** Only one signature

which performed well in the nasal dataset (Jacobsen_3) performed well in the blood dataset. Full

AUC results are in **Supplementary Table 5 and 6.**

Figure 5: Boxplots of published blood diagnostic signatures that reached statistical significance (p value <0.05) and had a

bootstrapped AUC of >0.7 in distinguishing TB from controls using ssGSEA in the A) nasal and B) blood datasets and using

GSVA methods in the C) nasal and D) blood datasets.

*For blood, only the top ten (based on AUC) published blood diagnostic signatures are shown.

GSVA= gene set variation analysis; ssGSEA= single-sample gene set enrichment analysis.

Discussion

In a cohort of treatment-naive PLHIV with presumptive TB in Uganda, we identified 44 DEGs in nasal

cells between adults with microbiologically confirmed TB disease compared with individuals without

TB. These 44 DEGs yielded a promising AUC of > 0.87 in predicting TB across six models in cross-

validation within our dataset. Amongst those with TB disease, we also found a blunted inflammatory

nasal response, with downregulation of several innate immune pathways. Compartmental

differences were seen, with contrasting patterns of inflammation in TB between the nose and blood

and little overlap in nasal and blood DEGs. Our proof-of-concept study highlights that there are

differences in nasal gene expression associated with TB disease that are distinct from blood.

There are no published nasal transcriptomic studies in TB; however, half of the top ten nasal DEGs

have been associated with TB in blood. *SPIB,* which ranked amongst the most important predictive

genes across all models, is important for plasmacytoid dendritic cell function[47] and memory B cell

maintenance[48]. *SPIB* downregulation was noted in active TB patients in multiple studies[49-51]. *CLC,*

which is linked to neutrophil recruitment[52], was part of a four-marker blood signature for

differentiating adults with TB and non-TB febrile infection[53]. *ITLN,* which has a role in pathogen

recognition, was downregulated in newly diagnosed adults with TB, increasing in expression with

treatment[49]. Changes in expression of T cell markers *PTGDR2* and *CD69* have also been associated

with early signs of pulmonary TB[54] and TB infection, respectively[55]. Top DEGs not previously linked to

TB include *CEACAM21*, which facilitates opsonin-independent recognition and clearance of specific

gram-negative bacterial pathogens[56]. *CST1, CLC, FETUB* and *VSMT1* were upregulated in asthma and

allergic diseases in nasal studies[57-59], which could reflect overrepresentation of allergic diseases in

our non-TB control group.

GSEA, ORA and IPA suggested that TB was associated with suppressed nasal innate immune

responses. IFN-γ activity was the top enriched downregulated pathway from the Hallmark gene-sets

using fGSEA . IFN-γ is an important cytokine for immunity against mycobacteria. It facilitates the host

defense to *Mtb* through stimulating macrophage activity and producing reactive nitrogen species[60],

with ineffective IFN-γ activity associated with disseminated mycobacterial infection[61]. Neutrophils

were the top downregulated REACTOME enriched pathway. Whilst excessive neutrophil activation

can cause tissue damage, they have a protective role in early stages[62] and adult contacts with higher

blood neutrophil counts were shown to be less likely to develop TB infection[63]. Chemokine signaling

was the most enriched downregulated KEGG pathway. Single nucleotide polymorphisms in

chemokines involved in cell migration in TB, were shown to increase susceptibility to active TB[64,65].

The nose is the first line of defense for airborne pathogens and early control of respiratory infection

is dependent on robust nasal mucosal immune responses[66]. Dampening of IFN-γ activity,

neutrophils, and chemokines in addition to other enriched nasal pathways like complement,

phagocytosis, and toll-like receptor signaling, may underlie failure of the early protective nasal

response to *Mtb* in TB disease.

The nasal host response in TB differed from blood. Few published TB blood signatures could classify

TB in our nasal sequencing dataset. Moreover, downregulated nasal innate immune responses in TB

were accompanied by upregulation of inflammatory pathways in blood. Unlike blood, nasal samples

have more heterogeneity in cell composition, consisting of both immune and epithelial

components[14]. We could not show that nasal immune pathways were driven by immune cell

composition. However, neutrophils in the blood were higher in PLHIV with TB versus those without

TB, consistent with our observation of increased neutrophil enrichment in blood.

Our finding of upregulated innate immune pathways in TB blood signatures confirms previously

published studies in PLHIV[10,67,68]. Similar to the downregulation of immune nasal pathways in TB in

our study, Mick and colleagues found that toll-like receptor signaling, inflammasomes, neutrophil

degranulation and chemokine signaling were attenuated in the nasopharyngeal response in SARS-

CoV-2 patients compared to other viral infections[16]. Hurst and colleagues also noted that systemic

symptoms in SARS-CoV-2-infected individuals were correlated with suppressed nasal innate and

adaptive immune pathways and upregulation of these same pathways in matched blood samples[69].

In contrast, two studies comparing nasal and blood samples in children with viral infections,

identified upregulation in immune activity and strong innate responses in both compartments[14,15].

Although our cross-sectional study cannot infer causality, we can hypothesize that reduced mucosal

immune responses and inadequate control of *Mtb* in the nose could promote progression to lung

disease and consequent systemic inflammation. Longitudinal studies using paired nasal and blood

samples are needed to understand the temporal relationship between the nasal and blood immune

response and TB disease.

A streamlined nasal gene signature based on variable importance (NASAL_4) met the WHO criteria

for a TB triage test (>90% sensitivity and 70% specificity) in cross-validation. Three genes within

NASAL_4 relate to immune activity: *SPIB* is linked to interferon production and proinflammatory

cytokines[47], *CD1b* is associated with antigen presentation[70] and *TESPA* relates to T cell maturation[71].

Whilst the AUC for nasal models based on all DEGs was slightly lower than blood, most of the top ten

nasal DEGs were not ISGs, unlike blood. This is promising given that the performance of many ISG-

dominant RNA blood signatures in differentiating TB from viral infections or in PLHIV has been

inadequate[8,10]. Moreover, adults with TB missed by blood models were classified correctly by nasal

models and vice versa, suggesting potential value of combined nasal-blood diagnostic biomarkers.

All adults in our study had HIV. Studies of the nasal transcriptome in HIV are limited, however HIV

causes mucosal immune dysfunction[72], which will shape nasal transcriptional changes,

independently of TB. Based on flow cytometry and single cell sequencing, Phiri and colleagues found

increased nasal neutrophilic abundance and inflammation in PLHIV compared with non-infected

individuals and that nasal neutrophils had transcriptional signatures of stress and senescence and

nasal T cells had signatures of exhaustion and apoptosis[73]. Differing numbers of nasal CD3 and CD4

cells based on HIV status have also been noted[74]. Further studies looking at the nasal TB response

among individuals without HIV would help explore the extent to which nasal gene profiles in our

study are driven by HIV.

Our work has limitations. The reliance on sputum Ultra as the TB reference standard in PLHIV means

paucibacillary TB disease could be missed in the control group. We could not follow up controls to

identify later development of TB disease and were unable to evaluate the role of nasal signatures in

predicting TB progression. Secondly, our ART-naive cohort was at high risk of opportunistic infections

and nasopharyngeal bacterial colonization. However, we were unable to ascertain and adjust for

these as well as any alternative diagnoses in non-TB controls that would affect gene expression. Case

numbers were too small to study the impact of CD4 count on nasal signatures, and other covariates

that impact gene expression, like malnutrition and TB infection via TST or IGRA status, were

unavailable. Finally, nasal DEG-based classifiers were trained and evaluated on the same dataset

since the sample size was too small to allocate a separate test set, and there are no other published

nasal transcriptomic datasets for TB. Although we employed k-fold cross-validation to reduce

overfitting, the need for external validation of nasal signatures on independent datasets remains.

Despite these limitations, these are the first data interrogating the nasal transcriptome in TB disease.

Whilst developing a nasal diagnostic test will require a more accessible technology than sequencing,

detecting gene expression differences is the first step. Our focus on ART-naïve PLHIV who have an

elevated risk of TB addresses a major gap in HIV-TB studies and by only studying symptomatic adults,

we remove the bias of using healthy controls. Although outside the scope of this work, sequencing

nasal samples enable the concurrent study of the respiratory microbiome and opportunity to

develop integrated host-microbiome diagnostic biomarkers. Although we focused on adults

diagnosed with Ultra on expectorated sputum as our microbiological reference standard, a nasal

biomarker could still have additional value for TB diagnosis given the ease of sampling compared

with sputum. Future work to confirm the utility of nasal signatures in TB requires discovery and

external validation of nasal biomarkers across larger, independent, geographically distinct cohorts,

that include adults with and without HIV across the spectrum of TB disease, as well as other difficult

to diagnose groups, such as young children or those without microbiologically confirmed TB.

Conclusion

In conclusion, our exploratory study using minimally invasive nasal samples suggests that host nasal

transcriptomics is feasible and informative for the diagnosis of TB, whereby elucidating gene

expression profiles at the initial pathogen encounter reveals novel insights into host-TB interaction.

We found distinct nasal gene expression patterns associated with TB disease in PLHIV that were not

captured in blood, with DEGs able to distinguish TB from non-TB controls. Our findings support

further study of the nose for non-sputum-based TB diagnostic biomarkers, which is an urgent

research priority to reduce global TB-associated mortality.

<u>Additional tables greater than one A4 in length</u>

Table 2: The top ten enriched pathways in the Hallmark gene sets, KEGG and Reactome pathways based on fGSEA in nasal and blood samples

NASAL			BLOOD		
	Adjusted p-value	Normalized enrichment score		Adjusted p-value	Normalized enrichment score
Hallmark gene set					
interferon gamma response*	4.58E-33	-3.1391781	heme metabolism	1.44E-17	2.69075782
interferon alpha response*	1.51E-23	-3.1566226	interferon gamma response*	1.95E-17	2.65410796
allograft rejection	8.15E-20	-2.7435101	EF2 targets	1.18E-12	-2.504435
inflammatory response*	5.64E-19	-2.667695	complement*	4.29E-11	2.33214589
TNF-A signaling via NF-kB*	1.45E-12	-2.3558235	IL6 JAK STAT3 signaling*	1.47E-09	2.50731015
IL2 STAT5 signaling	1.49E-10	-2.2420619	MYC targets	1.47E-09	-2.2858968
complement*	1.00E-08	-2.1158445	inflammatory response*	1.71E-08	2.15816136
IL6 JAK STAT3 signaling*	1.52E-08	-2.3716069	hypoxia	5.38E-08	2.11394868
bile acid metabolism	1.77E-05	2.04062867	interferon alpha response*	5.38E-08	2.34472265
cholesterol homeostasis	4.66E-05	2.04477918	TNF-A signaling via NF-kB*	7.37E-08	2.10746927
REACTOME pathway					
neutrophil degranulation*	1.22E-20	-2.3753505	neutrophil degranulation*	4.60E-48	3.03136454
cilium assembly	2.14E-19	2.82675326	platelet activation signaling and aggregation	1.65E-09	2.24160937
infraglabellar transport	5.73E-18	3.28715505	signaling by interleukins*	3.23E-09	1.99216824
organelle biogenesis and maintenance	2.01E-16	2.50485971	rrna processing	5.40E-09	-2.5587675
interferon alpha beta signalling	9.70E-14	-2.878876	antigen processing cross presentation	1.41E-07	2.36613708
immunoregulatory interactions between lymphoid and a non-lymphoid cell	1.25E-13	-2.6368349	interferon gamma signaling	1.41E-07	2.41404798
signaling by interleukins*	1.07E-12	-2.0969441	trna processing	9.24E-07	-2.335618
interferon signaling	1.34E-11	-2.262785	rrna modification in the nucleus and cytosol	1.10E-06	-2.5765856
leishmania infection	3.79E-09	-2.2903078	response to elevated platelet cytosolic ca2	2.32E-06	2.18883777
cholesterol biosynthesis	2.46E-08	2.8140458	diseases of signal transduction by growth factor receptors and second messengers	3.68E-06	1.80933376
KEGG pathway					
chemokine signaling pathway	1.24E-13	-2.5264553	regulation of actin cytoskeleton	9.20E-08	2.15502842
fc gamma r mediated phagocytosis*	9.01E-10	-2.4900559	leukocyte transendothelial migration	7.01E-06	2.20647871
cytokine cytokine receptor interaction	9.21E-09	-2.1236459	fc gamma r mediated phagocytosis*	1.67E-05	2.17411173
natural killer cell mediated cytotoxicity	9.21E-09	-2.3593142	lysosome	2.77E-05	2.05385563
toll like receptor signaling pathway*	9.21E-09	-2.4708031	focal adhesion	3.76E-05	1.92835864
leishmania infection*	2.48E-07	-2.3806651	leishmania infection*	6.30E-05	2.13257101

JAK STAT signaling pathway	1.45E-06	-2.1495304	acute myeloid leukemia	8.44E-05	2.17502533
B cell receptor signaling pathway	1.96E-06	-2.2891477	complement and coagulation cascades	8.44E-05	2.2165221
T cell receptor signaling pathway	1.96E-06	-2.1668356	neurotrophin signaling pathway	8.44E-05	1.9836064
MAPK signaling pathway	2.04E-06	-1.8908071	toll like receptor signaling pathway*	8.86E-05	2.05311921

575   *Negative normalized enrichment scores indicate the pathway is downregulated in TB compared to non-TB controls, and positive enrichment scores indicate the pathway is upregulated in TB.*

576   *Enriched pathways which are shared in both nasal and blood samples are italicized and have an asterisk (*).*

List of abbreviations

ART: antiretroviral therapy

AUC: area under the curve

DEG: differentially expressed genes

GSEA: gene set enrichment analysis

GSVA: gene set variation analysis

IPA: ingenuity pathway analysis

KNN: k nearest neighbors

ORA: over representation analysis

PCA: principal component analysis

PLHIV: people living with HIV

PLS: partial least squares

RF: random forest

ROC: receiver operating characteristic curve

SVM: support vector machine

ssGSEA: single sample gene set enrichment analysis

TB : tuberculosis

WHO: world health organization

<u>Declarations</u>

Ethics approval and consent to participate

This study used human data. Informed consent was obtained from all participants. The study

protocol and informed consent forms was approved, with ethics approval given by institutional

review boards of all partnering institutions (Boston University Medical Campus, Joint Clinical

Research Centre and Uganda National Council for Science and Technology). This study did not use

any animal data.

Consent for publication

Not applicable

Availability of data and materials

The datasets analyzed during the current study are available in the SRA repository

(PRJNA1379661) on https://www.ncbi.nlm.nih.gov/bioproject/PRJNA1379661

Computational code and datasets are also available at:

https://github.com/nisreenkhambati/uganda_nasal_cells.

Competing interests

The authors declare that they have no competing interests

Funding

This work was supported by grants from the National Institute of Allergy and Infectious Diseases of

the National Institutes of Health: UO1AI52084 (JE), R01AI152159 (PS), R21AI154387 (WEJ),

R01GM127430 (WEJ), R01AI175315 (WEJ) and the PROVIDENCE/BOSTON CFAR (P30AI042853).

Author contributions

Conceptualization: JE, LN, WEJ, SG

Methodology: NK, KLT, LN, WEJ

Resources: LN, WEJ, PS

Project administration: LN, WEJ

Investigation: EN, LN, KLT, NK

Software: NK

Data curation: EN, NK

Formal analysis: NK (except IPA done by AVV)

Validation: KLT, SL, JKA, WEJ

Writing–original draft: NK

Writing –review of final manuscript: all authors

Visualisation- NK

Supervision: RS, AJP, EMB, DOC, WEJ

Funding acquisition: JE, LN, WEJ, SG

Acknowledgements

We would like to thank all the participants in Uganda who were part of this study, as well as all local

clinical, lab and data teams involved.

References

1.   Meintjes G, Maartens G. HIV-Associated Tuberculosis. N Engl J Med. 2024;391(4):343-55.
doi:10.1056/NEJMra2308181
2.   The World Health Organization. Global tuberculosis report 2024. Geneva: The WHO; 2024.
3.   Gupta RK, Lucas SB, Fielding KL, Lawn SD. Prevalence of tuberculosis in post-mortem studies
of HIV-infected adults and children in resource-limited settings: a systematic review and meta-
analysis. AIDS. 2015;29(15):1987-2002. doi:10.1097/QAD.0000000000000802
4.   The World Health Organization. High-priority target product profiles or new tuberculosis
diagnostics: report of a consensus meeting. Geneva; 2014.
5.   Warsinske H, Vashisht R, Khatri P. Host-response-based gene signatures for tuberculosis
diagnosis: A systematic comparison of 16 signatures. PLoS Med. 2019;16(4):e1002786.
doi:10.1371/journal.pmed.1002786
6.   Muwanga VM, Mendelsohn SC, Leukes V, Stanley K, Mbandi SK, Erasmus M, et al. Blood
transcriptomic signatures for symptomatic tuberculosis in an African multicohort study. Eur Respir J.
2024;64(2). doi:10.1183/13993003.00153-2024
7.   Gupta-Wright A, Ha H, Abdulgadar S, Crowder R, Emmanuel J, Mukwatamundu J, et al.
Evaluation of the Xpert MTB Host Response assay for the triage of patients with presumed
pulmonary tuberculosis: a prospective diagnostic accuracy study in Viet Nam, India, the Philippines,
Uganda, and South Africa. Lancet Glob Health. 2024;12(2):e226-e34. doi:10.1016/S2214-
109X(23)00541-7
8.   Mann T, Gupta RK, Reeve BWP, Ndlangalavu G, Chandran A, Krishna AP, et al. Blood RNA
biomarkers for tuberculosis screening in people living with HIV before antiretroviral therapy
initiation: a diagnostic accuracy study. Lancet Glob Health. 2024;12(5):e783-e92. doi:10.1016/S2214-
109X(24)00029-9
9.   Mendelsohn SC, Verhage S, Mulenga H, Scriba TJ, Hatherill M. Systematic review of
diagnostic and prognostic host blood transcriptomic signatures of tuberculosis disease in people
living with HIV. Gates Open Res. 2023;7:27. doi:10.12688/gatesopenres.14327.2
10.   Esmail H, Lai RP, Lesosky M, Wilkinson KA, Graham CM, Horswell S, et al. Complement
pathway gene activation and rising circulating immune complexes characterize early disease in HIV-
associated tuberculosis. Proc Natl Acad Sci U S A. 2018;115(5):E964-E73.
doi:10.1073/pnas.1711853115
11.   Kicic A, de Jong E, Ling KM, Nichol K, Anderson D, Wark PAB, et al. Assessing the unified
airway hypothesis in children via transcriptional profiling of the airway epithelium. J Allergy Clin
Immunol. 2020;145(6):1562-73. doi:10.1016/j.jaci.2020.02.018
12.   Perez Rogers JF. Shared Gene Expression Alterations in Nasal and Bronchial Epithelium for
Lung Cancer Detection. J Natl Cancer Inst. 2017;109(7). doi:10.1093/jnci/djw327
13.   Gower AC, Steiling K, Brothers JF, 2nd, Lenburg ME, Spira A. Transcriptomic studies of the
airway field of injury associated with smoking-related lung disease. Proc Am Thorac Soc.
2011;8(2):173-9. doi:10.1513/pats.201011-066MS
14.   Yu J, Peterson DR, Baran AM, Bhattacharya S, Wylie TN, Falsey AR, et al. Host Gene
Expression in Nose and Blood for the Diagnosis of Viral Respiratory Infection. J Infect Dis.
2019;219(7):1151-61. doi:10.1093/infdis/jiy608
15.   Do LAH, Pellet J, van Doorn HR, Tran AT, Nguyen BH, Tran TTL, et al. Host Transcription Profile
in Nasal Epithelium and Whole Blood of Hospitalized Children Under 2 Years of Age With Respiratory
Syncytial Virus Infection. J Infect Dis. 2017;217(1):134-46. doi:10.1093/infdis/jix519
16.   Mick E, Kamm J, Pisco AO, Ratnasiri K, Babik JM, Castaneda G, et al. Upper airway gene
expression reveals suppressed immune responses to SARS-CoV-2 compared with other respiratory
viruses. Nat Commun. 2020;11(1):5854. doi:10.1038/s41467-020-19587-y

17.   Ziegler CGK, Miao VN, Owings AH, Navia AW, Tang Y, Bromley JD, et al. Impaired local intrinsic
immunity to SARS-CoV-2 infection in severe COVID-19. Cell. 2021;184(18):4713-33 e22.
doi:10.1016/j.cell.2021.07.023
18.   Hurst JH, Mohan AA, Dalapati T, George IA, Aquino JN, Lugo DJ, et al. Differential host
responses within the upper respiratory tract and peripheral blood of children and adults with SARS-
CoV-2 infection. medRxiv. 2023:2023.07.31.23293337. doi:10.1101/2023.07.31.23293337
19.   Jochems SP, Marcon F, Carniel BF, Holloway M, Mitsi E, Smith E, et al. Inflammation induced
by influenza virus impairs human innate immune control of pneumococcus. Nat Immunol.
2018;19(12):1299-308. doi:10.1038/s41590-018-0231-y
20.   Castro-Nallar E, Shen Y, Freishtat RJ, Perez-Losada M, Manimaran S, Liu G, et al. Integrating
microbial and host transcriptomics to characterize asthma-associated microbial communities. BMC
Med Genomics. 2015;8:50. doi:10.1186/s12920-015-0121-1
21.   Pandya R, He YD, Sweeney TE, Hasin-Brumshtein Y, Khatri P. A machine learning classifier
using 33 host immune response mRNAs accurately distinguishes viral and non-viral acute respiratory
illnesses in nasal swab samples. Genome Med. 2023;15(1):64. doi:10.1186/s13073-023-01216-0
22.   Yahya M, Rulli M, Toivonen L, Waris M, Peltola V. Detection of Host Response to Viral
Respiratory Infection by Measurement of Messenger RNA for MxA, TRIM21, and Viperin in Nasal
Swabs. J Infect Dis. 2017;216(9):1099-103. doi:10.1093/infdis/jix458
23.   Andrews S. FastQC: A Quality Control Tool for High Throughput Sequence Data 2010
[Available from: https://www.bioinformatics.babraham.ac.uk/projects/fastqc/. Accessed: 6th January
2026
24.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics. 2014;30(15):2114-20. doi:10.1093/bioinformatics/btu170
25.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal
RNA-seq aligner. Bioinformatics. 2013;29(1):15-21. doi:10.1093/bioinformatics/bts635
26.   GENCODE. Human Release 44 (GRCh38.p14) 2022 [Available from:
https://www.gencodegenes.org/human/release_44.html. Accessed: 6th January 2026
27.   Durinck S, Spellman PT, Birney E, Huber W. Mapping identifiers for the integration of genomic
datasets with the R/Bioconductor package biomaRt. Nat Protoc. 2009;4(8):1184-91.
doi:10.1038/nprot.2009.97
28.   Gu Z, Eils R, Schlesner M. Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. Bioinformatics. 2016;32(18):2847-9.
doi:10.1093/bioinformatics/btw313
29.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. Genome Biol. 2014;15(12):550. doi:10.1186/s13059-014-0550-8
30.   Korotkevich G, Sukhov V, Sergushichev A. Fast gene set enrichment analysis. bioRxiv. 2019.
31.   Kuhn M. Building Predictive Models in R Using the caret Package. Journal of Statistical
Software. 2008;28(5):1 - 26. doi:10.18637/jss.v028.i05
32.   Channon-Wells S, Habgood-Coote D, Vito O, Galassini R, Wright VJ, Brent AJ, et al. Integration
and validation of host transcript signatures, including a novel 3-transcript tuberculosis signature, to
enable one-step multiclass diagnosis of childhood febrile disease. J Transl Med. 2024;22(1):802.
doi:10.1186/s12967-024-05241-4
33.   Bobak CA, Titus AJ, Hill JE. Comparison of common machine learning models for classification
of tuberculosis using transcriptional biomarkers from integrated datasets. Applied Soft Computing.
2019;74:264-73. doi:https://doi.org/10.1016/j.asoc.2018.10.005
34.   Anderson ST, Kaforou M, Brent AJ, Wright VJ, Banwell CM, Chagaluka G, et al. Diagnosis of
childhood tuberculosis and host RNA expression in Africa. N Engl J Med. 2014;370(18):1712-23.
doi:10.1056/NEJMoa1303657
35.   Ding S, Yi X, Gao J, Huang C, Zhou Y, Yang Y, et al. Combining bioinformatics and machine
learning to identify diagnostic biomarkers of TB associated with immune cell infiltration. Tuberculosis
(Edinb). 2024;149:102570. doi:10.1016/j.tube.2024.102570

36.   Turner CT, Gupta RK, Tsaliki E, Roe JK, Mondal P, Nyawo GR, et al. Blood transcriptional
biomarkers for active pulmonary tuberculosis in a high-burden setting: a prospective, observational,
diagnostic accuracy study. Lancet Respir Med. 2020;8(4):407-19. doi:10.1016/S2213-2600(19)30469-
2
37.   John CR. MLeval: Machine Learning Model Evaluation 2020 [Computer software version 0.3].
Available from: https://cran.r-project.org/web/packages/MLeval/index.html. Accessed: 10th January
2026
38.   Fluss R, Faraggi D, Reiser B. Estimation of the Youden Index and its associated cutoff point.
Biom J. 2005;47(4):458-72. doi:10.1002/bimj.200410135
39.   Rusinova I, Forster S, Yu S, Kannan A, Masse M, Cumming H, et al. Interferome v2.0: an
updated database of annotated interferon-regulated genes. Nucleic Acids Res. 2013;41(Database
issue):D1040-6. doi:10.1093/nar/gks1215
40.   Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, et al. clusterProfiler 4.0: A universal enrichment tool
for interpreting omics data. Innovation (Camb). 2021;2(3):100141. doi:10.1016/j.xinn.2021.100141
41.   Broad Institute. Human MSigDB Collections: Broad Institute, Inc., Massachusetts Institute of
Technology, and Regents of the University of California; 2023 [Available from: https://www.gsea-
msigdb.org/gsea/msigdb/collections.jsp. Accessed: 6th January 2026
42.   Gene Ontology C, Aleksander SA, Balhoff J, Carbon S, Cherry JM, Drabkin HJ, et al. The Gene
Ontology knowledgebase in 2023. Genetics. 2023;224(1). doi:10.1093/genetics/iyad031
43.   Kramer A, Green J, Pollard J, Jr., Tugendreich S. Causal analysis approaches in Ingenuity
Pathway Analysis. Bioinformatics. 2014;30(4):523-30. doi:10.1093/bioinformatics/btt703
44.   Newman AM, Liu CL, Green MR, Gentles AJ, Feng W, Xu Y, et al. Robust enumeration of cell
subsets from tissue expression profiles. Nat Methods. 2015;12(5):453-7. doi:10.1038/nmeth.3337
45.   Newman AM, Steen CB, Liu CL, Gentles AJ, Chaudhuri AA, Scherer F, et al. Determining cell
type abundance and expression from bulk tissues with digital cytometry. Nat Biotechnol.
2019;37(7):773-82. doi:10.1038/s41587-019-0114-2
46.   Johnson WE, Odom A, Cintron C, Muthaiah M, Knudsen S, Joseph N, et al. Comparing
tuberculosis gene signatures in malnourished individuals using the TBSignatureProfiler. BMC Infect
Dis. 2021;21(1):106. doi:10.1186/s12879-020-05598-z
47.   Sasaki I, Hoshino K, Sugiyama T, Yamazaki C, Yano T, Iizuka A, et al. Spi-B is critical for
plasmacytoid dendritic cell function and development. Blood. 2012;120(24):4733-43.
doi:10.1182/blood-2012-06-436527
48.   Horiuchi S, Koike T, Takebuchi H, Hoshino K, Sasaki I, Fukuda-Ohta Y, et al. SpiB regulates the
expression of B-cell-related genes and increases the longevity of memory B cells. Front Immunol.
2023;14:1250719. doi:10.3389/fimmu.2023.1250719
49.   Cliff JM, Lee JS, Constantinou N, Cho JE, Clark TG, Ronacher K, et al. Distinct phases of blood
gene expression pattern through tuberculosis treatment reflect modulation of the humoral immune
response. J Infect Dis. 2013;207(1):18-29. doi:10.1093/infdis/jis499
50.   Bah SY, Forster T, Dickinson P, Kampmann B, Ghazal P. Meta-Analysis Identification of Highly
Robust and Differential Immune-Metabolic Signatures of Systemic Host Response to Acute and
Latent Tuberculosis in Children and Adults. Front Genet. 2018;9:457. doi:10.3389/fgene.2018.00457
51.   Natarajan S, Ranganathan M, Hanna LE, Tripathy S. Transcriptional Profiling and Deriving a
Seven-Gene Signature That Discriminates Active and Latent Tuberculosis: An Integrative
Bioinformatics Approach. Genes (Basel). 2022;13(4). doi:10.3390/genes13040616
52.   Gevaert E, Delemarre T, De Volder J, Zhang N, Holtappels G, De Ruyck N, et al. Charcot-
Leyden crystals promote neutrophilic inflammation in patients with nasal polyposis. J Allergy Clin
Immunol. 2020;145(1):427-30.e4. doi:10.1016/j.jaci.2019.08.027
53.   Roe JK, Thomas N, Gil E, Best K, Tsaliki E, Morris-Jones S, et al. Blood transcriptomic diagnosis
of pulmonary and extrapulmonary tuberculosis. JCI Insight. 2016;1(16):e87238.
doi:10.1172/jci.insight.87238

54.   Shaukat SN, Eugenin E, Nasir F, Khanani R, Kazmi SU. Identification of immune biomarkers in
recent active pulmonary tuberculosis. Sci Rep. 2023;13(1):11481. doi:10.1038/s41598-023-38372-7
55.   Chen ZY, Wang L, Gu L, Qu R, Lowrie DB, Hu Z, et al. Decreased Expression of CD69 on T Cells
in Tuberculosis Infection Resisters. Front Microbiol. 2020;11:1901. doi:10.3389/fmicb.2020.01901
56.   Pils S, Gerrard DT, Meyer A, Hauck CR. CEACAM3: an innate immune receptor directed
against human-restricted bacterial pathogens. Int J Med Microbiol. 2008;298(7-8):553-60.
doi:10.1016/j.ijmm.2008.04.005
57.   Xu Z, Forno E, Sun Y, Manni ML, Han YY, Kim S, et al. Nasal epithelial gene expression and
total IgE in children and adolescents with asthma. J Allergy Clin Immunol. 2024;153(1):122-31.
doi:10.1016/j.jaci.2023.09.014
58.   Klingler AI, Stevens WW, Tan BK, Peters AT, Poposki JA, Grammer LC, et al. Mechanisms and
biomarkers of inflammatory endotypes in chronic rhinosinusitis without nasal polyps. J Allergy Clin
Immunol. 2021;147(4):1306-17. doi:10.1016/j.jaci.2020.11.037
59.   Tomizawa H, Yamada Y, Arima M, Miyabe Y, Fukuchi M, Hikichi H, et al. Galectin-10 as a
Potential Biomarker for Eosinophilic Diseases. Biomolecules. 2022;12(10).
doi:10.3390/biom12101385
60.   Shanmuganathan G, Orujyan D, Narinyan W, Poladian N, Dhama S, Parthasarathy A, et al.
Role of Interferons in Mycobacterium tuberculosis Infection. Clin Pract. 2022;12(5):788-96.
doi:10.3390/clinpract12050082
61.   Dorman SE, Holland SM. Mutation in the signal-transducing chain of the interferon-gamma
receptor and susceptibility to mycobacterial infection. J Clin Invest. 1998;101(11):2364-9.
doi:10.1172/JCI2901
62.   de Waal AM, Hiemstra PS, Ottenhoff TH, Joosten SA, van der Does AM. Lung epithelial cells
interact with immune cells and bacteria to shape the microenvironment in tuberculosis. Thorax.
2022;77(4):408-16. doi:10.1136/thoraxjnl-2021-217997
63.   Martineau AR, Newton SM, Wilkinson KA, Kampmann B, Hall BM, Nawroly N, et al.
Neutrophil-mediated innate immune resistance to mycobacteria. J Clin Invest. 2007;117(7):1988-94.
doi:10.1172/JCI31097
64.   Feng WX, Flores-Villanueva PO, Mokrousov I, Wu XR, Xiao J, Jiao WW, et al. CCL2-2518 (A/G)
polymorphisms and tuberculosis susceptibility: a meta-analysis. Int J Tuberc Lung Dis.
2012;16(2):150-6. doi:10.5588/ijtld.11.0205
65.   Sanchez-Castanon M, Baquero IC, Sanchez-Velasco P, Farinas MC, Ausin F, Leyva-Cobian F, et
al. Polymorphisms in CCL5 promoter are associated with pulmonary tuberculosis in northern Spain.
Int J Tuberc Lung Dis. 2009;13(4):480-5.
66.   Zhou X, Wu Y, Zhu Z, Lu C, Zhang C, Zeng L, et al. Mucosal immune response in biology,
disease prevention and treatment. Signal Transduct Target Ther. 2025;10(1):7. doi:10.1038/s41392-
024-02043-4
67.   Verma S, Du P, Nakanjako D, Hermans S, Briggs J, Nakiyingi L, et al. "Tuberculosis in advanced
HIV infection is associated with increased expression of IFNγ and its downstream targets". BMC
Infect Dis. 2018;18(1):220. doi:10.1186/s12879-018-3127-4
68.   Kulkarni V, Queiroz ATL, Sangle S, Kagal A, Salvi S, Gupta A, et al. A Two-Gene Signature for
Tuberculosis Diagnosis in Persons With Advanced HIV. Front Immunol. 2021;12:631165.
doi:10.3389/fimmu.2021.631165
69.   Hurst JH, McCumber AW, Aquino JN, Rodriguez J, Heston SM, Lugo DJ, et al. Age-Related
Changes in the Nasopharyngeal Microbiome Are Associated With Severe Acute Respiratory
Syndrome Coronavirus 2 (SARS-CoV-2) Infection and Symptoms Among Children, Adolescents, and
Young Adults. Clin Infect Dis. 2022;75(1):e928-e37. doi:10.1093/cid/ciac184
70.   Shahine A. The intricacies of self-lipid antigen presentation by CD1b. Mol Immunol.
2018;104:27-36. doi:10.1016/j.molimm.2018.09.022

71. Wang D, Zheng M, Lei L, Ji J, Yao Y, Qiu Y, et al. Tespa1 is involved in late thymocyte
development through the regulation of TCR-mediated signaling. Nat Immunol. 2012;13(6):560-8.
doi:10.1038/ni.2301
72. Paiardini M, Frank I, Pandrea I, Apetrei C, Silvestri G. Mucosal immune dysfunction in AIDS
pathogenesis. AIDS Rev. 2008;10(1):36-46.
73. Phiri JA, Sibale LL, Kapira G, Mlogoti L, Nyirenda JT, Mitole N, et al. Persistent pneumococcal
colonisation in antiretroviral-treated HIV infection is associated with nasal inflammation. Nat
Commun. 2025. doi:10.1038/s41467-025-67258-7
74. Moss RB, Scott TA, Goldrich M, Pitale M, Daniel A, Lebovics R, et al. Nasal mucosal cells in
human immunodeficiency virus type 1-seropositive patients with sinusitis. J Clin Lab Anal.
1996;10(6):418-22. doi:10.1002/(SICI)1098-2825(1996)10:6<418::AID-JCLA18>3.0.CO;2-#