A Methodological Framework for Passive Visual Mirror Exposure in Children with CVI and an Atelic Profile

Author

Edwige Smague[1]

Affiliations:

[1]N.E.T.A: Neuronal Epileptotranscriptomic, Sydney, Australia

Contact:

edwige@netadiscovery.org

ORCID ID: 0009-0002-8182-7888

Abstract

Children with cerebral visual impairment (CVI) presenting an atelic profile constitute a population that is largely excluded from neurovisual research, primarily due to the absence of reliable behavioural responses to standard assessment or intervention paradigms. This exclusion often leads to the assumption that a lack of observable response reflects an absence of visual processing, despite converging evidence from neuroscience indicating that neural processing may persist in the absence of overt behavioural expression.

This article proposes a pre-empirical methodological framework describing **a** passive visual mirror exposure paradigm intended to inform the design of future feasibility studies in children with CVI and an atelic profile. The visual mirror is defined as the repeated

presentation of a highly predictable, coherent visual experience, intended to reduce perceptual instability and cognitive load associated with typical visual environments, without aiming to restore or train visual function.

The paradigm relies on passive visual exposure delivered via a large screen, or, where appropriate and tolerated, a virtual reality headset configured in a strictly non-interactive mode, without requiring effort, task engagement, or voluntary responses from the child. The proposed framework is grounded in a precautionary ethical rationale specifically adapted to behaviourally non-responsive paediatric populations, defined here as children presenting absent voluntary behavioural responses to sensory or task-based instructions.

The framework outlines predefined safety criteria and non-verbal physiological indicators, including heart rate and autonomic markers, as illustrative examples of feasibility and tolerability monitoring, rather than behavioural performance.

This manuscript does not report or initiate research involving human participants. It is intended to characterise the ethical acceptability, conceptual implementability, and methodological considerations of a passive visual mirror paradigm in this vulnerable population, and to inform future research

46 considerations contingent upon formal ethical approval, without

47 implying any obligation or commitment to subsequent studies.

49 **Keywords:** Atelic profile; cerebral visual impairment (CVI);

50 cerebral plasticity; visual mirror exposure; non-verbal children;

51 passive virtual reality; vision rehabilitation.

1. Introduction

Cerebral visual impairment (CVI) (1) is now recognised as one of

the leading causes of paediatric visual disability. It results from

damage to or dysfunction of the cerebral pathways involved in

visual processing, independent of ocular integrity (2). Visual

difficulties associated with CVI are heterogeneous and may

include impairments in gaze orienting, motion perception, object

recognition, sustained visual attention, or visuomotor coordination.

Within this population, children presenting an atelic profile

constitute a particularly marginalised group. Atelicity is defined

here as a persistent inability to initiate, maintain, or orient

behaviour towards a goal, even in simple sensory contexts (3).

Children with atelic profiles are typically unable to follow

instructions or produce reliable behavioural responses, which

effectively excludes them from conventional visual assessment and

rehabilitation paradigms that rely on task engagement or
observable performance.

This exclusion has created a significant methodological and ethical
gap. The absence of empirical data in this subgroup makes it
difficult to determine whether these children are genuinely non-
responsive to visual stimuli or whether they are simply
inaccessible to existing assessment methods. In this context,
behavioural non-responsiveness cannot be assumed to reflect an
absence of visual processing, as converging evidence from
neuroscience indicates that perceptual and neural processing may
persist in the absence of overt behavioural expression.

The present paper addresses this gap by proposing a feasibility-
informed methodological framework designed to conceptually
examine physiological responses to controlled visual stimuli in the
absence of voluntary behaviour. Rather than aiming to assess
therapeutic efficacy, clinical benefit, or functional visual
improvement, the framework is explicitly limited to outlining
methodological and ethical conditions under which a passive visual
mirror exposure paradigm could be evaluated in children with CVI
and an atelic profile, while allowing the acquisition of stable, non-
behavioural physiological signals.

Accordingly, the primary objective of this work is not to report an

implemented study, but to define and formalise the methodological

and ethical considerations relevant to determining whether

repeated sessions of passive visual mirror exposure may be

feasible and tolerable under controlled conditions in behaviourally

non-responsive paediatric populations. The central feasibility

question is therefore presented as a prospective and conceptual

inquiry, rather than as an empirically tested outcome.

2. The Atelic Child and Visual Experience

2.1 Atelicity as a Disruption of the Perception–Prediction-
Action Loop

In children presenting an atelic profile, the absence of goal-

directed behaviour should not be interpreted as an absence of

perception. Evidence from cognitive and neuroscientific research

indicates that perceptual and memory-related processing can occur

in the absence of voluntary control or overt behavioural expression

(4). Accordingly, limited behavioural responsiveness does not

preclude the presence of underlying sensory processing.

Atelicity can be described as a functional difficulty in initiating,

maintaining, or organising behaviour in response to sensory

information. In this context, the child may receive sensory input

but be unable to use it reliably to guide action, exploration, or sustained engagement with the environment (5). This description does not imply a deficit in sensory reception itself, but rather a disruption in the translation of sensory information into observable, goal-directed behaviour.

2.2 Visual Experience in Atelic Children with Cerebral Visual Impairment

In atelic children with cerebral visual impairment, visual experience is often reported by clinicians and caregivers as unstable and difficult to interpret (6). Commonly described features include:

- **Image Instability**: where the visual scene may appear fragmented or inconsistent;
- **Excessive Perceptual Detail**: resulting in difficulty prioritising or organising visual information;
- **Difficulty Anticipating Visual Change**: such that new objects or changes in lighting may be experienced as abrupt or unpredictable;
- **Impaired Recognition of Visual Stability**: for example, difficulty recognising that an object remains the same despite minor changes in position or context.

At a functional level, these characteristics are associated with increased perceptual uncertainty and elevated cognitive load

during visual processing. Over time, visual engagement may

become effortful or aversive, leading to reduced spontaneous

visual exploration or apparent disengagement from visual stimuli

(7).

This section provides a conceptual and descriptive framework to

support the design choices of the proposed protocol. It does not

constitute a testable theoretical model and does not imply specific

mechanisms of impairment. Rather, it serves to justify the use of

simplified, predictable, and non-interactive visual stimuli within a

pre-empirical methodological framework involving behaviourally

non-responsive paediatric populations.

3. Research Question and Objectives

3.1 Primary Feasibility Question

The visual mirror exposure paradigm described in this article is

conceptually informed by principles derived from mirror-based

approaches originally developed in post-stroke motor rehabilitation

(8), while differing fundamentally in its intended function and

mechanism. Rather than relying on visual comparison or illusion,

the present paradigm is designed to provide a coherent, stable, and

highly predictable visual stimulus that may temporarily substitute

for unstable or cognitively costly visual environments in children

with cerebral visual impairment, consistent with sensory substitution principles (9) and descriptive models of predictive processing (7).

In the context of cerebral visual impairment and atelic profiles, the paradigm is explicitly designed for behaviourally non-responsive paediatric populations and does not require task engagement, instruction following, or voluntary behavioural responses.

The primary feasibility question articulated in this manuscript is therefore formulated as a prospective and conceptual inquiry, as follows: Would it be feasible to deliver repeated sessions of passive visual mirror exposure to children with cerebral visual impairment and an atelic profile while maintaining physiological tolerability and obtaining stable, interpretable non-behavioural physiological signals in behaviourally non-responsive paediatric populations?

3.2 Primary Objective

The primary objective of this work is to define and formalise the feasibility and physiological tolerability criteria relevant to a passive visual mirror exposure protocol in the context of future empirical studies involving children with cerebral visual impairment and an atelic profile. Feasibility is defined in terms of

the theoretical implementability of the protocol without inducing

sustained physiological distress, and while enabling the potential

acquisition of usable autonomic physiological measures in a

population for whom active, task-based paradigms are not

accessible (7).

3.3 Secondary Methodological Objectives

The secondary objectives outlined here are strictly methodological

and illustrative. They are intended to characterise the types of

protocol performance indicators and measurement properties that

may be relevant rather than to assess therapeutic efficacy or

clinical benefit. Specifically, the objectives are:

- to assess whether non-behavioural physiological measures,

including heart rate and heart rate variability, could in principle be

reliably recorded during passive visual mirror exposure sessions in

this population;

- to examine the intra-individual stability and variability of these

physiological signals as a methodological consideration under

conditions of highly controlled and predictable visual stimulation

(7);

- to outline potential rates of session completion, interruption, or

early termination due to indicators of non-verbal dissent or

physiological distress, in accordance with precautionary ethical

principles;

•   to explore whether protocol tolerability may reasonably be

expected to vary across individual neurovisual profiles within the

atelic population, without making claims of group-level inference.

3.4 Reference to Feasibility, Tolerability and Safety Criteria

To ensure the highest standard of participant safety, the present

manuscript specifies feasibility and tolerability criteria as a pre-

empirical governance framework. These predefined thresholds and

decision rules, detailed in Table 1, constitute a conceptual safety

and oversight structure intended to guide future protocol

implementation.

Their primary purpose is to characterise the reliability, tolerability,

and ethical acceptability of the protocol in children with atelic

profiles, and to ensure that any future empirical study remains

grounded in continuous monitoring of participant well-being and

protocol stability. These criteria are not intended to demonstrate

therapeutic efficacy but to guide ethical conduct and interpretation

within a feasibility-focused study.

Table 1. Illustrative feasibility, tolerability, and safety indicators with predefined decision thresholds (pre-empirical
framework).

Category	Indicator	Feasibility Thresholds and Interpretation

216 Note. HR: Heart Rate; HRV: Heart Rate Variability.

217 The indicators and thresholds presented in this table are provided as illustrative, pre-empirical examples intended to inform the
218 methodological design and ethical governance of future feasibility studies. They do not correspond to data collection, participant
219 monitoring, or decision-making in an implemented study.

Protocol Implementation	Session Completion Rate	≥ 75% of planned sessions would be expected to be completed without premature termination across a minimum sample of 10 participants.
Technical Reliability	Usable Physiological Data	≥ 80% of sessions would be expected to yield usable HR and HRV data, defined as less than 20% signal artefact, at the individual level.
Physiological Tolerability	Acute Distress Signal	HR would be expected to remain within less than a 20% increase from the individual baseline for at least 90% of exposure time for each participant.
Autonomic Stability	HRV Stability	HRV stability would be characterised by the absence of a persistent decrease sustained over three consecutive sessions within an individual participant.
Ethical Safety	Dissent Response	Ethical safety would require immediate session termination upon detection of predefined non verbal dissent markers, with full compliance with this stopping rule.
Attrition	Participant Withdrawal	Feasibility would be supported if participant withdrawal due to intolerance would not exceed 20%, corresponding to no more than two out of ten participants.
Informational Potential	Signal Sensitivity	Informational potential would be indicated by the presence of measurable physiological modulation during controlled novelty phases in at least 50% of participants within the predefined feasibility sample.

Table 1 summarises the primary feasibility question, objectives,
indicators, and predefined feasibility and tolerability criteria as part
of a pre-empirical methodological framework. All criteria are
specified a priori to illustrate how ethical conduct, safety
monitoring, and outcome interpretation could be structured in
future feasibility studies, rather than to report procedures applied in
an implemented study. The table links each research objective to
concrete physiological or procedural indicators and to explicit
decision rules intended to guide prospective protocol design and
ethical governance, rather than protocol continuation or early
termination in an ongoing study. Importantly, these criteria are not
intended to demonstrate therapeutic efficacy or clinical benefit.
Rather, they are designed to conceptually examine whether the
proposed visual mirror exposure protocol may be considered
ethically manageable and methodologically implementable under
controlled conditions in children with cerebral visual impairment
and an atelic profile, while enabling the potential acquisition of
interpretable non-behavioural physiological data. This structure
supports transparency, ethical oversight, and methodological
clarity at the protocol design stage, without implying any
obligation, initiation, or commitment to subsequent research.

4. Methods

4.1 Participants

The target population comprises children with cerebral visual impairment (CVI) presenting an atelic profile, defined as a persistent inability to initiate, maintain, or orient behaviour towards a goal, even in simple sensory contexts. This population is characterised as behaviourally non-responsive, which currently precludes participation in conventional visual assessment or rehabilitation protocols that rely on task engagement or observable performance.

No participants are recruited as part of the present manuscript. Any future empirical implementation of the proposed protocol would require prior approval from an appropriate Human Research Ethics Committee and would be conducted within an institutional framework.

Given the conceptual and pre-empirical scope of this work, no formal sample size calculation is performed. A minimum target sample of 10 participants is specified here illustratively, as a commonly accepted benchmark in feasibility study design, to indicate the scale at which protocol feasibility, tolerability, and measurement reliability could be evaluated in this rare and heterogeneous population using an intra-individual design.

Inclusion Criteria

265   In a prospective implementation scenario, participants would be

266   expected to meet the following criteria:

267   • A clinical diagnosis of cerebral visual impairment, established

268   by a qualified clinician and independent of ocular pathology;

269   • A behavioural profile consistent with atelicity, operationalised

270   as the absence of reliable goal-directed responses to visual or

271   task-based instructions;

272   • Age within the paediatric range appropriate for non-invasive

273   physiological monitoring, as would be determined by the host

274   institution's ethics committee;

275   • Medical stability permitting participation in brief passive visual

276   exposure sessions without contraindication;

277   • Informed consent would be obtained from a parent or legal

278   guardian.

279   **Exclusion Criteria**

280   In a future empirical context, participants would be expected to be

281   excluded if any of the following criteria were met:

282   • Uncontrolled epilepsy or recent seizure activity judged to pose

283   an increased risk during visual stimulation;

284   • Severe medical instability or acute illness at the time of

285   participation;

•   Known intolerance to screen-based or head-mounted visual

exposure;

•   Conditions preventing reliable acquisition of physiological

signals, such as severe movement disorders incompatible with

sensor placement;

•   Any clinical judgement indicating that participation would not

be in the child's best interest.

Ethical Justification

The ethical justification for the present protocol lies in its

methodological design rather than solely in the general

vulnerability of the target population. The proposed visual mirror

exposure paradigm is strictly passive and non-invasive, and does

not require instruction following, task engagement, or voluntary

behavioural responses. As such, it is specifically adapted to

children for whom conventional assessment or intervention

procedures are not ethically appropriate due to their limited

capacity to provide reliable behavioural assent.

Given the reduced capacity for verbal or behavioural

communication in behaviourally non-responsive children with an

atelic profile, the absence of overt response would not be

interpreted as consent. Participant protection would be

ensured through continuous monitoring of predefined

physiological and behavioural indicators of potential non-verbal

dissent. These indicators and their associated thresholds are

specified a priori and operationalised in Table 1 as part of a pre-

empirical ethical governance framework. In particular, session

termination would be triggered by predefined autonomic criteria

aligned with Table 1, including a sustained heart rate increase

beyond the 20% threshold relative to the individual baseline, a

persistent reduction in heart rate variability across consecutive

sessions, and/or the emergence of behavioural markers indicative

of distress or refusal.

Any detection of such indicators would result in immediate

termination of the session, in accordance with the predefined

stopping rules. This precautionary approach prioritises participant

safety and minimises risk exposure while allowing conceptual

evaluation of protocol feasibility under controlled conditions.

The present manuscript does not report or initiate a feasibility

study. It does not involve therapeutic intent, does not seek to

induce functional change, and does not withhold standard care. Its

purpose is limited to defining the ethical and methodological

conditions under which the protocol could be implemented safely

and tolerably in this population, based on explicit, predefined

safety thresholds, without implying any obligation or commitment

to subsequent research.

4.2 Setting and Equipment

In a future implementation, the visual mirror exposure

paradigm would be delivered in a controlled clinical or research

setting designed to minimise extraneous sensory stimulation and to

ensure participant safety. Sessions would take place in a quiet

room with reduced ambient noise and simplified visual

surroundings, in order to limit non-specific sensory load and

potential sources of distraction. The protocol is designed to be

implementable across clinical and research settings without

requiring site-specific adaptation.

Modes of Visual Presentation

Visual stimuli would be presented using one of two non-interactive

modalities, selected on the basis of participant tolerance, comfort,

and clinical judgement:

• A large screen positioned directly in front of the child, which

constitutes the default mode of visual presentation and would

be used whenever head-mounted devices are not tolerated or

are deemed inappropriate.

• A virtual reality headset, configured in a strictly passive, non-

interactive mode, providing full-field visual exposure and

reducing peripheral visual input. Use of a head-mounted device

would be optional and contingent upon individual tolerance; it

would be discontinued immediately if signs of discomfort or

non-verbal dissent are observed.

Both modalities are designed to deliver identical visual content and

temporal parameters. No interactive features, task demands, or

response contingencies are included in either modality. The

protocol is conceptually evaluated for feasibility across

presentation modalities rather than for equivalence or comparative

performance between modalities.

Controlled Conditions

Across both modes of visual presentation, the following

parameters would be controlled and standardised:

• Stimulus size, contrast, luminance, colour, and motion

characteristics.

• Duration of visual exposure blocks and inter-block intervals.

• Viewing distance in the screen condition, adjusted to promote

central visual exposure while limiting peripheral engagement.

• Absence of additional visual, auditory, or tactile stimulation during

exposure.

The purpose of this controlled environment is not to create an

immersive or engaging experience, but to ensure precise regulation

of visual input and to reduce non-specific sensory stimulation. This

configuration is intended to approximate a mirror-like condition in

which the child would be exposed to a single, isolated, and

predictable visual stimulus under conditions of minimal sensory

competition.

Equipment for Physiological Monitoring

Physiological data would be acquired using non-invasive sensors

suitable for paediatric populations. Monitoring would focuses on

autonomic measures, including heart rate and heart rate variability,

that could be recorded continuously during baseline, exposure, and

post-exposure periods.

All physiological monitoring devices would be required to be

certified for human use and comply with applicable regulatory

standards (e.g. CE marking and/or TGA or FDA approval,

depending on device availability and site requirements). Detailed

device specifications would be provided in supplementary

documentation or appendices where required by the reviewing

ethics committee.

Sensor placement and data acquisition procedures would be

selected to minimise discomfort and to allow reliable signal

recording without requiring voluntary movement, verbalisation, or

task engagement from the child. Signal quality would be assessed

in real time to identify artefacts resulting from excessive

movement, sensor displacement, or technical failure. Periods of

data affected by such artefacts would be flagged and excluded

from analysis according to predefined criteria.

Safety Monitoring and Personnel

In a future implementation scenario, continuous monitoring during

all sessions would be conducted by the researcher delivering the

protocol, who would be trained in the recognition of physiological

and behavioural indicators of distress in paediatric

neurodevelopmental populations, with immediate access to a

clinician affiliated with the host institution for safety escalation.

All personnel responsible for session monitoring would be required

to have received specific training in the recognition of

physiological and behavioural indicators of distress, including

autonomic changes, non-verbal dissent markers, and signs of

sensory overload in behaviourally non-responsive

children.Monitoring personnel would be authorised to interrupt or

terminate sessions immediately if predefined safety

thresholds were met or if any indication of discomfort or

distress were observed, in accordance with the predefined stopping

rules outlined in Table 1 and Section 4.5 as part of a pre-empirical

safety governance framework.

4.3 Visual Mirror Exposure Protocol

The visual mirror exposure protocol described in this manuscript consists of brief, repeated, and strictly controlled sessions of passive visual stimulation. The protocol is designed to expose the child to a simple, coherent, and highly predictable visual environment, without requiring instruction following, task engagement, or voluntary behavioural responses. It is specifically adapted to behaviourally non-responsive paediatric populations.

Session Structure

In a prospective implementation, each session would comprise the following sequential phases, delivered in a fixed and predefined order:

1. **Baseline Phase**

A resting baseline would be recorded prior to visual exposure. During this phase, no visual stimulus would be presented. Physiological signals would be recorded to establish the individual resting autonomic state.

2. **Stabilisation Phase**

A static, high-contrast visual stimulus would be presented against a uniform background. This phase is intended to allow exposure to a

stable visual signal without the need to process motion, change, or

novelty.

3. **Continuity Phase**

The visual stimulus would undergo slow, continuous, and regular

motion, such as linear translation at a constant velocity. No

acceleration, abrupt transitions, or unpredictable changes would be

introduced.

4. **Controlled Novelty Phase**

Limited and predictable modifications would be introduced to the

stimulus, such as gradual changes in colour, size, or trajectory.

These modifications would be predefined, infrequent, and

temporally constrained, and would be intended to probe autonomic

sensitivity while minimising the risk of sensory overload.

All phases would be delivered in a fixed sequence. No adaptive or

contingent modifications would be applied within a session.

Stimulus Parameters

Across all sessions, visual stimuli would be characterised by:

• A very limited number of objects presented simultaneously;

• High contrast between stimulus and background;

•   Slow and continuous motion when movement is present;

•   Absence of faces, social content, symbolic material, or rapid visual

events.

Stimulus parameters, including duration, luminance, contrast,

colour range, and motion characteristics, would be

predefined and would remain identical across participants, except

where minor adjustments might be required to ensure tolerability.

Temporal Progression

In a future empirical context, the protocol would follow a

predefined temporal progression across multiple weeks. Gradual

increases in perceptual complexity would be introduced only after

successful completion of earlier phases, as defined by the absence

of predefined physiological or behavioural stopping criteria.

Early sessions would prioritise static and highly predictable visual

stimuli. Subsequent sessions would introduce slow motion and,

later, controlled and limited novelty in a structured and non-

adaptive manner. This progression is intended to minimise

perceptual load and prediction error while allowing repeated

exposure under stable and controlled conditions.

The full session schedule, phase composition, and temporal

progression would   be   specified a   priori   and would   be

summarised in Table 2 as an illustrative implementation example.

Table 2. Illustrative Visual Mirror Exposure Protocol: Session Structure, Stimulus Parameters, and Temporal Progression
(Pre-Empirical Framework).

Phase	Visual content	Motion characteristics	Duration (minutes)	Predictability level	Purpose
Baseline	No stimulus	None	3	Not applicable	Illustratively intended to establish an individual autonomic resting state.
Stabilisation	Single high-contrast static object	None	4	Maximal	Illustratively intended to promote visual stability and reduce sensory load.
Continuity	Same object	Slow linear motion at constant velocity	4	High	Illustratively intended to support temporal predictability and sustained engagement.
Controlled novelty	Same object	Predefined gradual change (colour, size, or trajectory)	2	Moderate, predefined	Illustratively intended to probe autonomic sensitivity without sensory overload.

Week range	Phases included	Novelty introduction
Weeks 1 to 2	Baseline, Stabilisation	None
Weeks 3 to 4	Baseline, Stabilisation, Continuity	None
Weeks 5 to 6	All phases	Controlled novelty only

Note. The session structure, stimulus parameters, and temporal progression presented in this table are provided as illustrative, pre-

empirical examples intended to inform the methodological design of future feasibility studies. They do not correspond to sessions

delivered, participants exposed, or procedures implemented in an ongoing study.

Table 2 summarises the structure of the visual mirror exposure
protocol as an illustrative, pre-empirical framework, including
session phases, stimulus parameters, and temporal progression. All
durations, stimulus characteristics, and phase sequences are
specified a priori to illustrate protocol stability and safety
considerations, rather than to describe procedures applied in an
implemented study. The protocol is intentionally brief to minimise
physiological burden and to ensure tolerability in a behaviourally
non-responsive paediatric population. Temporal progression across
weeks is fixed at the design level and is not intended to adapt to
observed responses, except conceptually when predefined safety
stopping rules would be triggered in a future implementation. This
table provides the operational specification at the protocol design
stage required for methodological transparency and ethical
oversight.

Protocol Integrity

Within a prospective implementation, all sessions would be
delivered according to the predefined protocol without
modification based on observed physiological responses, except in
cases where stopping rules would be triggered. Any deviation from
the planned session structure, including early termination, would
be documented and considered in feasibility assessment.

4.4 Outcomes and Measures

The present manuscript does not report an implemented feasibility study and does not aim to assess therapeutic efficacy or functional visual improvement.

All outcomes are conceptual, methodological, and exploratory, and are selected to define feasibility, tolerability, and measurement reliability criteria relevant to the visual mirror paradigm in a behaviourally non-responsive paediatric population.

Outcome measures are limited to non-behavioural physiological signals, defined as autonomic physiological measures that could be recorded without requiring voluntary motor, verbal, or task-based responses from the child. Autonomic physiological measures are not interpreted as specific markers of visual processing, but as non-behavioural indicators of physiological regulation, tolerability, and phase-dependent sensitivity to controlled visual input. These measures are chosen to allow conceptual examination of physiological engagement and tolerability during passive visual exposure.

Primary Outcomes

The primary outcomes defined here relate to physiological tolerability of the protocol. Tolerability is operationalised as the

absence of sustained autonomic destabilisation indicative of distress during or following exposure. The following measures are identified as relevant indicators and would be recorded continuously throughout baseline, exposure, and post-exposure phases in a future empirical study:

- Heart rate, used to detect acute or sustained autonomic activation relative to individual baseline.
- Heart rate variability, used as an index of autonomic flexibility and parasympathetic regulation.

Predefined thresholds for physiological alert and stopping rules are specified at the design stage and detailed in Section 4.5.

Secondary Outcomes

Secondary outcomes are methodological and descriptive in nature. They include:

- The feasibility of acquiring continuous, interpretable physiological signals across all session phases.
- The presence or absence of non-verbal dissent markers, defined as behavioural or physiological signals that would be interpreted as refusal or discomfort, including abrupt increases in heart rate, sustained reduction in heart rate variability, sudden motor stiffening, agitation, or atypical vocalisations.

• The proportion of sessions that would be expected to be completed

as planned versus interrupted or terminated early due to safety

criteria.

• Intra-individual stability and variability of physiological signals

across repeated sessions.

No outcome relies on behavioural performance, visual choice, task

completion, or explicit communication from the child.

Measurement Windows

Physiological measures would be analysed within predefined

temporal windows corresponding to each session phase. Baseline

measures would serve as the individual reference state. Changes

observed during stabilisation, continuity, and controlled novelty

phases would be interpreted relative to this baseline, without

inference of cognitive or perceptual content.

Periods affected by signal artefact, defined as loss or distortion of

the physiological signal rendering data uninterpretable due to

excessive movement or sensor displacement, would be flagged and

excluded according to predefined criteria.

All outcomes and indicators are summarised in Table 3 as part of

the proposed methodological framework.

Table 3. Illustrative outcome domains, measures, indicators, and interpretation framework (pre-empirical).

Outcome domain	Measure	Phase(s)	Indicator	Interpretation purpose
Physiological Tolerability	HR	All phases	Stability relative to baseline	Illustratively intended to inform the assessment of autonomic tolerability of exposure without inference of therapeutic effect.
Physiological Tolerability	HRV	All phases	Maintenance of autonomic flexibility	Illustratively intended to inform the detection of physiological stress or destabilisation as safety indicators only.
Feasibility of Measurement	Signal Acquisition Quality	All phases	Proportion of usable signal	Illustratively intended to inform the feasibility of physiological monitoring in a non-responsive population.
Non-Verbal Dissent	Behavioural and Physiological Markers	All phases	Presence of predefined indicators	Illustratively intended to define conditions under which predefined stopping rules would be triggered as an ethical safeguard, not as an outcome measure.
Session Completion	Session Status	Whole session	Completed versus interrupted	Illustratively intended to inform the assessment of procedural feasibility independent of clinical change.
Measurement Reliability	Intra-individual signal patterns	Across sessions	Signal consistency	Illustratively intended to inform the evaluation of methodological robustness without group-level inference.

Note: HR: Heart Rate; HRV: Heart Rate Variability

The outcomes, measures, indicators, and interpretation purposes presented in this table are provided as illustrative, pre-empirical elements intended to inform the design and interpretation framework of future feasibility studies. They do not correspond to data collected, analyses performed, or outcomes evaluated in an implemented study.

Table 3 summarises the predefined outcomes, measures, and

indicators as part of a pre-empirical methodological

framework intended to inform the evaluation of feasibility,

tolerability, and measurement reliability in future feasibility

studies. All outcomes are specified a priori and are strictly non-

behavioural in nature. The structure is designed to conceptually

examine whether stable and interpretable physiological data could

be collected during passive visual mirror exposure, while ensuring

participant safety. These outcomes are not intended to support

claims regarding visual perception, cognitive processing, or

therapeutic efficacy, but to establish the methodological

preconditions required for future mechanistic or interventional

studies.

4.5 Safety and Stopping Rules

In a prospective implementation, participant safety would be

ensured through predefined physiological and behavioural stopping

rules, operationalised in Table 1 and applied as part of a pre-

empirical safety governance framework. The visual mirror

exposure protocol would be governed by these predefined safety

thresholds and stopping rules, which are designed to prioritise

physiological and behavioural indicators of potential discomfort or

distress in a behaviourally non-responsive paediatric population.

Safety Monitoring Procedures

In a future empirical context, safety monitoring would be conducted continuously during all session phases by trained personnel, with immediate authority to interrupt the session if predefined stopping criteria were met. Monitoring would be performed in real time throughout baseline, exposure, and post-exposure periods.

The absence of overt behavioural response would not be interpreted as consent or tolerance. Instead, safety assessment would be based on objective physiological changes and observable behavioural signs that may indicate non-verbal dissent.

Figure 1 provides an illustrative, pre-empirical example of how real-time physiological safety monitoring could be implemented during a single visual mirror exposure session, including the definition of an individual baseline and the application of predefined safety thresholds. This figure is provided for conceptual clarification only and does not represent empirical data.

Figure 1. Illustrative, pre-empirical example of real-time physiological safety monitoring during a single visual mirror exposure session (mock-up).

This illustrative mock-up depicts how physiological safety

monitoring could be implemented during a passive visual mirror

exposure session. The solid grey horizontal line represents the

individual resting heart rate baseline. The dashed red line indicates

the predefined safety threshold corresponding to a +20% increase

relative to baseline. The shaded area represents the predefined

physiological safety zone. The solid line represents a simulated

example of heart rate variation over time across session phases,

including a controlled novelty phase. In this illustrative scenario,

physiological modulation remains within predefined safety limits,

and no stopping criteria would be triggered. This figure is provided

for explanatory purposes only and does not represent empirical

data.

Physiological Stopping Criteria

In a prospective implementation, sessions would be immediately terminated if any of the following physiological criteria were met:

- A sudden and sustained increase in heart rate exceeding approximately 20% above the individual resting baseline, persisting beyond transient fluctuations;

- A marked and persistent reduction in heart rate variability relative to baseline levels, indicative of reduced autonomic flexibility;

- Any pattern of autonomic destabilisation suggesting sustained physiological activation rather than transient orienting responses.

These thresholds are defined as operational precautionary criteria, rather than diagnostic markers. They are specified a priori to prevent prolonged exposure to potential stress in the absence of subjective reporting.

Behavioural Stopping Criteria

In parallel with physiological monitoring, sessions would be terminated immediately if any of the following behavioural indicators were observed:

- Sudden motor stiffening or freezing not present at baseline;

- Marked agitation, repetitive movements, or abrupt changes in posture;

- Atypical vocalisations or changes in breathing pattern suggestive of distress;

- Any behavioural change that would be judged by the supervising clinician or researcher to reflect discomfort or refusal.

Behavioural indicators would be interpreted conservatively. When uncertainty exists, the decision rule would systematically favour immediate session termination.

Stopping Procedures and Documentation

If a stopping criterion were met, visual stimulation would be ceased immediately, and the child would be returned to a neutral resting state. No attempt would be made to resume the session on the same day.

All instances of early termination, including the triggering criterion and session phase, would be documented and would be incorporated into feasibility and tolerability considerations. Early termination would not be considered a protocol deviation but a predefined indicator relevant to feasibility assessment.

Ethical positioning

These stopping rules operationalise a precautionary ethical

governance approach in which participant protection supersedes

data completeness. These stopping procedures are not intended to

infer subjective emotional states, but to provide clear, auditable

documentation of discontinuation decisions based on observable

biological and behavioural signals.

The presence, frequency, and context of stopping events are

defined here as conceptual feasibility indicators, intended to

inform the ethical viability of future empirical investigations using

this paradigm.

4.6 Procedure timeline

In a future empirical context, the procedure would follow a

predefined and fixed timeline designed to evaluate feasibility,

tolerability, and measurement reliability of the visual mirror

paradigm across repeated exposures. The timeline is structured to

minimise sensory load, ensure participant safety, and

allow conceptual examination of intra-individual physiological

stability over time.

Each participant would follow the same temporal progression,

independent of physiological responses, except when predefined

stopping rules were triggered. No adaptive modification of

stimulus parameters would be performed based on interim

observations. No interim analyses or protocol adaptations would be

planned.

The protocol would be organised into three successive phases

spanning six weeks. Progression from one phase to the next would

follow a predefined fixed schedule and would not be adapted based

on observed responses, except in cases where predefined stopping

rules would result in session termination. Successful

completion would be defined retrospectively for feasibility

considerations and would not alter the planned timeline.

Early sessions would prioritise maximal predictability and

perceptual stability. Subsequent sessions would introduce

controlled and limited novelty in order to probe autonomic

sensitivity while maintaining a low cognitive and sensory burden.

All sessions would be scheduled with sufficient temporal spacing

to avoid cumulative fatigue or sensitisation effects. Session

frequency and duration would be identical across participants.

The full procedure timeline, including session content and

duration, is specified a priori at the design stage and is

summarised in Table 4 as an illustrative implementation example.

Table 4. Illustrative procedure timeline and visual mirror

exposure progression (pre-empirical framework).

Phase	Weeks	Session duration
Initial exposure (static stimuli)	Weeks 1–2	**10 to 15 minutes per session**
Predictable motion exposure	Weeks 3–4	**10 to 15 minutes per session**
Controlled novelty exposure	Weeks 5–6	**10 to 15 minutes per session**

Note. The procedure timeline and session durations presented in

this table are provided as illustrative, pre-empirical examples

intended to inform the design of future feasibility studies. They do

not correspond to sessions delivered or participants enrolled in an

implemented study.

Table 4 summarises the predefined temporal structure of the visual

mirror exposure protocol as an illustrative, pre-empirical

framework. Each phase is designed to incrementally modulate

perceptual complexity while preserving high stimulus

predictability.

Session durations are intentionally brief to reduce physiological

and sensory burden. The progression from static to dynamic

stimuli, followed by controlled novelty, reflects the theoretical aim

of gradually engaging visual predictive mechanisms without

inducing overload.

This timeline does not represent a therapeutic dose escalation and

is not intended to induce functional change. Rather, it is intended

to illustrate how protocol feasibility, tolerability, and physiological

signal stability could be examined within a predefined, feasibility-

focused design framework, prior to any empirical implementation.

5. Data Analysis Plan

This section outlines a predefined, pre-empirical descriptive

analysis framework intended to inform the evaluation of

feasibility, tolerability, and measurement reliability of the visual

mirror paradigm in future feasibility studies. The analyses focus

exclusively on non-behavioural physiological signals, predefined

temporal windows corresponding to protocol phases, and explicit

decision rules specified a priori. No hypothesis testing, efficacy

evaluation, or inferential analysis is proposed at this stage.

The complete data analysis plan, including signals, temporal

windows, indicators, and decision thresholds, is specified at the

protocol design stage and is summarised in Table 5 as an

illustrative analytical framework.

 Table 5. Illustrative data analysis framework: signals, temporal windows, indicators, and decision rules (pre-empirical).

Analysis domain	Physiological signal	Temporal window	Descriptive metric	Indicator	Decision rule
Baseline Physiological State	HR	Pre-exposure	Mean HR	Stable autonomic baseline	Baseline would be considered valid if the signal were continuous and artefact-free at the individual level.
Baseline Physiological State	HRV	Pre-exposure	Time-domain indices	Interpretable baseline	Baseline indices would be retained if they were computable at the individual level.
Exposure Tolerability	HR	Stabilisation	Mean and peak HR	Absence of sustained activation	Session continuation would be supported if HR were to remain within predefined thresholds at the individual level
Exposure Tolerability	HRV	Stabilisation	Qualitative pattern	Autonomic stability	Session interruption would be considered if a sustained reduction indicative of distress were to be observed.
Sensitivity to Controlled Novelty	HR	Controlled novelty phase	Peak and pattern relative to baseline	Autonomic reactivity	Data inclusion would be considered if signal quality were to meet the criteria specified in Table 1; session termination would be considered if predefined non-verbal dissent markers (Section 4.5) were to be detected.
Non-Verbal Dissent	Combined Signals	All phases	Presence of markers	Ethical safety	Immediate termination would be considered upon detection.
Measurement Reliability	HR and HRV	Across sessions	Intra-individual consistency	Signal stability	Descriptive assessment would be conducted at the intra-individual level across repeated sessions.
Feasibility	Procedural	Entire protocol	Completion rate	Implementability	Overall feasibility would be evaluated descriptively at the

Assessment	Indicators				protocol design level, using an illustrative feasibility sample size (e.g. n = 10), without group-level inference.

Note: HR: Heart Rate; HRV: Heart Rate Variability

The analysis domains, signals, temporal windows, indicators, and decision rules presented in this table are provided as illustrative,

pre-empirical elements intended to inform the analytical framework of future feasibility studies. They do not correspond to analyses

conducted, decisions applied, or data generated in an implemented study.

Table 5 summarises the predefined data analysis plan as part of a

pre-empirical methodological framework, linking each analytical

domain to specific physiological signals, temporal windows,

descriptive metrics, and explicit decision rules. In line with the

objectives of a feasibility-focused study, the table specifies how

physiological data could be examined descriptively at the

individual level, without hypothesis testing or group-level

inference. The decision rules outlined in the table define a priori

the conditions under which protocol feasibility, physiological

tolerability, and measurement reliability would be

considered achieved, partially achieved, or not achieved. By

specifying these analytical procedures at the protocol design stage,

the table ensures transparency, prevents analytical flexibility, and

establishes a clear interpretive structure for interpreting outcomes

strictly in terms of feasibility rather than therapeutic efficacy.

5.1 General Analytical Structure

This manuscript does not report an implemented feasibility study.

It is not designed to test therapeutic efficacy, causal effects, or

group-level hypotheses.

All analyses described here are pre-specified at the protocol design

level and are intended to inform future feasibility studies. They are

descriptive, exploratory, and strictly limited to the evaluation of

feasibility, tolerability, and measurement reliability of the visual

mirror paradigm.

Analyses would focus on intra-individual trajectories rather than

between-participant comparisons. Each child would serve as their

own reference, and no inferential statistics aimed at generalisation

would be planned.

No hypothesis testing, effect size estimation, or correction for

multiple comparisons is proposed.

5.2 Data Preprocessing and Signal Quality Control

Physiological signals would be visually and algorithmically

inspected prior to analysis to identify artefacts related to

movement, sensor displacement, or technical failure.

Segments affected by signal artefact would be flagged and

excluded according to predefined criteria. Only periods with

interpretable physiological data would be retained for analysis.

Baseline values would be computed individually for each

participant using resting-state segments recorded prior to visual

exposure in a future implementation.

5.3 Descriptive Analysis of Physiological Signals

Heart rate and heart rate variability would be summarised

descriptively within predefined temporal windows corresponding

to protocol phases.

Analyses would examine:

• baseline resting periods,

• stabilisation phases,

• continuity phases,

• controlled novelty phases.

For each participant, temporal evolution of physiological

signals would be visualised across sessions using within-subject

plots and time-series representations.

No aggregation across participants would be performed.

5.4 Assessment of Feasibility and Tolerability

Feasibility outcomes would be evaluated using procedural

indicators, including:

• number of sessions initiated,

• number of sessions completed,

• frequency and timing of early session termination,

• occurrence of protocol deviations.

Tolerability would be assessed through the absence or presence of

sustained physiological activation exceeding predefined thresholds,

and through documentation of non verbal dissent markers

triggering session interruption.

These outcomes would be summarised descriptively at the

individual level.

5.5 Assessment of Measurement Reliability

Measurement reliability would be examined by evaluating:

• stability of physiological signals across repeated sessions within

individuals,

• consistency of signal patterns across comparable protocol phases,

• proportion of sessions yielding usable physiological data.

Reliability assessment does not assume stationarity or normality of

signals and would remain limited to descriptive pattern

identification.

5.6 Interpretation Structure Specific to Feasibility Assessment

Interpretation would rely on within-session and within-individual

contrasts across predefined protocol phases, rather than on absolute

autonomic values.

Differential autonomic patterns would be used to characterise the

internal structure and interpretability of physiological signals.

They would not be used to infer magnitude of response, functional

visual processing, or therapeutic effect.

Outcomes would be interpreted as indicating:

• feasibility achieved,

• feasibility partially achieved,

• feasibility not achieved,

Based on the combination of protocol completion rates,

physiological tolerability, and signal interpretability at the

individual level.

The predefined target sample size of 10 participants is specified as

an illustrative feasibility objective, rather than a rigid requirement.

No post hoc modification of decision thresholds would be

permitted.

Importantly, the protocol would be considered feasible even if

physiological stabilisation reflects habituation rather than

engagement, provided predefined tolerability and signal

interpretability criteria are met.

5.7 Transparency and Reproducibility

All analysis procedures, decision rules, and feasibility criteria are

defined a priori at the protocol design stage, prior to any data

collection, to minimise analytical flexibility.

Any deviation from the planned analytical procedures would be

documented and reported alongside feasibility considerations in a

future implementation.

6. Ethics, Oversight, and Regulatory Considerations

This manuscript describes a clinical feasibility study protocol and

does not involve the collection, analysis, or reporting of empirical

data from human participants.

No study procedures have been initiated, and no participants have

been recruited.

The proposed visual mirror paradigm is intended for future

empirical implementation in a highly vulnerable paediatric

population. Consequently, ethical considerations are integrated at

the protocol design level, prior to any data collection. Formal

ethical approval will be sought from the appropriate Human

Research Ethics Committee at the host institution before the

commencement of any study involving human participants.

The protocol is designed in accordance with the precautionary

principle. The visual mirror exposure is strictly passive, non-

invasive, and does not require instruction following, task

engagement, or voluntary behavioural responses. The absence of
overt behavioural response is not interpreted as consent.
Participant protection relies on continuous monitoring of
predefined physiological and behavioural indicators of potential
non-verbal dissent, including autonomic destabilisation and
observable signs of distress. Any indication of predefined non-
verbal dissent would result in immediate termination of the
session, in accordance with the stopping rules specified in Section
4.5.

No therapeutic intent is associated with this protocol. The study
does not aim to induce functional change, does not replace or
withhold standard care, and does not involve deception,
randomisation, or experimental manipulation beyond controlled
visual exposure. The sole purpose of the proposed protocol is to
determine whether the paradigm can be implemented safely,
tolerably, and with sufficient measurement reliability to justify
consideration of future mechanistic or efficacy studies.

Data management, storage, and analysis procedures for any future
implementation will comply with institutional, national, and
international regulations governing research involving children and
vulnerable populations. In a future empirical implementation, the
protocol could be discontinued if predefined feasibility, safety, or
tolerability criteria are not met. Such an outcome would be

considered ethically appropriate and informative for future
research planning.

7. Discussion

The present manuscript describes a clinical feasibility study
protocol designed to evaluate a visual mirror paradigm in children
with cerebral visual impairment (CVI) presenting an atelic profile.
As no empirical data have been collected, this discussion does not
interpret outcomes or effects, but instead examines the
methodological validity, scope, and limits of the proposed
protocol. The primary methodological value of this protocol lies in
its explicit adaptation to a population for whom conventional
visual assessment or rehabilitation approaches are neither ethically
nor practically applicable. By relying exclusively on passive visual
exposure and non-behavioural autonomic physiological measures,
the protocol avoids any requirement for voluntary responses, task
engagement, or instruction following. This design directly
addresses a major methodological gap in the study of children with
severe CVI and atelic profiles, while remaining compatible with
stringent ethical constraints.

A central strength of the proposed approach is its exclusive focus
on feasibility, tolerability, and measurement reliability, rather than
on clinical efficacy or functional visual processing. The use of

930   predefined session phases and within-session, within-individual

931   contrasts allows evaluation of whether physiological signals

932   exhibit structured and interpretable organisation in response to

933   controlled visual input, without reliance on absolute autonomic

934   values. This structure is specifically intended to determine whether

935   the proposed paradigm is sufficiently stable, interpretable, and

936   ethically manageable to justify future empirical implementation.

937   The protocol also incorporates conservative safety monitoring and

938   stopping rules designed to protect participants who may be unable

939   to express discomfort through overt behavioural or verbal means.

940   By clearly distinguishing between analytic inclusion criteria and

941   ethical termination criteria, the design prioritises participant

942   protection while preserving methodological transparency. These

943   safeguards are essential when working with highly vulnerable

944   paediatric populations and constitute a core component of the

945   feasibility assessment.

946   Several limitations inherent to the proposed design must be

947   acknowledged. The absence of behavioural outcomes, group-level

948   comparisons, or functional performance measures necessarily

949   restricts the interpretive scope of the data that may be generated by

950   future implementation. In addition, autonomic physiological

951   signals are not interpreted as specific markers of visual processing,

but rather as indirect indicators of physiological regulation,

tolerability, and signal stability. These constraints are intentional

and reflect the ethical and practical boundaries associated with

research involving children with severe cerebral visual impairment

and atelic profiles, rather than methodological shortcomings.

The predefined feasibility, tolerability, and measurement reliability

criteria are intended to guide both the ethical conduct and

interpretation of the study. Should these criteria not be met, early

discontinuation of the protocol would constitute an ethically

appropriate and informative outcome. Conversely, if the criteria

are satisfied, the findings may inform future methodological or

clinical research considerations, without implying any obligation

or commitment to subsequent studies.

This discussion therefore addresses the validity and limits of the

proposed methodological approach itself, rather than interpreting

outcomes that have not yet been observed. The manuscript should

be read as a methodological protocol evaluated prior to data

collection, establishing clear ethical, procedural, and analytical

boundaries for any future empirical investigation.

Declarations

Ethics Statement: This manuscript presents a conceptual and

methodological protocol developed within an independent research

974	programme. No data involving human participants were collected,
975	generated, or analysed as part of the present work.
976	The proposed protocol is intended to inform future empirical
977	research involving vulnerable paediatric populations. Formal
978	ethical approval will be sought from the appropriate Human
979	Research Ethics Committee (HREC) at the host institution prior to
980	the commencement of any study involving human participants.
981	The protocol has been designed in accordance with the
982	precautionary principle. Given the limited capacity for verbal
983	assent in the target population, the absence of a behavioural
984	response is not interpreted as consent. The protocol explicitly
985	integrates physiological and behavioural indicators as markers of
986	potential non-verbal dissent, and any such indicators would lead to
987	immediate termination of the session in future implementations.
988	Generative AI tools were used during the preparation of the
989	manuscript to facilitate linguistic revision from French to academic
990	English. This tool was not used to generate scientific content,
991	interpret data, or draw conclusions. All content has been reviewed,
992	verified, and approved by the author.
993	**Data and Materials Availability**
994	No empirical data were collected for this study. The protocol
995	materials and methodological documentation have been archived

996 in a private repository and can be shared with reviewers or editors

997 upon request.

998 **Data Availability Statement**

999 Not applicable. No data were generated or analysed in the current

1000 study.

1001

1002 1. Smague E. Literature Review of the Influence of

1003 Neurovisual Disorders on the Diagnosis of Autism in Australian

1004 Children. Current Developmental Disorders Reports. 2024;12(1):3.

1005 2. Chokron S, Bernard JM, Imbert M. Length representation

1006 in normal and neglect subjects with opposite reading habits studied

1007 through a line extension task. Cortex. 1997;33(1):47–64.

1008 3. Hacohen A, editor The psychological reality of the

1009 telic/atelic distinction: Evidence from adult and child Hebrew.

1010 Proceedings of the 21st Annual Meeting of the Israel Associations

1011 for Theoretical Linguistics,(unnumbered); 2006.

1012 4. Derrien D, Garric C, Sergent C, Chokron S. The nature of

1013 blindsight: implications for current theories of consciousness.

1014 Neurosci Conscious. 2022;2022(1):niab043.

1015 5. Friston K, Mattout J, Kilner J. Action understanding and

1016 active inference. Biol Cybern. 2011;104(1-2):137–60.

1017 6. Dutton GN, Jacobson LK. Cerebral visual impairment in

1018 children. Semin Neonatol. 2001;6(6):477–85.

7.   Friston K. The free-energy principle: a unified brain
theory? Nat Rev Neurosci. 2010;11(2):127–38.

8.   Ramachandran VS, Rogers-Ramachandran D. Synaesthesia
in phantom limbs induced with mirrors. Proc Biol Sci.
1996;263(1369):377–86.

9.   Bach-y-Rita P, S WK. Sensory substitution and the human-
machine interface. Trends Cogn Sci. 2003;7(12):541–6.

1026