Visual Mirror Exposure for Children with Cerebral Visual Impairment and an Atelic Profile: A Clinical Feasibility Study Protocol.

Author

Edwige Smague[1]

Affiliations:

[1]N.E.T.A: Neuronal Epileptotranscriptomic, Sydney, Australia

Contact:

edwige@netadiscovery.org

ORCID ID: 0009-0002-8182-7888

Abstract

Children with cerebral visual impairment (CVI) presenting an atelic profile constitute a population that is largely excluded from neurovisual research, primarily due to the absence of reliable behavioural responses to standard assessment or intervention paradigms. This exclusion often leads to the assumption that a lack of observable response reflects an absence of visual processing, despite converging evidence from neuroscience indicating that neural processing may persist in the absence of overt behavioural expression.

This protocol describes a clinical feasibility study designed to evaluate the feasibility, tolerability, and measurement reliability of a passive visual mirror exposure paradigm in children with CVI

and an atelic profile. The visual mirror is defined as the repeated

presentation of a highly predictable, coherent visual experience,

intended to reduce perceptual instability and cognitive load

associated with typical visual environments, without aiming to

restore or train visual function.

The paradigm relies on passive visual exposure delivered via a

large screen, or, where appropriate and tolerated, a virtual reality

headset configured in a strictly non-interactive mode, without

requiring effort, task engagement, or voluntary responses from the

child. The study is embedded within a precautionary ethical

framework specifically adapted to behaviourally non-responsive

paediatric populations, defined here as children presenting absent

voluntary behavioural responses to sensory or task-based

instructions.

Feasibility and tolerability are assessed using predefined safety

criteria and non-verbal physiological indicators, including heart

rate and autonomic markers, rather than behavioural performance.

This feasibility study is designed to characterise the ethical

acceptability, implementability, and physiological tolerability of

the protocol in this vulnerable population, and to inform future

research considerations without implying any obligation or

commitment to subsequent studies.

Keywords: Atelic profile; cerebral visual impairment (CVI); cerebral plasticity; visual mirror exposure; non-verbal children; passive virtual reality; vision rehabilitation.

1. Introduction

Cerebral visual impairment (CVI) (1) is now recognised as one of the leading causes of paediatric visual disability. It results from damage to or dysfunction of the cerebral pathways involved in visual processing, independent of ocular integrity (2). Visual difficulties associated with CVI are heterogeneous and may include impairments in gaze orienting, motion perception, object recognition, sustained visual attention, or visuomotor coordination.

Within this population, children presenting an atelic profile constitute a particularly marginalised group. Atelicity is defined here as a persistent inability to initiate, maintain, or orient behaviour towards a goal, even in simple sensory contexts (3). Children with atelic profiles are typically unable to follow instructions or produce reliable behavioural responses, which effectively excludes them from conventional visual assessment and rehabilitation paradigms that rely on task engagement or observable performance.

This exclusion has created a significant methodological and ethical gap. The absence of empirical data in this subgroup makes it

difficult to determine whether these children are genuinely non-
responsive to visual stimuli or whether they are simply
inaccessible to existing assessment methods. In this context,
behavioural non-responsiveness cannot be assumed to reflect an
absence of visual processing, as converging evidence from
neuroscience indicates that perceptual and neural processing may
persist in the absence of overt behavioural expression.

The present study addresses this gap by proposing a feasibility-
focused methodological approach designed to evaluate
physiological responses to controlled visual stimuli in the absence
of voluntary behaviour. Rather than aiming to assess therapeutic
efficacy, clinical benefit, or functional visual improvement, the
protocol is explicitly limited to evaluating whether a passive visual
mirror exposure paradigm can be implemented safely and tolerably
in children with CVI and an atelic profile, while allowing the
acquisition of stable, non-behavioural physiological signals.

The primary objective of this feasibility study is to determine
whether repeated sessions of passive visual mirror exposure can be
delivered under controlled conditions without inducing sustained
physiological distress, and while enabling the collection of
interpretable autonomic markers. Accordingly, the primary
feasibility question addressed in this study is as follows: Is it

feasible to deliver repeated passive visual mirror exposure sessions

to children with CVI and an atelic profile while maintaining

physiological tolerability in behaviourally non-responsive

paediatric populations?

2. The Atelic Child and Visual Experience

2.1 Atelicity as a Disruption of the Perception–Prediction-
Action Loop

In children presenting an atelic profile, the absence of goal-

directed behaviour should not be interpreted as an absence of

perception. Evidence from cognitive and neuroscientific research

indicates that perceptual and memory-related processing can occur

in the absence of voluntary control or overt behavioural expression

(4). Accordingly, limited behavioural responsiveness does not

preclude the presence of underlying sensory processing.

Atelicity can be described as a functional difficulty in initiating,

maintaining, or organising behaviour in response to sensory

information. In this context, the child may receive sensory input

but be unable to use it reliably to guide action, exploration, or

sustained engagement with the environment (5). This description

does not imply a deficit in sensory reception itself, but rather a

disruption in the translation of sensory information into observable, goal-directed behaviour.

2.2 Visual Experience in Atelic Children with Cerebral Visual Impairment

In atelic children with cerebral visual impairment, visual experience is often reported by clinicians and caregivers as unstable and difficult to interpret (6). Commonly described features include:

- **Image Instability**: where the visual scene may appear fragmented or inconsistent;
- **Excessive Perceptual Detail**: resulting in difficulty prioritising or organising visual information;
- **Difficulty Anticipating Visual Change**: such that new objects or changes in lighting may be experienced as abrupt or unpredictable;
- **Impaired Recognition of Visual Stability**: for example, difficulty recognising that an object remains the same despite minor changes in position or context.

At a functional level, these characteristics are associated with increased perceptual uncertainty and elevated cognitive load during visual processing. Over time, visual engagement may become effortful or aversive, leading to reduced spontaneous

visual exploration or apparent disengagement from visual stimuli

(7).

This section provides a conceptual and descriptive framework to

support the design choices of the proposed protocol. It does not

constitute a testable theoretical model and does not imply specific

mechanisms of impairment. Rather, it serves to justify the use of

simplified, predictable, and non-interactive visual stimuli in a

feasibility-focused study involving behaviourally non-responsive

paediatric populations.

3. Research Question and Objectives

3.1 Primary Feasibility Question

The visual mirror exposure paradigm evaluated in this study is

conceptually informed by principles derived from mirror-based

approaches originally developed in post-stroke motor rehabilitation

(8), while differing fundamentally in its intended function and

mechanism. Rather than relying on visual comparison or illusion,

the present paradigm is designed to provide a coherent, stable, and

highly predictable visual stimulus that may temporarily substitute

for unstable or cognitively costly visual environments in children

with cerebral visual impairment, consistent with sensory

substitution principles (9) and descriptive models of predictive

processing (7).

In the context of cerebral visual impairment and atelic profiles, the

paradigm is explicitly designed for behaviourally non-responsive

paediatric populations and does not require task engagement,

instruction following, or voluntary behavioural responses.

The primary feasibility question addressed in this study is therefore

as follows: Is it feasible to deliver repeated sessions of passive

visual mirror exposure to children with cerebral visual impairment

and an atelic profile while maintaining physiological tolerability

and obtaining stable, interpretable non-behavioural physiological

signals in behaviourally non-responsive paediatric populations?

3.2 Primary Objective

The primary objective of this clinical feasibility study is to

evaluate the feasibility and physiological tolerability of a passive

visual mirror exposure protocol in children with cerebral visual

impairment and an atelic profile. Feasibility is defined in terms of

the ability to implement the protocol as planned, without inducing

sustained physiological distress, and while enabling the acquisition

of usable autonomic physiological measures in a population for

whom active, task-based paradigms are not accessible (7).

3.3 Secondary Methodological Objectives

The secondary objectives of this study are strictly methodological and exploratory. They are intended to characterise protocol performance and measurement properties rather than to assess therapeutic efficacy or clinical benefit. Specifically, the objectives are:

- to assess whether non-behavioural physiological measures, including heart rate and heart rate variability, can be reliably recorded during passive visual mirror exposure sessions in this population;

- to examine the intra-individual stability and variability of these physiological signals across repeated sessions under conditions of highly controlled and predictable visual stimulation (7);

- to document rates of session completion, interruption, or early termination due to indicators of non-verbal dissent or physiological distress, in accordance with precautionary ethical principles;

- to explore whether protocol tolerability varies across individual neurovisual profiles within the atelic population, without making claims of group-level inference.

3.4 Reference to Feasibility, Tolerability and Safety Criteria

To ensure the highest standard of participant safety, feasibility and

tolerability criteria are defined a priori. These predefined

thresholds and decision rules, detailed in Table 1, constitute a

mandatory safety and governance framework for protocol

management.

Their primary purpose is to characterise the reliability, tolerability,

and ethical acceptability of the protocol in children with atelic

profiles, and to ensure that study conduct remains grounded in

continuous monitoring of participant well-being and protocol

stability. These criteria are not intended to demonstrate therapeutic

efficacy but to guide ethical conduct and interpretation within a

feasibility-focused   study.

Table 1. Feasibility, tolerability, and safety indicators with predefined decision thresholds.

Category	Indicator	Feasibility Thresholds and Interpretation
Protocol Implementation	Session Completion Rate	≥ 75% of planned sessions are completed without premature termination across a minimum sample of 10 participants.
Technical Reliability	Usable Physiological	≥ 80% of sessions yield usable HR and HRV data (defined as < 20% signal artefact) at the individual level.

206   Note. HR: Heart Rate; HRV: Heart Rate Variability.
207

	Data	
Physiological Tolerability	Acute Distress Signal	HR remains within < 20% increase from individual baseline for at least 90% of exposure time for each participant.
Autonomic Stability	HRV Stability	Absence of a persistent decrease in HRV sustained over 3 consecutive sessions within participant.
Ethical Safety	Dissent Response	100% compliance with the immediate termination upon detection of non-verbal dissent markers.
Attrition	Participant Withdrawal	Feasibility is supported if participant withdrawal due to intolerance does not exceed 20% (i.e. no more than 2 out of 10 participants).
Informational Potential	Signal Sensitivity	Presence of measurable physiological modulation during controlled novelty phases in $\geq$ 50% of participants within the predefined feasibility sample.

Table 1 summarises the primary feasibility question, objectives,

indicators, and predefined feasibility and tolerability criteria for

this clinical feasibility study. All criteria are specified a priori, prior

to data collection, to guide ethical conduct, safety monitoring, and

interpretation of the study outcomes. The table links each research

objective to concrete physiological or procedural indicators and to

explicit decision rules governing protocol continuation or early

termination. Importantly, these criteria are not intended to

demonstrate therapeutic efficacy or clinical benefit. Rather, they

are designed to determine whether the proposed visual mirror

exposure protocol can be implemented safely and tolerably under

controlled conditions in children with cerebral visual impairment

and an atelic profile, while enabling the acquisition of interpretable

non-behavioural physiological data. This structure supports

transparency, ethical oversight, and methodological clarity within a

feasibility-focused study, without implying any obligation or

commitment to subsequent research.

4. Methods

4.1 Participants

The target population comprises children with cerebral visual

impairment (CVI) presenting an atelic profile, defined as a

persistent inability to initiate, maintain, or orient behaviour

towards a goal, even in simple sensory contexts. This population is characterised as behaviourally non-responsive, which currently precludes participation in conventional visual assessment or rehabilitation protocols that rely on task engagement or observable performance.

Participants will be recruited prospectively following approval from the appropriate Human Research Ethics Committee, using the present protocol without modification.

Given the exploratory and feasibility-focused nature of this clinical feasibility study, no formal sample size calculation is performed. A minimum target sample of 10 participants is specified a priori as sufficient to evaluate protocol feasibility, tolerability, and measurement reliability in this rare and heterogeneous population, using an intra-individual design.

Inclusion Criteria

Participants must meet the following criteria:

- a clinical diagnosis of cerebral visual impairment, established by a qualified clinician and independent of ocular pathology;
- a behavioural profile consistent with atelicity, operationalised as the absence of reliable goal-directed responses to visual or task-based instructions;

• age within the paediatric range appropriate for non-invasive

physiological monitoring, as determined by the host

institution's ethics committee;

• medical stability permitting participation in brief passive visual

exposure sessions without contraindication;

• informed consent obtained from a parent or legal guardian.

Exclusion Criteria

Participants meeting any of the following criteria will be excluded:

• uncontrolled epilepsy or recent seizure activity judged to pose

an increased risk during visual stimulation;

• severe medical instability or acute illness at the time of

participation;

• known intolerance to screen-based or head-mounted visual

exposure;

• conditions preventing reliable acquisition of physiological

signals, such as severe movement disorders incompatible with

sensor placement;

• any clinical judgement indicating that participation would not

be in the child's best interest.

Ethical Justification

The ethical justification for the present protocol lies in its

methodological design rather than solely in the general

vulnerability of the target population. The proposed visual mirror

exposure paradigm is strictly passive and non-invasive, and does

not require instruction following, task engagement, or voluntary

behavioural responses. As such, it is specifically adapted to

children for whom conventional assessment or intervention

procedures are not ethically appropriate due to their limited

capacity to provide reliable behavioural assent.

Given the reduced capacity for verbal or behavioural

communication in behaviourally non-responsive children with an

atelic profile, the absence of overt response is not interpreted as

consent. Participant protection is ensured through continuous

monitoring of predefined physiological and behavioural indicators

of potential non-verbal dissent. These indicators and their

associated thresholds are specified a priori and operationalised in

Table 1. In particular, session termination is triggered by

predefined autonomic criteria aligned with Table 1, including a

sustained heart rate increase beyond the 20% threshold relative to

the individual baseline, a persistent reduction in heart rate

variability across consecutive sessions, and/or the emergence of

behavioural markers indicative of distress or refusal.

Any detection of such indicators results in immediate termination

of the session, in accordance with the predefined stopping rules.

This precautionary approach prioritises participant safety and

minimises risk exposure while allowing evaluation of protocol

feasibility under controlled conditions.

This feasibility study does not involve therapeutic intent, does not

seek to induce functional change, and does not withhold standard

care. Its purpose is limited to determining whether the protocol can

be implemented safely and tolerably in this population, based on

explicit, predefined safety thresholds, without implying any

obligation or commitment to subsequent research.

4.2 Setting and Equipment

The visual mirror exposure paradigm is delivered in a controlled

clinical or research setting designed to minimise extraneous

sensory stimulation and to ensure participant safety. All sessions

take place in a quiet room with reduced ambient noise and

simplified visual surroundings, in order to limit non-specific

sensory load and potential sources of distraction. The protocol is

designed to be implementable across clinical and research settings

without requiring site-specific adaptation.

Modes of Visual Presentation

Visual stimuli are presented using one of two non-interactive

modalities, selected on the basis of participant tolerance, comfort,

and clinical judgement:

• A large screen positioned directly in front of the child, which

constitutes the default mode of visual presentation and is used

whenever head-mounted devices are not tolerated or are

deemed inappropriate.

• A virtual reality headset, configured in a strictly passive, non-

interactive mode, providing full-field visual exposure and

reducing peripheral visual input. Use of a head-mounted device

is optional and contingent upon individual tolerance; it is

discontinued immediately if signs of discomfort or non-verbal

dissent are observed.

Both modalities are designed to deliver identical visual content and

temporal parameters. No interactive features, task demands, or

response contingencies are included in either modality. The

protocol is evaluated for feasibility across presentation modalities

rather than for equivalence or comparative performance between

modalities.

Controlled Conditions

Across both modes of visual presentation, the following

parameters are controlled and standardised:

- Stimulus size, contrast, luminance, colour, and motion characteristics.
- Duration of visual exposure blocks and inter-block intervals.
- Viewing distance in the screen condition, adjusted to promote central visual exposure while limiting peripheral engagement.
- Absence of additional visual, auditory, or tactile stimulation during exposure.

The purpose of this controlled environment is not to create an immersive or engaging experience, but to ensure precise regulation of visual input and to reduce non-specific sensory stimulation. This configuration is intended to approximate a mirror-like condition in which the child is exposed to a single, isolated, and predictable visual stimulus under conditions of minimal sensory competition.

Equipment for Physiological Monitoring

Physiological data are acquired using non-invasive sensors suitable for paediatric populations. Monitoring focuses on autonomic measures, including heart rate and heart rate variability, recorded continuously during baseline, exposure, and post-exposure periods. All physiological monitoring devices used in the study are certified for human use and comply with applicable regulatory standards (e.g. CE marking and/or TGA or FDA approval, depending on device availability and site requirements). Detailed device

specifications are provided in supplementary documentation or

appendices where required by the reviewing ethics committee.

Sensor placement and data acquisition procedures are selected to

minimise discomfort and to allow reliable signal recording without

requiring voluntary movement, verbalisation, or task engagement

from the child. Signal quality is assessed in real time to identify

artefacts resulting from excessive movement, sensor displacement,

or technical failure. Periods of data affected by such artefacts are

flagged and excluded from analysis according to predefined

criteria.

Safety Monitoring and Personnel

Continuous monitoring during all sessions will be conducted by

the researcher delivering the protocol, who is trained in the

recognition of physiological and behavioural indicators of distress

in paediatric neurodevelopmental populations, with immediate

access to a clinician affiliated with the host institution for safety

escalation. All personnel responsible for session monitoring have

received specific training in the recognition of physiological and

behavioural indicators of distress, including autonomic changes,

non-verbal dissent markers, and signs of sensory overload in

behaviourally non-responsive children.

380   Monitoring personnel are authorised to interrupt or terminate

381   sessions immediately if predefined safety thresholds are met or if

382   any indication of discomfort or distress is observed, in accordance

383   with the predefined stopping rules outlined in Table 1 and Section

384   4.5.

385   **4.3 Visual Mirror Exposure Protocol**

386   The visual mirror exposure protocol consists of brief, repeated, and

387   strictly controlled sessions of passive visual stimulation. The

388   protocol is designed to expose the child to a simple, coherent, and

389   highly predictable visual environment, without requiring

390   instruction following, task engagement, or voluntary behavioural

391   responses. It is specifically adapted to behaviourally non-

392   responsive paediatric populations.

393   **Session Structure**

394   Each session comprises the following sequential phases, delivered

395   in a fixed and predefined order:

396   1. **Baseline Phase**

397   A resting baseline is recorded prior to visual exposure. During this

398   phase, no visual stimulus is presented. Physiological signals are

399   recorded to establish the individual resting autonomic state.

2. **Stabilisation Phase**

A static, high contrast visual stimulus is presented against a uniform background. This phase is intended to allow exposure to a stable visual signal without the need to process motion, change or novelty.

3. **Continuity Phase**

The visual stimulus undergoes slow, continuous, and regular motion, such as linear translation at a constant velocity. No acceleration, abrupt transitions, or unpredictable changes are introduced.

4. **Controlled Novelty Phase**

Limited and predictable modifications are introduced to the stimulus, such as gradual changes in colour, size, or trajectory. These modifications are predefined, infrequent and temporally constrained, and are intended to probe autonomic sensitivity while minimising the risk of sensory overload.

All phases are delivered in a fixed sequence. No adaptive or contingent modifications are applied within a session.

Stimulus Parameters

Across all sessions, visual stimuli are characterised by:

• A very limited number of objects presented simultaneously;

• High contrast between stimulus and background;

• Slow and continuous motion when movement is present;

• Absence of faces, social content, symbolic material, or rapid visual

events.

Stimulus parameters, including duration, luminance, contrast,

colour range, and motion characteristics, are predefined and remain

identical across participants, except where minor adjustments are

required to ensure tolerability.

Temporal Progression

The protocol follows a predefined temporal progression across

multiple weeks. Gradual increases in perceptual complexity are

introduced only after successful completion of earlier phases, as

defined by the absence of predefined physiological or behavioural

stopping criteria.

Early sessions prioritise static and highly predictable visual

stimuli. Subsequent sessions introduce slow motion and, later,

controlled and limited novelty in a structured and non-adaptive

manner. This progression is intended to minimise perceptual load

and prediction error while allowing repeated exposure under stable

and controlled conditions.

The full session schedule, phase composition, and temporal

progression are specified a priori and summarised in Table 2.

Table 2. Visual Mirror Exposure Protocol: session structure, stimulus parameters, and temporal progression.

Phase	Visual content	Motion characteristics	Duration (minutes)	Predictability level	Purpose
Baseline	No stimulus	None	3	Not applicable	Establish individual autonomic resting state
Stabilisation	Single high-contrast static object	None	4	Maximal	Promote visual stability and reduce sensory load
Continuity	Same object	Slow linear motion at constant velocity	4	High	Support temporal predictability and sustained engagement
Controlled novelty	Same object	Predefined gradual change (colour, size, or trajectory)	2	Moderate, predefined	Probe autonomic sensitivity without sensory overload

Week range	Phases included	Novelty introduction
Weeks 1 to 2	Baseline, Stabilisation	None
Weeks 3 to 4	Baseline, Stabilisation, Continuity	None
Weeks 5 to 6	All phases	Controlled novelty only

Table 2 summarises the structure of the visual mirror exposure
protocol, including session phases, stimulus parameters, and
temporal progression. All durations, stimulus characteristics, and
phase sequences are predefined a priori to ensure protocol stability
and participant safety. The protocol is intentionally brief to
minimise physiological burden and to ensure tolerability in a
behaviourally non-responsive paediatric population. Temporal
progression across weeks is fixed and does not adapt to observed
responses, except when predefined safety stopping rules are
triggered. This table provides the operational specification required
for protocol replication and ethical oversight.

Protocol Integrity

All sessions are delivered according to the predefined protocol
without modification based on observed physiological responses,
except in cases where stopping rules are triggered. Any deviation
from the planned session structure, including early termination, is
documented and considered in feasibility assessment.

4.4 Outcomes and Measures

The present feasibility study does not aim to assess therapeutic
efficacy or functional visual improvement.

All outcomes are strictly methodological and exploratory, and are selected to evaluate feasibility, tolerability, and measurement reliability of the visual mirror paradigm in a behaviourally non-responsive paediatric population.

Outcome measures are limited to non-behavioural physiological signals, defined as autonomic physiological measures recorded without requiring voluntary motor, verbal, or task-based responses from the child. Autonomic physiological measures are not interpreted as specific markers of visual processing, but as non-behavioural indicators of physiological regulation, tolerability, and phase-dependent sensitivity to controlled visual input. These measures are chosen to allow indirect assessment of physiological engagement and tolerability during passive visual exposure.

Primary Outcomes

The primary outcomes relate to physiological tolerability of the protocol. Tolerability is operationalised as the absence of sustained autonomic destabilisation indicative of distress during or following exposure. The following measures are recorded continuously throughout baseline, exposure, and post-exposure phases:

- Heart rate, used to detect acute or sustained autonomic activation relative to individual baseline.

•   Heart rate variability, used as an index of autonomic flexibility and

parasympathetic regulation.

Predefined thresholds for physiological alert and stopping rules are

specified *a priori* and detailed in Section 4.5.

Secondary Outcomes

Secondary outcomes are methodological and descriptive in nature.

They include:

•   The feasibility of acquiring continuous, interpretable physiological

signals across all session phases.

•   The presence or absence of non-verbal dissent markers, defined as

behavioural or physiological signals interpreted as refusal or

discomfort, including abrupt increases in heart rate, sustained

reduction in heart rate variability, sudden motor stiffening,

agitation, or atypical vocalisations.

•   The proportion of sessions completed as planned versus interrupted

or terminated early due to safety criteria.

•   Intra-individual stability and variability of physiological signals

across repeated sessions.

No outcome relies on behavioural performance, visual choice, task

completion, or explicit communication from the child.

Measurement Windows

Physiological measures are analysed within predefined temporal windows corresponding to each session phase. Baseline measures serve as the individual reference state. Changes observed during stabilisation, continuity, and controlled novelty phases are interpreted relative to this baseline, without inference of cognitive or perceptual content.

Periods affected by signal artefact, defined as loss or distortion of the physiological signal rendering data uninterpretable due to excessive movement or sensor displacement, are flagged and excluded according to predefined criteria.

All outcomes and indicators are summarised in Table 3.

519 **Table 3. Outcomes, Measures, Indicators, and Interpretation Structure**

Outcome domain	Measure	Phase(s)	Indicator	Interpretation purpose
Physiological Tolerability	HR	All phases	Stability relative to baseline	Assess autonomic tolerability of exposure without inference of therapeutic effect.
Physiological Tolerability	HRV	All phases	Maintenance of autonomic flexibility	Detect physiological stress or destabilisation as safety indicators only.
Feasibility of Measurement	Signal Acquisition Quality	All phases	Proportion of usable signal	Determine feasibility of physiological monitoring in a non-responsive population.
Non-Verbal Dissent	Behavioural and Physiological Markers	All phases	Presence of predefined indicators	Trigger predefined stopping rules as an ethical safeguard, not as an outcome measure.
Session Completion	Session Status	Whole session	Completed versus interrupted	Assess procedural feasibility independent of clinical change
Measurement Reliability	Intra-individual signal patterns	Across sessions	Signal consistency	Evaluate methodological robustness without group-level inference.

Note: HR: Heart Rate; HRV: Heart Rate Variability

Table 3 summarises the predefined outcomes, measures, and

indicators used to evaluate feasibility, tolerability, and

measurement reliability in this clinical feasibility study. All

outcomes are specified *a priori* and are strictly non-behavioural in

nature. The structure is designed to determine whether stable and

interpretable physiological data can be collected during passive

visual mirror exposure, while ensuring participant safety. These

outcomes are not intended to support claims regarding visual

perception, cognitive processing, or therapeutic efficacy, but to

establish the methodological preconditions required for future

mechanistic or interventional studies.

4.5 Safety and Stopping Rules

Participant safety is ensured through predefined physiological and

behavioural stopping rules, operationalised in Table 1 and applied

uniformly across all sessions. The visual mirror exposure protocol

is governed by these predefined safety thresholds and stopping

rules, which are designed to prioritise physiological and

behavioural indicators of potential discomfort or distress in a

behaviourally non-responsive paediatric population.

Safety Monitoring Procedures

Safety monitoring is conducted continuously during all session

phases by trained personnel, with immediate authority to interrupt

the session if predefined stopping criteria are met. Monitoring is

performed in real time throughout baseline, exposure, and post-

exposure periods.

The absence of overt behavioural response is not interpreted as

consent or tolerance. Instead, safety assessment is based on

objective physiological changes and observable behavioural signs

that may indicate non-verbal dissent.

Figure 1 provides an illustrative example of how real-time

physiological safety monitoring is implemented during a single

visual mirror exposure session, including the definition of an

individual baseline and the application of predefined safety

thresholds. This figure is provided for conceptual clarification only

and does not represent empirical data.

Figure 1. Illustrative example of real-time physiological safety
monitoring during a single visual mirror exposure session
(mock-up).

This illustrative mock-up shows how physiological safety is

monitored during a passive visual mirror exposure session. The

solid grey horizontal line represents the individual resting heart

rate baseline. The dashed red line indicates the predefined safety

threshold corresponding to a +20% increase relative to baseline.

The shaded area represents the predefined physiological safety

zone. The solid line represents heart rate variation over time across

session phases, including a controlled novelty phase. In this

example, physiological modulation remains within predefined

safety limits, and no stopping criteria are triggered. This figure is

provided for explanatory purposes only and does not represent

empirical data.

Physiological Stopping Criteria

Sessions are immediately terminated if any of the following

physiological criteria are met:

• a sudden and sustained increase in heart rate exceeding

approximately 20% above the individual resting baseline,

persisting beyond transient fluctuations;

• a marked and persistent reduction in heart rate variability

relative to baseline levels, indicative of reduced autonomic

flexibility;

• any pattern of autonomic destabilisation suggesting

sustained physiological activation rather than transient

orienting responses.

These thresholds are operational precautionary criteria rather than

diagnostic markers. They are defined a priori to prevent prolonged

exposure to potential stress in the absence of subjective reporting.

Behavioural Stopping Criteria

In parallel with physiological monitoring, sessions are terminated

immediately if any of the following behavioural indicators are

observed:

• sudden motor stiffening or freezing not present at baseline.

• Marked agitation, repetitive movements, or abrupt changes in

posture;

• atypical vocalisations or changes in breathing pattern suggestive of

distress;

•   any behavioural change judged by the supervising clinician or

researcher to reflect discomfort or refusal.

Behavioural indicators are interpreted conservatively. When

uncertainty exists, the decision rule systematically favours

immediate session termination.

Stopping Procedures and Documentation

If a stopping criterion is met, visual stimulation is ceased

immediately, and the child is returned to a neutral resting state. No

attempt is made to resume the session on the same day.

All instances of early termination, including the triggering criterion

and session phase, are systematically documented and incorporated

into feasibility and tolerability analyses. Early termination is not

considered a protocol deviation but a predefined outcome relevant

to feasibility assessment.

Ethical positioning

These stopping rules operationalise a precautionary ethical

governance approach in which participant protection supersedes

data completeness. These stopping procedures are not intended to

infer subjective emotional states, but to provide clear, auditable

documentation of discontinuation decisions based on observable

biological and behavioural signals.

The presence, frequency, and context of stopping events constitute

key feasibility outcomes in this feasibility study and directly

inform the ethical viability of future empirical investigations using

this paradigm.

4.6 Procedure timeline

The procedure follows a predefined and fixed timeline designed to

evaluate feasibility, tolerability, and measurement reliability of the

visual mirror paradigm across repeated exposures. The timeline is

structured to minimise sensory load, ensure participant safety, and

allow assessment of intra individual physiological stability over

time.

Each participant follows the same temporal progression,

independent of physiological responses, except when predefined

stopping rules are triggered. No adaptive modification of stimulus

parameters is performed based on interim observations. No interim

analyses or protocol adaptations will be performed during the

study.

The protocol is organised into three successive phases spanning six

weeks. Progression from one phase to the next follows a

predefined fixed schedule and is not adapted based on observed

responses, except in cases where predefined stopping rules result in

session termination. Successful completion is defined

retrospectively for feasibility evaluation and does not alter the

planned timeline.

Early sessions prioritise maximal predictability and perceptual

stability. Subsequent sessions introduce controlled and limited

novelty in order to probe autonomic sensitivity while maintaining a

low cognitive and sensory burden.

All sessions are scheduled with sufficient temporal spacing to

avoid cumulative fatigue or sensitisation effects. Session frequency

and duration are identical across participants.

The full procedure timeline, including session content and

duration, is specified *a priori* and summarised in Table 4.

Table 4. Procedure timeline and visual mirror exposure progression

Phase	Weeks	Session duration
Initial exposure (static stimuli)	Weeks 1–2	**10 to 15 minutes per session**
Predictable motion exposure	Weeks 3–4	**10 to 15 minutes per session**
Controlled novelty exposure	Weeks 5–6	**10 to 15 minutes per session**

654   Table 4 summarises the predefined temporal structure of the visual

655   mirror exposure protocol. Each phase is designed to incrementally

656   modulate perceptual complexity while preserving high stimulus

657   predictability.

658   Session durations are intentionally brief to reduce physiological

659   and sensory burden. The progression from static to dynamic

660   stimuli, followed by controlled novelty, reflects the theoretical aim

661   of gradually engaging visual predictive mechanisms without

662   inducing overload.

663   This timeline does not represent a therapeutic dose escalation and

664   is not intended to induce functional change. Its sole purpose is to

665   support evaluation of protocol feasibility, tolerability, and

666   physiological signal stability within a predefined, feasibility-

667   focused study framework.

668   **5. Data Analysis Plan**

669   This section outlines the predefined descriptive analysis structure

670   used to evaluate feasibility, tolerability, and measurement

671   reliability of the visual mirror paradigm. The analyses focus

672   exclusively on non-behavioural physiological signals, predefined

673   temporal windows corresponding to protocol phases, and explicit

674   decision rules specified a priori. No hypothesis testing or efficacy

675   evaluation is planned in this feasibility study. The complete data

676   analysis plan, including signals, temporal windows, indicators, and

677   decision thresholds, is summarised in Table 5.

678   **Table 5. Data analysis plan: signals, windows, indicators, and decision rules**

Analysis domain	Physiological signal	Temporal window	Descriptive metric	Indicator	Decision rule
Baseline Physiological State	HR	Pre-exposure	Mean HR	Stable autonomic baseline	Baseline valid if signal is continuous and artefact-free for each participant.
Baseline Physiological State	HRV	Pre-exposure	Time-domain indices	Interpretable baseline	Retained if indices are computable at the individual level.
Exposure Tolerability	HR	Stabilisation	Mean and peak HR	Absence of sustained activation	Session continues if HR remains within predefined thresholds for that participant.
Exposure Tolerability	HRV	Stabilisation	Qualitative pattern	Autonomic stability	Session interrupted if sustained reduction indicating distress is observed.
Sensitivity to Controlled Novelty	HR	Controlled novelty phase	Peak and pattern relative to baseline	Autonomic reactivity	Data point included in intra-individual analysis if signal quality meets Table 1 criteria; session terminated if predefined non-verbal dissent markers (Section 4.5) are detected.
Non-Verbal Dissent	Combined Signals	All phases	Presence of markers	Ethical safety	Immediate termination upon detection.
Measurement Reliability	HR and HRV	Across sessions	Intra-individual consistency	Signal stability	Assessed descriptively within participants across repeated sessions
Feasibility Assessment	Procedural Indicators	Entire protocol	Completion rate	Implementability	Overall feasibility evaluated across the predefined feasibility sample (minimum n = 10), without group-level inference

679   Note: HR: Heart Rate; HRV: Heart Rate Variability

Table 5 summarises the predefined data analysis plan for this feasibility study, linking each analytical domain to specific physiological signals, temporal windows, descriptive metrics, and explicit decision rules. In line with the objectives of a feasibility-focused study, the table specifies how physiological data will be examined descriptively at the individual level, without hypothesis testing or group-level inference. The decision rules outlined in the table define a priori the conditions under which protocol feasibility, physiological tolerability, and measurement reliability will be considered achieved, partially achieved, or not achieved. By specifying these analytical procedures before data collection, the table ensures transparency, prevents analytical flexibility, and establishes a clear interpretive structure for interpreting outcomes strictly in terms of feasibility rather than therapeutic efficacy.

5.1 General Analytical Structure

This feasibility study is not designed to test therapeutic efficacy, causal effects, or group-level hypotheses.

All analyses are therefore descriptive, exploratory, and strictly limited to the evaluation of feasibility, tolerability, and measurement reliability of the visual mirror paradigm.

Analyses focus on intra-individual trajectories rather than between-

participant comparisons. Each child serves as their own reference,

and no inferential statistics aimed at generalisation are planned.

No hypothesis testing, effect size estimation, or correction for

multiple comparisons will be conducted.

5.2 Data Preprocessing and Signal Quality Control

Physiological signals are visually and algorithmically inspected

prior to analysis to identify artefacts related to movement, sensor

displacement, or technical failure.

Segments affected by signal artefact are flagged and excluded

according to predefined criteria. Only periods with interpretable

physiological data are retained for analysis.

Baseline values are computed individually for each participant

using resting state segments recorded prior to visual exposure.

5.3 Descriptive Analysis of Physiological Signals

Heart rate and heart rate variability are summarised using

descriptive statistics within predefined temporal windows

corresponding to protocol phases.

Analyses examine:

•   baseline resting periods,

•   stabilisation phases,

•   continuity phases,

•   controlled novelty phases.

For each participant, temporal evolution of physiological signals is
visualised across sessions using within subject plots and time
series representations.

No aggregation across participants is performed.

5.4 Assessment of Feasibility and Tolerability

Feasibility outcomes are evaluated using procedural indicators,
including:

•   number of sessions initiated,

•   number of sessions completed,

•   frequency and timing of early session termination,

•   occurrence of protocol deviations.

Tolerability is assessed through the absence or presence of
sustained physiological activation exceeding predefined thresholds,
and through documentation of non verbal dissent markers
triggering session interruption.

These outcomes are summarised descriptively at the individual

level.

5.5 Assessment of Measurement Reliability

Measurement reliability is examined by evaluating:

• stability of physiological signals across repeated sessions within

individuals,

• consistency of signal patterns across comparable protocol phases,

• proportion of sessions yielding usable physiological data.

Reliability assessment does not assume stationarity or normality of

signals and is limited to descriptive pattern identification.

5.6 Interpretation Structure Specific to Feasibility Assessment

Interpretation relies on within-session and within-individual

contrasts across predefined protocol phases, rather than on absolute

autonomic values. Differential autonomic patterns observed

between baseline, stabilisation, continuity, and controlled novelty

phases are used to characterise the internal structure and

interpretability of physiological signals. These contrasts inform

decision rules by allowing distinction between phase-dependent

physiological organisation and random fluctuation or non-specific

global activation. They are not used to infer magnitude of

response, functional visual processing, or therapeutic effect.

Outcomes are interpreted as indicating:

• feasibility achieved,

• feasibility partially achieved,

• feasibility not achieved,

Based on the combination of protocol completion rates,

physiological tolerability, and signal interpretability at the

individual level.

The predefined target sample size of 10 participants represents an a

priori feasibility objective rather than a rigid requirement. In the

event that recruitment is limited due to population rarity or ethical

constraints, feasibility interpretation will be conducted

transparently based on the achieved sample, with explicit reporting

of recruitment limitations and without post hoc modification of

feasibility criteria.

No post hoc modification of decision thresholds is permitted.

Importantly, the protocol is considered feasible even if

physiological stabilisation reflects habituation rather than

engagement, provided predefined tolerability and signal

interpretability criteria are met.

5.7 Transparency and Reproducibility

All analysis procedures, decision rules, and feasibility criteria are

defined a priori, prior to any data collection, to minimise analytical

flexibility and to prevent post hoc reinterpretation. Any deviation

from the planned analytical procedures, including unexpected data

quality constraints or session level interruptions triggered by

predefined stopping rules, will be documented transparently and

reported alongside the feasibility and tolerability results. No post

hoc changes to decision thresholds or outcome definitions will be

introduced after data collection begins.

This analytical structure establishes the methodological conditions

required to justify the consideration of future mechanistic or

efficacy studies, contingent upon feasibility, tolerability, and

measurement reliability being demonstrated.

6. Ethics, Oversight, and Regulatory Considerations

This manuscript describes a clinical feasibility study protocol and

does not involve the collection, analysis, or reporting of empirical

data from human participants.

No study procedures have been initiated, and no participants have
been recruited.

The proposed visual mirror paradigm is intended for future
empirical implementation in a highly vulnerable paediatric
population. Consequently, ethical considerations are integrated at
the protocol design level, prior to any data collection. Formal
ethical approval will be sought from the appropriate Human
Research Ethics Committee at the host institution before the
commencement of any study involving human participants.

The protocol is designed in accordance with the precautionary
principle. The visual mirror exposure is strictly passive, non-
invasive, and does not require instruction following, task
engagement, or voluntary behavioural responses. The absence of
overt behavioural response is not interpreted as consent.
Participant protection relies on continuous monitoring of
predefined physiological and behavioural indicators of potential
non-verbal dissent, including autonomic destabilisation and
observable signs of distress. Any indication of predefined non-
verbal dissent will result in immediate termination of the session,
in accordance with the stopping rules specified in Section 4.5.

No therapeutic intent is associated with this protocol. The study
does not aim to induce functional change, does not replace or

withhold standard care, and does not involve deception,

randomisation, or experimental manipulation beyond controlled

visual exposure. The sole purpose of the proposed protocol is to

determine whether the paradigm can be implemented safely,

tolerably, and with sufficient measurement reliability to justify

consideration of future mechanistic or efficacy studies.

Data management, storage, and analysis procedures for any future

implementation will comply with institutional, national, and

international regulations governing research involving children and

vulnerable populations. As a feasibility study involving a

vulnerable paediatric population, the protocol may be discontinued

if predefined feasibility, safety, or tolerability criteria are not met.

Such an outcome would be considered ethically appropriate and

informative for future research planning.

7. Discussion

The present manuscript describes a clinical feasibility study

protocol designed to evaluate a visual mirror paradigm in children

with cerebral visual impairment (CVI) presenting an atelic profile.

As no empirical data have been collected, this discussion does not

interpret outcomes or effects, but instead examines the

methodological validity, scope, and limits of the proposed

protocol. The primary methodological value of this protocol lies in

its explicit adaptation to a population for whom conventional

visual assessment or rehabilitation approaches are neither ethically

nor practically applicable. By relying exclusively on passive visual

exposure and non-behavioural autonomic physiological measures,

the protocol avoids any requirement for voluntary responses, task

engagement, or instruction following. This design directly

addresses a major methodological gap in the study of children with

severe CVI and atelic profiles, while remaining compatible with

stringent ethical constraints.

A central strength of the proposed approach is its exclusive focus

on feasibility, tolerability, and measurement reliability, rather than

on clinical efficacy or functional visual processing. The use of

predefined session phases and within-session, within-individual

contrasts allows evaluation of whether physiological signals

exhibit structured and interpretable organisation in response to

controlled visual input, without reliance on absolute autonomic

values. This structure is specifically intended to determine whether

the proposed paradigm is sufficiently stable, interpretable, and

ethically manageable to justify future empirical implementation.

The protocol also incorporates conservative safety monitoring and

stopping rules designed to protect participants who may be unable

to express discomfort through overt behavioural or verbal means.

By clearly distinguishing between analytic inclusion criteria and

ethical termination criteria, the design prioritises participant

protection while preserving methodological transparency. These

safeguards are essential when working with highly vulnerable

paediatric populations and constitute a core component of the

feasibility assessment.

Several limitations inherent to the proposed design must be

acknowledged. The absence of behavioural outcomes, group-level

comparisons, or functional performance measures necessarily

restricts the interpretive scope of the data that may be generated by

future implementation. In addition, autonomic physiological

signals are not interpreted as specific markers of visual processing,

but rather as indirect indicators of physiological regulation,

tolerability, and signal stability. These constraints are intentional

and reflect the ethical and practical boundaries associated with

research involving children with severe cerebral visual impairment

and atelic profiles, rather than methodological shortcomings.

The predefined feasibility, tolerability, and measurement reliability

criteria are intended to guide both the ethical conduct and

interpretation of the study. Should these criteria not be met, early

discontinuation of the protocol would constitute an ethically

appropriate and informative outcome. Conversely, if the criteria

885   are satisfied, the findings may inform future methodological or

886   clinical research considerations, without implying any obligation

887   or commitment to subsequent studies.

888   This discussion therefore addresses the validity and limits of the

889   proposed methodological approach itself, rather than interpreting

890   outcomes that have not yet been observed. The manuscript should

891   be read as a methodological protocol evaluated prior to data

892   collection, establishing clear ethical, procedural, and analytical

893   boundaries for any future empirical investigation.

894   **Declarations**

895   **Ethics Statement:** This manuscript presents a conceptual and

896   methodological protocol developed within an independent research

897   programme. No data involving human participants were collected,

898   generated, or analysed as part of the present work.

899   The proposed protocol is intended to inform future empirical

900   research involving vulnerable paediatric populations. Formal

901   ethical approval will be sought from the appropriate Human

902   Research Ethics Committee (HREC) at the host institution prior to

903   the commencement of any study involving human participants.

904   The protocol has been designed in accordance with the

905   precautionary principle. Given the limited capacity for verbal

906   assent in the target population, the absence of a behavioural

907   response is not interpreted as consent. The protocol explicitly

integrates physiological and behavioural indicators as markers of potential non-verbal dissent, and any such indicators would lead to immediate termination of the session in future implementations.

Data and Materials Availability

No empirical data were collected for this study. The protocol materials and methodological documentation have been archived in a private repository and can be shared with reviewers or editors upon request.

Data Availability Statement

Not applicable. No data were generated or analysed in the current study.

1.   Smague E. Literature Review of the Influence of Neurovisual Disorders on the Diagnosis of Autism in Australian Children. Current Developmental Disorders Reports. 2024;12(1):3.

2.   Chokron S, Bernard JM, Imbert M. Length representation in normal and neglect subjects with opposite reading habits studied through a line extension task. Cortex. 1997;33(1):47–64.

3.   Hacohen A, editor The psychological reality of the telic/atelic distinction: Evidence from adult and child Hebrew. Proceedings of the 21st Annual Meeting of the Israel Associations for Theoretical Linguistics,(unnumbered); 2006.

4.   Derrien D, Garric C, Sergent C, Chokron S. The nature of

blindsight: implications for current theories of consciousness.

Neurosci Conscious. 2022;2022(1):niab043.

5.   Friston K, Mattout J, Kilner J. Action understanding and

active inference. Biol Cybern. 2011;104(1-2):137–60.

6.   Dutton GN, Jacobson LK. Cerebral visual impairment in

children. Semin Neonatol. 2001;6(6):477–85.

7.   Friston K. The free-energy principle: a unified brain

theory? Nat Rev Neurosci. 2010;11(2):127–38.

8.   Ramachandran VS, Rogers-Ramachandran D. Synaesthesia

in phantom limbs induced with mirrors. Proc Biol Sci.

1996;263(1369):377–86.

9.   Bach-y-Rita P, S WK. Sensory substitution and the human-

machine interface. Trends Cogn Sci. 2003;7(12):541–6.