SHAPE-AI: Development and Expert Validation of a Survey for Human–AI Performance
Evaluation in Healthcare

Mark V Mai, MD, MHS[1,2], Mustafa Ozkaynak[3], PhD, Jenna L. Marquard [4], PhD, Richard J.
Simonson[5,6] PhD, Richard J. Holden, PhD,[8] Hanna J. Barton, PhD[9], Ken Catchpole PhD[10],
Melinda Jamil, MS[11], Sarah Brauer, MS, RN, NI-BC[12], & Swaminathan Kandaswamy, PhD,
FAMIA[1]

1 - Emory University, Department of Pediatrics, Atlanta, GA

2 - Children's Healthcare of Atlanta, Atlanta, GA

3 - University of Colorado | Anschutz Medical Campus, College of Nursing, Aurora, CO

4 – University of Minnesota School of Nursing, Minneapolis, MN

5- Children's Mercy Hospital, Kansas City, MO

6 – University of Missouri Kansas City School of Medicine, Kansas City, MO

7 – St. Luke's Health System, Boise, ID

8 – Indiana University, Bloomington, IN

9 – BerbeeWalsh Department of Emergency Medicine, University of Wisconsin – Madison

10- Medical University of South Carolina, College of Medicine, Charleston, SC

11 – Advocate Health, Charlotte, NC

12 – St Lukes Health System, Boise, ID

Running head: SHAPE-AI instrument development

Article type: Original Research

Exact word count of text: 3634

Corresponding Author:
Swaminathan Kandaswamy
Assistant Professor
Director of Human Factors in Information System Design
Emory University School of Medicine, Department of Pediatrics
Email: Swaminathan.Kandaswamy@emory.edu
Address: 2015 Uppergate Drive, Atlanta, GA 30322

Funding: This work was supported by the Agency for Healthcare Research and Quality
(AHRQ) and the National Center for Advancing Translational Sciences (NCATS) of the National
Institutes of Health (NIH) grant numbers 5R03HS029417-02, UL1TR002378, and
KL2TR002381.

Key Words: Human-AI Teaming, Psychometrics, Trust in Automation, Clinical Decision
Support, Situation Awareness

Abstract

Objective

To develop and content-validate a brief, expert-informed Survey for Human–AI

Performance Evaluation (SHAPE-AI) for near-real-time assessment of how clinical AI

affects human performance.

Background

AI-enabled clinical decision support can improve outcomes only when aligned with

clinician workflows, and cognitive demands. Existing evaluations measure technical

performance and adoption , providing limited assessment of how AI shapes human

performance. There is a lack of concise, operationally feasible instruments to measure

AI impact on these human factors outcomes in clinical settings.

Method

We used a construct-driven, multi-stage development process. A literature review and

prior qualitative work with users of a deployed sepsis prediction tool identified core

human performance constructs. Preliminary items were created and iteratively refined

through two expert panels. Six clinical informatics experts evaluated representativeness

and clarity using content validity indices (CVI). Seven human factors experts then

refined constructs, item wording, and response formats through ratings and focus

groups, emphasizing discriminant validity, cognitive bias mitigation, and feasibility for

deployment within 24 hours of AI use.

Results

A concise 10-item instrument was created, comprising perceived impact, interpretability,

agreement with the AI's findings, agreement with the AI's recommendations, trust,

workload, provider–patient and provider–team relationships, unexpected outcomes.

Conclusion

The SHAPE-AI instrument is a theoretically grounded, operationally feasible tool to

monitor human performance as relates to AI use.

Application

Health care organizations can deploy SHAPE-AI as a rapid, standardized probe to

detect workflow misalignment, mis-calibrated reliance, communication disruptions, and

unintended consequences of AI, informing safer design, implementation, and

optimization of clinical AI tools.

Précis: SHAPE-AI is a brief, expert-validated survey designed to capture clinicians'

near-real-time perceptions of how AI-enabled decision support affects human

performance such as their situational awareness, decision-making, workload, and trust.

SHAPE-AI offers health systems a practical, standardized way to monitor and

understand the impact of AI on human performance.

INTRODUCTION

Artificial intelligence (AI)–enabled tools are increasingly deployed in healthcare to improve clinical outcomes.[1] However, good outcomes are only possible when these tools fit clinician workflows and the broader sociotechnical context in which they are embedded. A model may be mathematically robust yet still fail clinically if its outputs are hard to interpret, appear at the wrong time, or erode user trust.[2,3] In this sense, technical accuracy is necessary but not sufficient for a safe, high quality care delivery.

Despite this reality, the implementation of clinical AI is currently hindered by three systemic failures: an over-emphasis on technical accuracy, a neglect of workflow integration, and a failure to account for the complexities of human cognition

First, technical accuracy describes how well the model predicts or classifies clinical events, often measured by discrimination and calibration.[3] These metrics are the focus of most AI evaluations. They tell us how the model performs with data, but not whether clinicians will see, understand, or use its outputs in real time.

Second, a neglect of workflow integration often renders even accurate models unhelpful; an alert that fires too often, appears at a busy moment, or is buried in a crowded interface can disrupt situational awareness rather than enhance it .[4–6] The same vendor sepsis prediction model has produced very different outcomes in different health systems: one reported increased alert fatigue,[4] while another reported reduced mortality.[7] These contrasting results highlight how local workflows, organizational context, and implementation decisions shape whether AI tools are ultimately beneficial.

Finally, human cognition refers to how clinicians perceive, interpret, and act on

information from AI tools. AI does not operate in isolation; it becomes part of a joint

human–AI system in which clinicians must combine model outputs with their own

expertise, patient preferences, and resource constraints.

Established Human Factors Constructs

Human factors research provides the necessary constructs to explain why AI succeeds

or fails at the point of care. One key construct is situation awareness - what information

clinicians notice, how they make sense of it, and how they anticipate what will happen

next.[8] An AI tool might improve situation awareness by highlighting a subtle risk that

would otherwise be missed, or it might degrade it by distracting attention from more

important cues.

Another key construct is trust, the extent to which clinicians are willing to rely on the AI

tool when making decisions.[9,10] If clinicians under-trust the tool, they may ignore useful

recommendations. If they over-trust-, they may follow incorrect advice without sufficient

scrutiny (over-reliance or automation bias).[9–11] Communicating limitations and

uncertainty about the tool can help calibrate trust and reduce automation bias,

supporting more balanced decision making. [12]

A related construct is interpretability, which we use here to mean how understandable

and usable the AI output is to the clinician in context.[13] A technically "explainable" model

is not necessarily interpretable at the bedside. If clinicians cannot see why a patient is

being flagged, they may either disregard the alert or accept it uncritically. Another

important idea is human–AI agreement: the extent to which clinicians' judgments align

with the tool's findings and recommendations, and when they reasonably diverge.

Workload and cognitive burden are also critical constructs. AI tools are often promoted

as ways to reduce burden, but they can also add work by generating more alerts,

requiring extra checks, or prompting new documentation tasks. [6,14–17] An accurate alert

that appears at a high stress- moment can increase cognitive load and distraction. Over

time, this can contribute to alert fatigue and burnout.

AI tools can further affect team communication and relationships, how AI reshapes

interactions among clinicians, patients, and the technology itself. [14] An AI generated risk

score might prompt more timely huddles, or it might replace direct communication with a

silent alert in the electronic health record. These changes influence not only efficiency

and coordination, but also clinicians' perceived safety of using the tool in -real-world-

practice.

The Need for Real-Time Measurement

Despite the importance of these human and workflow factors, most evaluations of

clinical AI still rely on technical metrics and coarse adoption outcomes, such as alert

adherence or override rates.[3] These measures show whether clinicians followed an

alert, but not *why* they did so, or how the AI tool affected their situation awareness,

decision-making, workload, or team interactions. In other words, we lack practical ways

to measure the "missing link" between model performance and clinical outcomes.

Human factors principles are crucial for understanding not only how clinicians use AI

when it works well, but also how they detect, interpret, and recover from AI--related-

errors.

Clinicians are often the first to sense when an AI tool is misaligned with real-world

practice. By evaluating clinicians' near-real-time experience of AI at the point of care,

health systems can detect latent failures and integration problems before they lead to

harm. In this way, clinicians can act as "canaries in the coal mine" for post-deployment

performance drops, workflow mismatches, and potential bias in AI tools. There is also a

growing risk that differences between AI recommendations and clinicians' final decisions

will be interpreted simply as noncompliance, rather than as appropriate clinical

judgment in the face of imperfect tools. Without a clear view of human–AI interaction,

clinicians may be unfairly blamed for overriding AI in situations where the model's

recommendation is poorly timed, poorly calibrated, or misaligned with patient needs.

Despite recognition of the need to measure human performance in relation to AI

systems,[18] there remains a paucity of validated, operationally feasible instruments

designed to capture these human factors constructs in near-real---time. Existing

measures tend to focus on general attitudes toward technology or require lengthy

administration that is impractical in busy clinical settings.

Study Objective

The objective of this study was therefore to develop and content-validate a brief,

expert--informed survey instrument to assess the near--real--time impact of AI on

human performance in clinical settings, termed SHAPE---AI (**S**urvey for **H**uman-**A**I

Performance **E**valuation). Prior work by our group using a deployed sepsis prediction

alert identified several key human performance constructs, including changes in

situational awareness, diagnostic reasoning, treatment planning, team communication,

and response behavior.[19] Building on that foundation, SHAPE-AI is designed as a

concise, generalizable measurement instrument that can be deployed shortly after AI

use across different clinical AI and clinical decision support tools. It captures clinicians'

perceptions of impact, agreement with AI findings and recommendations, interpretability,

trust, workload, sociotechnical relationships, and unintended consequences. Such an

instrument can support rapid---cycle evaluation, guide implementation adaptations, and

provide a standardized lens on human–AI teaming that complements technical

performance metrics and outcome data.

By utilizing a construct-driven process that integrated prior empirical work, literature

review, and sequential input from clinical informatics and human factors experts, we

sought to operationalize core domains of human–AI interaction. Our goal was to

develop a tool that is both practical for routine deployment and conceptually aligned with

contemporary human factors theory, enabling health systems to systematically monitor

how AI systems shape clinician decision-making, workload, team dynamics, and

perceived safety.

METHODS

Developing measurement constructs

Following the survey scale design process described by AMEE (An International

Association for Medical Education) Guide No. 87, [20] we first conducted a literature

review to identify foundational measurement constructs.[18] As described in our previous

published work,[21] we conducted semi-structured interviews with users of a live sepsis AI

system to elucidate how they interacted with the AI tool and understand the

operationalization of these human performance constructs in the context of acute care

settings. To minimize recall bias, we identified and interviewed providers and nurses

within 72 hours of them seeing the AI sepsis tool in production. After synthesizing these

qualitative learnings, we developed preliminary survey items.

Validation and refinement of preliminary survey items

We used the content validity index (CVI) to evaluate how well the items represented

each human performance construct being measured. In line with psychometric

standards, a CVI of >=0.8 was established as the threshold for item retention and

refinement.[22] We conducted sequential expert validation of the survey instrument with

two distinct expert panels to ensure the tool achieved both clinical relevance and human

factors rigor.

Phase 1: Clinical Informatics Expert Panel

To establish preliminary validity, six board certified clinical informaticians with

experience in AI and clinical decision support (CDS) design, implementation, and

evaluation were recruited from the Pediatric CDS Collaborative, a national research

collaborative composed of clinical informaticists from nearly 20 pediatric institutions in

the US. Each expert rated each survey item assessing how representative and clear the

items were with respect to the construct of interest. For example, for the workload

construct participants were asked these two question(s):

a. For the overall care of this patient, on a scale of 0 (Low workload)-100

202   (High Workload) please rate your over all workload

203   Please rate how representative the above question is to understand the

204   workload. (The subjective demands of a particular task).

205   i. Item is not representative

206   ii. Item needs major revisions to be representative

207   iii. Item needs minor revision to be representative

208   iv. Item is representative

209   b. For the overall care of this patient, on a scale of 0 (Low workload)-100

210   (High Workload) please rate your overall workload

211   Please rate how representative the above question is to understand the

212   workload. (The subjective demands of a particular task).

213   i. Item is not clear

214   ii. Item needs major revisions to be clear

215   iii. Item needs minor revision to be clear

216   iv. Item is clear

217 Panelists also provided qualitative feedback on redundancy, the clinical interpretability

218 of the instrument items, and the potential for adapting the instrument to diverse CDS

219 interventions.

Phase 2: Human Factors Expert Panel

Seven experts with formal training in human factors who have studied human

performance related to healthcare were recruited nationally via e-mail. Survey items

revised after Phase 1 were first rated individually by these experts using the same type

of survey question used in Phase 1 for clarity and representativeness/alignment with

established human performance constructs. The panel then participated in four

facilitated 90-minute focus group discussions (two for each round) addressing: (1) the

scope of the tool (e.g., distinguishing between specific alerts vs broader AI "findings"

and "recommendations"), (2) the adequacy and discriminant validity of constructs such

as agreement, interpretability, trust, workload, sociotechnical relationships, and

unintended consequences, and (3) item wording specifically designed to mitigate

cognitive biases, scaling, and response-option structure. After each round, the updated

instrument was re-rated for clarity and feasibility until consensus was reached.

Producing the SHAPE-AI Survey

The final step of the development process involved a thematic synthesis of the

qualitative feedback and quantitative ratings collected during the two expert validation

phases. Discrepancies between the clinical informatics requirements for operational

feasibility (Phase 1) and the human factors requirements for theoretical rigor (Phase 2)

were resolved through iterative review by the core research team (authors MVM, MO,

SK). Final decisions were made to ensure the instrument could capture the nuance of

human-AI teaming across diverse clinical AI applications

Ethics:

242 This research complied with the American Psychological Association Code of Ethics and was

243 approved by the Institutional Review Board at Children's Healthcare of Atlanta STUDY00001383

Results

Identification of Human Performance Constructs

The initial literature review identified seven core human performance constructs potentially impacted by AI in healthcare: situational awareness, explainability, trust, automation bias, human-computer agreement, workload, and the impact on staff relationships (Table 1). Based on our previously published work, including the initial survey review of human performance constructs and subsequent semi-structured interviews,[21] we expanded this list to include differential diagnosis, treatment planning, team communication, tool satisfaction, and clinician response behavior. These constructs formed the basis of the initial 10-item survey instrument subjected to expert validation.

Table 1: Initial Constructs and Survey Items

Construct(s)	Definition	Initial survey items
Situation Awareness / Impact	Dynamic understanding of ongoing situation through the lens of perception, comprehension and projection.[8]	Did the alert lead to (multiselect) a. more timely care for the patient b. change in your diagnosis or differential c. change in treatment for the patient d. change in staffing or level of care for this patient e. improvement in team communication f. situational awareness
Impact	Changes in diagnosis, efficiency, team communication, or confidence	
Workload	Subjective demands of a particular task.[23]	For the overall care of this patient, on a scale of 0 (Low workload)-100 (High Workload) please rate your overall workload
Explainability	Characteristics of AI that enable users to know logic	a. How well do you understand why this alert is appearing at the time when it appeared?

	behind decisions and parameters used to make the decisions. [23]	Not at all b. Slightly c. Somewhat d. Mostly e. Completely

Human-Computer Agreement	User agreement with the AI system[24]	How surprised are you that this alert showed for the patient?
		    a. Not at all
		    b. Slightly
		    c. Somewhat
		    d. Mostly
		    e. Completely
		Do you agree with alert that this patient was at high risk for \<clinical condition\>?
		    a. Not at all
		    b. Slightly
		    c. Somewhat
		    d. Mostly
		    e. Completely
		When the alert appeared for this patient, how appropriate was the \<alert\> ?
		    a. Not at all
		    b. Slightly
		    c. Somewhat
		    d. Mostly
		    e. Completely

Automation Bias	Tendency to overly rely on automation leading to errors of commission or omission [25,26]	N/A (unable to measure directly from clinician responses)

Trust	Trust is the attitude that an AI system will help achieve an individual's goals in a situation characterized by uncertainty and vulnerability.[9]
	
	Overtrust: Trust exceeds system capabilities, leading to misuse.[9,10]
	
	Distrust: Trust falls short of system capabilities, leading to disuse.[9,10]
		In general, how much do you trust this alert ?
		    a. Not at all
		    b. Slightly
		    c. Somewhat
		    d. Mostly
		    e. Completely

Relationship between staff and patients	The way in which people regard and behave toward each other.	How adversely did the alert/huddle impact your relationship with the patient?
		    a. Not at all
		    b. Slightly
		    c. Somewhat
		    d. Mostly
		    e. Completely

Satisfaction	The overall contentment, engagement, experience, and happiness.	In general, how satisfied are you with the alert?
		    a. Not at all
		    b. Slightly
		    c. Somewhat
		    d. Mostly
		    e. Completely

Alert Performance	The effectiveness and efficiency in delivering relevant, timely, and actionable information to the right clinician through the right channel to improve patient outcomes and safety, without causing undue burden or "alert fatigue	How often do you ignore this alert because of how frequently it is incorrect
		    a. Almost never
		    b. Once in a while
		    c. Sometimes
		    d. Often
		    e. Almost always

Phase 1: Clinical Informatics Expert Panel

The Clinical Informatics expert panel evaluated the initial instrument (Table 1) for

content validity and clarity. The Clinical Informatics panel rated most items > 0.8 on the

CVI, except for four constructs – Impact (CVI = 1.0 [Representativeness]; 0.7 [Clarity]),

Human-Computer Agreement (CVI = 0.6 [Representativeness]; 0.8 [Clarity]), Workload

(CVI = 0.7 [Representativeness]; 0.7 [Clarity]), and Patient-Staff Relationships (CVI =

0.5 [Representativeness]; 0.7 [Clarity]). Panel feedback highlighted three primary

concerns:

- Redundancy in Agreement: Experts noted that the two separate items measuring "human-computer agreement" were conceptually indistinct and recommended merging them into a single assessment of agreement.
- Ambiguity in Workload Measurement: Experts questioned the validity of a single-item workload measure. They argued that respondents may struggle to distinguish the incremental cognitive load of the AI tool (e.g., interpretation and decision-making) from their baseline clinical workload (e.g., clinical complexity).
- Generalizability: The panel recommended replacing AI-specific terminology with broader CDS language to ensure the instrument could evaluate diverse algorithmic interventions (e.g., non-AI rules-based alerts).

Phase 2: Human Factors Expert Panel

The Human Factors expert panel reviewed the revised instrument Feedback centered on refining construct validity, managing potential cognitive bias, optimizing measurement design, and clarifying the operational scope.

Construct Refinement and Measurement Challenges

A primary outcome of Phase 2 was the intentional decoupling of core constructs to mitigate ambiguity and improve discriminant validity, especially for near-real-time assessment.

Experts universally agreed with breaking "Human-Computer Agreement" into two distinct components, helping to identify cases where a clinician agrees with the AI's finding or risk assessment but disagrees with its suggested action.

1. **Agreement with the AI's Finding or Risk Assessment:** Evaluating the underlying model's identification of risk or condition
2. **Agreement with the AI's Recommendation or Suggested Action:** Evaluating the proposed clinical course

The panel noted that questions about understanding the AI rationale often confuse the technical feature of explainability (the system interface stating how it works) with the cognitive outcome of interpretability (the user's ability to make sense of the output). Experts cautioned that relying on self-reported understanding risked measuring the persuasiveness of the interface (sometimes referred to as "storytelling"). Given that perceived transparency is highly variable and can potentially affect clinician interaction with the AI system, the need for subjective assessment via the survey was noted as a valuable leading indicator.

To address the ambiguity identified in Phase 1, the panel recommended a simplified scaling approach focused on the specific outcome of the interaction. Rather than attempting to measure absolute cognitive load, the final question was refined to capture the relative change in burden: whether the AI system created more work, reduced work, or resulted in no additional work.

Discussion highlighted that teaming (which includes communication, coordination, collaboration, and trust) is complex and context-dependent, often requiring specialized

306   methodologies like observation to measure accurately. Panelists agreed that assessing

307   the direct impact on human-to-human team communication and coordination was

308   necessary, particularly in scenarios where the AI agent might intervene or replace

309   existing communication channels. This resulted in items focused on the AI's influence

310   on provider-patient and provider-team interactions

311   Cognitive Biases and Item Formatting

The survey instrument was iteratively modified to address potential respondent bias

inherent in self-reporting.

Experts noted that items requiring clinicians to report their cognitive state (e.g., degree

of surprise or initial suspicion) were susceptible to recall and confirmation ("saving

face") bias, wherein a respondent might retrospectively alter their pre-decision state to

align with the correct AI outcome. To mitigate this, items were reframed to evaluate

system performance (e.g., "How clearly did the tool indicate...") rather than implying a

potential failure in user judgment.

For concepts like trust, a graded continuum was recommended over binary (Yes/No)

options. Furthermore, panelists noted difficulty in differentiating subtle differences, such

as "slightly" versus "somewhat," on some scales. For scales related to influence or

impact, the inclusion of a neutral option, such as "No Effect," placed centrally between

positive and negative poles, was recommended to avoid forcing a response where the

AI was not perceived as having an impact.

To ensure the instrument's longevity, the panel advocated for broadening the

terminology beyond "alert" to accommodate diverse CDS modalities, including passive

alerts, predictive risk scores, and other AI interventions that offer findings or

recommendations. Operational constraints (i.e. interruption to patient care and

challenges in administering surveys immediately after AI system interactions) required

the survey to be deployed within 24 hours of exposure to the AI tool. Questions focus on

a specific patient interaction to minimize recall bias.

333 Final Instrument Domains

334 Following this iterative process, the expert panel reached a final consensus on the

335 SHAPE-AI instrument. The validated domains and a summary of the survey items are

336 presented in Table 2, while the complete survey instrument is provided as a standalone

337 resource in the Appendix 1.

338 *Table 2: Final Constructs and Survey Items*

Construct(s)	Description	Final survey items

Perceived Impact	Changes in diagnosis, efficiency, team communication, or confidence	1. Did the AI have an impact on any of the following (multiselect)
		    a. Situational awareness (e.g. seeing new information and/or new understanding of patient risk, anticipate actions you need to take)
		    b. Timely care for the patient
		    c. Patient diagnosis or differential
		    d. Treatment for the patient
		    e. Staffing or level of care for this patient
		    f. Team communication and/or co-ordination
		    g. Confidence in your clinical decisions
		    h. None of the above
		Free Text Optional

Interpretability	Cognitive clarity of the output rationale	2. How clearly did the AI system explain the reason for its finding? (Interpretability)
		    a. Very Clearly
		    b. Somewhat Clearly
		    c. Not Clearly at All

		Free Text Optional

Human-Computer Agreement	Assessment of the finding vs. the recommendation.	**3. Did you agree with the AI's finding [risk assessment]? (Agreement)**
		    a. Yes
		    b. No
		Free Text (optional)
		4. (If "No" in question 3) Which of the following best explains why you did not fully agree with the AI's finding? (Select all that apply)
		    a. The AI's finding was based on inaccurate or outdated information.
		    b. The AI's finding did not account for important clinical context or patient factors.
		    c. The AI's findings did not match my own clinical reasoning.
		    d. Other (please specify)
		5. Did you agree with the AI's recommendation?
		    a. Yes
		    b. No
		Free Text (optional)
		6. (If "No" to question 5) Please explain why you didn't agree with the AI recommendation.
		    a. Evidence to the recommendation is weak
		    b. There are other equally good alternative actions
		    c. The action recommendation doesn't meet patient condition.
		    d. The action was not feasible to perform in this clinical setting.
		    e. The action was not specific enough to be useful.
		    f. Other (Please specify)

Workload	Relative change in clinical burden.	7. How did this AI system affect your work? (Workload)
		    a. Created more work
		    b. Reduced my work
		    c. No additional work

Trust	Sentiment toward specific recommendations.	8. To what extent did you trust this <AI system>? (Trust)
		    a. Fully trusted
		    b. Somewhat trusted
		    c. Neutral/Unsure
		    d. Did not trust

Unexpected outcomes	Open-ended capture of leading indicators for safety, both positive and negative.	9. Did or could this <AI System> lead to potential unexpected outcomes (reduced cognitive burden, unintended clinician impact or patient safety events/harm, patient/family interactions, health equity etc.)? Multi select
		    a. Positive Unexpected Outcome(s)
		    b. Negative Unexpected Outcome(s)

		c. No Unexpected Outcome Optional description for each option **Free Text (optional)**

Relationship between staff and patients	Influence on provider-patient and provider-team interactions.	10. Compared to baseline, how did the AI system affect your interaction with the patient/family?
		    a. Positively
		    b. Negatively
		    c. No effect at all
		Free Text (optional)

Automation Bias	Tendency to overly rely on automation leading to errors of commission or omission [25,26]	Optional:
		11. Did you verify the AI's finding with other clinical resource (e.g., labs, vitals) or another clinician or your own training/knowledge before acting? (Automation Bias)
		    a. Yes
		    b. No
		    c. Not necessary

DISCUSSION

This study developed and iteratively refined a concise survey tool to characterize the

near-real-time impact of AI on human performance in clinical settings, explicitly

grounding the work in human factors and sociotechnical systems framework. By starting

from constructs identified in the literature[18] and then subjecting a 10-item tool to review

by clinical informatics and human factors experts, the work moves beyond ad hoc

usability questions toward a more systematic representation of how AI systems shape

situational awareness, decision-making, workload, and team relationships. The final

domains – perceived impact, agreement with findings and recommendations,

interpretability, trust, workload, sociotechnical relationships, and unintended

consequences – align with the "Joint Cognitive System" (JCS) framework for human-AI

teaming in healthcare, which emphasize that performance is an emergent property of

the human-AI partnership rather than the technical output of the model alone.[27,28]

A key contribution of this work is the explicit separation of agreement with the AI's

finding from agreement with its recommended action. Prior studies of AI-enabled

decision support often treat acceptance or override as a single behavioral endpoint,

obscuring situations where clinicians endorse the risk assessment but deliberately

choose a different course of action based on patient preference, comorbidities, or

resource constraints.[29,30] By decoupling these constructs, the instrument serves as a

diagnostic tool for identifying specific breakdowns in the human-AI control loop,

distinguishing between model miscalibration (poor findings) and clinical misalignment

(poor recommendations). Similarly, distinguishing interpretability (clarity of the output

rationale in context) from technical explainability acknowledges that human

performance depends not on the existence of a rationale, but on the clinician's ability to

apply that rationale to safe and efficient decision-making at the point of care.[13,31]

At the same time, expert feedback highlighted the inherent difficulty of measuring

higher-order cognitive constructs such as trust, explainability, and automation bias via

brief self-report immediately after a clinical encounter. Consistent with recent reviews,

the panels emphasized that trust in AI is dynamic and multidimensional;[32] perceived

trust captured on a rating scale may diverge substantially from "calibrated trust," in

which clinicians align their reliance on the tool with its true performance characteristics.

While we do not directly measure calibrated trust, this can be approximated by linking

participant response Item 8 with participant alignment in the AI system

recommendations and with clinical outcomes. For example, if the model positive

predictive value is only 10% but clinicians follow through with treatment decisions for

50% of AI recommendations there is overtrust. Likewise, post-hoc explanations can

increase user confidence even when they are driven by spurious features, meaning that

self-reported "understanding" may sometimes reflect interface persuasiveness rather

than genuine insight into model behavior.[31,33,34] In response, the final version of the

instrument focuses on user sentiment (e.g., perceived interpretability and trust) as

leading indicators and deliberately avoids over-claiming them as direct measures of

safety or error detection, reinforcing the need to triangulate survey responses with log

data, model performance, and outcome measures.

The panels also surfaced important design trade-offs between brevity and construct

coverage. To make the instrument deployable within 24 hours of an AI interaction and

feasible for routine use, several potentially important constructs – such as detailed error

identification and recovery behaviors, training adequacy, and equity impacts – could not

be fully elaborated. Instead, they are partially captured through broader items on

unintended consequences and free-text responses, which can be used to trigger deeper

qualitative inquiry where needed. This design choice aligns with a growing recognition

that frontline clinicians are already experiencing substantial cognitive and

documentation burden,[35] and that any additional measurement must be tightly scoped

to avoid exacerbating the very workload and alert fatigue issues it aims to study.[36] The

resulting instrument is therefore best understood as a rapid, situational probe that can

be layered with observational studies, think-aloud protocols, or ethnographic methods

when organizations need richer insight into human-AI teaming.

Several limitations suggest directions for future work. First, the current validation relied

on expert panels rather than large-scale psychometric evaluation with end users, so

further testing is needed to assess reliability, sensitivity to change, and construct validity

across diverse AI tools, use cases, clinical roles, and settings. Second, the instrument

was designed primarily around predictive AI and CDS-style interventions; although

terminology was broadened to encompass a wider range of AI modalities, additional

adaptation will likely be required for generative systems that support documentation,

summarization, or patient communication. Third, because the survey captures

subjective, scenario-specific perceptions, it cannot on its own quantify calibrated trust,

automation bias, or safety outcomes;[34] these require linkage to event logs, override

patterns, error reports, and patient-level outcomes. Finally, while the tool attends to

team communication and patient–family relationships, more work is needed to integrate

it with complementary methods that capture team cognition and organizational context.

Addressing these gaps will be essential to realizing the full potential of the instrument as

part of a multi-method toolkit for evaluating and improving human performance with AI in

healthcare.

CONCLUSION

SHAPE-AI represents a significant step toward the systematic and proactive monitoring

of human-AI teaming in clinical practice. This concise, expert-validated 10-item

instrument is designed to measure the human factors impact of clinical AI in near-real-

time. By prioritizing brevity and clarity, the tool is operationally viable for deployment

within the high-stakes environment of clinical workflows without inducing survey fatigue.

While limitations in self-reporting trust and higher-order cognitive processes remain

inherent to survey methodologies, this instrument provides a standardized

sociotechnical foundation for assessing how AI interventions shape clinician judgment,

workload, and team dynamics. Ultimately, SHAPE-AI allows health systems to detect

and mitigate latent failures in their human-AI joint cognitive system. Future work should

focus on pairing this subjective measure with technical logs, clinician performance

metrics, and patient outcomes to build a holistic and multimodal view of safety, utility,

and human-AI collaboration in healthcare.

Key points:

- SHAPE-AI is a brief, 10-item survey designed to capture clinicians' near-real-time

experience of AI-enabled clinical decision support at the point of care.

- The instrument was developed through a construct-driven process integrating

literature review, prior qualitative work with sepsis AI users, and sequential input

from clinical informatics and human factors experts.

- SHAPE-AI measures key human factors domains: perceived impact,

interpretability, agreement with findings, agreement with recommendations, trust,

workload, provider–patient and provider–team relationships, unexpected

outcomes, and optional automation bias.

- SHAPE-AI is intended to complement technical performance metrics and

outcome data, offering health systems a standardized, low-burden tool to detect

workflow misalignment, mis-calibrated reliance, and unintended consequences of

AI in clinical practice.

References

1. Tighe, P., Mossburg, S. & Gale, B. Artificial Intelligence and Patient Safety: Promise and Challenges. *Artif. Intell. Patient Saf. Promise Chall.* https://psnet.ahrq.gov/perspective/artificial-intelligence-and-patient-safety-promise-and-challenges (2024).

2. Parasuraman, R. & Manzey, D. H. Complacency and Bias in Human Use of Automation: An Attentional Integration. *Hum. Factors* **52**, 381–410 (2010).

3. Nicolson, A., Bradburn, E., Gal, Y., Papageorghiou, A. T. & Noble, J. A. The human factor in explainable artificial intelligence: clinician variability in trust, reliance, and performance. *NPJ Digit. Med.* **8**, 658 (2025).

4. Wong, A. *et al.* External Validation of a Widely Implemented Proprietary Sepsis Prediction Model in Hospitalized Patients. *JAMA Intern. Med.* **181**, 1065–1070 (2021).

5. Cabitza, F., Rasoini, R. & Gensini, G. F. Unintended Consequences of Machine Learning in Medicine. *JAMA* **318**, 517–518 (2017).

6. Hudson, T. J. *et al.* Impact of Ambient Artificial Intelligence Documentation on Cognitive Load. *Mayo Clin. Proc. Digit. Health* **3**, 100193 (2025).

7. Tarabichi, Y. *et al.* Improving Timeliness of Antibiotic Administration Using a Provider and Pharmacist Facing Sepsis Early Warning System in the Emergency Department Setting: A Randomized Controlled Quality Improvement Initiative∗. *Crit. Care Med.* **50**, 418–427 (2022).

8. Endsley, M. R. Toward a Theory of Situation Awareness in Dynamic Systems: *https://doi.org/10.1518/001872095779049543* **37**, 32–64 (2016).

9. Lee, J. D. & See, K. A. Trust in automation: Designing for appropriate reliance. *Hum.*

Factors **46**, 50–80 (2004).

10. Parasuraman, R. & Riley, V. Humans and Automation: Use, Misuse, Disuse, Abuse.

Hum. Factors J. Hum. Factors Ergon. Soc. **39**, 230–253 (1997).

11. Giorgetti, C., Contissa, G. & Basile, G. Healthcare AI, explainability, and the human-

machine relationship: a (not so) novel practical challenge. *Front. Med.* **12**, 1545409

(2025).

12. Goddard, K., Roudsari, A. & Wyatt, J. C. Automation bias: a systematic review of

frequency, effect mediators, and mitigators. *J. Am. Med. Inform. Assoc.* **19**, 121–127

(2012).

13. Rudin, C. Stop explaining black box machine learning models for high stakes

decisions and use interpretable models instead. *Nat. Mach. Intell.* **1**, 206–215

(2019).

14. Wekenborg, M. K., Gilbert, S. & Kather, J. N. Examining human-AI interaction in

real-world healthcare beyond the laboratory. *NPJ Digit. Med.* **8**, 169 (2025).

15. Stults, C. D. *et al.* Evaluation of an Ambient Artificial Intelligence Documentation

Platform for Clinicians. *JAMA Netw. Open* **8**, e258614 (2025).

16. Asgari, E. *et al.* Impact of Electronic Health Record Use on Cognitive Load and

Burnout Among Clinicians: Narrative Review. *JMIR Med. Inform.* **12**, e55499 (2024).

17. Evan Orenstein *et al.* Alert burden in pediatric hospitals: a cross-sectional analysis

of six academic pediatric health systems using novel metrics. *J. Am. Med. Inform.*

Assoc. https://doi.org/10.1093/jamia/ocab179 (2021) doi:10.1093/jamia/ocab179.

18. Sujan, M., Pool, R. & Salmon, P. Eight human factors and ergonomics principles for

healthcare artificial intelligence. *BMJ Health Care Inform.* **29**, e100516 (2022).

19. Kandaswamy, S. *et al.* Human performance evaluation of a pediatric AI sepsis

model. *J. Am. Med. Inform. Assoc.*

20. Artino, A. R., La Rochelle, J. S., Dezee, K. J. & Gehlbach, H. Developing

questionnaires for educational research: AMEE Guide No. 87. *Med. Teach.* **36**, 463–

474 (2014).

21. Kandaswamy, S. *et al.* Human performance evaluation of a pediatric artificial

intelligence sepsis model. *J. Am. Med. Inform. Assoc.* ocaf106 (2025)

doi:10.1093/jamia/ocaf106.

22. Rubio, D., Berg-Weger, M., Tebb, S., Lee, E. & Rauch, S. Objectifyng content

validity: Conducting a content validity study in social work research. *Soc. Work Res.*

27, 94–104 (2003).

23. Hussain, F., Hussain, R. & Hossain, E. Explainable Artificial Intelligence (XAI): An

Engineering Perspective. https://doi.org/10.48550/arxiv.2101.03613 (2021)

doi:10.48550/arxiv.2101.03613.

24. Vasey, B. *et al.* Reporting guideline for the early-stage clinical evaluation of decision

support systems driven by artificial intelligence: DECIDE-AI. *Nat. Med. 2022 285* **28**,

924–933 (2022).

25. Parasuraman, R. & Manzey, D. H. Complacency and bias in human use of

automation: an attentional integration. *Hum. Factors* **52**, 381–410 (2010).

26. Goddard, K., Roudsari, A. & Wyatt, J. C. Automation bias: a systematic review of frequency, effect mediators, and mitigators. *J. Am. Med. Inform. Assoc. JAMIA* **19**, 121 (2012).

27. Berretta, S. *et al.* Defining human-AI teaming the human-centered way: a scoping review and network analysis. *Front. Artif. Intell.* **6**, (2023).

28. Hollnagel, E. & Woods, D. D. *Joint Cognitive Systems: Foundations of Cognitive Systems Engineering.* (CRC Press, Boca Raton, 2005). doi:10.1201/9781420038194.

29. Yu, Y. *et al.* Enhancing Clinician Trust in AI Diagnostics: A Dynamic Framework for Confidence Calibration and Transparency. *Diagnostics* **15**, 2204 (2025).

30. Rosenbacke, R., Melhus, Å. & Stuckler, D. False conflict and false confirmation errors are crucial components of AI accuracy in medical decision making. *Nat. Commun.* **15**, 6896 (2024).

31. Ghassemi, M., Oakden-Rayner, L. & Beam, A. L. The false hope of current approaches to explainable artificial intelligence in health care. *Lancet Digit. Health* **3**, e745–e750 (2021).

32. McGrath, M. J., Lack, O., Tisch, J. & Duenser, A. Measuring trust in artificial intelligence: validation of an established scale and its short form. *Front. Artif. Intell.* **8**, (2025).

33. Babic, B., Gerke, S., Evgeniou, T. & Cohen, I. G. Beware explanations from AI in health care. *Science* **373**, 284–286 (2021).

34. Byrne, R. M. J. Good Explanations in Explainable Artificial Intelligence (XAI): Evidence from Human Explanatory Reasoning. in *Proceedings of the Thirty-Second*

International Joint Conference on Artificial Intelligence 6536–6544 (International

Joint Conferences on Artificial Intelligence Organization, Macau, SAR China, 2023).

doi:10.24963/ijcai.2023/733.

35. National Academies of Sciences, Engineering, and Medicine; National Academy of

Medicine; Committee on Systems Approaches to Improve Patient Care by

Supporting Clinician Well-Being. *Taking Action Against Clinician Burnout: A Systems*

Approach to Professional Well-Being. (National Academies Press (US), Washington

(DC), 2019).

36. Chaparro, J. D. *et al.* Clinical Decision Support Stewardship: Best Practices and

Techniques to Monitor and Improve Interruptive Alerts. *Appl. Clin. Inform.* **13**, 560–

568 (2022).

541