Machine learning and burden analyses highlight novel genes in Parkinson's Disease

Sitki Cem Parlar, BSc[1, 2, 3], Eric Yu, PhD[1, 2, 3], Sajanth Kanagasingam[1, 2, 3], Meredith Zhang, MSc[1, 2, 3], Lang Liu, MSc[1, 2], Morvarid Ghamgosar Shahkhali BSc[1, 2, 3], Cloe Chantereault[1, 2, 3], Nathan Karpilovsky, BSc[3, 5], Donovan Worrall, BSc[3, 5], Rhalena A Thomas, PhD[3, 5], Konstantin Senkevich, PhD[2, 3, 6], Edward A Fon, MD[2, 3, 5], Ziv Gan-Or, PhD[1, 2, 3]*

Abstract

Background Genome-wide association studies (GWAS) have identified numerous risk loci for Parkinson's disease, yet identifying causal genes and mechanisms remains challenging due to non-coding associations and complex linkage disequilibrium.

Methods We prioritized genes within 147 GWAS loci using an XGBoost machine-learning model trained on 285 multi-omic features, including brain-specific eQTLs and single-cell expression. Prioritized candidates underwent gene- and domain-level rare variant burden analysis via optimal sequenced kernel association test (SKAT-O) across the Accelerating Medicines Partnership for Parkinson's Disease and UK Biobank cohorts (N case = 6,435, N proxy = 13,889, N control = 343,160).

Findings The model prioritized 406 genes with 48 features contributing to the prioritization. Meta-analysis of rare variants at the gene and domain levels replicated established associations (*GBA1*, *LRRK2*) and identified six novel potential risk genes *ANKRD27* driven by p.Arg21Cys, *UBXN2A*, *FAM171A1*, *ERCC8*, *BNC2*, and *ADNP*. Domain-level analysis uniquely uncovered a significant association in the zinc-finger domain of *ADNP*, which was masked in gene-level tests. Additionally, at cohort-level, one novel gene was nominated: *LRRC45* driven by p.Met607ArgfsTer57.

Interpretation Integrating machine-learning prioritization with gene- and domain-level burden testing identified novel genes potentially involved in Parkinson's disease, with further validation needed to elucidate causality and mechanisms.

Funding This study was supported by grants from the Galen and Hilary Weston Foundation, the Michael J. Fox Foundation, the Canadian Consortium on Neurodegeneration in Aging (CCNA), the Canada First Research Excellence Fund (CFREF). Lastly, the study received contributions from the G-Can (GBA1-Canada) Initiative. G-Can is supported by The Hilary and Galen Weston Foundation, Silverstein Foundation, and J. Sebastian van Berkom and Ghislaine Saucier. S.C.P. is supported by a Canadian Institutes of Health Research (CIHR) Canada Graduate Scholarship – Doctoral (CGS-D) award. Z.G.-O. holds a Fonds de recherche du Québec - Santé (FRQS) Chercheurs-boursiers award and is a William Dawson Scholar.

Author Affiliations:

[1]*Department of Human Genetics, McGill University, Montréal, Québec, Canada*

[2]*Montreal Neurological Institute-Hospital (The Neuro), McGill University, Montréal, Québec, Canada*

[3]*Department of Neurology and Neurosurgery, McGill University, Montréal, Québec, Canada*

[4]*DataTecnica LLC, Washington, DC, USA*

[5]*Early Drug Discovery Unit (EDDU), Montreal Neurological Institute-Hospital (The Neuro), Montreal, QC, Canada*

[6]*Department of Specialized Medicine, Division of Medical Genetics, McGill University Health Centre, Montreal, Québec, Canada.*

Correspondence to: Ziv Gan-Or

1033 Pine Avenue West, Room 310, Montréal (Québec), Canada, H3A 1A1

ziv.gan-or@mcgill.ca

Running title: ML prioritization and burden in PD

Keywords: Parkinson's disease, Machine learning, Rare variants, Burden analysis, Gene prioritization

Abbreviations: AMP-PD: Accelerating Medicines Partnership for Parkinson's Disease eQTL: Expression quantitative trait loci GWAS: Genome-wide association studies ML: Machine learning PD: Parkinson's disease SKAT-O: Optimal sequenced kernel association test UKBB: UK Biobank

Introduction

Parkinson's disease (PD) is the fastest-growing neurodegenerative disorder globally, affecting millions of individuals and imposing a substantial burden on healthcare systems.[1] While approximately 15% of cases have a family history of PD, the genetic architecture of the disease remains largely elusive, with a significant portion of heritability unexplained.[2] Large-scale genome-wide association studies (GWAS) have been instrumental in mapping the genetic landscape of PD, revealing numerous risk loci. However, the vast majority of these signals map to non-coding regions, making the identification of the specific causal variants, genes and mechanisms within these loci a major challenge.

To bridge the gap between statistical association and biological function, novel computational strategies are required. Recently, we demonstrated that machine learning approaches integrating multi-omic data can effectively prioritize high-confidence candidate genes within GWAS loci.[3] By leveraging tissue-specific transcriptomic and epigenetic features, these models move beyond simple genomic distance to identify drivers of disease risk. Furthermore, GWAS signals primarily capture common variants with small effect sizes, potentially neglecting the contribution of rare, highly penetrant variants.

Due to limited statistical power, rare variants are conventionally analyzed via burden testing, which aggregates variants into a single "gene-level" unit. While effective for well-established genes such as *GBA1*, *LRRK2*, and *SNCA*, this approach lacks functional resolution. Pathogenic mechanisms are often specific to distinct structural regions of a protein, such as the kinase domain of *LRRK2*,[4] meaning that aggregating variants across the entire gene may dilute potent signals with benign noise. Consequently, the current list of validated PD targets remains small, limiting our ability to uncover novel therapeutic targets.

In this study, we combined machine-learning-based gene prioritization on the most recent European PD GWAS study, which is the largest of its kind, with a high-resolution rare variant burden analysis. By demarcating proteins into their constituent functional domains, we aimed to increase the specificity of genetic association testing. This dual approach allows us to not only refine the list of prioritized candidates from European GWAS loci but also to uncover novel, domain-specific pathogenic mechanisms that are missed by traditional gene-level analyses.

Methods

Machine Learning and Prioritization Framework

To capture a comprehensive and high-yield map of the genetic architecture underlying PD, we constructed a composite set of risk loci by integrating the two largest available GWAS meta-analyses.[5,6] We incorporated summary statistics from the recent European GWAS and manually reinstated 13 unique loci from the previous meta-analysis by Nalls et al. (2019) that were absent in the newer release, resulting in a final dataset of 147 unique PD risk loci. To ensure broad capture of potential causal candidates, all protein-coding genes (N = 3105) located within a +/- 1mb window of the independent GWAS lead variants within each locus were extracted for prioritization.

Our prioritization strategy relied on an XGBoost[7], a gradient boosting machine-learning model trained to distinguish high-confidence risk genes from non-causal candidates. The training set was constructed using a "positive" class of 12 genes, comprising the established PD genes *GBA1*, *LRRK2*, *SNCA*, *GCH1*, *MAPT*, *TMEM175*, and *VPS13C*,[2] alongside the less established yet likely genes *SCARB2*, *CTSB*, *GRN*, *GALC*, and *GPNMB*.[8-11] All other protein-coding genes residing within the risk loci of these positive labelled genes (N = 191) were defined as the "negative" training set. To address the significant imbalance between the small number of positive labels and the large number of negative examples, we applied a specific weighting correction during training. We scaled the weight of the positive class by the ratio of negative-to-positive samples, ensuring the model optimizes for sensitivity towards the minority risk-gene class.

We characterized each gene using a high-dimensional set of 285 multi-omic features (Supplementary Table 1) derived from genomic, transcriptomic, and epigenomic datasets (Supplementary Table 2). Crucially, to account for heterogeneity across loci, rather than relying on absolute values, we applied a "neighbourhood scoring" method to standardize features relative to the maximum value observed within each specific locus. Key features included genomic distance metrics (log-transformed), variant severity scores, and brain-specific regulatory evidence from eQTLs, Hi-C, and snRNA-seq data. Training proceeded in a two-stage process:

first, recursive feature elimination was applied to remove redundant features and to reduce the dimensionality of the dataset, followed by hyperparameter tuning using 5-fold cross-validation on the refined feature set. The fully detailed model training process is provided in the Supplementary Methods (pp 1-2).

Gene-Level and Domain-Level Rare Variant Burden

Study Population and Sequencing

Our study population included whole-genome sequencing (WGS) data from the Accelerating Medicines Partnership for Parkinson's Disease (AMP-PD)[12] and the UK Biobank (UKBB).[13] To balance phenotypic rigor with statistical power, we performed two distinct meta-analyses. First, to prioritize clinical specificity, we performed a "limited" meta-analysis (Total N = 6,435 cases, 37,504 controls). This included the AMP-PD cohort and a subset of the UKBB containing PD cases identified strictly via clinical diagnosis, self-report, or death registry records, matched with 10x randomly sampled controls. Second, to maximize statistical power, we performed a "full" meta-analysis (Total N = 20,324 cases, 343,160 controls). This combined AMP-PD with the full UKBB dataset that included both PD cases and "proxy" cases defined as participants self-reporting a parental or sibling history of PD.

Sequencing data were aligned to the hg38 reference genome. We applied rigorous quality control at both the sample level (removing relatedness and non-European ancestry) and genetic level (filtering for depth and genotype quality). Quality control steps and software used are detailed in the Supplementary Methods (pp 3-4).

Gene and Domain Demarcation

A key innovation of our approach is the resolution of burden analysis at the functional domain level. For gene-level definitions, we utilized canonical coordinates and prioritized MANE Select transcripts as a biologically relevant reference.[14] To demarcate functional domains, we developed an automated pipeline integrating the Ensembl REST API[15] with InterPro.[16] We extracted protein features exclusively from Pfam-A, SMART, and PROSITE Profiles, filtering for features with validated InterPro Accessions to ensure consensus biological signatures.

Discontinuous fragments were collapsed into consensus ranges, and overlaps were resolved via a strict hierarchy (Pfam > SMART > PROSITE). Finally, the amino acid boundaries were mapped to hg38 genomic coordinates. The full gene and domain demarcation pipelines are detailed in the Supplementary Methods (pp 4-5).

Burden Analysis and Statistical Methods

Rare variants were annotated using Ensembl Variant Effect Predictor (VEP).[17] Further, we utilized the Combined Annotation Dependent Depletion (CADD) framework,[18] which integrates diverse genomic features to predict deleteriousness via a unified Phred-scaled score. For analysis, the variants were categorized into three functional groups: Non-synonymous (missense, in-frame indels, splice region), CADD > 20 (top 1% of predicted deleterious), and Loss-of-Function (LoF) (stop-gain, frameshift, splice acceptor/donor). Domain-level analysis was limited to non-synonymous category variants.

To evaluate individual variant associations, Odds Ratios (OR) and 95% Confidence Intervals (CI) were calculated using 2x2 contingency tables with a Haldane-Anscombe correction (adding 0.5 to all cells) to handle zero counts. P-values for variant-level tests were derived using the Wald test.

Rare variant burden testing was performed using the optimal sequence kernel association test (SKAT-O),[19] adjusting for sex, age, and the first five principal components. Results were meta-analyzed using Stouffer's Z-score method weighted by case counts. P-values were corrected using the False Discovery Rate (FDR) method, applied separately for gene-level and domain-level tests. Detailed statistical procedures are provided in the Supplementary Methods (p 5).

Role of the funding source

The funders of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report.

Results

Machine Learning Prioritizes 406 Genes for Parkinson's Disease Association

To capture a comprehensive and high-yield map of the genetic architecture underlying PD, we constructed a composite set of risk loci by integrating the two largest available GWAS meta-analyses. We incorporated summary statistics from the recent European GP2 study and manually reinstated unique loci from the previous meta-analysis that were absent in the newer release, resulting in a final dataset of 147 unique PD risk loci. To ensure broad capture of potential causal candidates, all protein-coding genes located within a +/- 1mb window of the independent lead variants within each locus were extracted for prioritization.

We prioritized the top three scoring genes in each locus. This yielded a total of 406 unique genes prioritized across the 147 risk loci (See Supplementary Table 3 and Figure 1). 23 genes were nominated multiple times due to overlapping neighbouring loci. Notable overlaps included *LSM7*, *SPPL2B*, and *TMPRSS9*, which were prioritized in three separate loci (122, 146, and 147). Other recurrent nominations included *BIRC6*, *SDHC*, *MED12L*, *AS3MT*, *TOX3*, *ADARB1*, *IQCB1*, and *ZSCAN31*, each emerging as the top-ranked candidate in paired loci. We assessed the distribution of probability scores among the 406 prioritized genes. A high probability score (>0.75) was assigned to 33 genes (8.1%), including more established genes such as *SNCA* (0.86), *LRRK2* (0.86), *GCH1* (0.84), and *CTSB* (0.91) as well as novel candidates like *ZYG11B* (0.92), *PXK* (0.92), *LMBRD1* (0.93). A further 142 genes (35.0%) received moderate probability scores between 0.50 and 0.75.

Conversely, the majority of the prioritized list showed lower confidence, with 231 genes (56.9%) scoring below 0.50. Notably, 123 genes (30.3%) had scores <0.25, indicating that for many loci, even the second or third-ranked candidates had weak predictive evidence compared to the top nomination. For example, while *CLCN3* (0.82) and *MFAP3L* (0.80) in locus 43 both showed strong support, other loci such as 51 (*E2F3*, 0.29) and 131 (*DSCAM*, 0.64; *MX1*, 0.10) showed a steep drop-off in probability after the primary candidate (See Supplementary Table 3).

Feature Contribution and Predictive Drivers

To interpret the machine learning model, we utilized Shapley Additive exPlanations (SHAP)[20] values to quantify the contribution of each feature to the final probability score. The instance-level contribution of these features is visualized in Figure 2, while the global hierarchy of feature importance is summarized in Figure 3.

Consistent with established GWAS biology,[21] genomic distance features (e.g., DIST_MIN, DIST_MEAN) were the dominant drivers of the model. As illustrated in the top rows of Figure 2, proximity to the locus risk variants provided a high baseline probability for the majority of nominated genes. However, the model demonstrated the capacity to override physical proximity when functional evidence was sufficiently strong. In 27 of the 147 risk loci (18.4%), the top-ranked candidate was not the gene with the lowest mean distance to the risk variants (See Supplementary Table 3). Beyond this distance-based prior, the model identified specific functional genomic signals as critical discriminators. As shown in Figure 3, the next most predictive features were cis-eQTL effect sizes derived from large-scale brain meta-analyses (BrainMeta) and the PsychENCODE consortium. This indicates that when candidates are equidistant from a risk variant, the model prioritizes genes with strong regulatory evidence in relevant tissue types. Additionally, we observed a distinct, high-specificity contribution from single-cell gene expression profiles. The model prioritized expression within specific *CALB1*-positive neuronal subpopulations (e.g., CALB1_CRYM_CCDC68 and CALB1_GEM). While these features had a lower global impact than distance, Figure 2 demonstrates that they contributed strongly to the scoring of specific gene subsets, suggesting they capture cell-type-specific vulnerabilities that broad tissue expression metrics miss.

Burden analyses reveal novel targets at both gene- and domain-level

To further validate the nominated genes and to identify novel rare variant burden associations, we performed rare variant burden analyses via SKAT-O in AMP-PD, UKBB, and UKBB with proxy cohorts (Table 1). The analysis included the 406 genes nominated via our machine-learning prioritization framework and 767 domain-level targets identified by our domain demarcation process (See Supplementary Table 4). Post quality control steps, the analysis

included a total of 27,777 unique rare variants from AMP-PD, 49,177 from UKBB, 155,399 from UKBB with proxy.

We subsequently performed two meta-analyses: (1) a limited meta-analysis with AMP-PD and UKBB (without proxies) and (2) a full meta-analysis with AMP-PD and UKBB (including proxies). The complete list of burden results is detailed in the Supplementary Tables 5 and 6 for gene- and domain-level analyses, respectively. The complete list of variants with detailed functional annotations, population frequencies, and OR calculations have been made accessible at: https://doi.org/10.5281/zenodo.18285356. The subset of findings reaching nominal significance ($P < 0.01$) in the full meta-analysis is presented in Tables 2 (gene-level) and 3 (domain-level). For these prioritized targets, detailed variant-level statistics across all functional categories are stratified by cohort in Supplementary Tables 7–9 (gene-level) and Supplementary Tables 10–12 (domain-level).

At gene-level, post-corrections, we identified significant associations ($P_{FDR} < 0.05$) in *GBA1* and *BNC2* in the limited meta-analysis, *GBA1*, *ANKRD27*, *FAM171A1*, *ERCC8*, and *UBXN2A* in the full meta-analysis, and *SNCA* and *LRCC45* genes in AMP-PD, which did not survive either meta-analysis. As expected, rare variants within *GBA1* and *SNCA*, two well-established PD risk genes showed significant associations: in the case of *GBA1,* the signal was across all three functional categories (non-synonymous, CADD > 20, and LoF) in the full meta-analysis, whereas, for the *SNCA* gene, the signal was limited to the AMP-PD cohort and the non-synonymous rare variants.

None of the novel genes identified in this analysis were the top-ranked candidate in their respective loci. *BNC2* (Locus 70) was prioritized as the third candidate. *ANKRD27* (Locus 124), *FAM171A1* (Locus 72), and *ERCC8* (Locus 45) were all ranked second in their respective loci, while *UBXN2A* (Locus 12) was the third candidate. Finally, *LRRC45* (Locus 119) was the second-highest ranked gene. Specific probability scores for each candidate are listed in Supplementary Table 3.

The sole novel signal in the limited meta-analysis, observed in *BNC2* (for variants with CADD score > 20), was driven by a significant cohort-level association in AMP-PD. This signal

involved 530 variants, mostly intronic, with no single variant emerging as a clear driver (See Supplementary Table 7), therefore this association should be considered with caution.

Among the novel hits from the full-meta-analysis, *ANKRD27* (Full meta-analysis P_{FDR} = 0.027 in the CADD > 20 category) was driven primarily by a single recurrent missense variant, p.Arg21Cys, which showed a consistent risk effect across all three cohorts (UKBB with proxy: OR = 1.20, 95% CI: 1.08–1.34, P = 1.03 x 10^{-3}; UKBB: OR = 1.29, 95% CI: 1.01–1.65, P =0.04; AMP-PD: OR = 1.21, 95% CI: 0.81–1.80, P = 0.36). The remaining three gene-level hits in the full-meta-analysis (*FAM171A1* and *ERCC8* in the LoF category; *UBXN2A* in the CADD > 20 category) did not implicate specific driver variants, appearing instead to result from the cumulative burden of multiple rare alleles (See Supplementary Tables 7–9).

In the case of *LRRC45*, significant signals were limited to the AMP-PD cohort in both the CADD > 20 (P_{FDR} = 6.28×10^{-4}) and LoF (P_{FDR} = 5.01×10^{-3}) categories. Both signals were driven by the same potentially protective frameshift variant p.Met607ArgfsTer57 found in 3 cases and 31 controls in AMP-PD (OR = 0.11, 95% CI: 0.04–0.38, P = 3.60 x 10^{-4}). This variant was not present in the UKBB cohort, which explains the limitation of signal only to AMP-PD.

At domain-level, post-corrections, non-synonymous rare variants in the *GBA1* 30 TIM-barrel and 30 beta-sandwich domains were found significantly associated, overlapping with the gene-level results. Furthermore, the *LRRK2* Kinase domain rare non-synonymous variants showed post-FDR significant association, which was not captured at gene-level for *LRRK2* (Variant level associations detailed in Supplementary Tables 10–12). Additionally, we identified a novel, significant domain-specific association in *ADNP* within its Zinc Finger domain. Importantly, *ADNP* did not reach significance in the gene-level burden test, demonstrating the ability of our domain-demarcation pipeline to dissect the structural localization of this burden. In the machine learning prioritization, *ADNP* was the second-ranked candidate in its locus, scoring below the top hit *PTPN1*. No clear variant-level association could be discerned as the association seemed to be driven by a cumulative effect of multiple variants within the domain (See Supplementary Tables 10–12).

Discussion

In this study, we leveraged the latest and largest European GWAS meta-analysis to systematically prioritize and interrogate the genetic architecture of PD. Starting from 147 unique risk loci, our machine-learning framework prioritized 406 candidate genes. We identified several recurrent nominations across adjacent loci (e.g., *LSM7*, *BIRC6*, *TOX3*), implying that these regions likely represent single risk signals driven by extensive linkage disequilibrium rather than independent associations. While our machine learning model relied heavily on genomic distance, it importantly integrated functional genomic features, such as brain-specific eQTLs and cell-type-specific expression, identifying candidates that were not always the closest to risk variants in the locus.

Through our rare variant burden analysis, which included the demarcation of 406 candidate genes into 767 functional domains, we verified seven novel genes prioritized by machine learning, one of which was validated exclusively through domain-level resolution. This complementary approach proved critical for these findings as the seven genes were prioritized as second and third candidates respectively based on the machine learning model. By using the framework to define a broad search space rather than relying on a single high-confidence prediction, we enabled the rare variant burden analysis to identify risk associations that would have been overlooked by the prioritization model alone. In the gene-level analysis, we identified robust associations in *BNC2* in the limited-meta-analysis and in *ANKRD27, FAM171A1, ERCC8*, and *UBXN2A* within the full meta-analysis. The signal in *BNC2* was mostly driven by intronic variants with high CADD scores exclusively in the AMP-PD cohort. Given the lack of replication and the non-coding nature of the drivers, it remains unclear if this signal reflects *BNC2* biology or regulatory effects on neighboring genes, necessitating caution in interpretation.

The association in *ANKRD27* is particularly compelling given its known role in endosomal trafficking and interaction with the retromer complex, a pathway heavily implicated in PD pathology containing a PD-associated *VPS35* gene.[22] This signal was driven by the recurrent missense variant p.Arg21Cys, which showed a consistent risk effect across all cohorts. Notably, a non-coding variant within *ANKRD27* (rs11672605) shows genome-wide significance (1.61×10^{-8}) in European GWAS[5] but is in high linkage disequilibrium $(r^2 > 0.8)$ with the top

locus hit (rs11669800), making it difficult to distinguish the causal driver via GWAS alone. Our rare variant finding suggests that *ANKRD27* and specifically the p.Arg21Cys variant may be the functional drivers of this locus.

The associations in *FAM171A1* and *ERCC8* were driven largely by LoF variants. *ERCC8* encodes a component of the ubiquitin ligase complex involved in transcription-coupled nucleotide excision repair, and its dysfunction causes Cockayne syndrome, a rare segmental progeroid disorder characterized by growth failure, severe photosensitivity, and progressive neurodegeneration.[23] Given the accumulation of DNA damage and mitochondrial dysfunction in PD, *ERCC8* represents a plausible link to genomic instability. *FAM171A1* offers a plausible mechanistic link to PD pathology; recent evidence suggests that along with its paralog *FAM171A2*, *FAM171A1* may regulate the levels of progranulin (*GRN*), a recognized PD candidate gene.[24] The role of *UBXN2A*, a ubiquitin-binding protein involved in proteasomal degradation,[25] remains less established in the context of neurodegeneration. As the signals concerning these novel genes were cumulative rather than variant-specific, functional studies targeting the expression of these genes are warranted to elucidate their role in PD pathobiology.

Conversely, our analysis of *LRRC45* in the AMP-PD cohort revealed a potentially protective LoF signal driven by the frameshift variant p.Met607ArgfsTer57. *LRRC45* is a centrosomal protein involved in ciliogenesis.[26] Since PD pathology has been linked to defects in primary cilia and centrosomal cohesion (notably involving *LRRK2*),[27] a protective LoF effect could imply that reduced *LRRC45* activity mitigates specific cellular deficits, or that the gene normally exerts a deleterious gain-of-function in the disease state.

Our results emphasize that functional specificity via domain-level analysis can uncover insights masked by global gene-level tests. A prime example is *LRRK2*; consistent with literature showing that the p.G2019S variant exerts its effect via the kinase domain,[4,28] our analysis successfully isolated the burden to this specific domain. Parallel to this, we identified a novel domain-specific association in *ADNP* restricted to its Zinc Finger domain. *ADNP* is a neuroprotective transcription factor essential for brain formation,[29] and while it failed to reach significance at the gene level, the domain-resolved signal highlights the power of this approach to filter out protein-wide noise and pinpoint potentially risk-influencing motifs.

Our study has several limitations. First, both the GWAS priors and rare variant burden analyses were restricted to populations of European ancestry. Larger, diverse datasets are needed for more global conclusions. Second, while quality control was rigorous, slight differences between the AMP-PD and UKBB pipelines may introduce minor heterogeneity. Finally, the inclusion of proxy cases in the UKBB meta-analysis warrants restraint in interpretation, as proxy-based phenotypes share only half the genetic liability of clinically diagnosed PD.

In conclusion, our study shows that integrating machine learning-based gene prioritization with domain-resolution burden analysis offers a robust framework for dissecting the complex genetic architecture of PD. By prioritizing candidates based on multi-omic features and proximity to non-coding GWAS hits, we successfully identified novel risk genes *BNC2*, *ANKRD27*, *FAM171A1*, *ERCC8*, and *LRRC45* and uncovered a domain-level signal in *ADNP* (Zinc Finger) that was masked at the gene level. These findings not only highlight specific biological processes, such as endosomal trafficking, DNA repair, and ciliary function, but also demonstrate that multi-omics based, high-resolution genomic interrogation is highly valuable for translating GWAS loci into actionable biological targets. Future experimental validation of these candidates is critical to fully decipher their mechanistic roles in disease pathogenesis.

Data sharing

The WGS cohorts accessed and analysed in this study for the purposes of the burden analysis are accessible (conditional to institutional approval) at: AMP-PD Knowledge Platform (https://www.amp-pd.org) and the UK Biobank Research Analysis Platform. All information directly relevant to the study is included in the text and the supplementary files of this article, except for the full range of analysed variants, which have been made accessible online at the link: https://doi.org/10.5281/zenodo.18285356. All the scripts used in the machine learning prioritization and burden analyses are shared at the GitHub repository link: https://github.com/gan-orlab/ML_and_Burden_PD.

Acknowledgements

We extend our deepest gratitude to the participants from all cohorts included in this study for their essential contributions to this work. This research was performed using the NeuroHub infrastructure and was supported in part by the Canada First Research Excellence Fund, awarded to McGill University for the Healthy Brains, Healthy Lives (HBHL) initiative, as well as by Calcul Québec and the Digital Research Alliance of Canada. This study also made use of data from the UK Biobank under Application Number 45551.

Data for this article were obtained on 10 June 2025 from the Accelerating Medicines Partnership® (AMP®) Parkinson's Disease (AMP PD) Knowledge Platform, release version 4.2. Further details on AMP-PD are available at https://www.amp-pd.org. The AMP® PD program operates as a public-private partnership managed by the Foundation for the National Institutes of Health (FNIH) and funded by the National Institute of Neurological Disorders and Stroke (NINDS) in collaboration with the Aligning Science Across Parkinson's (ASAP) initiative. Industry partners include Celgene Corporation (a subsidiary of Bristol-Myers Squibb Company), GlaxoSmithKline plc (GSK), The Michael J. Fox Foundation for Parkinson's Research (MJFF), Pfizer Inc., AbbVie Inc., Sanofi US Services Inc., and Verily Life Sciences. "ACCELERATING MEDICINES PARTNERSHIP" and "AMP" are registered service marks of the U.S. Department of Health and Human Services.

The clinical data and biosamples used in this article were derived from the following cohorts: (i) the BioFIND study (MJFF and NINDS); (ii) the Harvard Biomarkers Study (HBS) and the Stephen & Denise Adams Center for Parkinson's Disease Research at Yale School of Medicine (CPDR-Y); (iii) the National Institute on Aging (NIA) International Lewy Body Dementia Genetics Consortium Genome Sequencing in Lewy Body Dementia Case-control Cohort (LBD); (iv) the MJFF LRRK2 Cohort Consortium (LCC); (v) the NINDS Parkinson's Disease Biomarkers Program (PDBP); (vi) the MJFF Parkinson's Progression Markers Initiative (PPMI); (vii) the NINDS Study of Isradipine as a Disease-modifying Agent in Subjects With Early Parkinson Disease, Phase 3 (STEADY-PD3); and (viii) the NINDS Study of Urate Elevation in Parkinson's Disease, Phase 3 (SURE-PD3).

BioFIND is sponsored by MJFF with support from NINDS; current study information is available at michaeljfox.org/news/biofind. The BioFIND investigators did not participate in the review of this manuscript's data analysis or content. Genome sequence data for the Lewy body dementia case-control cohort were generated at the U.S. National Institutes of Health Intramural Research Program, supported in part by the NIA (program #: 1ZIAAG000935) and NINDS (program #: 1ZIANS003154). The Harvard Biomarker Study (HBS) is a collaboration of HBS investigators (listed at https://www.bwhparkinsoncenter.org/biobank/) and is funded through philanthropy, NIH, and non-NIH sources. The Stephen & Denise Adams Center for Parkinson's Disease Research of Yale School of Medicine is similarly funded through philanthropy, NIH, and non-NIH sources. Neither the HBS nor CPDR-Y investigators participated in reviewing the data analysis or content of this manuscript.

Data were obtained from the MJFF-sponsored LRRK2 Cohort Consortium (LCC). Current study information can be found at https://www.michaeljfox.org/biospecimens. The LCC investigators did not review the data analysis or manuscript content. PPMI is sponsored by MJFF and supported by a consortium of scientific partners (full list available at https://www.ppmi-info.org/about-ppmi/who-we-are/study-sponsors). For up-to-date study information, visit www.ppmi-info.org. PPMI investigators did not participate in the review of the data analysis or content of this manuscript. The Parkinson's Disease Biomarker Program (PDBP) consortium is supported by NINDS at the NIH. A full list of PDBP investigators is available at https://pdbp.ninds.nih.gov/policy. PDBP investigators did not review the data analysis or content

of this manuscript. STEADY-PD3 is funded by NINDS at the NIH with support from MJFF and the Parkinson Study Group. Additional information is available at https://clinicaltrials.gov/ct2/show/study/NCT02168842. The STEADY-PD3 investigators did not review the data analysis or content of this manuscript. SURE-PD3 is funded by NINDS at the NIH with support from MJFF and the Parkinson Study Group. Additional information is available at https://clinicaltrials.gov/ct2/show/NCT02642393. The SURE-PD3 investigators did not review the data analysis or content of this manuscript.

Funding

This work was supported by grants from the Galen and Hilary Weston Foundation, the Michael J. Fox Foundation, the Canadian Consortium on Neurodegeneration in Aging (CCNA), and the Canada First Research Excellence Fund (CFREF), awarded to McGill University for the Healthy Brains for Healthy Lives (HBHL) initiative. We also acknowledge contributions from the G-Can (GBA1-Canada) Initiative, an open-science collaboration dedicated to GBA1-associated neurodegeneration. G-Can is supported by the Galen and Hilary Weston Foundation, the Silverstein Foundation, and J. Sebastian van Berkom and Ghislaine Saucier. S.C.P. is supported by a Canadian Institutes of Health Research (CIHR) Canada Graduate Scholarship – Doctoral (CGS-D) award. Z.G.-O. holds a Fonds de recherche du Québec - Santé (FRQS) Chercheurs-boursiers award and is a William Dawson Scholar.

Author Contributions

Contributors S.C.P., E.Y., and Z.G.-O. conceived and designed the study. E.Y. developed the machine learning framework and performed the prioritization analysis. S.C.P. performed the domain demarcation and rare-variant burden analysis and wrote the first draft of the manuscript. S.K. and M.Z. assisted with data extraction and quality control. All authors contributed to the interpretation of the data and critical revision of the manuscript. Z.G.-O. obtained funding and supervised the study. E.Y., L.L., and R.A.T. verified the data underlying the machine learning analysis; S.C.P. and Z.G.-O. verified the underlying data for the burden analysis. All authors had full access to all the data in the study and had final responsibility for the decision to submit for publication.

Competing interests

Z. G.-O. received consultancy fees from Lysosomal Therapeutics Inc. (LTI), Idorsia, Prevail Therapeutics, Inceptions Sciences (now Ventus), Neuron23, Handl Therapeutics, UCB, Capsida, Vanqua Bio, Congruence Therapeutics, Ono Therapeutics, Denali, Bial Biotech, Bial, EG427, Takeda, Jazz Pharmaceuticals, Simcere, Guidepoint, Lighthouse, and Deerfield. K.S. received consultancy fees from Acurex.

Supplementary material

Supplementary material is provided in the supplementary tables and the methods Supplementary Methods.

References

1.   Dorsey ER, Elbaz A, Nichols E, et al. Global, regional, and national burden of Parkinson's disease, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *The Lancet Neurology* 2018; **17**(11): 939-53.

2.   Blauwendraat C, Nalls MA, Singleton AB. The genetic architecture of Parkinson's disease. *The Lancet Neurology* 2020; **19**(2): 170-8.

3.   Yu E, Larivière R, Thomas RA, et al. Machine learning nominates the inositol pathway and novel genes in Parkinson's disease. *Brain* 2024; **147**(3): 887-99.

4.   Parlar SC, Senkevich K, Yu E, et al. LRRK2 rare-variant per-domain genetic burden in Parkinson's Disease: association confined to the kinase domain. *npj Parkinson's Disease* 2025; **11**(1): 102.

5.   Leonard HL. Novel Parkinson's Disease Genetic Risk Factors Within and Across European Populations. *medRxiv* 2025.

6.   Nalls MA, Blauwendraat C, Vallerga CL, et al. Identification of novel risk loci, causal insights, and heritable risk for Parkinson's disease: a meta-analysis of genome-wide association studies. *The Lancet Neurology* 2019; **18**(12): 1091-102.

7.   Chen T. XGBoost: A Scalable Tree Boosting System. *Cornell University* 2016.

8.   Diaz-Ortiz ME, Seo Y, Posavi M, et al. GPNMB confers risk for Parkinson's disease through interaction with α-synuclein. *Science* 2022; **377**(6608): eabk0637.

9.   Jones-Tabah J, He K, Karpilovsky N, et al. The Parkinson's disease risk gene cathepsin B promotes fibrillar alpha-synuclein clearance, lysosomal function and glucocerebrosidase activity in dopaminergic neurons. *Molecular neurodegeneration* 2024; **19**(1): 88.

10.   Nalls MA, Blauwendraat C, Sargent L, et al. Evidence for GRN connecting multiple neurodegenerative diseases. *Brain Communications* 2021; **3**(2): fcab095.

11.   Robak LA, Jansen IE, Van Rooij J, et al. Excessive burden of lysosomal storage disorder gene variants in Parkinson's disease. *Brain* 2017; **140**(12): 3191-203.

12.   Iwaki H, Leonard HL, Makarious MB, et al. Accelerating medicines partnership: Parkinson's disease. Genetic resource. *Movement Disorders* 2021; **36**(8): 1795-804.

13.   Consortium UBW-GS. Whole-genome sequencing of 490,640 UK Biobank participants. *Nature* 2025; **645**(8081): 692.

14.   Morales J, Pujar S, Loveland JE, et al. A joint NCBI and EMBL-EBI transcript set for clinical genomics and research. *Nature* 2022; **604**(7905): 310-5.

15.   Martin FJ, Amode MR, Aneja A, et al. Ensembl 2023. *Nucleic acids research* 2023; **51**(D1): D933-D41.

16.   Paysan-Lafosse T, Blum M, Chuguransky S, et al. InterPro in 2022. *Nucleic acids research* 2023; **51**(D1): D418-D27.

17.   McLaren W, Gil L, Hunt SE, et al. The ensembl variant effect predictor. *Genome biology* 2016; **17**(1): 122.

18.   Schubach M, Maass T, Nazaretyan L, Röner S, Kircher M. CADD v1. 7: using protein language models, regulatory CNNs and other nucleotide-level scores to improve genome-wide variant predictions. *Nucleic acids research* 2024; **52**(D1): D1143-D54.

19.   Lee S, Emond MJ, Bamshad MJ, et al. Optimal unified approach for rare-variant association testing with application to small-sample case-control whole-exome sequencing studies. *The American Journal of Human Genetics* 2012; **91**(2): 224-37.

20.   Lundberg SM, Lee S-I. A unified approach to interpreting model predictions. *Advances in neural information processing systems* 2017; **30**.

21.   Mountjoy E, Schmidt EM, Carmona M, et al. An open approach to systematically prioritize causal variants and genes at all published human GWAS trait-associated loci. *Nature genetics* 2021; **53**(11): 1527-33.

22.   Zimprich A, Benet-Pagès A, Struhal W, et al. A mutation in VPS35, encoding a subunit of the retromer complex, causes late-onset Parkinson disease. *The American Journal of Human Genetics* 2011; **89**(1): 168-75.

23.   Chen J, Su W, Gao D, et al. Clinical and genetic analysis of ERCC8-Related cockayne syndrome: hepatic dysfunction as a biomarker, anhidrosis as a rare feature, and rehabilitation outcomes for ankle contractures. *Frontiers in Genetics* 2025; **16**: 1591551.

24.   Xu W, Han S-D, Zhang C, et al. The FAM171A2 gene is a key regulator of progranulin expression and modifies the risk of multiple neurodegenerative diseases. *Science Advances* 2020; **6**(43): eabb3063.

25.   Sane S, Srinivasan R, Potts RA, et al. UBXN2A suppresses the Rictor-mTORC2 signaling pathway, an established tumorigenic pathway in human colorectal cancer. *Oncogene* 2023; **42**(21): 1763-76.

26.   Kurtulmus B, Yuan C, Schuy J, et al. LRRC45 contributes to early steps of axoneme extension. *Journal of cell science* 2018; **131**(18): jcs223594.

27.   Khan SS, Jaimon E, Lin Y-E, et al. Loss of primary cilia and dopaminergic neuroprotection in pathogenic LRRK2-driven and idiopathic Parkinson's disease. *Proceedings of the National Academy of Sciences* 2024; **121**(32): e2402206121.

28.   Sosero YL, Gan☐Or Z. LRRK2 and Parkinson's disease: from genetics to targeted therapy. *Annals of clinical and translational neurology* 2023; **10**(6): 850-64.

29.   Pinhasov A, Mandel S, Torchinsky A, et al. Activity-dependent neuroprotective protein: a novel gene essential for brain formation. *Developmental Brain Research* 2003; **144**(1): 83-90.

Figure legends

Figure 1. Probability score of the Parkinson's disease genome-wide association study candidate genes. This figure shows the probability scores from the machine learning model for each locus in the Parkinson's disease genome-wide association study loci sorted in descending order. For each gene, the top non-distance feature was used to colour the data.

Figure 2. Heat map of feature importance. The heat map is generated using Shapley Additive exPlanations (SHAP) value for the top candidate gene in each locus. The plot at the top represents the probability score of each gene.

Figure 3. Feature importance for the Parkinson's disease genome-wide association study gene prioritization model. Bee-swarm plot of feature importance using Shapley Additive exPlanations values along with the distribution of genes based on feature value.

Tables

Table 1. Study population demographics for rare variant burden analysis

Cohort	Cases			Controls		
	N	Mean age ± SD	Males, %	N	Mean age ± SD	Males, %
AMP-PD	3,051	64.38 (9.74)	1,903 (62.4%)	3,667	67.91 (8.02)	1,713 (46.7%)
UKBB	3,384	62.78 (5.33)	2,157 (63.7%)	33,837	56.82 (8.00)	15,679 (46.3%)
UKBB with proxy	17,273	59.17 (7.07)	7,934 (45.9%)	339,493	56.81 (7.99)	157,225 (46.3%)

SD – standard deviation, AMP-PD – Accelerating Medicines Partnership – Parkinson Disease, UKBB – UK biobank.

Table 2. Gene-level rare variant burden analysis results nominally significant ($P < 0.01$) in the full meta-analysis

Gene	Variant Type	AMP-PD P_{FDR}	UKBB P_{FDR}	Limited Meta-analysis P_{FDR}	UKBB with proxy P_{FDR}	Full Meta-analysis P_{FDR}
GBA1	CADD>20	0.07	6.55×10^{-9}	1.70×10^{-10}	3.68×10^{-17}	9.94×10^{-20}
GBA1	Non-synonymous	0.64	1.61×10^{-18}	4.34×10^{-14}	6.81×10^{-15}	1.25×10^{-15}
GBA1	LoF	0.66	0.97	0.82	0.02	7.65×10^{-3}
ANKRD27	CADD>20	0.98	0.36	0.76	2.26×10^{-3}	0.03
UBXN2A	CADD>20	0.95	0.59	0.81	0.02	0.04
ERCC8	LoF	0.84	0.59	0.68	0.06	0.04
FAM171A1	LoF	0.94	0.59	0.76	0.02	0.04
LRRK2	CADD>20	0.92	0.59	0.81	0.03	0.06
LRRK2	Non-synonymous	0.97	0.59	0.88	0.02	0.12
UBXN2A	Non-synonymous	0.95	0.59	0.81	0.07	0.18
ADNP	Non-synonymous	0.95	0.85	0.88	0.11	0.24
LRRC45	LoF	6.28×10^{-4}	0.87	0.08	0.99	0.28
BNC2	CADD>20	0.04	0.59	7.69×10^{-3}	0.99	0.28
CACUL1	Non-synonymous	0.70	0.95	0.81	0.85	0.48
KCNMB3	CADD>20	0.81	0.88	0.76	0.83	0.52
SNCA	Non-synonymous	2.67×10^{-3}	0.95	0.37	0.99	0.52
MTMR7	CADD>20	0.84	0.67	0.76	0.74	0.52

NCAM1	CADD>20	0.90	0.80	0.81	0.57	0.52

AMP-PD – Accelerating Medicines Partnership – Parkinson Disease, UKBB – UK biobank, CADD – Combined Annotation Dependent Depletion.

Meta-analysis definitions: (1) Limited: AMP-PD + UKBB PD cases (ICD, self-report, death records; N = 6,435 cases, 37,504 controls); (2) Full: AMP-PD + UKBB PD cases and proxies (N = 20,324 cases, 343,160 controls).

Table 3. Domain-level rare variant burden analysis results nominally significant ($P < 0.01$) in the full meta-analysis

Gene	Domain	Variant Type	AMP-PD P_{FDR}	UKBB P_{FDR}	Limited Meta-analysis P_{FDR}	UKBB with proxy P_{FDR}	Full Meta-analysis P_{FDR}
GBA1	Glycosyl hydrolase family 30 TIM-barrel domain	Non-synonymous	0.95	1.13×10^{-15}	3.40×10^{-12}	3.02×10^{-12}	5.14×10^{-13}
LRRK2	Protein kinase domain	Non-synonymous	0.95	0.59	0.83	9.68×10^{-12}	3.01×10^{-10}
GBA1	Glycosyl hydrolase family 30 beta sandwich domain	Non-synonymous	0.95	2.10×10^{-4}	1.93×10^{-3}	3.40×10^{-5}	1.88×10^{-5}
ADNP	ADNP zinc finger domain	Non-synonymous	0.95	0.86	0.87	3.23×10^{-3}	7.88×10^{-3}
NOL7	U3 small nucleolar RNA-associated protein NOL7 C-terminal	Non-synonymous	0.95	0.70	0.83	0.38	0.29
TOX3	High mobility group box domain	Non-synonymous	0.95	0.86	0.83	0.38	0.29
BNC2	Zinc finger C2H2-type	Non-synonymou	0.95	0.81	0.83	0.45	0.40

		s					
UBXN2A	SEP domain	Non-synonymous	0.95	0.59	0.83	0.37	0.46
KLHL29	BTB/Kelch-associated	Non-synonymous	0.95	0.59	0.83	0.45	0.71
SNRNP70	RNA recognition motif domain	Non-synonymous	0.95	0.94	0.83	0.99	0.71

AMP-PD – Accelerating Medicines Partnership – Parkinson Disease, UKBB – UK biobank.

Meta-analysis definitions: (1) Limited: AMP-PD + UKBB PD cases (ICD, self-report, death records; N = 6,435 cases, 37,504 controls); (2) Full: AMP-PD + UKBB PD cases and proxies (N = 20,324 cases, 343,160 controls).