Differential association of body mass index with hypoglossal nerve stimulation efficacy by pharyngeal collapse pattern in obstructive sleep apnea

Daniel Vena[1], Eric J. Kezirian[2], Andrew Wellman[1], David Kent[3], Mark D'Agostino[4,5], Joao L.G.C. Monteiro[1], Azarbarzin A[1], Tom Chen[6], Ludovico Messineo[1], Neda Esmaeili[1], Scott A. Sands[1]*, Phillip Huyett[7]*.

[1]Division of Sleep and Circadian Disorders, Brigham and Women's Hospital and Harvard Medical School, Boston, MA.

[2]Department of Head and Neck Surgery, David Geffen School of Medicine at UCLA, Los Angeles, California.

[3]Department of Otolaryngology-Head and Neck Surgery, Vanderbilt University Medical Center, Nashville, Tennessee.

[4]Southern New England Ear, Nose and Throat Group, Middlesex Hospital, Middlesex, Connecticut.

[5]Department of Surgery, Yale University School of Medicine, New Haven, Connecticut.

[6]Department of Population Medicine, Harvard Medical School and Harvard Pilgrim Health Care Institute, Boston, MA

[7]Department of Otolaryngology, Massachusetts Eye and Ear, Boston, MA

*,[†]Equal contributions

Correspondence:

Daniel Vena

221 Longwood Ave., Boston MA, 02115

Division of Sleep and Circadian Disorders, Department of Medicine, Brigham & Women's Hospital & Harvard Medical School, Boston, Massachusetts.

Tel: +1-617-732-8483; E-mail: dvena@bwh.harvard.edu

Author Contributions:

Conception and design of the work: DV, SAS, PH

Data collection/management: DV, JLGCM, EK, PH, DK, MD

Statistical analysis: DV, SAS, TC

Data interpretation: DV, EK, AW, DK, MD, JLGCM, AA, SAS, PH

Drafting the article: DV, SAS, PH

Critical revision of the article: DV, EK, AW, DK, MD, JLGCM, TC, AA, SAS, PH

Final approval of the version to be published: DV, EK, AW, DK, MD, JLGCM, AA, TC, DPW, SAS, PH

EJK holds a grant from the National Institutes of Health (HL160993).

AW reports grants from the National Institutes of Health (HL102321 and HL128658).

SAS was supported by the American Thoracic Society (15SDG25890059) and NIH NHLBI (R01HL146697).

Disclosures:

DV receives personal fees as a consultant and grant funding from Inspire Medical Systems.

EJK is a consultant for Cryosa, Nyxoah, and Berendo Scientific. EJK is a co-inventor related to intellectual property for Magnap and Berendo Scientific. EJK has previously received research funding from Inspire Medical Systems.

AA reports grant support from Somnifix and serves as a consultant for Respicardia, Eli Lilly, Inspire, Cerebra and Apnimed. Apnimed is developing pharmacological treatments for Obstructive Sleep Apnea. AA's interests were reviewed by Brigham and Women's Hospital and Mass General Brigham in accordance with their institutional policies.

AW works as a consultant for Apnimed, Nox, Inspire, Mosanna, Takeda, and iNOS. He has received grants from Prosomnus. He also has a financial interest in Apnimed, Mosanna, and iNOS, which are developing therapies for sleep apnea. Dr. Wellman's interests were reviewed and are managed by Brigham and Women's Hospital and Mass General Brigham in accordance with their conflict-of-interest policies.

DK has received research support from Inspire Medical Systems, Inc. and Invicta Medical, Inc. DK serves as a consultant for Invicta Medical, Inc. and is a scientific advisory board member for Nyxoah SA. DK also has intellectual property interests and has received research support from Nyxoah SA.

DPW is a consultant for Apnimed, Philips, Nidra, Mosanna, Onera, Bairiton, Xtrodes, Cerebra Health, Cryosa, Resonea and SleepRes;

SS received grant support from Apnimed, Prosomnus, and Dynaflex, and has served as a consultant for Apnimed, Nox Medical, Inspire Medical Systems, Eli Lilly, Respicardia, LinguaFlex, and Achaemenid. He receives royalties for intellectual property pertaining to combination pharmacotherapy for sleep apnea via his Institution. He is also co-inventor of intellectual property pertaining to wearable sleep apnea phenotyping also via his Institution. He has received equity in Achaemenid, a company commercializing biosensor technology for monitoring oral appliance treatment efficacy. He is also co-inventor of intellectual property pertaining to wearable sleep apnea phenotyping also via his Institution. His industry interactions are actively managed by his Institution.

PH serves as an educational consultant for Inspire Medical Systems and receives research support from Inspire Medical Systems and Nyxoah

The funders were not involved in the study design, collection, the writing of this article or the decision to submit it for publication.

Running title: BMI and pharyngeal collapse pattern on HGNS efficacy.

Key words: Sleep Apnea, Obstructive; Drug-induced Sleep Endoscopy; Site of pharyngeal collapse

Word count: 3403

ABSTRACT

Background. Hypoglossal nerve stimulation (HGNS) is an established surgical therapy for obstructive sleep apnea (OSA), yet treatment response is variable and appears to be influenced by both body mass index (BMI) and pattern of upper airway collapse. Whether excess body weight differentially affects HGNS efficacy across pharyngeal collapse patterns remains unclear.

Methods. We combined data from two independent HGNS cohorts (n=760) to examine whether the association between BMI and HGNS efficacy depends on pharyngeal collapse pattern identified by drug-induced sleep endoscopy. Collapse was categorized as predominantly laterally directed or anteroposterior (AP). Multivariable regression models assessed HGNS efficacy—quantified primarily as percent reduction in apnea–hypopnea index (AHI) on titration polysomnography and secondarily as treatment success (≥50% AHI reduction to <15 events/h)—while testing for effect modification by collapse pattern and adjusting for baseline AHI, partial collapse, surgical center, follow-up sleep study type, and prior or concomitant pharyngeal surgery.

Results. Increasing BMI was associated with substantially lower HGNS efficacy among patients with laterally directed collapse, whereas the relationship between BMI and efficacy was attenuated in those with AP collapse. Specifically, each 5 kg/m² increase in BMI was associated with a −19.7% (95% CI: −33.2 to −6.2) greater reduction in efficacy in lateral collapse compared with −3.8% (−8.0 to 0.36) in AP collapse (interaction p=0.027). Higher BMI also corresponded to reduced odds of treatment success in lateral collapse (odds ratio 4.4 [95% CI: 1.4–14.3] per 5 kg/m²), with no significant association observed in AP collapse (1.1 [0.75–1.5]; interaction p=0.023).

Conclusions. The influence of BMI on HGNS treatment response differs meaningfully by pharyngeal collapse phenotype. Incorporating collapse pattern into BMI-based eligibility criteria may improve patient selection and optimize HGNS outcomes.

Word count: 250

INTRODUCTION

Unilateral hypoglossal nerve stimulation (HGNS; Inspire Medical Systems) has become an increasingly utilized therapy for obstructive sleep apnea (OSA) in patients who are unable to tolerate positive airway pressure[1]. Despite its growing use, HGNS demonstrates variable efficacy, with a substantial proportion of patients failing to achieve an adequate reduction in apnea–hypopnea index (AHI)[2-9].

Two factors that have been consistently associated with reduced HGNS efficacy are elevated body mass index (BMI)[10] and pharyngeal collapse patterns with a prominent lateral component, including complete concentric collapse (CCC) at the palate and complete oropharyngeal lateral wall (OLW) collapse[5,11-14]. Higher BMI is associated with increased upper airway collapsibility, which may reduce the ability of HGNS prevent upper airway collapse[15-18]. Similarly, lateral pharyngeal collapse may limit HGNS effectiveness, as stimulation primarily produces anteroposterior displacement of the tongue and adjacent structures, potentially leaving laterally directed collapse unresolved. However, the interaction between BMI and pharyngeal collapse pattern in determining HGNS efficacy has not been well characterized.

Clarifying this interaction has important implications for clinical decision-making. Patients with higher BMI may still respond favorably to HGNS when collapse occurs at sites that are more directly targeted by stimulation, such as the tongue base. Conversely, patients with laterally directed collapse may experience good treatment response in the setting of lower BMI. Although lateral collapse patterns are more common at higher BMI, obesity does not uniformly predict the presence of lateral collapse, underscoring the need to consider both factors together when evaluating HGNS candidacy[19]. This issue is particularly relevant given recent changes to regulatory guidance that have expanded BMI eligibility for HGNS despite limited efficacy data.

The present study evaluates the relationship between BMI and HGNS efficacy across different pharyngeal collapse patterns using pooled data from two independent HGNS cohorts. We hypothesized that greater BMI would be associated with reduced HGNS efficacy in collapse patterns that are less responsive to stimulation, such as CCC and complete OLW collapse, but would have a smaller effect in predominantly anteroposterior collapse patterns.

METHODS

Study Design and Cohorts

This study pooled data from two independent cohorts of patients treated with HGNS for OSA: a multicenter retrospective cohort (n = 343)[5] and a prospective, single-center observational cohort (n = 419) drawn from the Mass Eye and Ear Sleep Surgery Registry. Both cohorts included patient demographics, BMI, pharyngeal collapse characteristics assessed by drug-induced sleep endoscopy (DISE), and baseline and post-treatment measures of OSA severity. BMI data were missing for two patients in the retrospective cohort, resulting in a total analytic sample of 760 patients.

In the prospective cohort, 560 patients who underwent HGNS implantation between October 2019 and June 2024 were enrolled in a study evaluating factors associated with HGNS efficacy (IRB: 2020P000156). Of these, 419 patients had completed post-implantation sleep testing and were included in the present analysis. Patients who did not undergo follow-up sleep testing due to patient preference or insurance denial remained under routine clinical follow-up. Patients who underwent tonsillectomy (n = 35), functional expansion pharyngoplasty (n = 10), or partial uvulectomy (n = 3) either prior to or concomitant with HGNS implantation without repeat DISE were included, with adjustment for these procedures in statistical analyses as described below.

HGNS Implantation and Sleep Study Assessment

All patients underwent DISE prior to HGNS implantation. Baseline AHI was assessed using either a clinical home sleep apnea test or an in-laboratory polysomnogram (PSG) performed prior to DISE evaluation. Patients subsequently underwent unilateral HGNS implantation using a standard surgical approach (Inspire Medical Systems, Golden Valley, MN). Device activation occurred approximately one month after implantation, followed by post-activation sleep testing to assess and optimize therapeutic efficacy.

Most patients underwent an in-laboratory titration PSG (tPSG) after device activation. A subset of patients in the prospective cohort (n = 62) elected to undergo home sleep apnea testing following subjective optimization of therapy at home based on symptom improvement and tolerability. Across cohorts, post-activation care pathways varied over time, with tPSG initially recommended at one month after activation and later updated to approximately three months. For patients undergoing tPSG, AHI was calculated during periods of optimal stimulation.

In the prospective cohort, sleep studies were scored according to American Academy of Sleep Medicine guidelines, with hypopneas defined by a 3% oxygen desaturation criterion[20]. In the multicenter retrospective cohort, sleep study scoring was not standardized across sites; however, scoring definitions for follow-up studies were consistent with those used at baseline within each patient.

Assessment of Pharyngeal Collapse

Pharyngeal collapse patterns were assessed using DISE and classified according to the VOTE system[21]. In the prospective cohort, DISE was performed using standardized techniques, and VOTE classifications were assigned at the time of the procedure by the treating surgeon. In the multicenter cohort, DISE examinations were performed according to standard practice at each participating center, and consensus VOTE scores were generated by four physician reviewers who were blinded to treatment outcomes[5].

Categorization of Collapse Patterns

For analytic purposes, collapse patterns were grouped into two primary categories: lateral collapse and AP collapse. Lateral collapse included CCC and complete OLW collapse. AP collapse included complete collapse of the tongue base, as well as AP collapse of the velopharynx or epiglottis.

To account for multilevel collapse, patients with complete tongue base collapse were categorized as having AP collapse regardless of other collapse sites. Patients with lateral collapse in the absence of complete tongue base collapse were categorized as lateral, irrespective of velopharyngeal or epiglottic collapse pattern. These groupings were mutually exclusive. Patients demonstrating only partial collapse across sites were categorized separately as a partial collapse group (n = 178).

Outcome Definitions

HGNS efficacy was assessed as the percent reduction in AHI from baseline to post-treatment evaluation.

Treatment success was defined using the Sher15 criterion[22], consisting of a ≥50% reduction in AHI to <15 events per hour, consistent with prior HGNS outcome studies[13].

Statistical Analysis

Baseline characteristics and HGNS-only outcomes are reported as median values with interquartile ranges. HGNS-only outcomes excluded patients who underwent upper airway surgery (functional expansion pharyngoplasty, tonsillectomy, or partial uvulectomy) either prior to or concomitant with HGNS implantation without repeat DISE (n = 48). Outcomes were summarized within categories of AP-directed collapse, laterally directed collapse, and partial collapse, with further stratification by BMI using a median split (median BMI: 27.5 kg/m²).

The primary analysis tested the hypothesis that greater BMI is associated with a greater reduction in HGNS efficacy among patients with laterally directed collapse compared with those with AP-directed collapse. Multivariable regression was used to model HGNS efficacy as a continuous outcome, with BMI (continuous), lateral collapse pattern (binary), and their interaction as primary predictors. Models were adjusted for baseline AHI, a partial collapse × BMI interaction, surgical center, type of follow-up sleep study (tPSG vs home sleep test), and prior or concomitant upper airway surgery.

Secondary analyses repeated the primary model using absolute change in AHI as the outcome. Additional analyses evaluated treatment success as a dichotomous outcome using logistic regression. Sensitivity analyses using alternative definitions of treatment success ($\geq$50% AHI reduction to <10 events per hour[23,24] and $\geq$70% AHI reduction[25]) were also performed. All models were additionally adjusted for age and sex, with these results reported in the supplement. Statistical significance was defined as a two-sided P value < 0.05 for the primary interaction term.

RESULTS

Cohort Characteristics and Unadjusted Outcomes

Baseline characteristics for the pooled cohort, as well as stratified by cohort and pharyngeal collapse pattern, are summarized in Tables 1 and 2. HGNS efficacy outcomes overall, by cohort, and stratified by collapse pattern and body mass index (BMI) are summarized in Table 3. Baseline demographic and clinical characteristics were broadly similar between cohorts (Table 1). Patients with laterally directed collapse exhibited slightly higher baseline apnea–hypopnea index (AHI) and BMI compared with those with anteroposterior (AP) collapse patterns (Table 2).

Across cohorts, HGNS efficacy and treatment success rates were lower among patients with laterally directed collapse compared with those with AP directed collapse (Table 3), consistent with prior reports[5,12-14]. Within the lateral and partial collapse groups, response rates were lower at higher BMI levels, whereas this pattern was less evident among patients with AP directed collapse.

Primary Analysis: Interaction Between BMI and Collapse Pattern

In multivariable regression models evaluating HGNS efficacy as percent reduction in AHI, a statistically significant interaction was observed between BMI and the presence of lateral collapse (β [95% CI]: -15.9 [-30.0, -1.8]; Table 4). Stratified estimates indicated a steeper decline in HGNS efficacy with increasing BMI among patients with lateral collapse compared with those with AP collapse. Specifically, each 5 kg/m² increase in BMI was associated with a -19.7 [-33.2, -6.2]% reduction in efficacy in patients with lateral collapse, compared with a -3.8 [-8.0, 0.36]% reduction in patients with AP collapse.

When absolute change in AHI was used as the outcome, a similar BMI × collapse pattern interaction was observed (-5.2 [-10.2, -0.17] events/h; Table 4). Increasing BMI was associated with a greater loss of efficacy in patients with lateral collapse (-6.4 [-11.1, -1.6] events/h per 5 kg/m² increase in BMI) compared with those with AP collapse (-1.2 [-2.7, 0.30] events/h per 5 kg/m² increase in BMI).

Secondary Analyses: Treatment Success

In logistic regression models evaluating treatment success, defined as a ≥50% reduction in AHI to <15 events per hour, the interaction between BMI and laterally directed collapse remained statistically significant (odds ratio [95% CI]: 4.1 [1.2, 14.0]; Table 4). Increasing BMI was associated with reduced odds of treatment success among patients with lateral collapse, whereas BMI was not significantly associated with treatment success among patients with AP collapse. Results were consistent when alternative definitions of treatment success were applied (Table E1, Online Supplement).

Partial Collapse Analyses

Because the primary model included an interaction term between partial collapse and BMI, the association between BMI and HGNS efficacy in patients with partial collapse could also be evaluated. In this group, HGNS efficacy decreased moderately with increasing BMI (−9.1% [−15.4, −3.3%] per 5 kg/m² increase in BMI).

In supplemental analyses, patients with partial collapse demonstrated higher HGNS efficacy compared with those with lateral collapse, with an improvement of 8.9% [1.8, 15.2%] in percent AHI reduction. This difference was associated with a higher likelihood of treatment success (odds ratio [95% CI]: 3.3 [1.6, 6.5]). These findings were comparable to the differences observed between AP and lateral collapse patterns (8.5% [2.0, 14.4%] improvement and 2.7-fold [1.5, 5.1] higher likelihood of success, respectively).

Additional Analyses

All primary findings were similar after additional adjustment for age and sex (Table E2, Online Supplement). Age and sex did not demonstrate significant interactions with collapse pattern. When collapse sites were examined individually, the association between BMI and HGNS efficacy within each site mirrored that of the broader collapse category to which they were assigned (Table E3, Online Supplement). Specifically, increasing BMI had minimal impact on HGNS efficacy in patients with complete AP velum, tongue base, or epiglottic collapse, whereas efficacy declined with increasing BMI in patients with complete OLW collapse and CCC.

DISCUSSION

In this pooled analysis of two independent HGNS cohorts, we observed that the relationship between body mass index (BMI) and HGNS efficacy differed according to pharyngeal collapse pattern. Specifically, increasing BMI was associated with progressively reduced HGNS efficacy in patients with lateral collapse patterns, whereas BMI had comparatively little impact on treatment efficacy among patients with predominantly anteroposterior (AP) collapse. These findings indicate that the effect of BMI on HGNS outcomes is not uniform across collapse patterns and suggest that BMI and collapse site should be considered jointly when evaluating treatment response.

Previous studies have demonstrated that HGNS efficacy and treatment success decline with increasing BMI[10,26-28]. The present study extends these observations by demonstrating that this association is modified by the site and pattern of pharyngeal collapse. Among patients with lateral collapse, a 5 kg/m² increase in BMI was associated with a marked reduction in HGNS efficacy and a substantial decrease in the likelihood of treatment success. In contrast, among patients with AP collapse, increasing BMI was associated with only modest changes in efficacy and was not associated with reduced odds of treatment success. Together, these findings suggest that BMI functions as a risk factor for reduced HGNS response primarily in the presence of lateral collapse.

Importantly, the observed interaction also indicates that lateral collapse does not uniformly preclude favorable HGNS outcomes[5,12,14]. In patients with lower BMI, HGNS efficacy and success rates were similar between AP and lateral collapse patterns. This observation suggests that lateral collapse patterns commonly considered difficult to treat may remain amenable to HGNS in the setting of lower BMI.

Several mechanisms may explain why increasing obesity preferentially impairs HGNS efficacy in patients with lateral walls collapse compared to those with AP collapse. First, increased obesity is associated with greater volume of the parapharyngeal fat pads and adipose tissue in the base of the tongue[29-31], and is a well-established contributor to increased pharyngeal collapsibility[15,17]. Accordingly, we suggest that when the laterally collapsing airway is only mildly collapsible, as is more common in patients with lower BMI, AP directed tongue movement can generate sufficient traction to promote lateral airway opening. This mechanical coupling between the tongue and pharyngeal tissues (i.e., glossopharyngeal coupling) may play a role in mediating the effects of HGNS on lateral collapse; however, with greater obesity and collapsibility, this coupling may be compromised. In contrast, for patients with AP collapse, increased adiposity may still increase collapsibility, but the airway remains treatable with HGNS, as the therapy directly targets the tongue base.

In addition, obesity—particularly central adiposity—is associated with reduced lung volumes and diminished caudal traction on the upper airway[32-35]. Reduced lung volume–related traction may further impair the ability of HGNS-induced tongue movement to influence lateral airway dimensions, thereby amplifying the adverse effect of increasing BMI in patients with lateral collapse. While these mechanisms are consistent with established

physiological principles, further studies incorporating direct pattern of upper airway collapse, measures of upper airway collapsibility, and lung volume are needed to clarify their relative contributions.

Although not a primary focus of the study, analyses of patients with partial collapse provide additional context. In this group, HGNS efficacy declined moderately with increasing BMI, but overall treatment response remained favorable and comparable to that observed in patients with complete AP collapse. Partial collapse was associated with higher efficacy and success rates than lateral collapse, and these differences were similar in magnitude to those observed between AP and lateral collapse. Furthermore, the directional pattern of partial collapse (AP versus lateral) did not significantly influence HGNS efficacy. These findings suggest that partial collapse represents a collapse phenotype that is generally responsive to HGNS, with modest susceptibility to BMI-related reductions in efficacy.

Several limitations should be considered when interpreting these findings. First, the protocol for scoring DISE data differed between the retrospective and prospective cohorts. In the retrospective cohort, collapse patterns were assigned based on consensus scoring by multiple blinded reviewers[5], whereas in the prospective cohort, scoring was performed clinically at the time of DISE. However, effect sizes for the BMI × collapse interaction were similar across cohorts, and test–retest analyses in the retrospective cohort demonstrated good agreement for identification of complete lateral and complete AP collapse, suggesting that differences in DISE scoring approach did not materially affect the results (Online Supplement).

Second, treatment efficacy was primarily assessed using titration polysomnography, which estimates AHI during periods of optimal stimulation and may overestimate real-world efficacy compared with single-setting home sleep apnea testing[3]. However, titration studies are performed under controlled conditions, minimize confounding due to changes in body weight or therapy adherence, and reduce discontinuation bias. Although absolute efficacy estimates may be higher than those obtained from home testing, we do not expect this approach to influence the observed interaction between BMI and collapse pattern. Consistent with this, sensitivity analyses adjusting efficacy estimates to approximate home sleep testing values resulted in a uniform downward shift without altering the BMI-related differences across collapse patterns (Online Supplement).

Third, the number of patients with BMI values above 35 kg/m² was limited, restricting inference regarding HGNS efficacy at higher BMI levels. Larger studies will be needed to evaluate whether the observed interaction persists across the full range of BMI currently considered eligible for HGNS therapy.

Finally, sleep study scoring criteria were not uniform across centers in the multicenter cohort, which may introduce variability in absolute AHI values. However, because scoring definitions were consistent within individuals across baseline and follow-up studies, percent AHI reduction is unlikely to be substantially affected, and the interaction between BMI and collapse pattern were consistent between the cohorts (Online Supplement).

In summary, this study demonstrates that BMI modifies HGNS efficacy in a collapse pattern–specific manner, with increasing BMI associated with reduced treatment response in patients with lateral pharyngeal collapse but not in those with AP collapse. These findings underscore the importance of considering anatomical collapse patterns alongside BMI when evaluating HGNS outcomes and provide a framework for more refined assessment of factors influencing treatment response.

REFERENCES

1.   Strollo PJ, Jr., Soose RJ, Maurer JT, et al. Upper-airway stimulation for obstructive sleep apnea. *The New England journal of medicine*. Jan 9 2014;370(2):139-49. doi:10.1056/NEJMoa1308659

2.   Pordzik J, Ludwig K, Seifen C, et al. Real-world data on polysomnography-and patient-reported outcomes in hypoglossal nerve stimulation and auto-titrating positive airway pressure therapy for obstructive sleep apnea. *Respiratory Medicine*. 2024;232:107750.

3.   Kaffenberger TM, Sina EM, Hambach B, et al. How we measure hypoglossal nerve stimulator outcome matters: titration versus single amplitude efficacy sleep studies. *Journal of Clinical Sleep Medicine*. 2024:jcsm. 11328.

4.   Strollo PJ, Soose RJ, Maurer JT, et al. Upper-Airway Stimulation for Obstructive Sleep Apnea. *New England Journal of Medicine*. 2014-01-09 2014;370(2):139-149. doi:10.1056/NEJMoa1308659

5.   Huyett P, Kent DT, D'Agostino MA, et al. Drug-Induced Sleep Endoscopy and Hypoglossal Nerve Stimulation Outcomes: A Multicenter Cohort Study. *The Laryngoscope*. 2021 2021;131(7):1676–1682.

6.   Strollo Jr P, Gillespie M, Soose R, et al. Stimulation Therapy for Apnea Reduction (STAR) Trial Group. Upper airway stimulation for obstructive sleep apnea: durability of the treatment effect at 18 months. *Sleep*. 2015 2015;38(10):1593–8.

7.   Soose RJ, Woodson BT, Gillespie MB, et al. Upper airway stimulation for obstructive sleep apnea: self-reported outcomes at 24 months. *Journal of Clinical Sleep Medicine*. 2016;12(1):43-48.

8.   Woodson BT, Soose RJ, Gillespie MB, et al. Three-year outcomes of cranial nerve stimulation for obstructive sleep apnea: the STAR trial. *Otolaryngology–head and neck surgery*. 2016 2016;154(1):181–188.

9.   Woodson BT, Strohl KP, Soose RJ, et al. Upper airway stimulation for obstructive sleep apnea: 5-year outcomes. *Otolaryngology–Head and Neck Surgery*. 2018 2018;159(1):194–202.

10.   Patel RM, Wang HZ, Jamro EL, et al. Response to Hypoglossal Nerve Stimulation Changes With Body Mass Index and Supine Sleep. *JAMA Otolaryngology–Head & Neck Surgery*. 2024;150(5):421-428.

11.   Vanderveken OM, Maurer JT, Hohenhorst W, et al. Evaluation of Drug-Induced Sleep Endoscopy as a Patient Selection Tool for Implanted Upper Airway Stimulation for Obstructive Sleep Apnea. *Journal of Clinical Sleep Medicine*. 2013-5-15 2013;doi:10.5664/jcsm.2658

12.   Nord RS, Fitzpatrick IV T, Pingree G, Islam A, Chafin A. Should lateral wall collapse be a contraindication for hypoglossal nerve stimulation? *American Journal of Otolaryngology*. 2024;45(1):104053.

13.   Huyett P, Wellman A, Caruso V, et al. Combination Tonsillectomy and Hypoglossal Nerve Stimulation for Sleep Apnea Patients With Oropharyngeal Lateral Wall Collapse. *Otolaryngology–Head and Neck Surgery*. 2024;

14.   Vena D, Huyett P, Op De Beeck S, et al. Flow-shape-derived Site of Pharyngeal Collapse Predicts Response to Hypoglossal Nerve Stimulation Therapy. *D20 MECHANISTIC INSIGHTS IN SLEEP DISORDERED BREATHING*. American Thoracic Society; 2023:A6271-A6271.

15.   Genta PR, Schorr F, Eckert DJ, et al. Upper airway collapsibility is associated with obesity and hyoid position. *Sleep*. 2014;37(10):1673-1678.

16.   Isono S. Obesity and obstructive sleep apnoea: mechanisms for increased collapsibility of the passive pharyngeal airway. *Respirology*. 2012;17(1):32-42.

17.   Kirkness JP, Schwartz AR, Schneider H, et al. Contribution of male sex, age, and obesity to mechanical instability of the upper airway during sleep. *Journal of applied physiology*. 2008;104(6):1618-1624.

18.   Schwartz AR, Gold AR, Schubert N, Stryzak A. Effect of weight loss on upper airway collapsibility in obstructive sleep apnea1-3. *Am Rev Respir Dis*. 1991;144(3 pt 1):494-498.

19.   Lan MC, Liu SY, Lan MY, Modi R, Capasso R. Lateral pharyngeal wall collapse associated with hypoxemia in obstructive sleep apnea. *The Laryngoscope*. 2015;125(10):2408-2412.

20.   Berry RB, Brooks R, Gamaldo CE, Hardling S, Marcus C, Vaughn B. Rules for scoring respiratory events in sleep: Update of the 2007 AASM manual for the scoring of sleep and associated events. *Journal of Clinical Sleep Medicine*. 2012 2012;8(5):567-619.

21.   Kezirian EJ, Hohenhorst W, de Vries N. Drug-induced sleep endoscopy: the VOTE classification. *European Archives of Oto-Rhino-Laryngology*. 2011 2011;268(8):1233–1236.

22.   Sher AE, Schechtman KB, Piccirillo JF. The efficacy of surgical modifications of the upper airway in adults with obstructive sleep apnea syndrome. *Sleep*. Feb 1996;19(2):156-77.

23.   Vena D, Azarbarzin A, Marques M, et al. Predicting sleep apnea responses to oral appliance therapy using polysomnographic airflow. *Sleep*. 2020 2020;43(7):zsaa004.

24.   Op de Beeck S, Wellman A, Dieltjens M, et al. Endotypic mechanisms of successful hypoglossal nerve stimulation for obstructive sleep apnea. *American journal of respiratory and critical care medicine*. 2021 2021;203(6):746–755.

25.   Bamagoos AA, Cistulli PA, Sutherland K, et al. Polysomnographic endotyping to select patients with obstructive sleep apnea for oral appliances. *Annals of the American Thoracic Society*. 2019 2019;16(11):1422–1431.

26.   Corr F, Kilinç F, Oros J, et al. Increased body mass index correlates with less favorable postoperative outcomes after hypoglossal nerve stimulation for obstructive sleep apnea: a retrospective cohort study. *World Neurosurgery*. 2023;180:e210-e218.

27.   Renslo B, Virgen CG, Sawaf T, et al. Long-term trends in body mass index throughout upper airway stimulation treatment: does body mass index matter? *Journal of Clinical Sleep Medicine*. 2023;19(6):1061-1071.

28.   Suurna MV, Steffen A, Boon M, et al. Impact of body mass index and discomfort on upper airway stimulation: ADHERE registry 2020 update. *The Laryngoscope*. 2021;131(11):2616-2624.

29.   Kim AM, Keenan BT, Jackson N, et al. Tongue fat and its relationship to obstructive sleep apnea. *Sleep*. Oct 1 2014;37(10):1639-48. doi:10.5665/sleep.4072

30.   Li Y, Lin N, Ye J, Chang Q, Han D, Sperry A. Upper airway fat tissue distribution in subjects with obstructive sleep apnea and its effect on retropalatal mechanical loads. *Respiratory care*. 2012;57(7):1098-1105.

31.   Schwab RJ. Properties of tissues surrounding the upper airway. *Sleep*. 1996;19(suppl_10):S170-S174.

32.   Hoffstein V, Zamel N, Phillipson EA. Lung volume dependence of pharyngeal cross-sectional area in patients with obstructive sleep apnea. *American Review of Respiratory Disease*. 1984;130(2):175-178.

33.   Owens RL, Malhotra A, Eckert DJ, White DP, Jordan AS. The influence of end-expiratory lung volume on measurements of pharyngeal collapsibility. *Journal of applied physiology*. 2010;108(2):445-451.

34.   Schwartz AR, Patil SP, Squier S, Schneider H, Kirkness JP, Smith PL. Obesity and upper airway control during sleep. *Journal of Applied Physiology*. 2010;108(2):430-435.

35.   Stadler DL, McEvoy RD, Sprecher KE, et al. Abdominal compression increases upper airway collapsibility during sleep in obese male obstructive sleep apnea patients. *Sleep*. 2009;32(12):1579-1587.

TABLES

Table 1. Baseline characteristics of the overall sample, across cohorts, and across pharyngeal collapse patterns.

Characteristic	Overall	Multi-center	Prospective
N	760	341	419
Age, years	60 [53, 68]	61 [53, 69]	60 [53, 67]
Men, n (%)	553 (72.8)	259 (76.0)	294 (70.2)
BMI, kg/m^2	29.0 [26.6, 31.3]	29.5 [27.0, 31.7]	28.8 [26.5, 31.0]
AHI, events/h	31.6 [22.7, 43.3]	34.3 [23.9, 44.0]	30.3 [22.3, 42.6]
AP directed, n (%)	495 (65.1)	211 (61.9)	284 (67.8)
Laterally directed, n (%)	87 (11.4)	40 (11.7)	47 (11.2)
Partial, n (%)	178 (23.4)	90 (26.4)	88 (21.0)

AHI: apnea-hypopnea index; BMI: body-mass index

AP directed, Laterally directed, and Partial represent the patients categorized as having AP directed collapse, laterally directed collapse and partial collapse, respectively.

Table 2. Baseline characteristics across anteroposterior (AP), lateral and partial collapse patterns.

Characteristic	AP	Lateral	Partial
N	495	87	178
Age, years	61 [54, 69]	56 [48, 64]	60 [53, 68]
Men, n (%)	352 (71.1%)	75 (86.2%)	126 (70.8%)
BMI, kg/m^2	28.8 [26.2, 31.0]	30.1 [28.6, 32.0]	29.3 [27.2, 31.5]
AHI, events/h	31.0 [22.5, 42.3]	35.9 [25.9, 44.9]	32.2 [21.7, 44.7]

AHI: apnea-hypopnea index; BMI: body-mass index

Table 3. Hypoglossal nerve stimulation outcomes in pooled and individual cohorts, stratified by pharyngeal collapse pattern (anteroposterior, lateral, and partial).

Cohort	Parameter	All	AP	Lateral	Partial
Pooled	N	712	482	60	170
	Percent reduction AHI	84.7 [62.6, 94.8]	85.1 [64.1, 95.3]	68.8 [43.8, 88.8]	85.7 [65.9, 95.8]
	N Success (%)	547 (76.8%)	377 (78.2%)	34 (56.7%)	136 (80.0%)
Multi-center	Percent reduction AHI	80.9 [55.5, 92.6]	81.6 [57.3, 93.4]	67.9 [42.0, 89.4]	80.8 [61.4, 93.9]
	N Success (%)	244 (71.6%)	153 (72.5%)	23 (57.5%)	68 (75.6%)
Prospective	Percent reduction AHI	87.4 [70.5, 96.0]	87.7 [72.7, 96.3]	70.0 [46.3, 87.0]	89.1 [71.9, 96.5]
	N Success (%)	303 (81.7%)	224 (82.7%)	11 (55.0%)	68 (85.0%)

AHI: apnea-hypopnea index; BMI: body-mass index

AP directed, Laterally directed, and Partial represent the patients categorized as having AP directed collapse, laterally directed collapse and partial collapse, respectively.

Sample analyzed for this table only includes patients that underwent HGNS-only (n=712), i.e., excludes patients who underwent upper airway surgery (functional expansion pharyngoplasty, tonsillectomy, or partial uvulectomy) concomitant with or prior to HGNS without repeat DISE.

Table 4. Summary of the interaction between BMI and collapse pattern (lateral vs. AP) and the estimates (β coefficients or odds ratios) for the effect of BMI on HGNS outcomes within patients with lateral and AP collapse.

Outcome	Interaction [95%CI], p-value	Lateral Estimate [95%CI]	AP Estimate [95%CI]
Percent reduction in AHI	−15.9 [−30.0, −1.8], p=0.027	−19.7 [−33.2, −6.2]	−3.8 [−8.0, 0.36]
Absolute reduction in AHI	−5.2 [−10.2, −0.17], p=0.043	−6.4 [−11.1, −1.6]	−1.2 [−2.7, 0.30]
Success	4.1 [1.2, 14.0], p=0.023	1.1 [0.75, 1.5]	4.4 [1.4, 14.3]

AHI: apnea-hypopnea index; BMI: body-mass index.

Linear regression was used for continuous outcomes (percent and absolute AHI reduction) and logistic regression for binary outcomes (success). All models were adjusted for baseline AHI, the presence of partial collapse × BMI interaction, surgical center, type of follow-up sleep study, and previous or concomitant upper airway surgery.

Estimates for percent and absolution AHI reduction represents the additional reduction in HGNS efficacy outcome per 5 kg/m^2 increase in BMI

For the outcome 'Success,' estimates represent odds ratios for achieving treatment success (≥50% AHI reduction to <10 events/h) per 5 kg/m² increase in BMI..

The interaction represents the additional reduction in HGNS efficacy outcomes per 5 kg/m² increase in BMI that is observed in patients with lateral collapse compared to those with AP collapse.

Association between body mass index and unilateral hypoglossal nerve stimulation efficacy depends on the pattern of pharyngeal collapse in obstructive sleep apnea

Daniel Vena[1], Eric J. Kezirian[2], Andrew Wellman[1], David Kent[3], Mark D'Agostino[4,5], Joao L.G.C. Monteiro[1], Azarbarzin A[1], Tom Chen[6], Ludovico Messineo[1], Neda Esmaeili[1], Scott A. Sands[1]*, Phillip Huyett[7]*.

[1]Division of Sleep and Circadian Disorders, Brigham and Women's Hospital and Harvard Medical School, Boston, MA.

[2]Department of Head and Neck Surgery, David Geffen School of Medicine at UCLA, Los Angeles, California.

[3]Department of Otolaryngology-Head and Neck Surgery, Vanderbilt University Medical Center, Nashville, Tennessee.

[4]Southern New England Ear, Nose and Throat Group, Middlesex Hospital, Middlesex, Connecticut.

[5]Department of Surgery, Yale University School of Medicine, New Haven, Connecticut.

[6]Department of Population Medicine, Harvard Medical School and Harvard Pilgrim Health Care Institute, Boston, MA

[7]Department of Otolaryngology, Massachusetts Eye and Ear, Boston, MA

*,[†]Equal contributions

Correspondence:

Daniel Vena

221 Longwood Ave., Boston MA, 02115

Division of Sleep and Circadian Disorders, Department of Medicine, Brigham & Women's Hospital & Harvard Medical School, Boston, Massachusetts.

Tel: +1-617-732-8483; E-mail: dvena@bwh.harvard.edu

Interaction between BMI and pharyngeal collapse pattern on HGNS success using alternative definitions of success

To demonstrate consistency of the interaction effect between BMI and lateral collapse on HGNS success, we reran our secondary analysis with other commonly used definitions of success: 50% AHI reduction to <10 /h[1,2] (category 2) and 70% AHI reduction[3] (category 3). Results, summarized in Table E1 were consistent with the HGNS success criterion used in the main paper (Category 1: 50% AHI reduction to <15 /h).

Table E1. Interaction between BMI and laterally directed collapse on HGNS success defined across commonly used definitions of success

HGNS success category	Interaction	AP-directed	Laterally-directed
Category 1 (main paper)	4.1 [1.2, 14.0]	1.1 [0.75, 1.5]	4.4 [1.4, 14.3]
Category 2	2.6 [0.86, 7.9]	1.2 [0.79, 1.9]	3.1 [1.1, 9.1]
Category 3	3.3 [1.0, 10.4]	1.4 [0.99, 1.9]	4.5 [1.5, 13.7]

Adjustment of age and sex

The analysis evaluating the interaction between lateral collapse and BMI on HGNS efficacy, as described in the main manuscript, was repeated with adjustment for age and sex. Adjustment had no meaningful impact on the interaction term or on the estimates of BMI effects within the lateral and AP collapse groups (Table E2).

Table E2. Summary of the interaction between BMI and collapse pattern (lateral vs. AP) and adjusted estimates (β coefficients or odds ratios) for the effect of BMI on HGNS outcomes within lateral and AP collapse groups, with models adjusted for age and sex.

Outcome	Interaction [95%CI], p-value	Lateral Estimate [95%CI]	AP Estimate [95%CI]
Percent reduction in AHI	−15.6 [−29.7, −1.6], p=0.029	−19.4 [−32.8, −5.9]	−3.7 [−7.8, 0.42]
Absolute reduction in AHI	−5.1 [−10.1, −0.1], p=0.043	−6.3 [−11.1, −1.5]	−1.2 [−2.6, 0.32]
Success	4.1 [1.2, 14.1], p=0.023	4.3 [1.3, 14.0]	1.0 [0.73, 1.5]

AHI: apnea-hypopnea index; BMI: body-mass index.

Linear regression was used for continuous outcomes (percent and absolute AHI reduction) and logistic regression for binary outcomes (success). All models were adjusted for age, sex, baseline AHI, the presence of partial collapse × BMI interaction, surgical center, type of follow-up sleep study, and previous or concomitant upper airway surgery.

Estimates for percent and absolution AHI reduction represents the additional reduction in HGNS efficacy outcome per 5 kg/m^2 increase in BMI

For the outcome 'Success,' estimates represent odds ratios for achieving treatment success (≥50% AHI reduction to <15 events/h) per 5 kg/m² increase in BMI..

The interaction represents the additional reduction in HGNS efficacy outcomes per 5 kg/m² increase in BMI that is observed in patients with lateral collapse compared to those with AP collapse.

Age and sex interaction with BMI on HGNS efficacy

To examine whether age and sex were differentially associated with HGNS efficacy in patients with AP versus lateral collapse, we repeated the primary analysis from the main paper, substituting age and sex for BMI. Specifically, we conducted a multivariable mixed model regression to evaluate HGNS efficacy as a function of the presence of lateral collapse, age/sex and their interaction (age/sex × collapse pattern), and adjusted for covariates baseline AHI, the presence of partial collapse, surgical center, type of follow-up sleep study, and previous or concomitant upper airway surgery.

The interaction between age and laterally directed collapse was not statistically significant in the linear regression model evaluating HGNS efficacy (β [95% CI]: 6.2 [−0.64, 7.9]% per 10-year increase in age; p = 0.08). Nonetheless, there was a suggestive trend toward modestly greater efficacy with increasing age among patients with lateral collapse (5.1 [−1.2, 11.5]% per 10-year increase), a pattern not observed in patients with AP collapse (−1.1 [−3.5, 1.4]% per 10-year increase). Similarly, the interaction between sex and laterally directed collapse was not statistically significant (β [95% CI]: −3.1 [−23.9, 17.6]%; p = 0.76 for men compared with women).

Interaction between BMI and various pharyngeal site-of-collapse categories in predicting HGNS efficacy

We also evaluated the interaction between various combinations of different site-of-collapse categories (e.g., oropharyngeal lateral wall collapse combined with anteroposterior velum collapse) and BMI in explaining HGNS efficacy. To achieve this, we divided the analysis into five logical components. In the first component (analysis group A in Table E3), we examined the individual categories that constitute laterally-directed collapse: complete concentric collapse of the soft palate (CCC) and complete oropharyngeal lateral wall collapse (OLW). Our goal was to determine whether patients within each category exhibited substantial reductions in HGNS efficacy, similar to the parent category of lateral collapse. Indeed, lateral collapse defined by CCC alone, OLW alone, or a combination of CCC and OLW demonstrated substantial reductions in HGNS efficacy with increasing BMI (Table E3), with effect sizes comparable to or greater than those reported in the main paper.

Since the findings for the different site-of-collapse categories within lateral collapse were similar to those for lateral collapse overall, we combined them back into a single category of lateral collapse. We then assessed whether anteroposterior (AP) velum and epiglottic collapse, when occurring concurrently with laterally-directed collapse, influenced its interaction with BMI in explaining HGNS efficacy; as well as how this compared to isolated lateral collapse (i.e., without any accompanying collapse patterns). As shown in analysis group B results from Table E3, isolated lateral collapse and lateral collapse combined with AP velum demonstrated substantial reductions in HGNS efficacy with increasing BMI, with effect sizes comparable to or greater than those reported in the main paper. We note that the combination of lateral collapse with epiglottic collapse could not be evaluated due to an insufficient number of patients exhibiting this pattern.

Altogether, the findings from Table E3 demonstrate that patients with lateral collapse—whether defined as CCC, OLW collapse, or a combination of the two, and whether occurring in isolation or concurrently with AP velum or epiglottic collapse—exhibit substantial reduction in HGNS efficacy as BMI increases.

Next, we performed a similar analysis for AP collapse patterns in analysis groups C, D, and E (Table E3) to demonstrate that HGNS efficacy remains relatively stable with increasing BMI across the individual site-of-collapse patterns that fall within the category of AP collapse. In analysis group C we evaluate change in HGNS efficacy with increasing BMI within various categories of tongue collapse: isolated tongue collapse, tongue collapse concurrent with collapse of the AP velum, epiglottis, and then AP-directed collapse without any tongue involvement for comparison. As summarized in Table E3, HGNS efficacy varies minimally (or even increases) with increasing BMI across the various categories, with effect sizes comparable to that of the main paper. We note that results from concurrent tongue and epiglottic collapse were not reliable given the small sample size.

In analysis group D, we evaluated the change in HGNS efficacy with increasing BMI across various categories of AP velum collapse (Table E3). Only isolated AP velum collapse had a sufficient sample size for meaningful conclusions, and it exhibited results consistent with AP-directed collapse, showing minimal change in HGNS efficacy with increasing BMI.

Lastly, analysis group E examined the change in HGNS efficacy with increasing BMI in cases of isolated epiglottic collapse. Although the sample size was small, the effect size fell between those observed for AP-directed and laterally-directed collapse.

Table E3. The effect of BMI on HGNS efficacy within various site-of-collapse categories.

Analysis group	Collapse category	N	Difference in Efficacy per 5 unit increase in BMI. Beta [95%CI]	VOTE scores comprising collapse category
A	CCC without OLW	10	−63.8 [−132.1, −3.4]	V2c, V2cE2
	OLW without CCC	72	−15.3 [−30.4, −0.18]	O2, V2apO2, O2E2
	CCC + OLW	5	−17.6 [−62.1, 26.9]	V2cO2, V2cO2E2
B	Isolated lateral collapse	71	−17.1 [−33.6, −0.65]	V2c, O2, V2cO2
	Lateral + AP velum	15	−17.2 [−44.2, 9.9]	V2apO2
	Lateral + Epiglottis	1	N/A	O2E2

Analysis group	Collapse category	N	Change in Efficacy with BMI Beta [95%CI]	VOTE scores within collapse category
C	Isolated Tongue	154	−4.3 [−11.5, 2.9]	T2
	Tongue + AP velum	201	−4.4 [−10.8, 1.9]	V2apT2, V2apT2E2
	Tongue + Epiglottis	2	60.3 [204.8, 69.5]	T2E2
	AP collapse without Tongue	138	0.98 [−8.1, 10.1]	V2ap, V2apE2
D	Isolated AP velum	111	1.6 [−8.3, 11.6]	V2ap
	AP velum + Epiglottis	10	−12.1 [−57.5, 33.3]	V2apE2
	AP collapse without AP velum	10	−5.2 [−10.0, −0.48]	T2, T2E2
E	Isolated Epiglottis	17	−7.4 [−36.4, 21.7]	E2

Efficacy defined by the percent reduction in apnea-hypopnea index from baseline.

CCC: Complete concentric collapse of the soft palate; OLW: oropharyngeal lateral walls; AP: anteroposterior

Effect of partial pharyngeal collapse on HGNS efficacy

To study the effect of partial collapse on HGNS efficacy we used multivariable mixed model regression to evaluate HGNS efficacy (continuous dependent variable) as a function of the presence of partial pharyngeal collapse pattern (dichotomous independent variable) and AP-directed collapse (dichotomous), and adjusted for covariates baseline AHI (continuous), surgical center, type of follow-up sleep study (titration PSG or single-setting home sleep test), and previous or concomitant upper airway surgery (functional expansion pharyngoplasty, tonsillectomy, and partial uvulectomy). We repeated this analysis using successful treatment response as the outcome (dichotomous) and logistic regression instead of linear regression. These analyses quantified increased HGNS efficacy and success rate in patients with partial collapse relative to lateral collapse. For comparison, the increased HGNS efficacy and success in AP collapse was also analyzed and presented. As described in the main paper, HGNS efficacy in partial collapse was greater by 9.0 [0.33, 17.6]% over lateral collapse, with a corresponding 3-fold greater likelihood of a successful response (odds ratio [95% CI]: 3.3 [1.6, 6.5]), both comparable to the 8.5 [2.0, 14.4]% efficacy gain and 2.7-fold [1.5, 5.1] increased likelihood seen in AP collapse over lateral collapse.

We also evaluated the effect of increasing BMI on HGNS efficacy and success rate in patients with partial collapse. To do so, we used the same interaction model described in the main paper. The analysis showed that BMI had a moderate effect on HGNS outcomes, decreasing efficacy by -9.2 [-16.1, -2.3]% per 5 kg/m^2. Success rate also decreased by 1.6 times per 5 kg/m^2, but this did not reach statistical significance (odds ratio: 1.6 [0.83, 3.0]).

Lastly, we investigated the effect of pattern of *partial* collapse (i.e., AP vs. lateral) using a multivariable mixed model regression that evaluated HGNS efficacy as function of lateral *partial* collapse, adjusting out complete collapse, and adjusting for the same covariates described above. The coefficient for laterally-directed partial collapse quantifies the difference in HGNS efficacy of laterally-directed partial collapse compared to AP-directed partial collapse. The model showed that there was no significant difference in HGNS efficacy (4.8 [-6.5, 16.0]%) and success rate (odds ratio [95%CI]: 1.5 [0.55, 4.2]) between lateral and AP partial collapse. Overall, this suggests that the pattern of collapse is less important in the context of partial collapse.

Interaction between BMI and laterally directed collapse on HGNS outcomes, analyzed separately by cohort

We report the interaction between BMI and the presence of lateral collapse on HGNS efficacy and success rate separately for the multi-center and prospective cohorts. The interaction effect for HGNS efficacy was substantial in both cohorts, measuring -13.7% (95% CI: -31.6, 4.1) in the prospective cohort and somewhat larger in the multi-center cohort at -18.7% (95% CI: -41.1, 3.6). Similarly, the interaction effect for HGNS success rate (odds ratio [95%CI]) in the multi-center cohort was 3.4 [0.77, 14.7], and a comparable effect was

found in the prospective cohort at 3.9 [0.33, 28.9]. These findings suggest that single-reviewer scoring yielded consistent relationships between site of collapse, BMI, and HGNS efficacy, supporting its reliability for this analysis.

Inter-rater reliability for classifying pharyngeal collapse as lateral or anteroposterior

We assessed inter-rater reliability for categorizing airway collapse as lateral or not and as AP or not. To do so, we fit a logistic mixed model with random effects for video and rater (model equation: $\text{logit}\, P(Y_{ij} = 1 | \eta_i, \zeta_j) = \alpha + \eta_i + \zeta_j$ for the ith video and jth rater). The logistic-scale within-task intraclass correlation coefficient (ICC) was calculated based on the variance of the random effects $\sigma_\eta^2, \sigma_\zeta^2$ using the equation[4]: $\text{ICC}_{\text{Task}} = \frac{\sigma_\eta^2}{\sigma_\eta^2 + \sigma_\zeta^2 + \pi^2/3}$.

The model demonstrated a high ICC for lateral collapse (0.72) and a fairly strong ICC for AP collapse (0.50), indicating good agreement among raters for both categorizations. These findings again support the reliability of site-of-collapse assessments made by a single reviewer in this analysis.

Interaction between BMI and lateral collapse on HGNS efficacy adjusted to reflect efficacy from home sleep apnea test.

To address the limitation that HGNS efficacy was determined based on optimal setting AHI, which overestimates actual HGNS efficacy[5], we estimated HGNS efficacy as it would be if treatment AHI were measured using an HSAT. We leveraged data from patients who underwent a post-treatment HSAT (n=60) to compute the mean and standard deviation of HSAT-derived HGNS efficacy distribution. We subsequently applied a linear transformation to adjust the distribution of HGNS efficacy calculated based on the titration study at optimal stimulation to match the mean and standard deviation of HGNS efficacy derived from the HSAT. With the adjusted HGNS efficacy, we re-ran our primary multivariable regression analysis, evaluating estimated HSAT-derived HGNS efficacy as a function of laterally-directed pharyngeal collapse, BMI, and their interaction (BMI × collapse pattern), adjusting for the same covariates as in the main analysis.

This adjustment resulted in a uniform ~10% downward shift in overall efficacy (Figure E1). The model again identified a significant interaction between BMI and lateral collapse in predicting HGNS efficacy (β [95% CI]: −15.6 [−29.3, −1.9]%, p = 0.026). Specifically, lateral collapse was associated with a similar reduction in efficacy (−19.2 [−32.4, −6.1]% per 5 kg/m² increase in BMI) compared to AP collapse (−3.6 [−7.7, 0.41]% per 5 kg/m² increase in BMI) (Figure E1).

Figure E1. Impact of body mass index (BMI) on adjusted hypoglossal nerve stimulation (HGNS) efficacy (percent reduction AHI) across AP- and laterally directed collapse patterns. Histograms at the top of the figure illustrates the BMI distribution within each pattern of collapse. HGNS efficacy was adjusted to reflect efficacy determined from follow-up home sleep testing.

REFERENCES

1.   Vena D, Azarbarzin A, Marques M, et al. Predicting sleep apnea responses to oral appliance therapy using polysomnographic airflow. *Sleep*. 2020 2020;43(7):zsaa004.

2.   Op de Beeck S, Wellman A, Dieltjens M, et al. Endotypic mechanisms of successful hypoglossal nerve stimulation for obstructive sleep apnea. *American journal of respiratory and critical care medicine*. 2021 2021;203(6):746–755.

3.   Bamagoos AA, Cistulli PA, Sutherland K, et al. Polysomnographic endotyping to select patients with obstructive sleep apnea for oral appliances. *Annals of the American Thoracic Society*. 2019 2019;16(11):1422–1431.

4.   Wu S, Crespi CM, Wong WK. Comparison of methods for estimating the intraclass correlation coefficient for binary responses in cancer prevention cluster randomized trials. *Contemporary clinical trials*. 2012;33(5):869-880.

5.   Kaffenberger TM, Sina EM, Hambach B, et al. How we measure hypoglossal nerve stimulator outcome matters: titration versus single amplitude efficacy sleep studies. *Journal of Clinical Sleep Medicine*. 2024:jcsm.11328.