Re-classification and Weighting of Multiple Causes of Death: US Death Certificates 2003–2023

Michael Levitt[1], Ben Marten[2], Gal Oren[1,3], John P.A. Ioannidis[4,5,6,7]

[1]Department of Structural Biology, Stanford University School of Medicine, Stanford, California, USA; [2]Mortality.Watch, St. Petersburg, Florida, USA;[3]Computer Science Department, Technion Israel Institute of Technology, Haifa, Israel, [4]Meta-Research Innovation Center at Stanford (METRICS), Stanford University School of Medicine, Stanford, California, USA; [5]Stanford Prevention Research Center, Department of Medicine, Stanford University School of Medicine, Stanford, California, USA; [6]Department of Epidemiology and Population Health, Stanford University School of Medicine, Stanford, California, USA; [7]Department of Biomedical Data Science, Stanford University School of Medicine, California, USA

Funding: National Institutes of Health R35 GM122543 (ML) and NIEHS R01ES032470 (JPAI) and N000142412687 by the Office of Naval Research (JPAI).

Data and code availability: https://github.com/scientific-computing-user/Reclassification-and-Weighting-of-Multiple-Causes-of-Death-US-Death-Certificates-2003-2023/

Disclosures: The authors have no conflicts of interest.

Acknowledgments: None

Keywords: death certificate, mortality, COVID-19, cause of death, vital statistics

Abstract

Death certificates record causes of death as reported by certifiers (Entity Axis) and as standardized by mortality coding rules (Record Axis). Conventional mortality statistics then reduce these to a single underlying cause, ignoring other contributing conditions; weighting schemes can instead distribute the mortality burden across all listed causes. We evaluated the impact of re-classification from Entity to Record axis and weighting across all 56,986,831 US death certificates from 2003–2023, mapping ICD-10 codes to 14 broad disease categories and testing three weighting schemes: W1 (50% weight to the underlying cause, 50% shared equally among contributing causes), W2 (equal weighting across all causes), and W2A (equal weighting at the ICD-10 code level). Entity and Record Axes agreed on underlying cause in 48,313,403 deaths (84.8%) by broad disease category and in 39,260,709 deaths (68.9%) by ICD-10 code (Supplementary Tables S5A, S5B); concordance reached 70.4% when using 3-character rather than 4-character ICD-10 categories. Re-classification markedly increased COVID-19 deaths (+92%) and Transport deaths (+43%) while markedly decreasing Other External Causes (-54%). Weighting substantially altered death burden attribution: COVID-19 (−44–63%) and Falls (−46–66%) decreased, Other External causes more than tripled (+204–254%), and deviations from unweighted counts were more pronounced with W2 and W2A than with W1. Weighting also brought disease-category counts closer to Entity Axis values and restored Respiratory seasonality patterns suppressed during the pandemic years. Systematic differences between reported and re-classified causes of death, and the choice of weighting scheme, profoundly alter disease burden estimates for several causes, with major implications for resource allocation and public health priorities.

SIGNIFICANCE STATEMENT

Standardized re-classification processes using mortality coding rules recast the selected causes of death in many death certificates. Moreover, vital statistics typically isolate a single underlying cause, while for many deaths multiple causes jointly lead to demise. Analysis of ~57 million deaths in the USA (2003-2023) shows that a large proportion of deaths are re-classified by rules to different causes than those filled by original certifiers. Analyses that give weight not only to recorded underlying cause but also to other listed causes lead to markedly different

estimates of deaths from several diseases. For example, the footprint of COVID-19 fatalities during the pandemic years decreases by 44-63%. Re-classification and weighting schemes may have profound impact on disease burden estimates and policy decisions.

INTRODUCTION

Death certificates are widely known to have high rates of inaccuracies and inconsistencies (1-5), but they serve as the foundational data source for mortality statistics and for public health policies that depend on these statistics. Problems often arise at the level of filling out the certificate forms by physicians and other eligible health care practitioners who certify the causes of death. The certifiers may lack experience, make inadvertent errors, or have less than optimal knowledge of the complete death situation, circumstances, and past medical history of the deceased. Autopsies are uncommonly performed. To enhance standardization, for several decades now systems have been in place that process the crude reported death certificates and re-classify the causes of death with specific mortality coding rules (6,7). The certifier-entered information is termed Entity Axis, and the re-classified standardized information is termed Record Axis. The data of the Record Axis are the ones endorsed officially to represent the cleaned information on underlying causes of death and on other contributing causes. The mortality coding rules for the cleaning and re-classification process from the Entity to the Record Axis are laid out in detail (6,7). However, they are sometimes revised, when a new version of the International Classification of Diseases (ICD) is released and when new ICD codes are added or existing ones are modified. The introduction of COVID-19 coding in the ICD system during the pandemic was such an example of a prominent modification (8).

Even with the Record Axis standardization, major inaccuracies may persist if the original input at the Entity Axis is flawed in ways that cannot be amended by the re-classification rules. Studies of in-depth medical record audits demonstrate large discrepancies between death certificates and more appropriate determination of causes of death based on the in-depth assessments; this has been particularly prominent for deaths where multi-morbidity is involved

such as COVID-19 (9-11). A pervasive issue relates to the fundamental traditional rule allowing only one underlying cause. In reality, many deaths have substantial contributions from multiple causes (12-22). The preference for listing a specific cause as underlying rather than contributing may change over time, depending on physician awareness and on how lethal that cause is perceived to be relative to competing causes (16). Evolving perceptions may create artefacts on the increase or decrease of specific diseases as underlying causes of death. Misclassifications of underlying and contributing causes become a greater challenge with increasing multimorbidity (17). For people with co-existing morbidities that have eroded their health, it may be best to give weight not just to one cause but many. Absent in-depth medical record audits, the listing of additional causes in the death certificates may provide some indication of the broader background for each death. Several authors have proposed weighting schemes where multiple contributing causes get a share of the death attribution (19-22).

Here we aimed to analyze all death certificates in the USA between 2003 and 2023 to empirically assess how common are changes between Entity and Record Axes and whether specific major patterns exist in these re-classifications. Moreover, we aimed to assess whether different weighting schemes for multiple causes of death may substantially affect the estimated burden of death from specific causes. We examined evolving patterns over time during this extended period, and separately in elderly and non-elderly populations paying attention to the pandemic period 2020-2023 and to COVID-19 deaths. COVID-19 represented a major change in cause of death classifications with extensive debate about the over- or under-counting of COVID-19 deaths (23).

METHODS

Death certificate data

Raw US death certificate data covering the 21-year period from 2003 to 2023 were obtained from the publicly accessible Centers for Disease Control and Prevention (CDC) National Center for Health Statistics (NCHS) FTP server (24). The dataset consists of annual mortality files (`mort2003us.zip` through `mort2023us.zip`). These archives were decompressed to yield fixed-width text files (`VS23MORT.DUSMCPUB`) containing death certificate data in specified columns. For each death, we extracted the complete information on ICD codes from both the Entity Axis and the Record Axis. We then classified the ICD codes into 14 broad disease categories, 7 natural ones (Circulatory, Cancer, Respiratory, Endocrine/Nutritional/Metabolic, Digestive, COVID-19, and Other Natural) and 7 external ones (Drug Poisoning, Transport, Suicide, Alcohol-induced, Falls, Homicide, Other External), following the classification scheme of Wrigley-Field et al (25) that simplifies a previously proposed grouping of the National Academies of Science, Engineering and Medicine (26). Our grouping is the same as in (25) except that we also have Falls as a separate broad category rather than merged with Other External, since reported Falls deaths have rapidly increased in the 21st century and thus would be worth closer examination. For COVID-19 we used three ICD-10 codes (U07.1, U07.2, and U09.9) as done before (25). The few (n=518) occurrences of uncertain values (all were string "1") under ICD-10 codes in the Record Axis were ignored.

When several ICD codes were mapped to the same broad disease category, they were usually counted only once in all analyses that used the broad disease categories (W2A is the exception). We used the broad-level classification to focus on major shifts across major groups of causes.

Entity Axis versus Record Axis comparisons

We noted for each broad disease category how often it is listed as what might be assumed to be the underlying cause of death in the Entity Axis (i.e. appearing in the line in Part I of the death certificate filled out by certifying physician or other eligible health care practitioner) versus as underlying cause in the Record Axis. We also examined in each Axis for each broad disease category how often it was listed but not as underlying cause; how often it was listed at all (either as underlying cause or other); and how often it was listed as the only broad disease category in the death certificate. We then evaluated the underlying causes of death according to the two Axes and noted how often they agreed at the broad disease category level and at the specific ICD level. Focusing on the broad disease categories, we also recorded how often each cause was promoted to underlying cause of death from the re-classification of Entity Axis into Record Axis and how often it was demoted from underlying cause of death during that re-classification, the ratio of promotions to demotions, and the specific pairs of broad disease categories on which major shifts occurred. Specifically for COVID-19 we also examined the specific ICD categories involved in such promotion and demotion switches.

When there was more than one ICD code listed in the last line of Part I in the Entity Axis, we used the one listed last for the main analyses. As a sensitivity analysis, we examined the one listed first in the last line of Part I.

Of note, Entity Axis is not normally used for underlying cause reporting, therefore the comparison between Entity and Record Axis is done primarily to examine the effect of the re-classification process that generates the official cause data (Record Axis). Moreover, examination of the Entity Axis data allows to see whether weighted analyses considering multiple causes of death bring the death counts closer to the original Entity Axis entries.

Weighting of multiple causes of death

We generated data for death counts comparing different weighting schemes using information from the Record Axis: the baseline W0 (counting only the listed underlying cause of death in the Record Axis); a hierarchical weighting scheme W1, where the underlying cause of death is given 50% of the weight, and the remaining contributing causes of death share equally among them the other 50% (e.g. if there are 2 contributing causes, then each of them gets 25% of the weight and the underlying cause gets 50% of the weight); an equal weighting scheme W2 where all causes share equally the weight (e.g. with two contributing causes, each of them gets 33.33% of the weight and the underlying cause also gets 33.33%) and a variant of W2 called W2A where all the causes (including repeated occurrences in the same broad category that are normally excluded) share the weight equally. Similar weighting schemes have been previously proposed in analyses of multiple cause of death data (19-22) and have used different levels of grouping of causes. Hence, we explored both the high-level broad categories grouping (W1, and W2) and the more granular ICD-level analyses (W2A). We estimated the absolute and relative magnitude of change between W1, W2, and W2A versus the baseline W0 (which represents the traditionally presented mortality statistics) for the number of deaths attributed to each of the broad death categories.

We mapped the baseline versus the W1, W2 and W2A estimates for each of the broad death categories per month in the 2003-2023 period. We also performed similar subgroup analyses for age groups of elderly (greater than 65 years old, GE65) and non-elderly (less than 65 years old, LT65). The plotted estimates were population-adjusted using annual mid-year US population estimates from CDC Wonder available through the Human Mortality Database (https://www.mortality.org/File/GetDocument/hmd.v6/USA/STATS/Population5.txt), with 2023 as the reference year. Scaling was performed separately for the overall population and for each

age subgroup using overall and subgroup-specific population denominators, such that observed deaths were multiplied by the ratio of 2023 population to that year's population within each stratum.

For the weighted analyses, we also performed two sensitivity analyses using the underlying cause information from the Record Axis and as additional causes the Part II causes listed from the Entity Axis; and using the underlying cause information from the Entity Axis (last line on Part I) and as additional causes the Part II causes listed in the Entity Axis. These sensitivity analyses aim to avoid the potential immediate causes of deaths (for example, sepsis) that might have otherwise contaminated the listing of other causes in the Record Axis.**Software**

All analyses were performed using Python scripts developed iteratively with LLM assistance (Claude, Anthropic). The LLM served as a coding assistant – the investigators specified the analytical requirements, reviewed all generated code for correctness, and validated results through multiple cross-checks including verifying that weighted totals sum to total deaths, comparing results across independent implementations, spot-checking calculations manually on subsets of data, and confirming concordance with published CDC statistics where available. The final Python scripts are deterministic and reproducible; they are provided along with the key data in https://github.com/scientific-computing-user/Reclassification-and-Weighting-of-Multiple-Causes-of-Death-US-Death-Certificates-2003-2023/.

RESULTS

Entity versus Record Axis

56,986,831 U.S. death certificates were issued in 2003–2023. Comparison of Entity and Record Axis data (Table 1) shows substantial changes for some broad categories of death such as COVID-19 and Other External causes. In relative terms, re-classification from Entity Axis to

Record Axis almost doubles the number of COVID-19 deaths (+92%) and creates also major increases in the number of Falls deaths (+69%), Transport deaths (+43%), and modest increases in the number of Suicide (+25%), Homicide (+30%), Endocrine (+16%) and Cancer deaths (+12%). Re-classification leaves Circulatory, Digestive, Alcohol-related and Drug Poisoning deaths largely unchanged. Conversely, it causes major decreases in Other External causes (-54%), and modest decreases in Other Natural causes (-14%), and Respiratory deaths (-11%).

The availability of additional contributing causes of death more than doubles the total number of available causes (n=57,611,546 more) even with the clustering in broad categories. For both Axes, slightly over a third of the deaths (n=20,502,014 or 20,808,620) have only one of the broad disease categories represented in the death certificate. The relative ranking changes substantially for some categories such as COVID-19 and Cancer when any cause is considered instead of only the underlying cause. There is a large diversity across the broad categories on the ratio of their appearance as contributing causes rather than as underlying causes (ranging from near zero to 6.25, see Supplementary Table S1).

In 7,749,865 (13.6%) of the 56,986,831 deaths, the last line of Part I in the death certificate had multiple ICD codes listed. When assuming as underlying cause of death the first ICD listed in the last line of Part I instead of the last ICD listed in that line, the respective results and changes versus the main analysis are shown in Supplementary Tables S2 and S3.

Table 2 shows the specific transitions across the broad disease categories between the Entity Axis and the Record Axis for the underlying cause of death. Entity and Record Axes agreed on underlying cause (the Entity "underlying cause" is that reported in the lowest, last line of Part I) in 48,313,403 (84.8%) of the total deaths based on broad disease categories and in 39,260,709 (68.9%) based on ICD-10 codes (Supplementary Table S5A) and in 30,806,674

(70.4%) when we used 3-character ICD-10 categories instead of 4-characters (e.g. J44 instead of J449). Concordance was 85.7% and 70.1%, respectively, in pre-pandemic years (2003-2019) and 81.6% and 65.0%, respectively, during pandemic years (2020-2023). Discordance rates slowly increased between 2003 and 2019 (by about 15%) and then had a marked increase in 2020 and 2021 and partial recovery in 2022-2023.

In sensitivity analysis using the first ICD listed in the last line of Part I instead of the last listed one, concordance rates were slightly lower (82.3% for broad category and 67.7% for ICD level) (see Supplementary Table S5A) and 69.1% with ICD-10 at 3-character categories.

Focusing on the broad death categories in Table 2 that have the most striking relative increases (COVID-19, Transport) or relative decreases (Other External, Alcohol) shows that a few specific category transitions are mostly responsible for the changes. For COVID-19, 288,936 deaths (29% of the 1,004,208 total) in the Record Axis are imported from the Respiratory category in the Entity Axis and another 119,181 deaths (12%) are imported from the Other Natural deaths in the Entity Axis. Transport deaths in the Record Axis receive 266,638 deaths (29% of the 915,055 total in the Record Axis) from the Other External causes in the Entity Axis. Other External causes get often re-classified also as Falls (218,625, 33% of Falls total deaths of 654,100 in Record Axis) and Suicide receives 139,167 deaths (18.2% of its total of 762,109 in the Record Axis) from Other External causes in the Entity Axis. Alcohol deaths in the Entity Axis often get re-classified as Drug Poisoning (87,860 deaths) in the Record Axis.

For COVID-19 deaths, the number of promotions to underlying cause of death (n=502,461) outnumber demotions (n=20,765) by 24-fold in the Entity to Record Axis re-classifications. Promotions largely re-classify the underlying cause of death away from respiratory and infection syndromes (Table S4). Most prominently, the ICD-10 J189 code of

pneumonia represents 43.8% of promotions. The demotions are much rarer and shift the underlying cause of death toward very diverse chronic disease causes.

Weighted causes of death

Table 3, panel 1 shows the unweighted (W0) and W1-, W2-, and W2A-weighted counts for deaths in broad disease categories for 2003-2023 and separately for the pandemic years 2020-2023 and Table 4 does the same separately for age strata. The weighting results in major changes for the relative contribution of some disease categories to the overall death burden and usually changes are more prominent with W2 and W2A.

With weighting, COVID-19 (-44-63%), Transport (-50-58%), Suicide (-50-60%), Falls (-46-66%) and Homicide (-50-65%) death counts decrease drastically and there are also major 19-32% decreases for Cancer and Alcohol-related deaths. Conversely, Other External causes more than triple (+204-254%); Other Natural, Respiratory and Endocrine, Nutrition and Metabolic deaths increase by 4-35% in the 2003-2023 data and slightly more prominently (8-44%) in the pandemic years alone. There are small decreases in Drug Poisoning and Digestive deaths, while Circulatory deaths remain largely unchanged. The rank order of causes changes substantially in the weighted analyses. For all years, all three weighting schemes promote Other External causes from rank 12 to 6. At the same time Digestive, COVID-19, Drug Poisoning, Suicide and Transport causes are demoted by between 1 and 3 ranks.

Table 3, panel 2 shows the weighted analyses using the Record Axis underlying cause with contribution causes from Part II of the Entity Axis instead of the causes from Record Axis as presented above: results were largely similar. Moreover, Figure S1 shows how the percent changes to each broad disease category for the different weighting schemes (W1, W2 and W2A) are highly correlated with correlation coefficients for the entire period of 0.809, and of 0.824 for

the pandemic years) for calculations done using the causes on Part II of the Entity Axis instead of causes on the Record Axis. Table 3, panel 3 shows that the results were largely similar when only information from Entity Axis (Entity underlying cause from last line of Part I with contributing causes from Part II) was used in the weighting.

These patterns follow directly from the disparity in count between underlying and contributing causes of death already seen in Table 1 and Supplementary Table S1. If there are equal counts of the two categories, sensible weighting is expected to have little effect (e.g. Circulatory). If there are fewer counts in contributing causes of death, such weighting decreases the death counts (e.g. COVID-19, Transport, Suicide, Falls) and if there are more counts in contributing causes of death, such weighting increases the death counts (e.g. Respiratory, Endocrine, Other External). In the extreme increases with weighting that are seen for Other External causes, the key reasons are respiratory tract obstruction, fractures, and injuries (Supplementary Table S6), and many of these ICD-10 codes probably reflect the nature rather than the cause of the injury.

Among deaths in ages under 65 years (Table 4), Cancer deaths are the most common cause in the unweighted counts, but with weighting the Circulatory category becomes the most common one. With the weighting schemes W2 and W2A Cancer gets to third rank after Circulatory and Other Natural causes. Conversely, with weighting the relative increases in Respiratory deaths and deaths due to Other Natural causes are more prominent in this age stratum, especially in the pandemic years. With weights W2 and W2A COVID-19 deaths are cut the most (by 64% and 57%, respectively) in this age stratum. In the age stratum of age 65 years and older (also in Table 4), for most of the broad death categories the relative change in death counts with weighting are similar as in the non-elderly stratum. However, there are some

exceptions. Boosting of Other External causes is less pronounced while boosting of Respiratory, Endocrine and Other Natural causes is more pronounced than in the non-elderly.

The corrections to the Record Axis underlying death counts made by all three weighting schemes are such as to make the counts become more similar to what they were in the Entity Axis of the death certificate. Specifically, the Pearson correlation coefficients of the fractional changes caused by weighting and the fractional change between Entity and Record axes is between 0.96 and 0.97 (Supplementary Table S7).

Figure 2 shows the evolution of death numbers between 2003 and 2023 for each of the broad disease categories using the unweighted W0 death counts and the W1-weighted death counts. There is observed seasonality for several broad categories. For Respiratory, but also for Circulatory, Endocrine Nutrition and Metabolic, Falls, and Other Natural disease causes there are annual peaks in the colder seasons and troughs in the summer, while for Transport and Suicide deaths the opposite seasonality pattern is seen. In the baseline W0 counts, the seasonality is largely lost for Respiratory deaths during the pandemic years, but it is recovered with the weighting. Supplementary Figures S3 and S4 show separate data for the elderly and non-elderly age groups. Similar patterns are seen with W2 and W2A weights (Supplementary Figures S5 and S6, and age subgroups in Supplementary Figures S7, S8, S9, and S10), although the magnitude of the deviations from W0 tend to be bigger with W2 and W2A than with W1.

There are also robust linear and logarithmic increases over time for some categories and age-groups (Supplementary Table S8; Supplementary Figure S11), such as Drug Poisoning (linear trend 6.7% per year), Alcohol (4.7% per year) and Falls (4.3% per year) deaths and less prominently for Other Natural (1.9% per year) and Suicide (2.0% per year). For the elderly, there are gradual decreases over time for Circulatory (-2.5% per year) and Cancer (-1.9% per year for

ages 65 or over) deaths. Linear and logarithmic annual percentage changes are very similar and weighting has little clear effect on these rates of change.

DISCUSSION

Our analysis of almost 57 million death certificates in the USA over 21 years offers insights on the impact of re-classifications during processing and of weighting for multimorbidity with multiple causes rather than a single death cause shaping each fatality. One third of deaths change ICD code for the selected underlying cause upon processing from the crude reported certificate to the version used in official national statistics and about half of them involve a shift even to a different broad disease category. Shifts have become more common over time and substantially escalated in the pandemic years, especially in 2020-2021, with the introduction of COVID-19 coding. Re-classification almost doubled the number of deaths where COVID-19 was selected as the underlying cause. Conversely, consideration of multimorbidity with weighting for multiple causes of death cut about half to two-thirds of the COVID-19 burden. It also generated substantive changes in the absolute death counts and in the relative ranking of several other causes of death. Weighting brought again the death counts for different disease categories closer to the Entity Axis. Seasonality peaks for Respiratory deaths were affected by the use or not of weighting.

The relatively high, and increasing, rates of disagreement between Record and Entity Axis are worrisome. They may reflect a high, and increasing, level of inaccuracy in filling death certificates. The escalating lack of concordance during 2020-2021 may reflect lack of acquaintance of certifiers in the new COVID-19 coding; or a particular difficulty in describing COVID-19 deaths and properly assigning the underlying cause of death in the standard framework where only one underlying cause is allowed. COVID-19 is a classic example where

almost all deaths occurred in the presence of major co-morbidities, and where several co-morbidities had strong associations with fatal outcomes in SARS-CoV-2 infections (27-29). It is thus possible that the WHO rules which were embedded also in the USA re-classification mortality coding rules (much like in other countries) were markedly overestimating COVID-19 death burden (23). Record Axis COVID-19 deaths have included many deaths that had multiple other causes with substantive contribution to the fatal outcome. Even beyond COVID-19, deaths with high levels of multimorbidity are probably becoming more common over time posing a challenge for certifiers to select a single underlying cause. Additional analyses using weighting for multiple causes of deaths need to be conducted across diverse countries, with priority on those that have the most complete information from death certificates. Some of our findings may be more specific to the USA (e.g. drug use and homicide counts) while others, such as changes in recording and coding of COVID-19 were experienced in many countries around the world.

Specific non-COVID-19 re-classifications are heavily represented, especially those involving the shift from Other External causes to Transport, Suicide, and Falls deaths. Many such shifts may be appropriate to capture the more precise initiating event that led to death. Some re-classifications may be debatable, however. For example, Alcohol-related death counts were overall reduced by the re-classification, but this may be underestimating the contribution of alcohol to many other conditions that eventually kill people. Moreover, it is well documented that alcohol is already grossly under-reported in death certificates: for example, traffic census data show that in motor traffic crash deaths with detected high alcohol levels, only 16% of death certificates in the USA mention alcohol, with wide variation across states (2-81%) (30).

The weighting schemes attempt to correct for the multi-factorial nature of deaths, a scenario that has become extremely common, largely the norm rather than the exception, in the

21st century. The paradigm of listing only one underlying cause of death was quite appropriate for eras where previously healthy, young people were dying from acute infectious diseases or violence. In the current circumstances, this paradigm has very questionable validity. Most people die from the concerted impact of multiple causes that undermine their health over a variable range of time. Some causes act on a shorter time horizon than others. The exact extent to which causes other than the underlying cause should be considered is debated. We tried three weighting options inspired by previous work (19-22), extending them to consider other causes both at the level of broad categories and at ICD level.

The weighted analyses yielded largely similar results in terms of direction, although W2 and W2A tend to produce larger deviations from the baseline unweighted W0 than W1. Weighting tends to bring the corrected death counts closer to what certifiers had selected in the Entity Axis. When multimorbidity is extreme (like in COVID-19), the more prominently altered estimates may be more appropriate. A much lower than officially recorded number of COVID-19 deaths is commensurate also with the small loss of life years estimated for many of those fatalities (31). It also agrees with the findings of several in-depth audit evaluations of medical records in different countries where large percentages of previously officially proclaimed COVID-19 deaths are found to have other primary, underlying death causes (9-11). It is thus likely that the true pandemic burden of COVID-19 corresponded to only less than half or even just a third of the officially assigned COVID-19 deaths in the USA. Probably COVID-19 during 2020-2023 accounted cumulatively for ~100,000 deaths among people less than 65 years old, and ~300,000 deaths among people 65 years old and older.

Changes in the frequency of specific death causes over time may reflect genuine changes due to changing incidence (e.g. infectious disease waves) or improvements in diagnostic and

therapeutic management and outcomes. However, changes may also reflect an alteration of the perceived importance of a death cause among health care practitioners and in the community in general. This may affect whether a cause is listed at all and whether it is chosen to be listed as underlying or contributing. An example has been dissected in the case of Alzheimer's disease where an increase in its listing as underlying cause was accompanied by a decrease in its listing as contributing cause (16). In our data, perception issues may underlie the observed patterns for COVID-19. Apparently, in the very early phase of the pandemic, some COVID-19 deaths were not recorded, either because their etiology was entirely missed or because certifiers may have not considered COVID-19 as the underlying cause. At this early phase, substantial under-reporting of COVID-19 deaths may have been reflected in increased death rates from other listed causes (32). Later the inverse pattern probably predominated, with severe over-counting of COVID-19 deaths in an environment where the pandemic was a dominant pervasive theme in health, media and in society at large, excessive testing was performed (23).

Perception, testing, and recognition issues are most stark with COVID-19 but may affect also in more subtle and chronic ways several major causes of deaths from chronic disease. For example, other investigators have noted that perception issues may propel ischemic heart disease, rather than diabetes or hypertension or heart failure to be listed as the underlying cause of death (22). Weighting of multiple death causes may correct some of this injustice of all-or-none attribution. However, it has also been argued that weighting may be inappropriate for some causes that might be truly the main and only determinant of fatality, as for example for many cancers (22). Moreover, some additional listed external causes, such as injuries and fractures, and respiratory tract obstruction may reflect mostly the nature of an injury and death process rather than causes that should get a share of the weight. Furthermore, one could further question

whether the chronic trends (with or without weighting) reflect real changes or perception changes. This dilemma could affect even the most common causes of disease, i.e. circulatory and cancer deaths. Certifiers may be certifying causes according to the increase or decrease of the reputational presence of each disease in the environment where they practice. Perception biases could make death certificate data become self-fulfilling prophecies about the burden of specific diseases and the evolution of this burden over time.

Our work has several limitations. First, it should be acknowledged that the selected grouping of death causes can make a difference. Here we used primarily high-level, broad categories of death, to avoid the frequent dilemmas between different ICD-10 codes within the same system, as in ischemic heart disease and hypertension (19-22). This approach captures only the most prominent shifts in causal categorization. At the level of single ICD-10 codes, shifts both with re-classification and with weighting were documented to be even more extreme.

Second, while rational arguments can be made regarding whether re-classified or specific weighted options have more (or less) validity than crude reported numbers of deaths, the real gold standard validation of different estimates requires autopsy studies and in-depth audit of medical records. Error rates regarding causes of death in death certificates are so high that healthy skepticism is warranted. Illustratively, in Vermont, a medical record audit study found that the underlying cause of death reported in death certificates was wrong in 60% and errors in ICD codes appeared in 92% of certificates (33). There is substantial potential for errors even with detailed medical records. For example, a study of medical examiners (arguably, the best-trained group to certify deaths) in Australia found that when they attributed deaths based on detailed perusal of medical records, their selected cause of death was still wrong 28% of the time when compared to autopsy (34). Third, we provide separate results for two large age strata and

these already show that the impact of different weighting may vary for some death causes in these age groups. However, we cannot exclude that the patterns generated from weighting may deviate even further from the baseline in more granular substrata, e.g. defined by residence status (community versus nursing homes) or socioeconomic indicators. Documentation of multiple as opposed to single causes of death depends on a wide range of socioeconomic and other factors (35). These associations may reflect both genuine differences in multi-morbidity, as well as differences in the likelihood that multiple causes would be documented even if they exist. For example, multiple causes may be more likely to be documented when patients get better health care with continuity and when they are hospitalized rather than when they have no care and die at home (35).

Acknowledging these sources of complexity, we argue that re-classification patterns and multiple causes of death should be routinely considered in analyses of death certificate data. Different re-classification patterns may point to structural defects in death certification that may require targeted training and education of certifiers; or that specific mortality coding rules may be promoting misleading changes. Statistics based on single underlying causes of death may be grossly inappropriate, more so for some causes like COVID-19 than for others. The reliance on single underlying causes of death has long served to encourage condition-specific focus in service provision, medical training, research and lobbying. Public health and health statistics worldwide should move away from unilateral dependence on this concept of only relative value and embrace more widely approaches that consider multiple causes of death. Different patterns emerging from weighting may point out that the burden from specific causes and categories of causes may be much larger or smaller than traditional death statistics suggest. This can have profound implications for resource allocation for both research and healthcare. Incorporation of

weighted estimates may help shape more appropriate, evidence-based public health priorities. Additional in-depth validation assessments of death causes in diverse populations, including higher rates of autopsies, are also needed to better understand and more properly calibrate death certificate data.

REFERENCES

1. Alipour J, Payandeh A. Common errors in reporting cause-of-death statements on death certificates: a systematic review and meta-analysis. J Forensic Leg Med 2021;82:102220.

2. Johansson A, Björkenstam C, Westerling R. Unexplained differences between hospital and mortality data indicated mistakes in death certification: an investigation of 1,094 deaths in Sweden during 1995. J Clin Epidemiol. 2009;62:1202-9.

3. Mathers CD, Fat DM, Inoue M, Rao C, Lopez AD. Counting the dead and what they died from: an assessment of the global status of cause of death data. Bull World Health Organ 2005;83(3):171-7.

4. Johansson LA, Westerling R. Comparing Swedish hospital discharge records with death certificates: implications for mortality statistics. Int J Epidemiol 2000 Jul;29(3):495-502.

5. Smith Sehdev AE, Hutchins GM. Problems with proper completion and accuracy of the cause-of-death statement. Arch Intern Med 2001;161(2):277-84.

6. National Vital Statistics System. About the mortality medical data system. https://www.cdc.gov/nchs/nvss/mmds/about_mmds.htm, last accessed December 24, 2025.

7. National Vital Statistics System. ACME and TRANSAX. https://www.cdc.gov/nchs/nvss/mmds/acme_transax.htm, last accessed December 24, 2025.

8. National Vital Statistics System. Reporting and coding deaths due to COVID-19. https://www.cdc.gov/nchs/covid19/coding-and-reporting.htm, last accessed December 24, 2025.

9. Basoulis D, Logioti K, Papaodyssea I, Chatzopoulos M, Alexopoulou P, Mavroudis P, et al. Deaths "due to" COVID-19 and deaths "with" COVID-19 during the Omicron variant surge, among hospitalized patients in seven tertiary-care hospitals, Athens, Greece. Sci Rep. 2025;15(1):13728.

10. Spreco A, Andersson C, Sjödahl R, Timpka T. Concordance between COVID-19 mortality statistics derived from clinical audit and death certificates in Östergötland county, Sweden. Public Health. 2023;221:46–9.

11. White P, Crowe C, Bowe A, Brennan A, Bruton O, O'Sullivan MB, O'Mahony MT, Sheahan A, Barrett P. An audit of COVID-19 death reporting in counties Cork and Kerry, Ireland, winter 2021–2022. Irish J Medical Sci 2023;192:1589-94.

12. Désesquelles A, Gamboni A, Demuru E and the MultiCause network. We only die once… but from how many causes? Popul Soc. 2016;534.

13. Moreno-Betancur M, Sadaoui H, Piffaretti C, Rey G. Survival analysis with multiple causes of death: extending the competing risks model. Epidemiology. 2017;28(1):12–9.

14. Israel RA, Rosenberg HM, Curtin LR. Analytical potential for multiple cause-of-death data. Am J Epidemiol. 1986;124(2):161–79.

15. Desesquelles A, Salvatore M, Frova L, Pace M, Pappagallo M, Meslé F, et al. Revisiting the mortality of France and Italy with the multiple-cause-of-death approach. Demogr Res. 2010;23(28):771–806.

16. Adair T, Temple J, Anstey KJ, Lopez AD. Is the rise in reported dementia mortality real? Analysis of multiple-cause-of-death data for Australia and the United States. Am J Epidemiol. 2022;191(7):1270–9.

17. D'Amico M, Agozzino E, Biagino A, Simonetti A, Marinelli P. Ill-defined and multiple causes on death certificates--a study of misclassification in mortality statistics. Eur J Epidemiol. 1999 Feb;15(2):141–8.

18. Baneshi MR, Eynstone-Hinkins J, McElwee P, Mishra GD, Moran L, Waller M, Dobson A. What can death records tell us about multimorbidity? J Epidemiol Community Health. 2023;77(8):507-514.

19. Bishop K, Balogun S, Eynstone-Hinkins J, Moran L, Martin M, Banks E, et al. Analysis of Multiple Causes of Death: A Review of Methods and Practices. Epidemiology. 2023;34(3):333.

20. Bishop K, Moreno-Betancur M, Balogun S, Eynstone-Hinkins J, Moran L, Rao C, et al. Quantifying cause-related mortality in Australia, incorporating multiple causes: observed patterns, trends and practical considerations. Int J Epidemiol. 2023;52(1):284–94.

21. Piffaretti C, Moreno-Betancur M, Lamarche-Vadel A, Rey G. A method for measuring the level of cause-related mortality using weighted multiple cause of death data. Bull World Health Organ. 2016;94:870-879B.

22. Dobson A, McElwee P, Baneshi MR, Eynstone-Hinkins J, Moran L, Waller M. A new data driven method for summarising multiple cause of death data. BMC Med Res Methodol. 2023;23(1):83.

23. Ioannidis JPA. Over- and under-estimation of COVID-19 deaths. Eur J Epidemiol. 2021 Jun;36(6):581–8.

24. https://ftp.cdc.gov/pub/Health_Statistics/NCHS/Datasets/DVS/mortality/

25. Wrigley-Field E, Raquib RV, Berry KM, Morris KJ, Stokes AC. Mortality trends among early adults in the United States, 1999-2023. JAMA Network Open 2025;8:e2457538.

26. NASEM. High and rising mortality rates among working-age adults. National Academies Press, Washington D.C., 2021:576.

27. Ganaza-Domingues KLT, Ramos-Milaré ÁCFH, Lera-Nonose DSSL, Brustolin AÁ, de Oliveira LF, Rosa JS, Otofuji Inada AY, Dias Leme AL, Pinel BI, Perina BS, de Souza Terron M, da Silva Santos T, Demarchi IG, Lonardoni MVC, Teixeira JJV. Effect of comorbidities on the mortality of patients with COVID-19: a systematic review of reviews and meta-analyses. Rev Med Virol. 2025 Mar;35(2):e70024.

28. Thakur B, Dubey P, Benitez J, Torres JP, Reddy S, Shokar N, Aung K, Mukherjee D, Dwivedi AK. A systematic review and meta-analysis of geographic differences in comorbidities and associated severity and mortality among individuals with COVID-19. Sci Rep. 2021 Apr 20;11(1):8562.

29. Ssentongo P, Ssentongo AE, Heilbrunn ES, Ba DM, Chinchilli VM. Association of cardiovascular disease and 10 other pre-existing comorbidities with COVID-19 mortality: A systematic review and meta-analysis. PLoS One. 2020;15(8):e0238215.

30. Castle IJ, Yi HY, Hingson RW, White AM. State variation in underreporting of alcohol involvement on death certificates: motor vehicle traffic crash fatalities as an example. J Stud Alcohol Drugs. 2014 Mar;75(2):299-312.

31. Ioannidis JPA, Pezzullo AM, Cristiano A, Boccia S. Global estimates of lives and life-years saved by COVID-19 vaccination during 2020-2024. JAMA Health Forum. 2025 Jul 3;6(7):e252223.

32. Li Y, Li H, Adair T. The impact of the pandemic on non-COVID-19 causes of death in the united States: a multiple cause of death analysis. European Journal of Epidemiology. 2025 Apr;40(4):463-74. DOI: 10.1007/s10654-025-01214-z.

33. McGivern L, Shulman L, Carney JK, Shapiro S, Bundock E. Death certification errors and the effect on mortality statistics. Public Health Rep. 2017;132(6):669-675.

34. Nashelsky MB, Lawrence CH. Accuracy of cause of death determination without forensic autopsy examination. Am J Forensic Med Pathol. 2003 Dec;24(4):313-9.

35. Wall MM, Huang J, Oswald J, McCullen D. Factors associated with reporting multiple causes of death. BMC Med Res Methodol. 2005 Jan 17;5(1):4.

Figure legends

Figure 1. Population-normalized monthly deaths across 14 broad categories during 2003-2023 for all ages comparing unweighted W0 with W1 weighting. In this figure as well as in supplementary figures S3 to S10, unweighted is a colored line and weighted is a thick black line. Disease categories are grouped by fixed Y-axis scales to allow visual comparison across panels. The top band (COVID-19, Circulatory, Cancer, Other natural, Respiratory) uses 0-130,000 deaths/month; the middle band (Endocrine, Digestive, Drug poisoning, Other external) uses 0-25,000; the bottom band (Transport, Suicide, Alcohol-related, Falls, Homicide) uses 0-6,000. All values are population-scaled to 2023 levels.

Table 1. Causes of death classified in broad disease categories[++] in Entity Axis and Record Axis, USA 2003-2023 (when there was more than one ICD code listed in the last line of Part I in the Entity Axis, we used the one listed last).

| Disease | Entity Axis | | | | Record Axis | | | |
	Underlying (i.e. lowest line of Part I) (% change)	Contributing	Only	Any	Underlying	Contributing	Only[+]	Any[#]
Circulatory	17,793,268	13,582,667	7,521,247	31,375,935	17,796,602 (0%)	13,510,161	7,678,693	31,306,763
Cancer	11,214,012	2,817,512	5,476,807	14,031,524	12,576,466 (12%)	1,435,404	5,482,941	14,011,870
Other Natural	12,386,337	17,171,701	4,603,744	29,558,038	10,708,481 (-14%)	18,124,458	4,662,437	28,832,939
Respiratory	5,978,426	9,277,447	1,295,638	15,255,873	5,303,862 (-11%)	9,896,044	1,295,750	15,199,906
Endocrine*	2,209,113	7,136,922	124,334	9,346,035	2,560,585 (16%)	6,780,460	156,081	9,341,045
Digestive	1,675,880	2,692,069	346,111	4,367,949	1,731,012 (3%)	2,394,730	346,257	4,125,742
COVID-19	522,512	647,301	58,211	1,169,813	1,004,208 (92%)	165,605	58,208	1,169,813
Drug Poisoning	1,130,962	342,169	701,529	1,473,131	1,229,766 (9%)	241,136	701,531	1,470,902
Transport	637,778	296,760	852	934,538	915,055 (43%)	19,491	853	934,546
Suicide	609,437	155,860	662	765,297	762,109 (25%)	3,190	662	765,299
Alcohol-related	686,868	669,287	102,929	1,356,155	676,440 (-2%)	679,715	155,262	1,356,155
Other External	1,451,348	3,420,721	269,514	4,872,069	671,615 (-54%)	4,197,101	269,509	4,868,716
Falls	386,291	427,221	161	813,512	654,100 (69%)	159,415	161	813,515
Homicide	304,599	96,049	275	400,648	396,530 (30%)	4,118	275	400,648
Uncertain	0	0	0	0	0	518	0	518
Total	**56,986,831**	**58,733,686**	**20,502,014**	**115,720,517**	**56,986,831**	**57,611,546**	**20,808,620**	**114,598,377**

[++] The mapping of the ICD-10 codes onto broad disease categories is given in Table S9.

[+] Only means that there only one broad category cause of death for the particular axis. There are 309,574 death certificates with more than a single unique broad cause of disease for the record or the entity axis when the other axis has only a single cause. For the vast majority of certificates (308,090), it is the Entity Axis that has multiple causes.

[#] Any means that we code all broad disease categories.

*endocrine, nutritional and metabolic

Table 2. Transition matrix from Entity Axis (columns) to Record Axis (rows). When there was more than one ICD code listed in the last line of Part I in the Entity Axis, we used the one listed last.

ENTITY RECORD	Circulatory	Cancer	Other Natural	Respiratory	Endocrine*	Digestive	COVID-19	Drug Poisoning	Transport	Suicide	Alcohol	Other External	Falls	Homicide	Total Record Axis
Circulatory	15,936,312	61,223	1,092,514	408,494	143,301	49,031	7,330	16,246	1,313	104	12,981	63,365	4,047	341	17,796,602
Cancer	344,036	11,020,772	598,117	360,314	109,320	104,079	3,591	1,323	135	26	14,196	19,895	640	22	12,576,466
Other Natural	628,963	53,307	9,350,816	406,374	130,961	60,841	4,473	7,028	714	68	16,553	45,311	2,840	232	10,708,481
Respiratory	297,054	29,430	506,545	4,346,838	57,638	17,118	2,539	2,832	239	14	4,870	37,738	966	41	5,303,862
Endocrine*	379,340	13,610	364,998	60,867	1,711,842	11,766	1,924	2,013	111	8	4,529	9,227	329	21	2,560,585
Digestive	68,911	14,182	187,179	54,731	15,488	1,372,198	338	1,017	159	8	2,568	13,926	257	50	1,731,012
COVID-19	54,475	10,168	119,181	288,936	21,328	5,017	501,747	204	48	5	567	2,251	271	10	1,004,208
Drug Poisoning	16,732	1,578	13,339	7,380	2,855	1,906	125	1,088,054	96	482	87,860	9,181	83	95	1,229,766
Transport	4,014	169	3,570	1,686	455	242	26	1,086	634,750	1	2,378	266,638	31	9	915,055
Suicide	531	443	9,809	359	121	78	14	1,943	5	608,669	966	139,167	3	1	762,109
Alcohol-related	17,325	2,231	48,391	11,891	8,062	48,038	87	3,809	50	31	530,942	5,339	233	11	676,440
Other External	29,504	4,740	63,407	22,983	5,584	4,561	89	4,721	107	14	4,980	530,516	391	18	671,615
Falls	15,552	2,144	27,326	7,253	1,974	929	229	426	39	2	3,402	218,625	376,199	0	654,100
Homicide	519	15	1,145	320	184	76	0	260	12	5	76	90,169	1	303,748	396,530
Total Entity Axis	17,793,268	11,214,012	12,386,337	5,978,426	2,209,113	1,675,880	522,512	1,130,962	637,778	609,437	686,868	1,451,348	386,291	304,599	56,986,831

*endocrine, nutritional and metabolic

Table 3. Underlying death counts for broad disease categories for 2003-2023 and for the pandemic years 2020-2023 with different weighting schemes (W0: no weighting, only the underlying cause is counted; W1: 50% weight for the underlying cause and the other 50% shared equally among other causes; W2: weight shared equally among all causes; W2A: weight shared equally among all causes but considering ICD-level causes rather than just broad categories). The upper panel weights the counts of ICD on the Record Axis. The middle panel weights the counts for the underlying ICD on the Record Axis with the contributing ICD on the Entity Axis. The lower panel weights the counts for the underlying ICD on the Entity Axis (last line of Part I) with the contributing ICD on the Entity Axis.

Counts and percentage relative to W0 for the counts of ICD causes from the Record Axis

Disease	All Years (2003-2023)				Pandemic Years (2020-2023)			
	W0	W1	W2	W2A	W0	W1	W2	W2A
Circulatory	17,797	17,960 (+1)	18,060 (+1)	17,045 (-4)	3,715	3,867 (+4)	3,942 (+6)	3,688 (-1)
Cancer	12,576	9,392 (-25)	8,546 (-32)	8,832 (-30)	2,494	1,827 (-27)	1,615 (-35)	1,678 (-33)
Other Natural	10,708	13,538 (+26)	14,424 (+35)	14,186 (+32)	2,468	3,131 (+27)	3,378 (+37)	3,286 (+33)
Respiratory	5,304	6,122 (+15)	6,287 (+19)	6,481 (+22)	1,046	1,388 (+33)	1,497 (+43)	1,511 (+44)
Endocrine*	2,561	2,653 (+4)	2,707 (+6)	3,404 (+33)	679	733 (+8)	766 (+13)	950 (+40)
Digestive	1,731	1,636 (-5)	1,565 (-10)	1,668 (-4)	394	365 (-7)	347 (-12)	375 (-5)
COVID-19	1,004	566 (-44)	370 (-63)	441 (-56)	1,004	566 (-44)	370 (-63)	441 (-56)
Drug Poisoning	1,230	1,140 (-7)	1,087 (-12)	988 (-20)	426	396 (-7)	377 (-11)	342 (-20)
Transport	915	461 (-50)	380 (-58)	437 (-52)	192	97 (-50)	81 (-58)	91 (-52)
Suicide	762	382 (-50)	308 (-60)	363 (-52)	174	87 (-50)	69 (-60)	83 (-52)
Alcohol-related	676	547 (-19)	512 (-24)	574 (-15)	202	162 (-20)	150 (-26)	170 (-16)
Other External	672	2,039 (+204)	2,376 (+254)	2,104 (+213)	145	454 (+212)	533 (+266)	482 (+231)
Falls	654	351 (-46)	225 (-66)	269 (-59)	180	97 (-46)	59 (-67)	72 (-60)
Homicide	397	200 (-50)	140 (-65)	195 (-51)	97	49 (-50)	34 (-66)	48 (-51)

Counts and percentage change relative to W0 of the underlying Record Axis cause with the contributing causes on Part II of the Entity Axis

Disease	All Years (2003-2023)				Pandemic Years (2020-2023)			
	W0	W1	W2	W2A	W0	W1	W2	W2A
Circulatory	17,816[+]	16,679 (-6)	16,594 (-7)	16,330 (-8)	3,715	3,556 (-4)	3,568 (-4)	3,484 (-6)
Cancer	12,576	10,789 (-14)	10,396 (-17)	10,514 (-16)	2,494	2,085 (-16)	1,971 (-21)	2,004 (-20)
Other Natural	10,692	13,742 (+29)	14,228 (+33)	13,961 (+31)	2,468	3,191 (+29)	3,336 (+35)	3,250 (+32)
Respiratory	5,308	4,794 (-10)	4,694 (-12)	4,880 (-8)	1,046	965 (-8)	952 (-9)	1,000 (-4)
Endocrine*	2,550	3,448 (+35)	3,706 (+45)	3,865 (+52)	679	920 (+35)	1,002 (+47)	1,046 (+54)
Digestive	1,731	1,662 (-4)	1,641 (-5)	1,695 (-2)	394	378 (-4)	372 (-6)	388 (-1)
COVID-19	1,004	784 (-22)	705 (-30)	733 (-27)	1,004	784 (-22)	705 (-30)	733 (-27)
Drug Poisoning	1,230	1,165 (-5)	1,142 (-7)	1,099 (-11)	426	403 (-5)	394 (-7)	380 (-11)
Transport	915	852 (-7)	839 (-8)	845 (-8)	192	178 (-8)	174 (-9)	176 (-9)
Suicide	762	444 (-42)	428 (-44)	431 (-43)	174	95 (-46)	91 (-48)	92 (-47)
Alcohol-related	680	665 (-2)	655 (-4)	677 (-0)	202	192 (-5)	187 (-8)	195 (-4)
Other External	672	1,246 (+85)	1,306 (+94)	1,273 (+90)	145	295 (+103)	310 (+113)	305 (+110)
Falls	654	478 (-27)	420 (-36)	445 (-32)	180	125 (-31)	105 (-42)	114 (-37)
Homicide	397	240 (-40)	235 (-41)	237 (-40)	97	53 (-46)	52 (-47)	52 (-46)

Counts and percentage change relative to W0 of the underlying Entity Axis (last line of Part 1) cause with the contributing causes on Part II of the Entity Axis

	All Years (2003-2023)	Pandemic Years (2020-2023)

Disease	W0	W1	W2	W2A	W0	W1	W2	W2A
Circulatory	17,793	16,726 (-6)	16,670 (-6)	16,386 (-8)	3,773	3,618 (-4)	3,631 (-4)	3,541 (-6)
Cancer	11,214	9,798 (-13)	9,484 (-15)	9,593 (-14)	2,208	1,882 (-15)	1,790 (-19)	1,821 (-18)
Other Natural	12,386	14,917 (+20)	15,271 (+23)	14,997 (+21)	2,960	3,525 (+19)	3,623 (+22)	3,536 (+19)
Respiratory	5,978	5,224 (-13)	5,049 (-16)	5,259 (-12)	1,392	1,206 (-13)	1,157 (-17)	1,215 (-13)
Endocrine*	2,209	3,225 (+46)	3,528 (+60)	3,713 (+68)	598	873 (+46)	969 (+62)	1,021 (+71)
Digestive	1,676	1,627 (-3)	1,615 (-4)	1,669 (-0)	388	374 (-4)	370 (-5)	386 (-0)
COVID-19	523	422 (-19)	382 (-27)	401 (-23)	523	422 (-19)	382 (-27)	401 (-23)
Drug Poisoning	1,131	1,120 (-1)	1,109 (-2)	1,072 (-5)	389	387 (-1)	382 (-2)	370 (-5)
Transport	638	653 (+2)	645 (+1)	651 (+2)	142	140 (-2)	138 (-3)	139 (-2)
Suicide	609	371 (-39)	359 (-41)	362 (-41)	140	82 (-41)	79 (-44)	80 (-43)
Alcohol-related	687	660 (-4)	650 (-5)	670 (-2)	208	190 (-9)	185 (-11)	192 (-8)
Other External	1,451	1,703 (+17)	1,709 (+18)	1,673 (+15)	307	385 (+25)	386 (+26)	382 (+24)
Falls	386	350 (-9)	330 (-15)	351 (-9)	111	92 (-17)	84 (-25)	91 (-18)
Homicide	305	190 (-38)	187 (-38)	189 (-38)	78	44 (-44)	43 (-45)	43 (-44)

*endocrine, nutritional metabolic

[+]There are small differences in W0 counts with the different classification scheme used in this panel.

Table 4. Death counts for broad disease categories for 2003-2023 and for the pandemic years 2020-2023 with different weighting schemes (W0: no weighting, only the underlying cause is counted; W1: 50% weight for the underlying cause and the other 50% shared equally among other causes; W2: weight shared equally among all causes; W2A: weight shared equally among all causes but considering ICD-level causes rather than just broad categories) separately for deaths under 65 years old and for deaths 65 years old or older. The causes weighted are all on the Record Axis.

Death counts for non-elderly, under 65 years old, in thousands (% of W0)

Disease	All Years (2003-2023)				Pandemic Years (2020-2023)			
	W0	W1	W2	W2A	W0	W1	W2	W2A
Circulatory	3,312	3,396 (+3)	3,425 (+3)	3,288 (-1)	695	721 (+4)	734 (+6)	703 (+1)
Cancer	3,635	2,744 (-25)	2,524 (-31)	2,576 (-29)	638	469 (-27)	417 (-35)	428 (-33)
Other Natural	2,086	2,890 (+39)	3,136 (+50)	3,068 (+47)	402	603 (+50)	678 (+69)	654 (+63)
Respiratory	808	1,074 (+33)	1,147 (+42)	1,181 (+46)	168	267 (+59)	302 (+79)	302 (+79)
Endocrine*	721	715 (-1)	713 (-1)	858 (+19)	180	189 (+5)	195 (+8)	232 (+29)
Digestive	532	561 (+5)	557 (+5)	579 (+9)	116	117 (+1)	115 (-1)	122 (+5)
COVID-19	248	138 (-44)	89 (-64)	106 (-57)	248	138 (-44)	89 (-64)	106 (-57)
Drug Poisoning	1,159	1,076 (-7)	1,027 (-11)	934 (-19)	396	368 (-7)	351 (-11)	318 (-20)
Transport	745	374 (-50)	311 (-58)	358 (-52)	153	77 (-50)	65 (-57)	73 (-52)
Suicide	622	312 (-50)	254 (-59)	297 (-52)	139	69 (-50)	56 (-60)	66 (-52)
Alcohol-related	526	426 (-19)	402 (-24)	450 (-15)	152	122 (-20)	113 (-25)	128 (-16)
Other External	356	1,278 (+259)	1,472 (+314)	1,303 (+267)	74	278 (+274)	322 (+333)	290 (+290)
Falls	100	52 (-48)	35 (-65)	42 (-58)	23	12 (-48)	8 (-65)	9 (-59)
Homicide	376	189 (-50)	132 (-65)	185 (-51)	92	46 (-50)	32 (-66)	46 (-51)

Death counts for elderly, 65 years old or older, in thousands (% of W0)

Disease	All Years (2003-2023)				Pandemic Years (2020-2023)			
	W0	W1	W2	W2A	W0	W1	W2	W2A
Circulatory	14,485	14,564 (+1)	14,635 (+1)	13,757 (-5)	3,021	3,147 (+4)	3,208 (+6)	2,986 (-1)
Cancer	8,942	6,648 (-26)	6,022 (-33)	6,257 (-30)	1,855	1,358 (-27)	1,197 (-35)	1,251 (-33)
Other Natural	8,623	10,648 (+23)	11,287 (+31)	11,119 (+29)	2,066	2,528 (+22)	2,701 (+31)	2,633 (+27)
Respiratory	4,496	5,048 (+12)	5,140 (+14)	5,299 (+18)	878	1,121 (+28)	1,196 (+36)	1,210 (+38)
Endocrine*	1,839	1,938 (+5)	1,994 (+8)	2,546 (+38)	499	544 (+9)	571 (+14)	718 (+44)
Digestive	1,199	1,076 (-10)	1,008 (-16)	1,089 (-9)	278	248 (-11)	232 (-17)	253 (-9)
COVID-19	756	427 (-43)	281 (-63)	334 (-56)	756	427 (-43)	281 (-63)	334 (-56)
Drug Poisoning	70	64 (-9)	60 (-15)	54 (-23)	30	28 (-8)	26 (-13)	24 (-21)
Transport	171	87 (-49)	69 (-60)	79 (-54)	39	20 (-49)	16 (-60)	18 (-54)
Suicide	140	70 (-50)	54 (-62)	66 (-53)	36	18 (-50)	13 (-62)	17 (-53)
Alcohol-related	150	121 (-20)	110 (-27)	124 (-17)	51	40 (-20)	37 (-28)	42 (-17)
Other External	316	760 (+141)	904 (+186)	801 (+153)	71	176 (+148)	210 (+196)	192 (+171)
Falls	554	299 (-46)	190 (-66)	227 (-59)	158	85 (-46)	51 (-67)	62 (-61)
Homicide[+]	20	11 (-48)	8 (-63)	10 (-51)	5	3 (-49)	2 (-64)	3 (-52)

*endocrine, nutritional metabolic

[+]The percentage changes using the complete counts not those rounded in thousands.

SUPPLEMENTARY MATERIAL
Supplementary Tables
Table S1. Causes of death classified in broad disease categories in Record Axis taken from Table 1 to show ratio of counts to Underlying.

Letter	Underlying	Contributing	Only	Anywhere	Contributing/ Underlying	Anywhere/ Underlying
All Ages						
Circulatory	17,796,602	13,510,161	7,678,693	31,306,763	0.76	1.76
Cancer	12,576,466	1,435,404	5,482,941	14,011,870	0.11	1.11
Other Natural	10,708,481	18,124,458	4,662,437	28,832,939	1.69	2.69
Respiratory	5,303,862	9,896,044	1,295,750	15,199,906	1.87	2.87
Endocrine*	2,560,585	6,780,460	156,081	9,341,045	**2.65**	**3.65**
Digestive	1,731,012	2,394,730	346,257	4,125,742	1.38	2.38
COVID-19	1,004,208	165,605	58,208	1,169,813	0.16	1.16
Drug Poisoning	1,229,766	241,136	701,531	1,470,902	0.20	1.20
Transport	915,055	19,491	853	934,546	0.02	1.02
Suicide	762,109	3,190	662	765,299	0.00	1.00
Alcohol-related	676,440	679,715	155,262	1,356,155	1.00	2.00
Other External	671,615	4,197,101	269,509	4,868,716	**6.25**	**7.25**
Falls	654,100	159,415	161	813,515	0.24	1.24
Homicide	396,530	4,118	275	400,648	0.01	1.01
Under 65						
Circulatory	3,311,687	2,870,974	1,427,809	6,182,661	0.87	1.87
Cancer	3,634,924	217,427	1,706,537	3,852,351	0.06	1.06
Other Natural	2,085,650	4,228,619	1,017,129	6,314,269	2.03	3.03
Respiratory	808,320	2,139,295	175,255	2,947,615	2.65	3.65
Endocrine*	721,155	1,548,828	66,023	2,269,983	**2.15**	**3.15**
Digestive	531,999	854,850	130,223	1,386,849	1.61	2.61
COVID-19	248,498	31,601	15,369	280,099	0.13	1.13
Drug Poisoning	1,159,365	205,731	673,252	1,365,096	0.18	1.18
Transport	744,515	10,074	704	754,589	0.01	1.01
Suicide	621,785	2,395	504	624,180	0.00	1.00
Alcohol-related	525,985	515,548	126,220	1,041,533	0.98	1.98
Other External	355,615	2,340,926	205,078	2,696,541	**6.58**	**7.58**
Falls	99,661	14,162	23	113,823	0.14	1.14
Homicide	376,112	2,709	256	378,821	0.01	1.01
65 Years or Older						
Circulatory	14,484,915	10,639,187	6,250,884	25,124,102	0.73	1.73
Cancer	8,941,542	1,217,977	3,776,404	10,159,519	0.14	1.14
Other Natural	8,622,831	13,895,839	3,645,308	22,518,670	1.61	2.61
Respiratory	4,495,542	7,756,749	1,120,495	12,252,291	1.73	2.73
Endocrine*	1,839,430	5,231,632	90,058	7,071,062	**2.84**	**3.84**
Digestive	1,199,013	1,539,880	216,034	2,738,893	1.28	2.28
COVID-19	755,710	134,004	42,839	889,714	0.18	1.18
Drug Poisoning	70,401	35,405	28,279	105,806	0.50	1.50
Transport	170,540	9,417	149	179,957	0.06	1.06
Suicide	140,324	795	158	141,119	0.01	1.01
Alcohol-related	150,455	164,167	29,042	314,622	1.09	2.09
Other External	316,000	1,856,175	64,431	2,172,175	**5.87**	**6.87**
Falls	554,439	145,253	138	699,692	0.26	1.26
Homicide	20,418	1,409	19	21,827	0.07	1.07

*endocrine nutritional and metabolic

Table S2. Causes of death classified in broad disease categories in Record Axis and Entity Axis, USA 2003-2023 using the sensitivity analysis approach (choosing the first listed ICD code in the last line of the Part I for the Entity Axis)

Disease	Record Axis				Entity Axis			
Disease	Underlying	Contributing	Only	Any	Underlying	Contributing	Only	Any
Circulatory	17,796,602 (0%)	13,510,161	7,678,693	31,306,763	17,770,736	13,605,199	7,521,247	31,375,935
Cancer	12,576,466 (12%)	1,435,404	5,482,941	14,011,870	11,182,712	2,848,812	5,476,807	14,031,524
Other Natural	10,708,481 (-13%)	18,124,458	4,662,437	28,832,939	12,342,764	17,215,274	4,603,744	29,558,038
Respiratory	5,303,862 (-11%)	9,896,044	1,295,750	15,199,906	5,979,056	9,276,817	1,295,638	15,255,873
Endocrine*	2,560,585 (19%)	6,780,460	156,081	9,341,045	2,159,635	7,186,400	124,334	9,346,035
Digestive	1,731,012 (3%)	2,394,730	346,257	4,125,742	1,674,443	2,693,506	346,111	4,367,949
COVID-19	1,004,208 (69%)	165,605	58,208	1,169,813	595,609	574,204	58,211	1,169,813
Drug Poisoning	1,229,766 (5%)	241,136	701,531	1,470,902	1,168,358	304,773	701,529	1,473,131
Transport	915,055 (430%)	19,491	853	934,546	172,795	761,743	852	934,538
Suicide	762,109 (555%)	3,190	662	765,299	116,440	648,857	662	765,297
Alcohol-related	676,440 (18%)	679,715	155,262	1,356,155	571,405	784,750	102,929	1,356,155
Other External	671,615 (-78%)	4,197,101	269,509	4,868,716	2,988,882	1,883,187	269,514	4,872,069
Falls	654,100 (178%)	159,415	161	813,515	235,675	577,837	161	813,512
Homicide	396,530 (1300%)	4,118	275	400,648	28,321	372,327	275	400,648
Uncertain	0	518	0	518	0	0	0	0
Total	56,986,831	57,611,546	20,808,620	114,598,377	56,986,831	58,733,686	20,502,014	115,720,517

*endocrine nutritional and metabolic

Table S3. Changes in causes of death classified in broad disease categories in Entity Axis when using the first listed ICD code rather than the last, in the last line of Part I. There is no change on the Record Axis.

BROAD CATEGORY	Entity Axis count using first ICD-10	Entity Axis count using last ICD-10	Change in count	Percent Change in count
Circulatory	17,770,736	17,793,268	-22,532	0%
Cancer	11,182,712	11,214,012	-31,300	0%
Other Natural	12,342,764	12,386,337	-43,573	0%
Respiratory	5,979,056	5,978,426	630	0%
Endocrine, Nutritional and Metabolic	2,159,635	2,209,113	-49,478	-2%
Digestive	1,674,443	1,675,880	-1,437	0%
COVID-19	595,609	522,512	73,097	14%
Drug Poisoning	1,168,358	1,130,962	37,396	3%
Transport	172,795	637,778	-464,983	-73%
Suicide	116,440	609,437	-492,997	-81%
Alcohol-related	571,405	686,868	-115,463	-17%
Other External	2,988,882	1,451,348	1,537,534	106%
Falls	235,675	386,291	-150,616	-39%
Homicide	28,321	304,599	-276,278	-91%

Table S4. The 30 most frequent ICD-10 codes shown as underlying cause in the Entity Axis replaced with COVID-19 (ICD-10 code U071) as underlying cause in the Record Axis.

ICD-10 code	Count	Cumulative percentage (%)	Description of cause
J189	220,367	43.8	Pneumonia, unspecified
B99	16,742	47.2	Other infectious diseases
A419	13,306	49.8	Sepsis, unspecified organism
J449	12,556	52.3	COPD, unspecified
J960	10,724	54.4	Acute respiratory failure
N179	10,579	56.5	Acute kidney failure
J969	10,470	58.6	Respiratory failure, unspecified
I10	10,300	60.7	Essential hypertension
F03	9,761	62.6	Dementia, unspecified
J80	7,590	64.1	ARDS
I500	6,199	65.4	Congestive heart failure
J129	6,099	66.6	Viral pneumonia, unspecified
I251	6,076	67.8	Atherosclerotic heart disease
E149	5,587	68.9	Diabetes, unspecified
B349	5,085	69.9	Viral infection, unspecified
E119	4,267	70.8	Type 2 diabetes
G309	4,245	71.6	Alzheimer disease
I48	4,170	72.4	Atrial fibrillation
R090	4,032	73.2	Asphyxia
N288	3,907	74.0	Other kidney disorders
I469	3,706	74.7	Cardiac arrest
N189	3,538	75.4	CKD, unspecified
I269	3,402	76.1	Pulmonary embolism
R688	3,351	76.8	Other specified symptoms
N185	3,076	77.4	CKD stage 5
I509	2,958	78.0	Heart failure, unspecified
J159	2,924	78.6	Bacterial pneumonia, unspecified
N19	2,575	79.1	Renal failure, unspecified
J690	2,468	79.6	Aspiration pneumonia
C349	2,112	80.0	Lung cancer

Table S5A. Changes between Record Axis and Entity Axis causes of death classified in broad disease categories when using the first listed ICD code in the last line of Part I of the Entity Axis versus using the last listed ICD code in the last line of Part I of the Entity Axis.

Year	Record Axis count	Entity Axis ICD-10 count with first degenerate Entity Axis code		Entity Axis ICD-10 count with last degenerate Entity Axis code		Entity Axis ICD-10 count with first degenerate Entity Axis 3 character code		Entity Axis ICD-10 count with last degenerate Entity Axis 3 character code		Entity Axis broad categories count with first degenerate Entity Axis code		Entity Axis broad categories count with first degenerate Entity Axis code	
	in 1000s	in 1000s	Entity / Record %	in 1000s	Entity / Record %	in 1000s	Entity / Record	in 1000s	Entity / Record	in 1000s	Entity / Record	in 1000s	Entity / Record
2003	2,448	1,759	71.8	1,733	70.8	1,781	72.8	1,755	71.7	2,131	87.0	2,078	84.9
2004	2,398	1,719	71.7	1,695	70.7	1,740	72.6	1,716	71.6	2,080	86.7	2,027	84.5
2005	2,448	1,762	72.0	1,737	71.0	1,784	72.9	1,759	71.8	2,123	86.7	2,067	84.5
2006	2,426	1,723	71.0	1,699	70.0	1,752	72.2	1,728	71.2	2,097	86.4	2,041	84.1
2007	2,424	1,726	71.2	1,703	70.3	1,749	72.1	1,725	71.2	2,090	86.2	2,031	83.8
2008	2,472	1,743	70.5	1,720	69.6	1,778	71.9	1,754	71.0	2,123	85.9	2,063	83.5
2009	2,437	1,721	70.6	1,698	69.7	1,755	72.0	1,732	71.0	2,094	85.9	2,035	83.5
2010	2,468	1,743	70.6	1,721	69.7	1,776	71.9	1,753	71.0	2,117	85.8	2,057	83.3
2011	2,515	1,765	70.2	1,742	69.3	1,805	71.7	1,781	70.8	2,155	85.7	2,094	83.2
2012	2,543	1,782	70.1	1,759	69.1	1,821	71.6	1,797	70.6	2,174	85.5	2,109	82.9
2013	2,597	1,816	69.9	1,791	69.0	1,860	71.6	1,834	70.6	2,221	85.5	2,155	83.0
2014	2,626	1,839	70.0	1,813	69.0	1,884	71.7	1,856	70.7	2,250	85.7	2,183	83.1
2015	2,713	1,888	69.6	1,860	68.6	1,934	71.3	1,904	70.2	2,319	85.5	2,247	82.8
2016	2,744	1,899	69.2	1,866	68.0	1,943	70.8	1,908	69.5	2,342	85.4	2,265	82.5
2017	2,814	1,921	68.3	1,885	67.0	1,968	70.0	1,930	68.6	2,386	84.8	2,309	82.1
2018	2,839	1,927	67.9	1,890	66.6	1,975	69.6	1,936	68.2	2,403	84.6	2,325	81.9
2019	2,855	1,933	67.7	1,893	66.3	1,981	69.4	1,939	67.9	2,418	84.7	2,337	81.9
2020	3,384	2,234	66.0	2,184	64.5	2,286	67.6	2,233	66.0	2,765	81.7	2,697	79.7
2021	3,464	2,203	63.6	2,149	62.0	2,253	65.0	2,196	63.4	2,747	79.3	2,694	77.8
2022	3,280	2,129	64.9	2,073	63.2	2,182	66.5	2,123	64.7	2,691	82.1	2,608	79.5
2023	3,091	2,031	65.7	1,977	64.0	2,083	67.4	2,026	65.6	2,586	83.7	2,495	80.7
Total 2003-23	56,987	39,261	68.9	38,588	67.7	40,091	70.4	39,385	69.1	48,313	84.8	46,920	82.3
Total 2003-19	43,768	30,663	70.1	30,205	69.0	31,287	71.5	30,807	70.4	37,523	85.7	36,425	83.2
Total 2020-23	13,219	8,598	65.0	8,383	63.4	8,804	66.6	8,578	64.9	10,790	81.6	10,495	79.4
Total 2003-23	56,986,831	39,260,709	68.9	38,588,288	67.7	40,090,579	70.4	39,384,860	69.1	48,313,403	84.8	46,919,850	82.3
Total 2003-19	43,768,050	30,663,135	70.1	30,205,010	69.0	31,286,621	71.5	30,806,674	70.4	37,523,391	85.7	36,425,349	83.2
Total 2020-23	13,218,781	8,597,574	65.0	8,383,278	63.4	8,803,958	66.6	8,578,186	64.9	10,790,012	81.6	10,494,501	79.4

Table S5B. Changes in causes of death classified in broad disease categories in Entity Axis when using the first listed ICD code in the last line of Part I versus using the last listed ICD code in the last line of Part I, showing results excluding COVID (U071).

Year	Record Axis count	Entity Axis ICD-10 count with first degenerate Entity Axis code		Entity Axis broad categories count with first degenerate Entity Axis code		Entity Axis ICD-10 count with first degenerate Entity Axis code; excluding U071		Entity Axis broad categories count with first degenerate Entity Axis code; excluding U071	
	in 1000s	in 1000s	Entity / Record %	in 1000s	Entity / Record %	in 1000s	Entity / Record %	in 1000s	Entity / Record %
2003	2,448	1,759	71.8	2,131	87.0	1,759	71.8	2,131	87.0
2004	2,398	1,719	71.7	2,080	86.7	1,719	71.7	2,080	86.7
2005	2,448	1,762	72.0	2,123	86.7	1,762	72.0	2,123	86.7
2006	2,426	1,723	71.0	2,097	86.4	1,723	71.0	2,097	86.4
2007	2,424	1,726	71.2	2,090	86.2	1,726	71.2	2,090	86.2
2008	2,472	1,743	70.5	2,123	85.9	1,743	70.5	2,123	85.9
2009	2,437	1,721	70.6	2,094	85.9	1,721	70.6	2,094	85.9
2010	2,468	1,743	70.6	2,117	85.8	1,743	70.6	2,117	85.8
2011	2,515	1,765	70.2	2,155	85.7	1,765	70.2	2,155	85.7
2012	2,543	1,782	70.1	2,174	85.5	1,782	70.1	2,174	85.5
2013	2,597	1,816	69.9	2,221	85.5	1,816	69.9	2,221	85.5
2014	2,626	1,839	70.0	2,250	85.7	1,839	70.0	2,250	85.7
2015	2,713	1,888	69.6	2,319	85.5	1,888	69.6	2,319	85.5
2016	2,744	1,899	69.2	2,342	85.4	1,899	69.2	2,342	85.4
2017	2,814	1,921	68.3	2,386	84.8	1,921	68.3	2,386	84.8
2018	2,839	1,927	67.9	2,403	84.6	1,927	67.9	2,403	84.6
2019	2,855	1,933	67.7	2,418	84.7	1,933	67.7	2,418	84.7
2020	3,384	2,234	66.0	2,765	81.7	2,026	66.9	2,556	84.5
2021	3,464	2,203	63.6	2,747	79.3	2,022	66.4	2,566	84.3
2022	3,280	2,129	64.9	2,691	82.1	2,041	66.1	2,603	84.3
2023	3,091	2,031	65.7	2,586	83.7	2,008	66.1	2,563	84.4
Total 2003-23	56,987	39,261	68.9	48,313	84.8	38,759	69.3	47,812	85.4
Total 2003-19	43,768	30,663	70.1	37,523	85.7	30,663	70.1	37,523	85.7
Total 2020-23	13,219	8,598	65.0	10,790	81.6	8,096	66.4	10,288	84.4
Total 2003-23	56,986,831	39,260,709	68.9	48,313,403	84.8	38,759,268	69.3	47,811,656	85.4
Total 2003-19	43,768,050	30,663,135	70.1	37,523,391	85.7	30,663,135	70.1	37,523,391	85.7
Total 2020-23	13,218,781	8,597,574	65.0	10,790,012	81.6	8,096,133	66.4	10,288,265	84.4

Table S6. Top five ICD-10 codes boosting the counts for Other External causes of death when contributing causes are included by weighting.

Count	Cumulative %	ICD-10	Explanation	Count	Cumulative %	ICD-10	Explanation
Under 65 years. Other External is not underlying cause of death				**Under 65 years. Other External is underlying cause of death**			
41026	3.5%	S099	Unspecified injury of head	4656	2.6%	R568	Other and unspecified convulsions
38155	6.8%	T179	Foreign body in respiratory tract, part unspecified	4033	4.8%	S099	Unspecified injury of head
31183	9.5%	T509	Poisoning: Other and unspecified drugs, medicaments and biological substances	2881	6.4%	I469	Cardiac arrest, unspecified
18045	11.0%	S069	Intracranial injury, unspecified	2768	7.9%	T68	Hypothermia
13391	12.1%	I469	Cardiac arrest, unspecified	2638	9.4%	T510	Toxic effect: Ethanol
65 years or over. Other External is not underlying cause of death				**65 years or over. Other External is underlying cause of death**			
125171	15.4%	T179	Foreign body in respiratory tract, part unspecified	7431	0.4%	J969	Respiratory failure, unspecified
67061	19.1%	S720	Fracture of neck of femur	6138	0.7%	R628	Other lack of expected normal physiological development
40565	21.3%	J969	Respiratory failure, unspecified	5065	1.0%	J449	Chronic obstructive pulmonary disease, unspecified
39975	23.5%	J449	Chronic obstructive pulmonary disease, unspecified	4877	1.3%	R13	Dysphagia
33167	25.3%	R13	Dysphagia	4433	1.5%	R688	Other specified general symptoms and signs

Table S7. Comparing the percentage changes in death counts between the Entity and Record axes with those between unweighted and weighted counts.

Disease	Death Counts					Death Counts normalized by W0				
	Record Underlying W0	Entity Underlying	W1	W2	W2A	Rec/W0	Ent/W0	W1/W0	W2/W0	W2A/W0
Circulatory	17,797	17,793	17,960	18,060	17,045	100.0%	100.0%	100.9%	101.5%	95.8%
Cancer	12,576	11,214	9,392	8,546	8,832	100.0%	89.2%	74.7%	68.0%	70.2%
Other Natural	10,708	12,386	13,538	14,424	14,186	100.0%	115.7%	126.4%	134.7%	132.5%
Respiratory	5,304	5,978	6,122	6,287	6,481	100.0%	112.7%	115.4%	118.5%	122.2%
Endocrine*	2,561	2,209	2,653	2,707	3,404	100.0%	86.3%	103.6%	105.7%	132.9%
Digestive	1,731	1,676	1,636	1,565	1,668	100.0%	96.8%	94.5%	90.4%	96.4%
COVID-19	1,004	523	566	370	441	100.0%	52.0%	56.4%	36.9%	43.9%
Drug Poisoning	1,230	1,131	1,140	1,087	988	100.0%	92.0%	92.7%	88.4%	80.3%
Transport	915	638	461	380	437	100.0%	69.7%	50.4%	41.5%	47.8%
Suicide	762	609	382	308	363	100.0%	80.0%	50.1%	40.4%	47.6%
Alcohol-related	676	687	547	512	574	100.0%	101.5%	80.9%	75.7%	84.9%
Other External	672	1,451	2,039	2,376	2,104	100.0%	216.1%	303.4%	353.6%	313.1%
Falls	654	386	351	225	269	100.0%	59.1%	53.7%	34.4%	41.1%
Homicide	397	305	200	140	195	100.0%	76.8%	50.4%	35.3%	49.1%
Correlation coefficient						1.000	1.000	0.969	0.974	0.960
RMS % Change						0.0%	37.9%	62.6%	78.9%	67.7%

*endocrine nutritional and metabolic

Table S8. Slopes of monthly death counts for different diseases, age buckets and weighting schemes. We only include those instances where the R2 for both the linear and logarithmic fit is greater than 0.6.

Disease	Bucket	Scheme	R2	B linear / y %	B log / y %	Disease	Bucket	Scheme	R2	B linear / y %	B log / y %
Circulatory	GE65	W0	0.694	-2.5	-2.4	Drug poisoning	LT65	W2A	0.833	6.8	6.8
Circulatory	GE65	W1	0.637	-2.1	-2.0	Suicide	ALL	W0	0.771	2.0	2.1
Circulatory	GE65	W2A	0.606	-2.0	-1.9	Suicide	ALL	W1	0.765	2.0	2.0
Cancer	ALL	W2	0.637	-0.7	-0.7	Suicide	ALL	W2	0.678	1.7	1.7
Cancer	ALL	W2A	0.646	-0.8	-0.7	Suicide	ALL	W2A	0.769	2.0	2.1
Cancer	GE65	W0	0.908	-1.9	-1.9	Suicide	LT65	W0	0.786	2.1	2.2
Cancer	GE65	W1	0.924	-2.1	-2.0	Suicide	LT65	W1	0.782	2.1	2.2
Cancer	GE65	W2	0.938	-2.3	-2.3	Suicide	LT65	W2	0.710	1.9	1.9
Cancer	GE65	W2A	0.938	-2.3	-2.3	Suicide	LT65	W2A	0.784	2.1	2.1
Cancer	LT65	W0	0.602	-0.9	-0.9	Alcohol-related	ALL	W0	0.856	4.7	4.7
Cancer	LT65	W1	0.762	-1.2	-1.2	Alcohol-related	ALL	W1	0.855	4.5	4.4
Cancer	LT65	W2	0.827	-1.5	-1.6	Alcohol-related	ALL	W2	0.849	4.2	4.2
Cancer	LT65	W2A	0.826	-1.5	-1.6	Alcohol-related	ALL	W2A	0.844	4.4	4.3
Other natural	ALL	W0	0.629	1.9	1.9	Alcohol-related	GE65	W0	0.818	4.3	4.2
Other natural	ALL	W1	0.617	1.8	1.9	Alcohol-related	GE65	W1	0.824	4.2	4.1
Other natural	ALL	W2	0.618	1.9	1.9	Alcohol-related	GE65	W2	0.816	4.0	3.9
Endocrine	LT65	W0	0.642	2.5	2.4	Alcohol-related	GE65	W2A	0.811	4.2	4.1
Endocrine	LT65	W1	0.625	3.0	2.9	Alcohol-related	LT65	W0	0.846	4.5	4.5
Endocrine	LT65	W2	0.609	3.3	3.1	Alcohol-related	LT65	W1	0.844	4.3	4.3
Endocrine	LT65	W2A	0.618	3.1	3.0	Alcohol-related	LT65	W2	0.838	4.1	4.1
Drug poisoning	ALL	W0	0.844	6.7	6.7	Alcohol-related	LT65	W2A	0.835	4.2	4.2
Drug poisoning	ALL	W1	0.84	6.7	6.7	Falls	ALL	W0	0.952	4.3	4.5
Drug poisoning	ALL	W2	0.837	6.7	6.7	Falls	ALL	W1	0.947	4.2	4.4
Drug poisoning	ALL	W2A	0.835	6.7	6.6	Falls	ALL	W2	0.940	3.7	3.8
Drug poisoning	GE65	W0	0.818	8.3	8.3	Falls	ALL	W2A	0.938	3.8	3.9
Drug poisoning	GE65	W1	0.818	8.5	8.6	Falls	GE65	W0	0.854	2.8	3.0
Drug poisoning	GE65	W2	0.816	8.7	8.7	Falls	GE65	W1	0.852	2.7	2.8
Drug poisoning	GE65	W2A	0.82	8.8	8.8	Falls	GE65	W2	0.773	2.0	2.1
Drug poisoning	LT65	W0	0.842	6.8	6.9	Falls	GE65	W2A	0.802	2.2	2.3
Drug poisoning	LT65	W1	0.838	6.9	6.9	Falls	LT65	W0	0.666	1.9	1.9
Drug poisoning	LT65	W2	0.835	6.9	6.8	Falls	LT65	W1	0.680	1.9	1.9

*endocrine nutritional and metabolic

Table S9 Mapping of ICD-10 codes to Broad disease categories as adapted from Wrigley-Field (Wrigley-Field E, Raquib RV, Berry KM, Morris KJ, Stokes AC. Mortality trends among early adults in the United States, 1999-2023. JAMA Network Open 2025:8e2457538).

ICD-10	Broad disease category	First_letter	Description
I169	Circulatory	I	I16.9 (Hypertensive crisis, unspecified)
K85	Digestive	K	K85 (Acute pancreatitis)
U099	COVID-19	U	U09.9 (Post COVID-19 condition)
E244	Alcohol-related	A	E24.4 (Alcohol-induced pseudo-Cushing syndrome)
F10	Alcohol-related	A	F10 (Mental and behavioural disorders due to use of alcohol)
G312	Alcohol-related	A	G31.2 (Degeneration of nervous system due to alcohol)
G621	Alcohol-related	A	G62.1 (Alcoholic polyneuropathy)
G721	Alcohol-related	A	G72.1 (Alcoholic myopathy)
I426	Alcohol-related	A	I42.6 (Alcoholic cardiomyopathy)
K292	Alcohol-related	A	K29.2 (Alcoholic gastritis)
K70	Alcohol-related	A	K70 (Alcoholic liver disease)
K852	Alcohol-related	A	K85.2 (Alcohol-induced acute pancreatitis)
K860	Alcohol-related	A	K86.0 (Alcohol-induced chronic pancreatitis)
R780	Alcohol-related	A	R78.0 (Finding of alcohol in blood)
X45	Alcohol-related	A	X45 (Accidental poisoning by and exposure to alcohol)
X65	Alcohol-related	A	X65 (Intentional self-poisoning by and exposure to alcohol)
Y15	Alcohol-related	A	Y15 (Poisoning by and exposure to alcohol, undetermined intent)
C00-D48	Cancer	C	C00-D48 (Neoplasms)
U071	COVID-19	C	U07.1 (COVID-19)
K00-K14	Digestive	D	K00-K14 (Diseases of oral cavity, salivary glands and jaws)
K20	Digestive	D	K20 (Oesophagitis)
K21	Digestive	D	K21 (Gastro-oesophageal reflux disease)
K22	Digestive	D	K22 (Other diseases of oesophagus)
K25	Digestive	D	K25 (Gastric ulcer)
K26	Digestive	D	K26 (Duodenal ulcer)
K27	Digestive	D	K27 (Peptic ulcer, site unspecified)
K28	Digestive	D	K28 (Gastrojejunal ulcer)
K290	Digestive	D	K29.0 (Acute haemorrhagic gastritis)
K291	Digestive	D	K29.1 (Other acute gastritis)
K293	Digestive	D	K29.3 (Chronic superficial gastritis)
K294	Digestive	D	K29.4 (Chronic atrophic gastritis)
K295	Digestive	D	K29.5 (Chronic gastritis, unspecified)
K296	Digestive	D	K29.6 (Other gastritis)
K297	Digestive	D	K29.7 (Gastritis, unspecified)
K298	Digestive	D	K29.8 (Duodenitis)
K299	Digestive	D	K29.9 (Gastroduodenitis, unspecified)
K30	Digestive	D	K30 (Dyspepsia)
K31	Digestive	D	K31 (Other diseases of stomach and duodenum)
K35-K38	Digestive	D	K35-K38 (Diseases of appendix)
K40-K46	Digestive	D	K40-K46 (Hernia)
K50-K52	Digestive	D	K50-K52 (Noninfective enteritis and colitis)
K55-K63	Digestive	D	K55-K63 (Other diseases of intestines)
K65-K66	Digestive	D	K65-K66 (Diseases of peritoneum)
K71	Digestive	D	K71 (Toxic liver disease)
K72	Digestive	D	K72 (Hepatic failure, not elsewhere classified)
K73	Digestive	D	K73 (Chronic hepatitis, not elsewhere classified)
K74	Digestive	D	K74 (Fibrosis and cirrhosis of liver)
K75	Digestive	D	K75 (Other inflammatory liver diseases)
K76	Digestive	D	K76 (Other diseases of liver)
K80	Digestive	D	K80 (Cholelithiasis)

K81	Digestive	D	K81 (Cholecystitis)
K82	Digestive	D	K82 (Other diseases of gallbladder)
K83	Digestive	D	K83 (Other diseases of biliary tract)
K850	Digestive	D	K85.0 (Idiopathic acute pancreatitis)
K851	Digestive	D	K85.1 (Biliary acute pancreatitis)
K853	Digestive	D	K85.3 (Drug-induced acute pancreatitis)
K858	Digestive	D	K85.8 (Other acute pancreatitis)
K859	Digestive	D	K85.9 (Acute pancreatitis, unspecified)
K861	Digestive	D	K86.1 (Other chronic pancreatitis)
K862	Digestive	D	K86.2 (Cyst of pancreas)
K863	Digestive	D	K86.3 (Pseudocyst of pancreas)
K868	Digestive	D	K86.8 (Other specified diseases of pancreas)
K869	Digestive	D	K86.9 (Disease of pancreas, unspecified)
K90-K92	Digestive	D	K90-K92 (Other diseases of the digestive system)
F11	Drug poisoning	D	F11 (Mental and behavioural disorders due to use of opioids)
F12	Drug poisoning	D	F12 (Mental and behavioural disorders due to use of cannabinoids)
F13	Drug poisoning	D	F13 (Mental and behavioural disorders due to use of sedatives or hypnotics)
F14	Drug poisoning	D	F14 (Mental and behavioural disorders due to use of cocaine)
F15	Drug poisoning	D	F15 (Mental and behavioural disorders due to use of other stimulants, including caffeine)
F16	Drug poisoning	D	F16 (Mental and behavioural disorders due to use of hallucinogens)
F19	Drug poisoning	D	F19 (Mental and behavioural disorders due to multiple drug use and use of other psychoactive substances)
X60	Drug poisoning	D	X60 (Intentional self-poisoning by and exposure to nonopioid analgesics, antipyretics and antirheumatics)
X61	Drug poisoning	D	X61 (Intentional self-poisoning by and exposure to antiepileptic, sedative-hypnotic, antiparkinsonism and psychotropic drugs, not elsewhere classified)
X62	Drug poisoning	D	X62 (Intentional self-poisoning by and exposure to narcotics and psychodysleptics [hallucinogens], not elsewhere classified)
X63	Drug poisoning	D	X63 (Intentional self-poisoning by and exposure to other drugs acting on the autonomic nervous system)
X64	Drug poisoning	D	X64 (Intentional self-poisoning by and exposure to other and unspecified drugs, medicaments and biological substances)
X85	Drug poisoning	D	X85 (Assault by drugs, medicaments and biological substances)
Y10	Drug poisoning	D	Y10 (Poisoning by and exposure to nonopioid analgesics, antipyretics and antirheumatics, undetermined intent)
Y11	Drug poisoning	D	Y11 (Poisoning by and exposure to antiepileptic, sedative-hypnotic, antiparkinsonism and psychotropic drugs, not elsewhere classified, undetermined intent)
Y12	Drug poisoning	D	Y12 (Poisoning by and exposure to narcotics and psychodysleptics [hallucinogens], not elsewhere classified, undetermined intent)
Y13	Drug poisoning	D	Y13 (Poisoning by and exposure to other drugs acting on the autonomic nervous system, undetermined intent)
Y14	Drug poisoning	D	Y14 (Poisoning by and exposure to other and unspecified drugs, medicaments and biological substances, undetermined intent)
X40	Drug poisoning	D	X40 (Accidental poisoning by and exposure to nonopioid analgesics, antipyretics and antirheumatics)
X41	Drug poisoning	D	X41 (Accidental poisoning by and exposure to antiepileptic, sedative-hypnotic, antiparkinsonism and psychotropic drugs, not elsewhere classified)
X42	Drug poisoning	D	X42 (Accidental poisoning by and exposure to narcotics and psychodysleptics [hallucinogens], not elsewhere classified)
X43	Drug poisoning	D	X43 (Accidental poisoning by and exposure to other drugs acting on the autonomic nervous system)
X44	Drug poisoning	D	X44 (Accidental poisoning by and exposure to other and unspecified drugs, medicaments and biological substances)
E00-E07	Endocrine	E	E00-E07 (Disorders of thyroid gland)
E10-E14	Endocrine	E	E10-E14 (Diabetes mellitus)
E15-E16	Endocrine	E	E15-E16 (Other disorders of glucose regulation and pancreatic internal secretion)
E20	Endocrine	E	E20 (Hypoparathyroidism)

E21	Endocrine	E	E21 (Hyperparathyroidism and other disorders of parathyroid gland)
E22	Endocrine	E	E22 (Hyperfunction of pituitary gland)
E23	Endocrine	E	E23 (Hypofunction and other disorders of pituitary gland)
E240	Endocrine	E	E24.0 (Pituitary-dependent Cushing disease)
E241	Endocrine	E	E24.1 (Nelson syndrome)
E242	Endocrine	E	E24.2 (Drug-induced Cushing syndrome)
E243	Endocrine	E	E24.3 (Ectopic ACTH syndrome)
E248	Endocrine	E	E24.8 (Other Cushing syndrome)
E249	Endocrine	E	E24.9 (Cushing syndrome, unspecified)
E25	Endocrine	E	E25 (Adrenogenital disorders)
E26	Endocrine	E	E26 (Hyperaldosteronism)
E27	Endocrine	E	E27 (Other disorders of adrenal gland)
E28	Endocrine	E	E28 (Ovarian dysfunction)
E29	Endocrine	E	E29 (Testicular dysfunction)
E30	Endocrine	E	E30 (Disorders of puberty, not elsewhere classified)
E31	Endocrine	E	E31 (Polyglandular dysfunction)
E32	Endocrine	E	E32 (Diseases of thymus)
E34	Endocrine	E	E34 (Other endocrine disorders)
E40-E46	Endocrine	E	E40-E46 (Malnutrition)
E50-E64	Endocrine	E	E50-E64 (Other nutritional deficiencies)
E65-E68	Endocrine	E	E65-E68 (Obesity and other hyperalimentation)
E70-E88	Endocrine	E	E70-E88 (Metabolic disorders)
X86	Homicide	H	X86 (Assault by corrosive substance)
X87	Homicide	H	X87 (Assault by pesticides)
X88	Homicide	H	X88 (Assault by gases and vapours)
X89	Homicide	H	X89 (Assault by other specified chemicals and noxious substances)
X90	Homicide	H	X90 (Assault by unspecified chemical or noxious substance)
X91	Homicide	H	X91 (Assault by hanging, strangulation and suffocation)
X92	Homicide	H	X92 (Assault by drowning and submersion)
X93	Homicide	H	X93 (Assault by handgun discharge)
X94	Homicide	H	X94 (Assault by rifle, shotgun and larger firearm discharge)
X95	Homicide	H	X95 (Assault by other and unspecified firearm discharge)
X96	Homicide	H	X96 (Assault by explosive material)
X97	Homicide	H	X97 (Assault by smoke, fire and flames)
X98	Homicide	H	X98 (Assault by steam, hot vapours and hot objects)
X99	Homicide	H	X99 (Assault by sharp object)
Y00	Homicide	H	Y00 (Assault by blunt object)
Y01	Homicide	H	Y01 (Assault by pushing from high place)
Y02	Homicide	H	Y02 (Assault by pushing or placing victim before moving object)
Y03	Homicide	H	Y03 (Assault by crashing of motor vehicle)
Y04	Homicide	H	Y04 (Assault by bodily force)
Y05	Homicide	H	Y05 (Sexual assault by bodily force)
Y06	Homicide	H	Y06 (Neglect and abandonment)
Y07	Homicide	H	Y07 (Other maltreatment syndromes)
Y08	Homicide	H	Y08 (Assault by other specified means)
Y09	Homicide	H	Y09 (Assault by unspecified means)
Y871	Homicide	H	Y87.1 (Sequelae of assault)
I10-I15	Circulatory	C	I10-I15 (Hypertensive diseases)
I00-I02	Circulatory	C	I00-I02 (Acute rheumatic fever)
I05-I09	Circulatory	C	I05-I09 (Chronic rheumatic heart diseases)
I20-I25	Circulatory	C	I20-I25 (Ischaemic heart diseases)
I26-I28	Circulatory	C	I26-I28 (Pulmonary heart disease and diseases of pulmonary circulation)
I30	Circulatory	C	I30 (Acute pericarditis)
I31	Circulatory	C	I31 (Other diseases of pericardium)
I33	Circulatory	C	I33 (Acute and subacute endocarditis)

I34	Circulatory	C	I34 (Nonrheumatic mitral valve disorders)
I35	Circulatory	C	I35 (Nonrheumatic aortic valve disorders)
I36	Circulatory	C	I36 (Nonrheumatic tricuspid valve disorders)
I37	Circulatory	C	I37 (Pulmonary valve disorders)
I38	Circulatory	C	I38 (Endocarditis, valve unspecified)
I40	Circulatory	C	I40 (Acute myocarditis)
I420	Circulatory	C	I42.0 (Dilated cardiomyopathy)
I421	Circulatory	C	I42.1 (Obstructive hypertrophic cardiomyopathy)
I422	Circulatory	C	I42.2 (Other hypertrophic cardiomyopathy)
I423	Circulatory	C	I42.3 (Endomyocardial (eosinophilic) disease)
I424	Circulatory	C	I42.4 (Endocardial fibroelastosis)
I425	Circulatory	C	I42.5 (Other restrictive cardiomyopathy)
I427	Circulatory	C	I42.7 (Cardiomyopathy due to drugs and other external agents)
I428	Circulatory	C	I42.8 (Other cardiomyopathies)
I429	Circulatory	C	I42.9 (Cardiomyopathy, unspecified)
I44	Circulatory	C	I44 (Atrioventricular and left bundle-branch block)
I45	Circulatory	C	I45 (Other conduction disorders)
I46	Circulatory	C	I46 (Cardiac arrest)
I47	Circulatory	C	I47 (Paroxysmal tachycardia)
I48	Circulatory	C	I48 (Atrial fibrillation and flutter)
I49	Circulatory	C	I49 (Other cardiac arrhythmias)
I50	Circulatory	C	I50 (Heart failure)
I51	Circulatory	C	I51 (Complications and ill-defined descriptions of heart disease)
I60-I69	Circulatory	C	I60-I69 (Cerebrovascular diseases)
I70-I78	Circulatory	C	I70-I78 (Diseases of arteries, arterioles and capillaries)
I80-I89	Circulatory	C	I80-I89 (Diseases of veins, lymphatic vessels and lymph nodes, not elsewhere classified)
I95-I99	Circulatory	C	I95-I99 (Other and unspecified disorders of the circulatory system)
B20-B24	Other natural	O	B20-B24 (Human immunodeficiency virus [HIV] disease)
F01-F09	Other natural	O	F01-F09 (Organic, including symptomatic, mental disorders)
F17	Other natural	O	F17 (Mental and behavioural disorders due to use of tobacco)
F18	Other natural	O	F18 (Mental and behavioural disorders due to use of volatile solvents)
F20-F29	Other natural	O	F20-F29 (Schizophrenia, schizotypal and delusional disorders)
F30-F39	Other natural	O	F30-F39 (Mood [affective] disorders)
F40-F48	Other natural	O	F40-F48 (Neurotic, stress-related and somatoform disorders)
F50-F59	Other natural	O	F50-F59 (Behavioural syndromes associated with physiological disturbances and physical factors)
F60-F69	Other natural	O	F60-F69 (Disorders of adult personality and behaviour)
F70-F79	Other natural	O	F70-F79 (Mental retardation)
F80-F89	Other natural	O	F80-F89 (Disorders of psychological development)
F90-F98	Other natural	O	F90-F98 (Behavioural and emotional disorders with onset usually occurring in childhood and adolescence)
F99-F99	Other natural	O	F99-F99 (Unspecified mental disorder)
A00-A09	Other natural	O	A00-A09 (Intestinal infectious diseases)
A16-A19	Other natural	O	A16-A19 (Tuberculosis)
A20-A28	Other natural	O	A20-A28 (Certain zoonotic bacterial diseases)
A30-A49	Other natural	O	A30-A49 (Other bacterial diseases)
A50-A64	Other natural	O	A50-A64 (Infections with a predominantly sexual mode of transmission)
A65-A69	Other natural	O	A65-A69 (Other spirochaetal diseases)
A70-A74	Other natural	O	A70-A74 (Other diseases caused by chlamydiae)
A75-A79	Other natural	O	A75-A79 (Rickettsioses)
A80-A89	Other natural	O	A80-A89 (Viral infections of the central nervous system)
A90-A99	Other natural	O	A90-A99 (Arthropod-borne viral fevers and viral haemorrhagic fevers)
B00-B09	Other natural	O	B00-B09 (Viral infections characterized by skin and mucous membrane lesions)
B15-B19	Other natural	O	B15-B19 (Viral hepatitis)
B25-B34	Other natural	O	B25-B34 (Other viral diseases)

B35-B49	Other natural	O	B35-B49 (Mycoses)
B50-B64	Other natural	O	B50-B64 (Protozoal diseases)
B65-B83	Other natural	O	B65-B83 (Helminthiases)
B85-B89	Other natural	O	B85-B89 (Pediculosis, acariasis and other infestations)
B90-B94	Other natural	O	B90-B94 (Sequelae of infectious and parasitic diseases)
B99-B99	Other natural	O	B99-B99 (Other infectious diseases)
D50-D89	Other natural	O	D50-D89 (Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism)
G00-G09	Other natural	O	G00-G09 (Inflammatory diseases of the central nervous system)
G10-G14	Other natural	O	G10-G14 (Systemic atrophies primarily affecting the central nervous system)
G20-G25	Other natural	O	G20-G25 (Extrapyramidal and movement disorders)
G30	Other natural	O	G30 (Alzheimer disease)
G310	Other natural	O	G31.0 (Circumscribed brain atrophy)
G311	Other natural	O	G31.1 (Senile degeneration of brain, not elsewhere classified)
G318	Other natural	O	G31.8 (Other specified degenerative diseases of nervous system)
G319	Other natural	O	G31.9 (Degenerative disease of nervous system, unspecified)
G35-G37	Other natural	O	G35-G37 (Demyelinating diseases of the central nervous system)
G40-G47	Other natural	O	G40-G47 (Episodic and paroxysmal disorders)
G50-G58	Other natural	O	G50-G58 (Nerve, nerve root and plexus disorders)
G60	Other natural	O	G60 (Hereditary and idiopathic neuropathy)
G61	Other natural	O	G61 (Inflammatory polyneuropathy)
G620	Other natural	O	G62.0 (Drug-induced polyneuropathy)
G622	Other natural	O	G62.2 (Polyneuropathy due to other toxic agents)
G628	Other natural	O	G62.8 (Other specified polyneuropathies)
G629	Other natural	O	G62.9 (Polyneuropathy, unspecified)
G64	Other natural	O	G64 (Other disorders of peripheral nervous system)
G70	Other natural	O	G70 (Myasthenia gravis and other myoneural disorders)
G71	Other natural	O	G71 (Primary disorders of muscles)
G720	Other natural	O	G72.0 (Drug-induced myopathy)
G722	Other natural	O	G72.2 (Myopathy due to other toxic agents)
G723	Other natural	O	G72.3 (Periodic paralysis)
G724	Other natural	O	G72.4 (Inflammatory myopathy, not elsewhere classified)
G728	Other natural	O	G72.8 (Other specified myopathies)
G729	Other natural	O	G72.9 (Myopathy, unspecified)
G80-G83	Other natural	O	G80-G83 (Cerebral palsy and other paralytic syndromes)
G90-G98	Other natural	O	G90-G98 (Other disorders of the nervous system)
H00-H57	Other natural	O	H00-H57 (Diseases of the eye and adnexa)
H60-H93	Other natural	O	H60-H93 (Diseases of the ear and mastoid process)
L00-L98	Other natural	O	L00-L98 (Diseases of the skin and subcutaneous tissue)
M00-M99	Other natural	O	M00-M99 (Diseases of the musculoskeletal system and connective tissue)
N00-N98	Other natural	O	N00-N98 (Diseases of the genitourinary system)
O00-O99	Other natural	O	O00-O99 (Pregnancy, childbirth and the puerperium)
P00-P96	Other natural	O	P00-P96 (Certain conditions originating in the perinatal period)
Q00-Q99	Other natural	O	Q00-Q99 (Congenital malformations, deformations and chromosomal abnormalities)
R00-R09	Other natural	O	R00-R09 (Symptoms and signs involving the circulatory and respiratory systems)
R10-R19	Other natural	O	R10-R19 (Symptoms and signs involving the digestive system and abdomen)
R20-R23	Other natural	O	R20-R23 (Symptoms and signs involving the skin and subcutaneous tissue)
R25-R29	Other natural	O	R25-R29 (Symptoms and signs involving the nervous and musculoskeletal systems)
R30-R39	Other natural	O	R30-R39 (Symptoms and signs involving the urinary system)
R40-R46	Other natural	O	R40-R46 (Symptoms and signs involving cognition, perception, emotional state and behaviour)
R47-R49	Other natural	O	R47-R49 (Symptoms and signs involving speech and voice)
R50-R68	Other natural	O	R50-R68 (General symptoms and signs)
R70	Other natural	O	R70 (Elevated erythrocyte sedimentation rate and abnormality of plasma viscosity)
R71	Other natural	O	R71 (Abnormality of red blood cells)

R72	Other natural	O	R72 (Abnormality of white blood cells, not elsewhere classified)
R73	Other natural	O	R73 (Elevated blood glucose level)
R74	Other natural	O	R74 (Abnormal serum enzyme levels)
R75	Other natural	O	R75 (Laboratory evidence of human immunodeficiency virus [HIV])
R76	Other natural	O	R76 (Other abnormal immunological findings in serum)
R77	Other natural	O	R77 (Other abnormalities of plasma proteins)
R781	Other natural	O	R78.1 (Finding of opiate drug in blood)
R782	Other natural	O	R78.2 (Finding of cocaine in blood)
R783	Other natural	O	R78.3 (Finding of hallucinogen in blood)
R784	Other natural	O	R78.4 (Finding of other drugs of addictive potential in blood)
R785	Other natural	O	R78.5 (Finding of psychotropic drug in blood)
R786	Other natural	O	R78.6 (Finding of steroid agent in blood)
R787	Other natural	O	R78.7 (Finding of abnormal level of heavy metals in blood)
R788	Other natural	O	R78.8 (Finding of other specified substances, not normally found in blood)
R789	Other natural	O	R78.9 (Finding of unspecified substance, not normally found in blood)
R79	Other natural	O	R79 (Other abnormal findings of blood chemistry)
R80-R82	Other natural	O	R80-R82 (Abnormal findings on examination of urine, without diagnosis)
R83-R89	Other natural	O	R83-R89 (Abnormal findings on examination of other body fluids, substances and tissues, without diagnosis)
R90-R94	Other natural	O	R90-R94 (Abnormal findings on diagnostic imaging and in function studies, without diagnosis)
R95-R99	Other natural	O	R95-R99 (Ill-defined and unknown causes of mortality)
U04	Other natural	O	U04 (Severe acute respiratory syndrome [SARS])
U070	Other natural	O	U07.0 (Vaping related disorder)
J00-J98	Respiratory	R	J00-J98 (Diseases of the respiratory system)
X66	Suicide	S	X66 (Intentional self-poisoning by and exposure to organic solvents and halogenated hydrocarbons and their vapours)
X67	Suicide	S	X67 (Intentional self-poisoning by and exposure to other gases and vapours)
X68	Suicide	S	X68 (Intentional self-poisoning by and exposure to pesticides)
X69	Suicide	S	X69 (Intentional self-poisoning by and exposure to other and unspecified chemicals and noxious substances)
X70	Suicide	S	X70 (Intentional self-harm by hanging, strangulation and suffocation)
X71	Suicide	S	X71 (Intentional self-harm by drowning and submersion)
X72	Suicide	S	X72 (Intentional self-harm by handgun discharge)
X73	Suicide	S	X73 (Intentional self-harm by rifle, shotgun and larger firearm discharge)
X74	Suicide	S	X74 (Intentional self-harm by other and unspecified firearm discharge)
X75	Suicide	S	X75 (Intentional self-harm by explosive material)
X76	Suicide	S	X76 (Intentional self-harm by smoke, fire and flames)
X77	Suicide	S	X77 (Intentional self-harm by steam, hot vapours and hot objects)
X78	Suicide	S	X78 (Intentional self-harm by sharp object)
X79	Suicide	S	X79 (Intentional self-harm by blunt object)
X80	Suicide	S	X80 (Intentional self-harm by jumping from a high place)
X81	Suicide	S	X81 (Intentional self-harm by jumping or lying before moving object)
X82	Suicide	S	X82 (Intentional self-harm by crashing of motor vehicle)
X83	Suicide	S	X83 (Intentional self-harm by other specified means)
X84	Suicide	S	X84 (Intentional self-harm by unspecified means)
Y870	Suicide	S	Y87.0 (Sequelae of intentional self-harm)
U01	Other external	O	U01 (Terrorism - Assault (homicide))
U02	Other external	O	U02 (Sequelae of terrorism)
U03	Other external	O	U03 (Terrorism Intentional (Suicide))
W00-W19	Falls	F	W00-W19 (Falls)
W20-W49	Other external	O	W20-W49 (Exposure to inanimate mechanical forces)
W50-W64	Other external	O	W50-W64 (Exposure to animate mechanical forces)
W65-W74	Other external	O	W65-W74 (Accidental drowning and submersion)
W75-W84	Other external	O	W75-W84 (Other accidental threats to breathing)

W85-W99	Other external	O	W85-W99 (Exposure to electric current, radiation and extreme ambient air temperature and pressure)
X00-X09	Other external	O	X00-X09 (Exposure to smoke, fire and flames)
X10-X19	Other external	O	X10-X19 (Contact with heat and hot substances)
X20-X29	Other external	O	X20-X29 (Contact with venomous animals and plants)
X30-X39	Other external	O	X30-X39 (Exposure to forces of nature)
X46	Other external	O	X46 (Accidental poisoning by and exposure to organic solvents and halogenated hydrocarbons and their vapours)
X47	Other external	O	X47 (Accidental poisoning by and exposure to other gases and vapours)
X48	Other external	O	X48 (Accidental poisoning by and exposure to pesticides)
X49	Other external	O	X49 (Accidental poisoning by and exposure to other and unspecified chemicals and noxious substances)
X50-X57	Other external	O	X50-X57 (Overexertion, travel and privation)
Y16	Other external	O	Y16 (Poisoning by and exposure to organic solvents and halogenated hydrocarbons and their vapours, undetermined intent)
Y17	Other external	O	Y17 (Poisoning by and exposure to other gases and vapours, undetermined intent)
Y18	Other external	O	Y18 (Poisoning by and exposure to pesticides, undetermined intent)
Y19	Other external	O	Y19 (Poisoning by and exposure to other and unspecified chemicals and noxious substances, undetermined intent)
Y20	Other external	O	Y20 (Hanging, strangulation and suffocation, undetermined intent)
Y21	Other external	O	Y21 (Drowning and submersion, undetermined intent)
Y22	Other external	O	Y22 (Handgun discharge, undetermined intent)
Y23	Other external	O	Y23 (Rifle, shotgun and larger firearm discharge, undetermined intent)
Y24	Other external	O	Y24 (Other and unspecified firearm discharge, undetermined intent)
Y25	Other external	O	Y25 (Contact with explosive material, undetermined intent)
Y26	Other external	O	Y26 (Exposure to smoke, fire and flames, undetermined intent)
Y27	Other external	O	Y27 (Contact with steam, hot vapours and hot objects, undetermined intent)
Y28	Other external	O	Y28 (Contact with sharp object, undetermined intent)
Y29	Other external	O	Y29 (Contact with blunt object, undetermined intent)
Y30	Other external	O	Y30 (Falling, jumping or pushed from a high place, undetermined intent)
Y31	Other external	O	Y31 (Falling, lying or running before or into moving object, undetermined intent)
Y32	Other external	O	Y32 (Crashing of motor vehicle, undetermined intent)
Y33	Other external	O	Y33 (Other specified events, undetermined intent)
Y34	Other external	O	Y34 (Unspecified event, undetermined intent)
Y86	Other external	O	Y86 (Sequelae of other accidents)
Y872	Other external	O	Y87.2 (Sequelae of events of undetermined intent)
Y88	Other external	O	Y88 (Sequelae with surgical and medical care as external cause)
Y89	Other external	O	Y89 (Sequelae of other external causes)
V01-V99	Transport	T	V01-V99 (Transport accidents)
Y85	Transport	T	Y85 (Sequelae of transport accidents)
T36-T50	Drug poisoning	T	T36-T50 (Poisoning by drugs and biological substances)
S00-S99	Other external	S	S00-S99 (Injuries to specific body regions)
T00-T35	Other external	T	T00-T35 (Injuries involving multiple/unspecified body regions and burns)
T51-T65	Other external	T	T51-T65 (Toxic effects of substances chiefly nonmedicinal as to source)
T66-T78	Other external	T	T66-T78 (Other and unspecified effects of external causes)
T79	Other external	T	T79 (Certain early complications of trauma)
T80-T89	Other external	T	T80-T89 (Complications of surgical and medical care NEC)
T90-T99	Other external	T	T90-T99 (Sequelae of injuries and other external causes)
V00	Transport	V	V00 (Pedestrian conveyance accident)
X58	Other external	X	X58 (Exposure to other specified factors)
X59	Other external	X	X59 (Exposure to unspecified factor)
Y35-Y39	Other external	Y	Y35-Y39 (Legal intervention and operations of war)
Y40-Y59	Other external	Y	Y40-Y59 (Drugs and biological substances causing adverse effects in therapeutic use)
Y60-Y69	Other external	Y	Y60-Y69 (Misadventures to patients during surgical and medical care)
Y70-Y82	Other external	Y	Y70-Y82 (Medical devices associated with adverse incidents)
Y83-Y84	Other external	Y	Y83-Y84 (Surgical and other medical procedures as cause of abnormal reaction)

| Y90-Y99 | Other external | Y | Y90-Y99 (Supplementary factors related to causes of morbidity and mortality) |

Supplementary Figures

Supplementary Figure S1. Showing how the percent changes of the broad disease category counts depend on the source of data. Along the x-axis we show data from the Record Axis. Along the y-axis we show data that combines the underlying cause-of-death from the Record Axis with the contributing causes from Part II of the Entity Axis.

Supplementary Figure S2. Increase in cause-of-death coding differences between death certifier-assigned and NCHS/ACME-assigned underlying cause of death over time and especially during the early years of the COVID-19 pandemic. Data shown for ICD-10 coding and broad disease categories with and excluding the key COVID-19 code. This data is shown in Table S5B.

Supplementary Figure S3 (LT65W1). Population-normalized monthly deaths across 14 broad categories during 2003-2023 for non-elderly people (less than 65 years old). The panels are in different scale. Light line: unweighted W0. Continuous thick line: W1 weights. Disease categories are grouped by fixed Y-axis scales to allow visual comparison across panels. The top band (COVID-19, Circulatory, Cancer, Other natural, Respiratory) uses 0-26,000 deaths/month; the middle band (Endocrine, Digestive, Drug poisoning, Other external) uses 0-12,000 deaths/month; the bottom band (Transport, Suicide, Alcohol-related, Falls, Homicide) uses 0-6,000 deaths/month. All values are population-scaled to 2023 levels.

Supplementary Figure S4 (GE65W1). Population-normalized monthly deaths across 14 broad categories during 2003-2023 for elderly people (age 65 years and over). The panels are in different scale. Light line: unweighted W0. Continuous thick line: W1 weights. Disease categories are grouped by fixed Y-axis scales to allow visual comparison across panels. The top band (COVID-19, Circulatory, Cancer, Other natural, Respiratory) uses 0-130,000 deaths/month; the middle band (Endocrine, Digestive, Drug poisoning, Other external) uses 0-25,000 deaths/month; the bottom band (Transport, Suicide, Alcohol-related, Falls, Homicide) uses 0-6,000 deaths/month.

Supplementary Figure S5 (ALLW2). Population-normalized monthly deaths across 14 broad categories during 2003-2023. The panels are in different scale. Light line: unweighted W0. Continuous thick line: W2 weights. Disease categories are grouped by fixed Y-axis scales to allow visual comparison across panels. The

top band (COVID-19, Circulatory, Cancer, Other natural, Respiratory) uses 0-130,000 deaths/month; the middle band (Endocrine, Digestive, Drug poisoning, Other external) uses 0-25,000 deaths/month; the bottom band (Transport, Suicide, Alcohol-related, Falls, Homicide) uses 0-6,000 deaths/month. All values are population-scaled to 2023 levels.

Supplementary Figure S6 (ALLW2A). Population-normalized monthly deaths across 14 broad categories during 2003-2023. The panels are in different scale. Light line: unweighted W0. Continuous thick line: W2A weights. Disease categories are grouped by fixed Y-axis scales to allow visual comparison across panels. The top band (COVID-19, Circulatory, Cancer, Other natural, Respiratory) uses 0-130,000 deaths/month; the middle band (Endocrine, Digestive, Drug poisoning, Other external) uses 0-25,000 deaths/month; the bottom band (Transport, Suicide, Alcohol-related, Falls, Homicide) uses 0-6,000 deaths/month. All values are population-scaled to 2023 levels.

Supplementary Figure S7 (LT65W2). Population-normalized monthly deaths across 14 broad categories during 2003-2023 for non-elderly people (less than 65 years old). The panels are in different scale. Light line: unweighted W0. Continuous thick line: W2 weights. Disease categories are grouped by fixed Y-axis scales to allow visual comparison across panels. The top band (COVID-19, Circulatory, Cancer, Other natural, Respiratory) uses 0-26,000 deaths/month; the middle band (Endocrine, Digestive, Drug poisoning, Other external) uses 0-12,000 deaths/month; the bottom band (Transport, Suicide, Alcohol-related, Falls, Homicide) uses 0-6,000 deaths/month. All values are population-scaled to 2023 levels.

Supplementary Figure S8 (LT65W2A). Population-normalized monthly deaths across 14 broad categories during 2003-2023 for non-elderly people (less than 65 years old). The panels are in different scale. Light line: unweighted W0. Continuous thick line: W2A weights. Disease categories are grouped by fixed Y-axis scales to allow visual comparison across panels. The top band (COVID-19, Circulatory, Cancer, Other natural, Respiratory) uses 0-26,000 deaths/month; the middle band (Endocrine, Digestive, Drug poisoning, Other external) uses 0-12,000 deaths/month; the bottom band (Transport, Suicide, Alcohol-related, Falls, Homicide) uses 0-6,000 deaths/month. All values are population-scaled to 2023 levels.

Supplementary Figure S9 (GE65W2). Population-normalized monthly deaths across 14 broad categories during 2003-2023 for elderly people (age 65 years and over). The panels are in different scale. Light line: unweighted W0. Continuous thick line: W2 weights.   Disease categories are grouped by fixed Y-axis scales to allow visual comparison across panels. The top band (COVID-19, Circulatory, Cancer, Other natural, Respiratory) uses 0-130,000 deaths/month; the middle band (Endocrine, Digestive, Drug poisoning, Other external) uses 0-25,000 deaths/month; the bottom band (Transport, Suicide, Alcohol-related, Falls, Homicide) uses 0-6,000 deaths/month. All values are population-scaled to 2023 levels.

Supplementary Figure S10 (GE65W2A). Population-normalized monthly deaths across 14 broad categories during 2003-2023 for elderly people (age 65 years and over). The panels are in different scale. Light line: unweighted W0. Continuous thick line: W2A weights.   Disease categories are grouped by fixed Y-axis scales to allow visual comparison across panels. The top band (COVID-19, Circulatory, Cancer, Other natural, Respiratory) uses 0-130,000 deaths/month; the middle band (Endocrine, Digestive, Drug poisoning, Other external) uses 0-25,000 deaths/month; the bottom band (Transport, Suicide, Alcohol-related, Falls, Homicide) uses 0-6,000 deaths/month. All values are population-scaled to 2023 levels.

Supplementary Figure S11. Linear and logarithmic fits to the population-normalized monthly deaths for the three age buckets (ALL, LT65 for people under 65 and GE65 for elderly people aged 65 years and over) across 14 broad categories during 2003-2023. Plots with unacceptable fits ($R2 < 0.6$, CC <. 0.77) have red plot boundaries. They are the majority occurring in 26 out of 42 cases. The other 16 plots show clear linear change over 21 years, with the possible exception of Drug poisoning, which grows somewhat exponentially.

Data 2003 to 2023. All years

$y = 0.4405x - 0.1387$
$R^2 = 0.8093$

Data 2020 to 2023. Pandemic years

$y = 0.4679x - 0.9536$
$R^2 = 0.8235$