1   **SGLT2 inhibitor use in type 2 diabetes in England: a population-based cohort**
2   **study of uptake of NICE guidance**
3

Dr Patrick Muller PhD,[1] Technical Adviser, ORCID: 0000-0002-6824-578X

Dr Jonathan Wray PhD,[1] Senior Analyst

Muksitur Rahman MSc,[1] Health Economist

James Hawkins MSc,[1] Health Economic Adviser

Dr Chirag Bakhai MBA,[2,3] GP and National Clinical Adviser for Diabetes, ORCID: 0009-
0008-4708-3305

Dr Daniel J Cuthbertson PhD,[4] Professor of Medicine, ORCID: 0000-0002-6128-0822

Dr Robert Willans PhD,[1] Technical Adviser

Dr Eleanor Yelland PhD,[1] Scientific Adviser, ORCID: 0000-0003-0148-8530

Shaun Rowark MSc,[1] Associate Director of Healthcare Data Analytics

Magdalena Watras MPharm,[1] Pharmacist Clinical Adviser

Dr Luke Sheridan Rains PhD,[1] Senior Technical Adviser, ORCID: 0000-0003-4905-4168

Dr Amanda I Adler PhD,[5] Professor of Diabetic Medicine and Health Policy, ORCID: 0000-
0002-2797-3385

Dr Lesley Owen PhD,[1] Health Economic Adviser

1. National Institute for Health and Care Excellence.
2. NHS Bedfordshire Luton and Milton Keynes Integrated Care Board.
3. NHS England.
4. Obesity and Endocrinology Research Group, Institute of Ageing and Chronic
Diseases, University of Liverpool.
5. Oxford Centre for Diabetes Endocrinology and Metabolism, University of Oxford.

Corresponding Author

Patrick Muller

National Institute for Health and Care Excellence, 2nd Floor 2 Redman Place,
London, E20 1JQ.

Email: **Patrick.muller@nice.org.uk**

Word count

Abstract: 315

Text: 4,342

Summary boxes: 269

ABSTRACT

Objectives An update to the NICE Type 2 diabetes (T2DM) guideline in February 2022 recommended an SGLT2 inhibitor be offered to people with cardiovascular disease (CVD) or heart failure (HF) as comorbidities and considered for people at high CVD risk. We report uptake of this guideline in England 18 months after its publication.

Design Observational cohort study.

Setting General practices contributing to the Clinical Practice Research Data Link, linked to hospital admission records.

Participants 587,826 people aged over 18 years with T2DM on 1st September 2023, stratified according to their CVD category (CVD only; HF only; CVD and HF; high CVD risk score; low CVD risk score) and chronic kidney disease (CKD) status, and further by age, gender, ethnicity, deprivation, and T2DM diagnosis duration.

Main outcome measures Percentage of patients with a current SGLT2 inhibitor prescription; odds ratios for association between patient characteristics and a current prescription.

Results In people with T2DM, the percentage with a current SGLT2 inhibitor prescription was 19.5% for people with CVD, 29.4% for people with HF, 30.5% for people with both CVD and HF, and 19.9% and 20.2% respectively for people at high and low CVD risk. In age-stratified analyses, uptake ordered from lowest to highest was as follows: low CVD risk score, high CVD risk score, CVD only, HF only, CVD and HF. In models adjusted for clinical and patient characteristics uptake was lower in people aged >60, women, Black people, and people living in areas of higher deprivation.

Conclusions Whilst prescribing of SGLT2 inhibitors continues to rise in England, an opportunity remains to increase uptake and to reduce inequalities in people with T2DM in 2026. We report inequalities by ethnicity and deprivation, and lower uptake for people with CVD without HF than people with HF, despite an equal guideline recommendation for these two groups. Additional evidence is needed on the effectiveness of SGLT2 inhibitors in frailer populations.

What is already known on this topic?

- In 2020 approximately 10% of people with type 2 diabetes (T2DM) and cardiovascular disease (CVD) and 14% of people with T2DM but without CVD in England had a current SGLT2 inhibitor prescription.

- In February 2022 NICE recommended that an SGLT2 inhibitor should be offered to people with T2DM with heart failure or CVD, and considered for people with T2DM at high risk of CVD; network meta-analyses have found 10% to 40% lower odds of cardiovascular mortality with treatment in these groups.

- Uptake of NICE guidelines in general practice has historically been variable, although higher when accompanied by pay-for-performance schemes such as the Quality and Outcomes Framework.

What this study adds

- By September 2023 the percentage of people with T2DM with a current SGLT2 inhibitor prescription had reached 19.5% in those with CVD as a comorbidity, 30.5% in those with heart failure, and 19.9% in those at high risk of CVD.

- Women, people of Black ethnicity, and people living in areas of high deprivation had lower odds of a current prescription in analyses adjusted for age, gender, cardiovascular comorbidity, and renal function.

How might these results change the focus of research or clinical practice?

- The results highlight the need for ongoing surveillance of uptake of NICE-recommended treatments for T2DM, and consideration of actions to address barriers to uptake. This is particularly important in the context of broader eligibility for SGLT2 inhibitor treatment in type 2 diabetes in England from 2026.

- These results support the development of initiatives and quality improvement programmes to improve evidence-based prescribing and address inequalities between clinical and demographic subgroups.

INTRODUCTION

The prevalence of type 2 diabetes (T2DM) has increased in England in recent decades, affecting approximately 7% of adults by March 2024.[1] The percentage of people with T2DM who are living with hypertension, renal disease, and cardiovascular disease as comorbidities has also risen in the same period.[2] Goals for treating T2DM have evolved from focusing on managing glycaemia to including primary and secondary prevention of cardiovascular and renal disease. Recent trial evidence on the cardiorenal benefits of SGLT2 inhibitors in people with T2DM has led to their addition to treatment strategies to achieve these goals.[3-5]

Between 2014 and 2023, the National Institute for Health and Care Excellence (NICE) in the UK published a range of guidance on SGLT2 inhibitors (Figure 1). In 2022, following Technology Appraisals (TAs) recommending SGLT2 inhibitor therapies for escalation of glucose-lowering treatment in T2DM, for symptomatic heart failure, and for chronic kidney disease (CKD), NICE updated its T2DM guideline.[6] The update expanded the T2DM population to be offered SGLT2 inhibitor therapy, recommending that an SGLT2 inhibitor with proven cardiovascular benefit be offered for people with heart failure or CVD as a comorbidity, and considered for people at high risk of CVD. The change was informed by network meta analyses indicating 10-40% lower odds of cardiovascular mortality associated with taking metformin and an SGLT2 inhibitor compared to metformin alone.[7]

NICE prioritises evaluation of the uptake of guidelines, with additional goals of supporting reductions in health inequalities according to protected characteristics and socioeconomic status.[8] The increasing availability of routinely collected electronic health and prescription records is facilitating evaluations of uptake and inequalities in uptake of medicines recommended by NICE in the populations in scope of guidance,[9] complementing the national surveillance of volumes of medicines prescribed in the NHS Innovation Scorecard.[10]

In the present study, we report variation in the percentage of people with T2DM who had a current SGLT2 inhibitor prescription on 1st September 2023 according to patients' comorbidity status, in a large and representative sample of English GP practices. We assess the evidence for inequalities in uptake according to age,

gender, deprivation, and ethnicity, separately for patients stratified according to their CVD risk (with CVD; with HF; with CVD and HF; high CVD risk score; low CVD risk score;) and by whether they have CKD. We consider the clinical and organisational factors that may explain the patterns in uptake of NICE guidance we report. Finally, we compare our findings to those in other high income countries, and discuss the policy implications of our results with reference the most recent update to the NICE T2DM guideline in 2026.

METHODS

Data sources

The cohort and outcomes for analysis were extracted from CPRD Aurum.[11] CPRD Aurum includes recorded symptoms, diagnoses, test results, and medication data for people at participating GP practices from 1995 to present, with approximately 24% coverage of the UK population by December 2023.[12] Patients in CPRD Aurum are broadly representative of the wider population with respect to their geographic distribution, deprivation, age, gender, and ethnicity.[13 14]

CPRD records of patients in England were linked to their Hospital Episode Statistics (HES) records to retrieve diagnoses and procedures recorded in NHS hospital inpatient admissions (with linkage up to March 2021 only),[15] and to their Index of Multiple Deprivation (IMD) quintile, categorised from 5 (least deprived) to 1 (most deprived). The IMD ranks small areas in England from most to least deprived using a single score based on 7 components of overall deprivation.[16] Linkages were done using an algorithm managed by NHS England.[17]

Lists of codes from the SNOMED (for CPRD) and ICD-10 and OPCS-4 (for HES) libraries were collated to select demographics, comorbidities, risk factors, and current medicines from the CPRD-HES dataset. Where relevant code lists were available from the General Practice Domain Set maintained by NHS England,[18] the Open Code Lists website maintained by the Bennett Institute,[19] or the CVDPrevent audit,[20] these were used. Otherwise, code lists were taken from the literature or created using the World Health Organisation ICD-10 codes library,[21] or free-text searches in the CPRD code browser. Each code list was reviewed at NICE by a

health economist, a T2DM guideline committee member, or both. Each code list with
details of its origin is available online (Appendix 1).

Cohort selection, stratification, exposure and outcome definitions

Patients were selected for the cross-sectional analysis of treatment uptake on 1
September 2023 (the "index date") through identifying patients with any record of
T2DM and no other form of diabetes in CPRD between 1st January 2000 and the
index date, and retaining those whose records could be linked to both HES and IMD
datasets, who were judged by CPRD to have a 'research-quality' record, and who
were aged 18 or over on the index date.[22] Patients were excluded if they died or de-
registered prior to the index date, or if they had less than one year registration on the
index date.

The selected people were categorised into mutually exclusive groups of CVD without
HF, HF without CVD, CVD with HF, high CVD risk score (QRISK2 score >10%[23], in
line with the standard UK threshold for treatment) without CVD or HF and low CVD
risk score (QRISK2 score <10%) without CVD or HF. These were further
subcategorised according to CKD status, with CKD defined as either an ACR > 3
mg/mmol or eGFR <60 ml/min/1.73m^2 in the most recent record of these in General
Practice within two years prior to the index date. QRISK2 was used instead of
QRISK3 as QRISK3 is poorly recorded in CPRD, and QRISK2 has similar
performance.[24] Most people did not have a recent QRISK2 recorded in CPRD
Aurum. This was instead estimated using an algorithm developed by researchers at
the London School of Hygiene and Tropical Medicine.[25] Appendix 2 describes the
implementation and validation of that algorithm for the present analysis.

Patients' IMD was divided into five groups defined by quintiles from least (5) to most
(1) deprived. For CVD risk factors, including laboratory results and BMI, the most
recent measurement within two years prior to the index date was used; if none were
available in this window the value was set to missing.

Presence of a current prescription of any medication was defined using the algorithm
developed by Farmer *et al.*[26] This estimates the days supply of a prescription using
any data available on the quantity of pills and the daily dose; or number of days
prescribed; or the average gap between previous prescriptions. If none of these are

available then a 28 day prescription length is assumed. If the most recent prescription date plus two times the estimated days supply falls after the index date, then the prescription is considered current. People with a current prescription for dapagliflozin, empagliflozin, canagliflozin, or ertugliflozin were counted as having a current SGLT2 inhibitor.

Descriptive and statistical analysis

The demographic and clinical characteristics of the prevalent population with T2DM in September 2023 were summarised using counts, percentages, and means, stratified by comorbidity (CVD risk score category, and CVD and HF status).

Uptake of SGLT2 inhibitors in each CVD group was defined the percentage of the population with a current prescription on the index date. Uptake was calculated overall, by comorbidity status, and further by age, gender, ethnicity, deprivation status, and duration of diagnosis.

The association between patient characteristics and odds of a current SGLT2 inhibitor prescription was then evaluated using logistic regression. First, one model was fitted to all people with either CVD or HF, with categorical variables for comorbidity status (CVD and CKD status) and all patient characteristics. Then, separate models were fitted for each CVD and CVD risk group (including people without CVD), each model including CKD status and all patient characteristics as covariables, to assess for heterogeneity in the association between patient characteristics and uptake between the different subpopulations. The analysis was conducted in STATA 16; all analysis code is available online (Appendix 1).

RESULTS

Baseline characteristics

Demographics

In total 587,826 people with prevalent T2DM in CPRD Aurum on 1[st] September 2023 met all the study inclusion criteria (Table 1). Most people were aged over 60 years (71.2%), 56.5% were male, and 75.0% were white. People with T2DM were more likely to live in deprived areas than the general population: 138,899 (23.6%) were living in the most deprived quintile in England compared to 95,425 (16.2%) living in the least deprived.

Cardiovascular disease and heart failure status

Cardiovascular disease was common: 129,306 (22.0%) had CVD without HF; 13,693 (2.3%) had HF without CVD; 35,266 (6.0%) had CVD and HF; 362,792 (61.7%) had neither CVD or HF but estimated 10 year CVD risk > 10%; and 46,769 (8.0%) had neither but CVD risk <10%.

Chronic kidney disease

In total 202,667 (34.5%) of the cohort had a record of comorbid CKD. Of these, 31,835 (15.7%) had stage 1 disease, 50,800 (25.1%) had stage 2 disease, 104,882 (51.8%) had stage 3 disease, and 15,150 (7.5%) had stage 4 or 5 disease (Appendix 3). A record of eGFR within 2 years prior to the index date was missing for 3.3% of the cohort.

Uptake of SGLT2 inhibitors

Overall uptake of SGLT-2 inhibitors for people with T2DM in General Practice was 20.7%, varying between 19.5% and 30.5% according to CVD group (Table 2). Uptake was higher in patients with heart failure (29.4% and 30.5% in patients with HF with and without CVD respectively) and lower in patients without heart failure (19.5%, 19.9%, and 20.2% in the patients without HF but with current CVD, high CVD risk, and low CVD risk, respectively). In the unadjusted results, uptake was markedly higher for people diagnosed with T2DM over 2 years prior to the analysis

index date compared to those diagnosed more recently; higher in men then women;

and higher in people aged under 70 years than those older than 70 years. Smaller

differences in unadjusted uptake were present according to CKD status, deprivation,

and ethnicity.

Table 1. Clinical and demographic characteristics of people in CPRD with prevalent

T2DM by cardiovascular disease comorbidity status, September 2023.

	All T2DM patients	CVD only	HF only	CVD + HF	High CVD risk, no HF	Low CVD risk, no HF*
Overall						
All patients	587,826 (100.0%)	129,306 (100.0%)	13,693 (100.0%)	35,266 (100.0%)	362,792 (100.0%)	46,769 (100.0%)
By CKD status						
No CKD**	365,549 (62.2%)	71,321 (55.2%)	5,167 (37.7%)	12,067 (34.2%)	240,048 (66.2%)	36,946 (79.0%)
Any CKD***	202,667 (34.5%)	55,529 (42.9%)	8,347 (61.0%)	22,778 (64.6%)	109,399 (30.2%)	6,614 (14.1%)
Missing	19,610 (3.3%)	2,456 (1.9%)	179 (1.3%)	421 (1.2%)	13,345 (3.7%)	3,209 (6.9%)
T2DM duration						
0-2 years	83,137 (14.1%)	13,145 (10.2%)	1,733 (12.7%)	3,049 (8.6%)	53,426 (14.7%)	11,784 (25.2%)
>2 years	504,689 (85.9%)	116,161 (89.8%)	11,960 (87.3%)	32,217 (91.4%)	309,366 (85.3%)	34,985 (74.8%)
Age (years)						
Mean age	67	72	74	76	67	50
18-39	13,894 (2.4%)	305 (0.2%)	103 (0.8%)	39 (0.1%)	2,267 (0.6%)	11,180 (23.9%)
40-49	43,494 (7.4%)	2,779 (2.1%)	395 (2.9%)	411 (1.2%)	20,056 (5.5%)	19,850 (42.4%)
50-59	112,272 (19.1%)	14,620 (11.3%)	1,309 (9.6%)	2,307 (6.5%)	79,429 (21.9%)	14,610 (31.2%)
60-69	156,682 (26.7%)	33,095 (25.6%)	2,622 (19.1%)	6,659 (18.9%)	113,179 (31.2%)	1,130 (2.4%)
70-79	153,988 (26.2%)	43,047 (33.3%)	4,164 (30.4%)	11,570 (32.8%)	95,205 (26.2%)	N/A
80-89	90,267 (15.4%)	29,532 (22.8%)	3,980 (29.1%)	11,405 (32.3%)	45,350 (12.5%)	-
90+	17,229 (2.9%)	5,928 (4.6%)	1,120 (8.2%)	2,875 (8.2%)	7,306 (2.0%)	-
Sex						
Male	332,160 (56.5%)	80,952 (62.6%)	6,950 (50.8%)	22,931 (65.0%)	203,181 (56.0%)	18,146 (38.8%)
Female	255,666 (43.5%)	48,354 (37.4%)	6,743 (49.2%)	12,335 (35.0%)	159,611 (44.0%)	28,623 (61.2%)
Deprivation (IMD)						
5 (Least deprived)	138,899 (23.6%)	30,890 (23.9%)	3,164 (23.1%)	8,616 (24.4%)	81,267 (22.4%)	14,962 (32.0%)
4	132,385 (22.5%)	28,417 (22.0%)	2,903 (21.2%)	7,834 (22.2%)	80,878 (22.3%)	12,353 (26.4%)
3	113,615 (19.3%)	24,850 (19.2%)	2,757 (20.1%)	6,670 (18.9%)	71,014 (19.6%)	8,324 (17.8%)
2	107,502 (18.3%)	24,118 (18.7%)	2,534 (18.5%)	6,579 (18.7%)	68,140 (18.8%)	6,131 (13.1%)
1 (Most deprived)	95,425 (16.2%)	21,031 (16.3%)	2,335 (17.1%)	5,567 (15.8%)	61,493 (16.9%)	4,999 (10.7%)
Ethnicity						
White	440,777 (75.0%)	103,630 (80.1%)	11,556 (84.4%)	29,534 (83.7%)	273,131 (75.3%)	22,926 (49.0%)
Mixed	7,266 (1.2%)	1,183 (0.9%)	122 (0.9%)	249 (0.7%)	4,235 (1.2%)	1,477 (3.2%)
Asian	87,284 (14.8%)	17,090 (13.2%)	987 (7.2%)	3,803 (10.8%)	54,790 (15.1%)	10,614 (22.7%)
Black	38,471 (6.5%)	5,324 (4.1%)	842 (6.1%)	1,292 (3.7%)	21,648 (6.0%)	9,365 (20.0%)
Chinese or Other	9,506 (1.6%)	1,644 (1.3%)	120 (0.9%)	329 (0.9%)	5,645 (1.6%)	1,768 (3.8%)
Not stated/known	4,522 (0.8%)	435 (0.3%)	66 (0.5%)	59 (0.2%)	3,343 (0.9%)	619 (1.3%)
*Counts rounded to nearest 10 where any cells have less than 10. **eGFR >59 & ACR <3/missing ***ACR > 3 or eGFR <60						

254   **Table 2.** Percentage of people with prevalent T2DM and a current prescription for an
255   SGLT-2 inhibitor, by cardiovascular disease status, September 2023.

	All T2DM patients	CVD, no HF	HF, no CVD	HF and CVD	High CVD risk	Low CVD risk
Overall						
All patients	20.7%	19.5%	29.4%	30.5%	19.9%	20.2%
CKD status						
No CKD	20.9%	20.7%	29.8%	31.6%	20.4%	20.1%
Any CKD	21.8%	18.5%	29.7%	30.3%	20.7%	27.8%
Missing	6.5%	7.6%	8.4%	9.0%	6.4%	5.4%
T2DM duration						
0-2 years	8.3%	7.8%	27.9%	25.5%	6.8%	8.6%
>2 years	22.8%	20.9%	29.7%	31.0%	22.2%	24.1%
Age (years)						
18-39	16.1%	23.9%	46.6%	43.6%	20.1%	14.7%
40-49	22.2%	29.0%	42.3%	48.2%	22.2%	20.4%
50-59	26.1%	30.3%	39.8%	44.0%	25.0%	24.1%
60-69	25.9%	27.4%	37.7%	41.4%	24.4%	20.3%
70-79	19.3%	18.9%	32.1%	34.1%	17.2%	N/A
80-89	10.5%	8.8%	21.6%	22.0%	7.6%	
90+	4.4%	2.4%	9.8%	11.0%	2.6%	
Sex						
Male	23.2%	22.2%	35.6%	34.9%	22.3%	19.1%
Female	17.5%	15.1%	23.1%	22.4%	17.0%	20.9%
Deprivation (IMD)						
5 (Least deprived)	20.9%	20.4%	28.7%	30.7%	20.0%	19.3%
4	20.7%	19.7%	30.1%	30.5%	20.0%	19.5%
3	20.5%	19.3%	28.5%	29.5%	19.8%	20.3%
2	20.9%	19.1%	30.8%	31.3%	20.1%	22.4%
1 (Most deprived)	20.6%	18.7%	29.2%	30.5%	19.9%	21.7%
Ethnicity						
White	21.0%	18.9%	29.5%	30.0%	20.3%	22.7%
Mixed	18.6%	21.0%	36.1%	34.5%	17.1%	17.0%
Asian	21.9%	24.5%	27.0%	34.8%	20.5%	19.3%
Black	16.5%	16.5%	29.8%	26.9%	15.5%	16.3%
Chinese or Other	18.6%	20.7%	35.8%	35.3%	17.0%	17.6%
Not stated / known	17.2%	15.4%	34.8%	30.5%	17.2%	15.7%

259 **Table 3.** Percentage of people with a current prescription for an SGLT-2 inhibitor, in
260 patients with T2DM and heart failure or CVD, by CKD status, September 2023.

	CVD only		HF only		CVD + HF	
	CKD	No CKD	CKD	No CKD	CKD	No CKD
Overall						
All patients	18.5%	20.7%	29.7%	29.8%	30.3%	31.6%
T2DM duration						
0-2 years	28.7%	47.5%	52.5%	52.6%	48.3%	48.0%
>2 years	19.3%	22.7%	29.8%	30.3%	30.6%	32.7%
Age (years)						
18-39	31.0%	23.1%	69.2%	39.7%	18.2%	60.0%
40-49	36.4%	27.9%	44.5%	41.3%	51.6%	49.6%
50-59	36.1%	29.5%	46.4%	37.6%	44.8%	44.8%
60-69	31.5%	26.2%	38.6%	37.9%	43.7%	40.4%
70-79	20.7%	17.7%	35.3%	27.5%	36.9%	29.6%
80-89	9.6%	7.8%	23.8%	14.9%	23.8%	16.3%
90+	2.6%	2.2%	10.8%	5.3%	11.6%	8.2%
Sex						
Male	21.8%	22.9%	36.2%	35.5%	35.1%	35.3%
Female	13.6%	16.8%	23.7%	22.7%	22.6%	22.4%
Deprivation (IMD)						
5 (Least deprived)	19.4%	21.6%	28.9%	29.0%	29.8%	33.1%
4	19.0%	20.8%	29.8%	31.4%	30.1%	31.9%
3	18.7%	20.2%	29.6%	27.3%	29.8%	29.6%
2	18.0%	20.4%	30.3%	32.3%	31.3%	31.7%
1 (Most deprived)	17.2%	20.2%	29.8%	28.8%	30.8%	30.6%
Ethnicity						
White	17.5%	20.3%	29.5%	30.0%	29.9%	30.9%
Mixed	22.2%	21.5%	31.5%	44.7%	34.4%	35.3%
Asian	25.7%	24.2%	28.0%	26.4%	34.2%	37.0%
Black	16.9%	16.8%	32.7%	25.8%	27.5%	27.8%
Chinese or Other	21.1%	21.5%	33.3%	42.0%	33.5%	39.7%
Not stated / known	20.6%	14.2%	34.2%	40.0%	31.3%	33.3%

When stratified by age group, a consistent association between CVD group and
SGLT2 inhibitor uptake was present, with uptake ordered from lowest to highest as
follows: low CVD risk score, high CVD risk score, CVD only, HF only, CVD and HF
(Table 2). In analysis stratified by CVD group and age, CKD was also consistently
associated with higher uptake of SGLT2 inhibitors for patients in each age category
with CVD or HF. In people aged 60-69 years with CVD, uptake was 26.2% in people

without CKD and 31.5% in people with CKD; similar differences were present in

other age groups.

Modelling analysis of uptake of SGLT2 inhibitors

All clinical patient characteristics were significantly (p<0.01) associated with

differences in the odds of having a current prescription for an SGLT2 inhibitor in a

model for people with CVD or HF which mutually adjusted for all other patient

characteristics, CVD group and CKD status (Table 4). CKD was associated with

higher odds of uptake (OR: 1.22, 95% CI: (1.19, 1.25)), as was HF (OR: 2.15 (2.06,

2.24)) and combined CVD and HF (OR: 2.15 (2.09, 2.22)). Patients aged 70-79

years had considerably lower odds than patients aged 18-39 (OR: 0.41 (0.33, 0.51)),

women had lower odds than men (OR: 0.67 (0.65, 0.68)), and people living in areas

in the highest quintile of deprivation had lower odds than those in the least deprived

(OR: 0.84 (0.81, 0.88)). Compared to people of White ethnicity, people of Asian

ethnicity had higher odds of a current prescription (OR: 1.07 (1.04, 1.11)) whilst

people Black ethnicity had lower odds (OR: 0.77 (0.72, 0.81)).

The same direction of association for each of the different patient characteristics in

the overall population was also generally observed within each CVD and CVD risk

group (Appendix 4, Figure 2). Higher age and female gender were consistently and

strongly associated with lower odds in all groups, except the low CVD risk group,

where odds were higher with higher age and only slightly lower for women (OR: 0.95

(0.90, 1.00)). Black ethnicity was associated with lower uptake in all groups, though

the difference was greater in patients without HF. Asian ethnicity was associated

with higher uptake in people with CVD. In patients with neither CVD nor HF, White

ethnicity was associated with considerably higher uptake than any other ethnicity.

CKD was associated with higher odds of a prescription in all subgroups (OR ranging

between 1.17 and 1.47), while more recent diagnosis was consistently associated

with lower odds except in patients without HF (Appendix Table 4).

Table 4. Logistic regression model for association between current SGLT2 inhibitor

prescription and patient clinical and demographic characteristics, people with T2DM

and either CVD or HF, September, 2023.

CKD		
No CKD	1.00	
CKD	1.22 (1.19, 1.25)	<0.01
Cardiovascular disease status		
CVD without HF	1.00	
HF without CVD	2.15 (2.06, 2.24)	<0.01
CVD and HF	2.15 (2.09, 2.22)	<0.01
Age group		
18-39	1.00	
40-49	0.98 (0.78, 1.23)	0.89
50-59	0.91 (0.74, 1.14)	0.41
60-69	0.71 (0.57, 0.88)	<0.01
70-79	0.41 (0.33, 0.51)	<0.01
80-89	0.19 (0.15, 0.24)	<0.01
90+	0.07 (0.06, 0.09)	<0.01
Gender		
Male	1.00	
Female	0.67 (0.65, 0.68)	<0.01
Deprivation		
5 (Least deprived)	1.00	
4	0.98 (0.94, 1.02)	0.38
3	0.91 (0.87, 0.95)	<0.01
2	0.89 (0.85, 0.92)	<0.01
1 (most deprived)	0.84 (0.81, 0.88)	<0.01
Ethnicity		
White	1.00	
Mixed	1.03 (0.91, 1.17)	0.60
Asian or Asian British	1.07 (1.04, 1.11)	<0.01
Black or Black British	0.77 (0.72, 0.81)	<0.01
Chinese or Other Ethnic Group	1.01 (0.91, 1.12)	0.88
Not stated/known	0.97 (0.78, 1.22)	0.80
Diagnosis recency		
0-1 years	1.00	
1-2 years	1.15 (1.05, 1.26)	<0.01
2-3 years	1.34 (1.22, 1.46)	<0.01
3-4 years	1.51 (1.38, 1.66)	<0.01
4-5 years	1.67 (1.53, 1.82)	<0.01
>5 years	3.08 (2.88, 3.29)	<0.01

Discussion

Principal findings

The percentage of people with T2DM with a current SGLT2 inhibitor prescription

remained relatively low by September 2023, 18 months after the 2022 NICE T2DM

guideline update. The percentage in the whole T2DM cohort was 20.7%. Uptake was

lower in people with comorbid CVD without HF (19.5%) and people at high CVD risk

without HF (19.9%) than for people with HF (29.4%). In age-stratified results, uptake

was higher with in people with comorbidities and greater CVD risk score: higher in

people with CKD than those without CKD; higher in people with CVD than people at

high CVD risk; and higher in people at high CVD risk than in those at low CVD risk.

In models which adjusted for patients' characteristics and comorbidity status, people

aged 70-79 had less than half the odds of having a current prescription than people

aged 18-39, and women had approximately two-thirds the odds of men. Consistent

ethnicity and deprivation inequalities in uptake were also observed in each of the

CVD and CVD risk groups analysed: people with Black ethnicity had lower odds of a

current prescription than people with White ethnicity, ranging from 8-40% lower in

the different groups, whilst people living in the most deprived areas had 9-10% lower

odds than those living in the least deprived areas.

Interpreting variation in uptake

Variation by comorbidity and NICE guidance type

Some of the variation in prescribing we report may be due to differences in the

speed and extent of implementation of different NICE guidance, including

Technology Appraisals on SGLT2 inhibitors for different indications between 2021

and 2023, and the 2022 T2DM guideline update (NICE NG28). Clinical

considerations may partly explain some of the other inequalities we report.

NICE NG28 does not differentiate between CVD and HF in the strength of its

recommendations, stating an SGLT2 inhibitor should be offered to people with T2DM

'if they have chronic heart failure or established atherosclerotic cardiovascular

disease'. However, we found prescribing was 50% higher in people with HF than in

people with CVD without HF. This may be due to prescribing to September 2023

being driven by Technology Appraisals for SGLT2 inhibitors for HF (2021-2023), in

addition to the NG28 (2022) guideline recommendations for their use for T2DM,

while no Technology Appraisals or regulatory approval existed specifically for the

use of SGLT2 inhibitors for CVD.[27] Higher uptake for people with T2DM and HF may

have been driven by continuation of prescribing initiated outside general practice,

due to the HF itself and independent of diabetes status, and may also reflect the

additional support for evidence-based prescribing provided by specialist HF and

community teams

Nonetheless, recommendations in NICE NG28 for groups with T2DM and not

covered by a relevant Technology Appraisal for another condition appear to have

been implemented to some extent in England. Specifically, in age-stratified results

prescribing of SGLT2 inhibitors was higher in people with T2DM with elevated CVD

risk scores than people with lower scores; and higher still for people with CVD. This

ordering mirrors the strength of recommendations in NG28 in these different groups.

However, further qualitative research is needed to understand the extent to which

variation in the strength of NICE recommendations can explain this variation.

The CPRD database analysed does not allow differentiation by where the

prescription was initiated, e.g. HF specialist or GP, or by primary reason for

prescribing, e.g. for cardiorenal benefit or glycaemic management. It is likely that

both reasons apply to varying extent in our population. However, the observation of

higher rates of prescribing for non-HF cohorts with longer durations of diagnosis,

compared to those diagnosed in the previous 2 years, suggest that glycaemic

management may have been a greater driver than cardiorenal protection in these

groups, as additional glucose-lowering medicines are typically required over time.

The increasing use of SGLT2 inhibitors in this period may also explain concurrent

downward trends in prescribing of glucose-lowering medicines without additional

cardiorenal benefits, such as sulphonylureas.[28]

Variation by age, gender, ethnicity, and deprivation

We report lower prescribing of SGLT2 inhibitors with higher age, particularly after

age 70 years. This could be partly due to concern about adverse outcomes relating

to SGLT2 inhibitor therapy, the avoidance of polypharmacy, and caution in the

context of frailty. NG28 recommends an SGLT2 inhibitor should be offered to

patients with T2DM under 40 if they have elevated lifetime CVD risk regardless of

their QRISK2 score, and the vast majority will have elevated lifetime risk. Differences

by gender may be at least partially due to adverse effects including genital mycotic

infections, recurrent UTIs being more common in women, and the need to avoid

SGLT2 inhibitors in women of child-bearing potential who are sexually active and not

using contraception. However, the extent to which the inequalities we report are due

to these factors is unclear. Seghezzo *et al* evaluated several drugs to treat HF in a

UK cohort with HF with and without T2DM, and reported similar age and gender

inequalities for SGLT2 inhibitors but also for drugs where UTIs and pregnancy are

not of concern.[29] The 2026 NICE T2DM guideline update cautions the use of SGLT2

inhibitors in people with T2DM and frailty, but highlights the sparsity of evidence for

this population,[30] while analysis of individual patient data indicates that frail patients

are underrepresented in RCTs and may have poorer outcomes.[31] One study in

Wales found similar profiles of comorbidity and adverse events between participants

in cardiovascular outcome trials of SGLT2 inhibitors and the real-world population

with a prescription,[32] and there is potential for further observational studies with

particular efforts to minimise confounding to inform guidance for frailer patient

groups.

Additional research is also needed to explore the differences in prescribing of SGLT2

inhibitors by ethnicity. The higher odds of a current prescription in Asian people may

reflect awareness of higher cardiovascular risk in this population. Although lower

prescribing in people with Black ethnicity may be partly due to concerns about DKA

with these medicines, and avoidance of use in those with a history of previous DKA

given that ketosis-prone diabetes is more prevalent in people of Black ethnicity,[33] it is

more likely that lower uptake in Black people, along with lower prescribing in areas of

higher deprivation, relate to systemic barriers in accessing healthcare including

differential provisional as well as financial, cultural and psychosocial factors.

Challenges to engagement with health services in minoritised groups and lack of

incentives to initiate SGLT2 inhibitors in primary care may both contribute.

Population health benefits of higher uptake of SGLT2 inhibitors

Network-meta analyses have demonstrated reductions in mortality, hospitalisation for heart failure, and myocardial infarctions associated with SGLT2 inhibitor therapy for people with T2DM and either CVD, HF, CKD, or high risk of CVD.[7] In the UK 92% of people with T2DM have at least one of these. With the loss of exclusivity of the patent for the most prescribed SGLT2 inhibitor in the UK, generic dapagliflozin has recently launched and acquisition costs are reducing, with an 82% price reduction in the NHS tariff between July 2025 and February 2026. The reductions imply costs of considerably less than £20,000 per additional quality-adjusted life year gained with SGLT2 inhibitor therapy, indicating substantial population health benefits can be gained from higher uptake whilst limiting displaced resources within the wider NHS.

Comparison to other studies on SGLT2 inhibitor uptake in the UK

To our knowledge no previous studies have reported uptake of SGLT2 inhibitors for T2DM up to 2023. Volumes of GP prescriptions for dapagliflozin (for any indication) increased in England approximately 80% between September 2023 and August 2025, from 495,000 individual units per month to 907,000.[34] Applying this increase to overall uptake in the present study (21%), implies it could have reached approximately 37% in people with T2DM by August 2025.

Farmer *et al* evaluated current prescriptions of people with T2DM with and without CVD, using CPRD GOLD up to 2019.[26] They reported that, in people without CVD, the percentage of people with a current SGLT-2 inhibitor prescription increased from 6.7% to 13.8% between end-2016 and end-2019. In people with CVD it increased from 4.3% to 9.8%. These results are compatible with our findings and suggest that, between 2019 and 2023, there was a modest rise in SGLT2i prescribing in those with T2DM without CVD and a greater increase in those with T2DM with CVD.

Seghezzo *et al* reported large increases in uptake of SGLT2 inhibitors in people with HF with reduced ejection from approximately 8% in January 2022 to over 25% by May 2024 in UK General Practice, with no indication of a plateau by 2024, and higher prescribing in people with comorbid T2DM.[29] Another large UK General Practice study on CKD up to December 2022 reported that 62.8% of people with CKD had guideline-directed indication for SGLT2 inhibitor, but uptake in that cohort

441   with CKD and T2DM was only 22%.[35] These results are consistent with those in our
442   subgroups of T2DM patients with comorbid HF and CKD.

443   Regarding the distribution of CVD risk scores in people with T2DM that we estimated
444   Young *et al* estimated than in 2020 59.6% of people with T2DM had high risk of CVD
445   and 6.9% had low risk,[36] compared to our estimates of 64% and 8% by 2023.

Comparison to other studies on SGLT2 inhibitor uptake internationally

Other comparable studies report lower uptake of SGLT2 inhibitors for T2DM and
CKD internationally compared to the UK, though there is sparse data after 2022.

Gagnon *et al* evaluated uptake of SGLT2 inhibitors in patients with T2DM and
comorbid HF, separately in a specialised heart function clinic and in all adults in
Alberta, Canada, between 2018 and 2022.[37] In the specialised clinic cohort it
reached 49.1% by March 2022, whilst in the whole province it reached 26.4% by
January 2022. In one population-based study of adults with T2DM hospitalised with
HF in New South Wales in Australia, uptake of SGLT2 inhibitors had reached 13.2%
by mid-2021, with lower uptake reported in older age groups, people with higher
frailty, and, in contrast to our analysis, people with CKD.[38] Another analysis of
114,000 adults with T2DM in primary care in Australia reported uptake of 13.6% in
people without CKD and 14.4% in people with CKD in 2020-2021.[39]

Uptake and inequalities in uptake of other NICE guidelines

There is evidence that including NICE indicators in the Quality and Outcomes
Framework has resulted in improved adherence to guidelines. [40] Otherwise,
however, there is evidence that adherence to NICE guideline recommendations in
General Practice is frequently low. Several small studies in General Practice settings
have evaluated uptake of NICE recommendations in areas including heart failure [41],
hypertension [42], T2DM and comorbid hypertension [43], and osteoarthritis [44]. They
have reported variable and low uptake.

Limitations

Evaluation of the QRISK2 algorithm we applied indicated that up to 20% of people
estimated as having a low CVD risk score (QRISK2 <10%) could have had a

QRISK2 of greater than 10% (Appendix 2). This imprecision may lead to some

under-estimation of the difference in uptake between people with high and low CVD

risk. Many people with CVD and HF may have been misclassified as not having

them, as recording of these in primary care can be incomplete,[45] and linked data

from HES was only available to March 2021. Such misclassification may result in

underestimation of uptake differences between people with and without

comorbidities.

We did not evaluate how many people had previously initiated but then stopped an

SGLT2 inhibitor, would be clinically unsuitable for treatment, or assess for

polypharmacy in our population. These factors will explain a proportion of the

patients not prescribed one. In the Canadian study of a cohort with T2DM and HF

Gagnon *et al* reported that 62.4% of people met all the eligibility criteria for trials

proving the effectiveness of SGLT2 inhibitors,[37] and in an analysis of uptake of

SGLT2 inhibitors for CKD in the UK, Forbes *et al* reported that 62.8% of patients with

T2DM and CKD were indicated for therapy with an SGLT2 inhibitor.[35] Further

research would be needed to identify a plausible ceiling for uptake for the people in

scope of NG28 in our analysis. However, if approximately a third of people were

ineligible for an SGLT2 inhibitor, our results would imply an opportunity to at least

double prescribing in people with HF, and triple prescribing in people with CVD

without HF.

Conclusion

By September 2023 prescribing of SGLT2 inhibitors remained relatively low for

people with T2DM recommended them in recent NICE technology appraisals and

guidelines in England. We report lower uptake in people with T2DM and CVD without

HF than people with T2DM and HF, despite an equal guideline recommendation for

both these groups, and inequalities by ethnicity and deprivation. SGLT2 inhibitor

prescribing increased substantially to September 2023 and afterwards to August

2025 nationally. However, our results show that progress has not been evenly

distributed, and that a substantial opportunity remains to increase prescribing in

2026.

A major update to the NICE T2DM guideline in February 2026 recommended that all

T2DM patients, regardless of their CVD risk, should be offered both metformin and

an SGLT2 inhibitor. This expands the eligible population to include people in all the

clinical subgroups considered in the present analysis. Action is now necessary to

support implementation of this guidance in all these groups, particularly in people

without CVD, CKD, or HF where uptake has been lower. Healthcare systems and

policy makers should explore interventions supporting case-finding and quality

improvement, such as pay-for-performance mechanisms, to ensure comprehensive

and equitable access to these cost-effective therapies. As well as qualitative

research on facilitators and barriers to SGLT2 inhibitor prescribing in primary care,

ongoing surveillance of T2DM care processes and prescribing is needed to monitor

uptake.

Contribution to authorship

The study was conceived by LO, PM, and designed by all the authors. PM and JW

did the data and statistical analysis with support from RW and EY; JW and MR

produced the code lists. All authors assisted with the interpretation of results. PM

wrote the manuscript with substantial input and revisions from all other authors. The

corresponding author attests that all listed authors meet authorship criteria and that

no other meeting the criteria have been omitted.

Acknowledgements, funding, and disclaimer

523   PM, JW, MR, JH, RW, EY, SR, MW, LSR and LO are employees of the National

524   Institute for Health and Care Excellence. AA is supported by funding from NIHR

525   Oxford Biomedical Research Centre. The authors are solely responsible for any

526   errors or omissions. The opinions expressed in this article are those of the authors

527   and do not necessarily reflect the position of their affiliations.

Disclosure of interests

DJC has received investigator-initiated grants from Astra Zeneca and Novo Nordisk, support for education from Perspectum and consultancy fees from Madrigal with any financial renumeration from pharmaceutical company consultation made to the University of Liverpool. All other authors declare no competing interests.

Sharing of data and results

No additional data are available; the authors are not the data controllers and do not have permission to share the data.

Transparency statement

PM had full access to all the data in the study, and affirms that the manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that there are no discrepancies between the study conducted and the study as originally planned.

Ethical approval

This study was based on data from the Clinical Practice Research Datalink obtained under licence from the UK Medicines and Healthcare products Regulatory Agency. The study was approved by the CPRD Research Data Governance process (ID: 23_003530).

REFERENCES

REFERENCES

1. Office for Health Improvement and disparities. Diabetes profile: statistical commentary, March
2025, 2025.
2. Pearson-Stuttard J, Holloway S, Polya R, et al. Variations in comorbidity burden in people with
type 2 diabetes over disease duration: A population-based analysis of real world evidence.
eClinicalMedicine 2022;52 doi: 10.1016/j.eclinm.2022.101584
3. Perkovic V, Jardine MJ, Neal B, et al. Canagliflozin and Renal Outcomes in Type 2 Diabetes and
Nephropathy. *New England Journal of Medicine* 2019;380(24):2295-306. doi:
doi:10.1056/NEJMoa1811744
4. Wiviott SD, Raz I, Bonaca MP, et al. Dapagliflozin and Cardiovascular Outcomes in Type 2 Diabetes.
New England Journal of Medicine 2019;380(4):347-57. doi: doi:10.1056/NEJMoa1812389
5. Ndumele CE, Rangaswami J, Chow SL, et al. Cardiovascular-Kidney-Metabolic Health: A
Presidential Advisory From the American Heart Association. *Circulation* 2023;148(20):1606-
35. doi: doi:10.1161/CIR.0000000000001184
6. National Institute for Health and Care Excellence. Type 2 diabetes in adults: management (NG28),
2024.
7. National Institute for Health and Care Excellence. Type 2 diabetes in adults: management
(Medicines update): Cost-effectiveness analysis: subsequent pharmacological therapy for
the management of type 2 diabetes (draft for consultation), 2025.
8. National Institute for Health and Care Excellence. What we do: NICE guidelines, 2024.
9. Curtis HJ, Walker AJ, Goldacre B. Impact of NICE guidance on tamoxifen prescribing in England
2011-2017: an interrupted time series analysis. *British journal of cancer* 2018;118(9):1268-
75. doi: 10.1038/s41416-018-0065-2 [published Online First: 2018/04/24]
10. NHS England. NICE Technology Appraisals in the NHS in England (Innovation Scorecard), 2023.
11. Wolf A, Dedman D, Campbell J, et al. Data resource profile: Clinical Practice Research Datalink
(CPRD) Aurum. *International Journal of Epidemiology* 2019;48(6):1740-40g. doi:
10.1093/ije/dyz034
12. Clinical Practice Research Datalink. CPRD Aurum December 2023 (Version 2023.12.001) [Data
set], 2023.
13. Herrett E, Gallagher AM, Bhaskaran K, et al. Data Resource Profile: Clinical Practice Research
Datalink (CPRD). *International Journal of Epidemiology* 2015;44(3):827-36. doi:
10.1093/ije/dyv098
14. Shiekh SI, Harley M, Ghosh RE, et al. Completeness, agreement, and representativeness of
ethnicity recording in the United Kingdom's Clinical Practice Research Datalink (CPRD) and
linked Hospital Episode Statistics (HES). *Population health metrics* 2023;21(1):3. doi:
10.1186/s12963-023-00302-0 [published Online First: 2023/03/16]
15. Boyd A, Cornish R, Johnson L, et al. Understanding Hospital Episode Statistics (HES). London, UK:
CLOSER, 2017.
16. Noble M, Wright G, Smith G, et al. Measuring Multiple Deprivation at the Small-Area Level.
Environment and Planning A: Economy and Space 2006;38(1):169-85. doi: 10.1068/a37168
17. Padmanabhan S, Carty L, Cameron E, et al. Approach to record linkage of primary care data from
Clinical Practice Research Datalink to other health-related patient data: overview and
implications. *European Journal of Epidemiology* 2019;34(1):91-99. doi: 10.1007/s10654-018-
0442-4
18. NHS England. Primary Care Domain reference sets 2023 [Available from:
https://isd.digital.nhs.uk/trud/user/guest/group/0/pack/8/subpack/659/releases.
19. OpenCodelists (Bennett Institute UoO. OpenCodelists 2024 [Available from:
https://www.opencodelists.org/.

20. Office for Health Improvement and Disparities and the NHS Benchmarking Network. The
Cardiovascular Disease Prevention Audit (CVDPREVENT) 2024 [Available from:
https://www.cvdprevent.nhs.uk/.
21. World Health Organisation. ICD-10 Version: 2010 2010 [Available from:
https://icd.who.int/browse10/2010/en.
22. Clinical Practice Research Datalink. CPRD: Denominator data 2024 [Available from:
https://www.cprd.com/denominator-data.
23. Hippisley-Cox J, Coupland C, Vinogradova Y, et al. Predicting cardiovascular risk in England and
Wales: prospective derivation and validation of QRISK2. *BMJ (Clinical research ed)*
2008;336(7659):1475-82. doi: 10.1136/bmj.39609.449676.25 [published Online First:
2008/06/25]
24. Hippisley-Cox J, Coupland C, Brindle P. Development and validation of QRISK3 risk prediction
algorithms to estimate future risk of cardiovascular disease: prospective cohort study. *BMJ*
(Clinical research ed) 2017;357:j2099. doi: 10.1136/bmj.j2099
25. Herrett E, Gadd S, Jackson R, et al. Eligibility and subsequent burden of cardiovascular disease of
four strategies for blood pressure-lowering treatment: a retrospective cohort study. *The*
Lancet 2019;394(10199):663-71. doi: 10.1016/S0140-6736(19)31359-5
26. Farmer RE, Beard I, Raza SI, et al. Prescribing in Type 2 Diabetes Patients With and Without
Cardiovascular Disease History: A Descriptive Analysis in the UK CPRD. *Clinical*
Therapeutics 2021;43(2):320-35. doi: 10.1016/j.clinthera.2020.12.015
27. Electronic Medicines Compendium. SMPC: Forxiga 10 mg film-coated tablets, 2024.
28. Bennett Institute for Applied Data Science. 6.1.2.1: Sulfonylureas: University of Oxford,.
29. Seghezzo G, Costello RE, Martineau F, et al. Impact of the COVID-19 Pandemic on Treatment and
Outcomes Among Patients with Heart Failure with Reduced Ejection Fraction: Nationwide
Studies Using the OpenSAFELY Platform. 2025
30. National Institute for Health and Care Excellence. Type 2 diabetes in adults: management
(medicines update) In development, 2025.
31. Wightman H, Butterly E, Wei L, et al. Frailty in randomized controlled trials of glucose-lowering
therapies for type 2 diabetes: An individual participant data meta-analysis of frailty
prevalence, treatment efficacy, and adverse events. *PLoS Med* 2025;22(4):e1004553. doi:
10.1371/journal.pmed.1004553 [published Online First: 2025/04/07]
32. Hanlon H, Wightman W, Sullivan S, et al. Assessing the representativeness of trials of Sodium-
glucose Cotransporter- 2 inhibitors in type 2 diabetes (preprint). 2025
33. Umpierrez GE, Smiley D, Kitabchi AE. Narrative review: ketosis-prone type 2 diabetes mellitus.
Annals of internal medicine 2006;144(5):350-7. doi: 10.7326/0003-4819-144-5-200603070-
00011 [published Online First: 2006/03/08]
34. OpenPrescribing.net. Dapagliflozin (0601023AG): Bennett Institute for Applied Data Science,
University of Oxford, 2025.
35. Forbes AK, Hinton W, Feher MD, et al. Implementation of chronic kidney disease guidelines for
sodium-glucose co-transporter-2 inhibitor use in primary care in the UK: a cross-sectional
study. *eClinicalMedicine* 2024;68 doi: 10.1016/j.eclinm.2024.102426
36. Young KG, Hopkins R, Shields BM, et al. Recent UK type 2 diabetes treatment guidance
represents a near whole population indication for SGLT2-inhibitor therapy. *Cardiovascular*
diabetology 2023;22(1):302. doi: 10.1186/s12933-023-02032-x [published Online First:
2023/11/03]
37. Gagnon LR, Hazra D, Perera K, et al. Uptake of SGLT2i and Outcomes in Patients with Diabetes
and Heart Failure: A Population-Based Cohort and a Specialized Clinic Cohort. *American*
heart journal 2024;274:11-22. doi: 10.1016/j.ahj.2024.04.007 [published Online First:
2024/04/27]
38. Lin J, Milder TY, Deakin CT, et al. Trends in use of sodium-glucose cotransporter 2 inhibitors
among people with type 2 diabetes following hospitalisation with heart failure: A

population-based study. *Diabetes research and clinical practice* 2025;224:112242. doi:
10.1016/j.diabres.2025.112242 [published Online First: 2025/05/11]

39. Wallace H, Wick J, Neuen BL, et al. Prevalence of SGLT2 inhibitor and GLP1 receptor agonist
prescriptions in type 2 diabetes patients with and without chronic kidney disease: Analysis of
an Australian primary care dataset. *Diabetes, Obesity and Metabolism* 2025;27(10):5599-
611. doi: https://doi.org/10.1111/dom.16608

40. Mendelson A, Kondo K, Damberg C, et al. The Effects of Pay-for-Performance Programs on
Health, Health Care Use, and Processes of Care: A Systematic Review. *Annals of internal*
medicine 2017;166(5):341-53. doi: 10.7326/m16-1881 [published Online First: 2017/01/24]

41. Paschalis T, Jones C. Plasma HbA1c in the investigation of suspected heart failure in general
practice: An audit of the 2018 NICE guidelines update. *Journal of family medicine and*
primary care 2020;9(2):1098-102. doi: 10.4103/jfmpc.jfmpc_917_19 [published Online First:
2020/04/23]

42. Heneghan C, Perera R, Mant D, et al. Hypertension guideline recommendations in general
practice: awareness, agreement, adoption, and adherence. *British Journal of General*
Practice 2007;57(545):948-52.

43. Kempegowda P, Chandan JS, Abdulrahman S, et al. Managing hypertension in people of African
origin with diabetes: Evaluation of adherence to NICE Guidelines. *Primary care diabetes*
2019;13(3):266-71. doi: 10.1016/j.pcd.2018.12.007 [published Online First: 2019/02/02]

44. Healey EL, Afolabi EK, Lewis M, et al. Uptake of the NICE osteoarthritis guidelines in primary care:
a survey of older adults with joint pain. *BMC musculoskeletal disorders* 2018;19(1):295. doi:
10.1186/s12891-018-2196-2 [published Online First: 2018/08/18]

45. Herrett E, Shah AD, Boggon R, et al. Completeness and diagnostic validity of recording acute
myocardial infarction events in primary care, hospital care, disease registry, and national
mortality records: cohort study. *BMJ (Clinical research ed)* 2013;346:f2350. doi:
10.1136/bmj.f2350 [published Online First: 2013/05/23]

FIGURE LEGENDS

Figure 1. Timeline of selected NICE Technology Appraisals (blue) and guideline
updates (orange) recommending SGLT2 inhibitors for different indications, 2014-
2023.

Figure 2. Odds ratios for association between patient characteristics and a current
SGLT2 inhibitor prescription, by cardiovascular disease status, models adjusted for
diagnosis recency and CKD stage, September 2023.

Mar 2022

TA773. Empagliflozin recommended for HF with reduced ejection fraction as add-on to optimised care.

Feb 2021

TA679. Dapagliflozin recommended for HF with reduced ejection fraction as add on to optimised care.

Mar 2022

TA775. Dapagliflozin recommended for CKD as add on to optimised care if eGFR in range (25, 75) ml/min/1.73 m^2 and T2DM or ACR>= 22.6 mg/mmol.

Jun 2023

TA902. Dapagliflozin recommended for CHF with preserved or mildly reduced ejection fraction as add on to optimised care.

2014-2019

TA315, TA 336, TA390, TA418, TA572, TA583. TAs recommend different SGLT2 inhibitors as part of triple therapy for glycaemic control in T2DM, or as monotherapy or dual therapy if specified other therapies are not appropriate.

Feb 2022

NG28. An SGLT2 inhibitor is recommended as part of first-line treatment for people with T2DM and comorbid HF or comorbid aCVD, for CVD risk reduction irrespective of HbA1c. Consider an SGLT2 inhibitor if T2DM and high CVD risk.

An SGLT2 inhibitor is recommended for people with T2DM and CKD if ACR >30mg/mmol as an add-on to optimised care.

1st Sep 2023

Analysis index date.