Randomized controlled trials claiming "personalized", "individualized" and "precision" interventions: characteristics, transparency and bias

Luigi Russo[1], Nicolò Lentini[1#], Luisa Soru[2], Roberta Pastorino[1,3], Stefania Boccia[1,3], John PA Ioannidis[4,5,6,7]

1. Section of Hygiene, Department of Life Sciences and Public Health, Università Cattolica del Sacro Cuore, Rome, Italy

2. Grant Office, Fondazione Policlinico Universitario A. Gemelli IRCCS, Rome, Italy

3. Department of Woman and Child Health and Public Health, Fondazione Policlinico Universitario A. Gemelli IRCCS, Rome, Italy

4. Stanford Prevention Research Center, Department of Medicine, Stanford University School of Medicine, Stanford, CA, USA.

5. Department of Epidemiology and Population Health, Stanford University School of Medicine, Stanford, CA, USA.

6. Department of Biomedical Data Science, Stanford University School of Medicine, Stanford, CA, USA.

7. Meta-Research Innovation Center at Stanford (METRICS), Stanford University, Stanford, CA, USA

#Corresponding Author

Nicolò Lentini MSc

Mail nicolo.lentini@unicatt.it

Section of Hygiene, University Department of Life Sciences and Public Health, Università Cattolica del Sacro Cuore, 00168 Rome, Italy

Abstract

The terms personalized, individualized and precision medicine are increasingly used to describe health interventions, yet their operational meaning in clinical research remains unclear. Despite extensive conceptual discussion, there is limited empirical evidence on how these labels are applied in randomized controlled trials (RCTs) and whether such trials meet standards of transparency and methodological rigor. We systematically examined 262 RCTs published between 2020 and 2022 that used the terms "personalized", "individualized", or "precision" in the title to describe an intervention. The term "personalized" was used most frequently (49.2%), followed by "individualized" (45.8%) and "precision" (5.0%). In most trials, personalization involved behavioral, digital, or pharmacological interventions, with few studies employing –omics approaches. Personalization was most often based on individual lifestyle factors, psychological characteristics, or disease classification. We also found that in most trials, personalization consisted of tailoring a single intervention to individuals (82.8%), often through individualized dosage (73.2%). Most included RCTs were judged to be at high risk of bias and showed limited transparency with respect to data and code sharing. Our study suggests that, in contemporary RCTs, the labels "personalized", "individualized", and "precision" are applied interchangeably to a wide range of heterogeneous interventions that are predominantly non-genomic. Greater conceptual clarity and stronger methodological standards are needed to ensure that claims of personalization in clinical research are empirically meaningful and reliable.

Introduction

Personalized, individualized, and precision medicine represent rapidly evolving and frequently touted domains within healthcare. The main principle is to tailor prevention, diagnosis and treatment to the unique characteristics of individuals, drawing on insights from genomics, behavior, lifestyle, and environmental factors. Intervention strategies are to be informed by the specific measurements, attributes, and responses of single participants taking advantage of the latest technological advancements.

However, the interpretation of the terms "personalized", "individualized", and "precision" may vary widely across disciplines, and even within the same field, researchers may use them inconsistently. Definitions from authoritative bodies highlight this ambiguity. For instance, the National Human Genome Research Institute [1] defines personalized medicine narrowly, as the use of an individual"s genetic profile to guide disease prevention, diagnosis, and treatment. By contrast, the International Consortium for Personalized Medicine (ICPerMed) [2] adopts a broader perspective, viewing it as a model that integrates both phenotypic and genotypic information (e.g., molecular profiling, imaging, lifestyle data) to inform therapy or prevention.

These concepts have permeated numerous medical specialties, illustrating their versatility but also the lack of consensus on their exact meaning. In oncology, for example, precision medicine often refers to genetic profiling that identifies mutations and guides targeted therapies [3,4]. In cardiology, personalized risk assessments combining genetic and lifestyle factors aim to inform individualized prevention strategies, such as interventions based on lifetime genetic risk scores [5,6]. Beyond genetics, the term "personalized" has been applied to very diverse interventions. Illustratively, these include tailored physiotherapy after surgery, customized text-message reminders for adherence [7], and music therapy for dementia patients [8].

The existing literature reflects this diversity of interpretation. Editorials, commentaries, and reviews have examined, debated, and promoted (often with some hype) the origins, evolution, and

future directions of personalized, individualized and precision medicine. Broadly, these writings fall into two groups: those that conceive of the concepts as an integration of genetic, social, and environmental features [9], and those that focus primarily on biological and genomic applications [10]. Ethical and social concerns [11], along with differences in public understanding [12], add further complexity. Despite the growing use of these terms in published clinical trials, operational clarity remains limited. Different articles may emphasize definitional debates [13], argue over which features should be included, or speculate about future developments [14,15], while the real-world implications of the terminology remain underexplored.

Clarifying how these terms are applied in clinical research is therefore essential. To address this gap, we surveyed randomized controlled trials (RCTs) published between 2020 and 2022 that describe interventions as "personalized", "individualized", or "precision". The aim was to map the range of meanings attached to these terms and to analyze how they are applied across different clinical contexts. Having assembled this literature corpus, we also aimed to assess whether these trials are transparent and whether they have low or high risk of bias.

Methods

We previously registered the protocol for this study on Open Science Framework [16]. The protocol aimed to generate a literature corpus of RCTs that prominently use in their titles the terms of interest to describe assessed interventions. Once the eligible trials were assembled and we could have a better sense of what they pertained to, we further extended the protocol to characterize better these trials and to explore features of transparency and bias. The full list of trials in available in Supplementary File 1 for further methodological research.

Search strategy

We searched for articles published between 2020 and 2022, in English, and indexed in Medline. We focused on RCTs that used the terms of interest to prominently describe the intervention in their title. We applied the following search string:

(precision [ti] OR personalized [ti] OR personalised [ti] OR individualized [ti] OR individualised [ti]) Filter[RandomizedControlledTrial]

After conducting the literature search, we evaluated each article against the following eligibility criteria: (1) any RCT on humans, including parallel arm, cross-over, cluster randomized or any other randomized design; and (2) randomization of the participants to an intervention considered "personalized", "individualized", or "precision". Both articles reporting the results of RCTs and those reporting protocols of RCTs were deemed eligible.

We excluded articles that did not report either the protocol or the results of an RCT (e.g., reviews, editorials, perspectives). We also excluded articles in which the relevant terms were not used to describe a randomized intervention, but rather for other purposes, as well as articles that did not address an actual or possible clinical condition and those not related to medicine and health.

Screening

Identified articles were uploaded into the Rayyan software. Two reviewers (L.R., A.P.) independently performed screening based on title and abstract, and disagreements were resolved

through consensus. Full texts of the identified articles were retrieved and reviewed independently by two researchers (L.R., L.S.), and disagreements were resolved by consensus. Articles meeting the eligibility criteria were subsequently entered into data extraction.

Data extraction

Two reviewers (L.R., L.S.) independently performed data extraction. From each included randomized controlled trial, we extracted:

General information: title, first author, DOI, journal of publication, year of publication, country, type of paper (protocol, trial).

Study characteristics: medical specialty (according to European Union directive [17]), specific disease, number of arms, total sample size, and study group. Studies were classified into two main categories:

Group 1: the personalized/individualized/precision intervention is compared to a non-personalized/individualized/precision intervention. This group was further subdivided post hoc into:

- ☐ Group 1a: studies with two or more intervention arms but only one personalized/individualized/precision;

- ☐ Group 1b: studies including more than one personalized/individualized/precision intervention arms

Group 2: the personalized/individualized/precision intervention is compared to another personalized/individualized/precision intervention.

Furthermore, for articles presenting trial results, we extracted whether the abstract conclusion was favorable to the personalized intervention, had mixed conclusions, or was unfavorable compared with the control. For articles in group 2 and for protocols this classification was not applicable.

Personalized, individualized, or precision intervention characteristics: we extracted which term was used; the type of application (i.e., primary prevention, secondary prevention, therapy); and the type of intervention (digital, omics, behavioral, dietary, medication, surgery, physical exercises,

radiotherapy, anesthesia, physiotherapy, rehabilitation, psychological). We evaluated whether the intervention provided was the same for all participants in the intervention arm; if it was not, we extracted how it differed (schedule of therapy, dosage of therapy, follow-up/visits, testing, drug used, anatomical target (e.g., radiotherapy applied to different body areas/organs; prostheses applied to different parts of the same bone or tissue). For dosage or schedule of therapy, we considered not only pharmacological treatments but any type of intervention, including, for example, diets with different caloric intake or different intensity in physical activity.

We extracted a description of the intervention, including the specific features used to personalize it, the category of individual features on which the intervention was based (as reported below), and the control arm intervention.

The individual features upon which the intervention is framed were classified in the following main categories and subcategories: demographics (age, sex, place of birth, domicile), socioeconomic (financial status, cultural and social, education), anthropometrics (height, weight, BMI), genetic (genetic testing, monogenic disorder), lifetime risk, omics (microbiome, polygenic risk score, metabolomics, nutrigenomics, proteomics), lifestyle and behavioral, psychological (patient preferences, motivation), vital signs (blood pressure, temperature, pulse rate, respiration rate), imaging tools (radiologic, magnetic resonance, scintigraphy, ECG, PET, ultrasound, etc.), tests on biological material (blood, urine, biopsy, other), complex scores (specify), physical or functional capacity, disease classification or other (specify).

As part of post hoc data extraction, we extracted from each study the number of personalized/individualized/precision interventions present and classified them as 1, 2, 3, 4, or multiple (>4). Moreover, we extracted from each study how the intervention content was tailored across participants, defined as the number of different ways in which the personalized intervention was delivered, and classified this as 1, 2, 3, 4, multiple (>4), or different for each patient. Finally, for studies that included differences in dosage, we extracted the number of dosage levels present (1, 2, 3, 4, multiple [>4], or for each patient), whereas studies without dosage differences were

recorded as "no dose involved." This approach allowed us to capture studies in which a single personalized intervention (e.g., a diet) was conceptually one intervention but varied at the individual level in terms of content, dose, or both.

Data extraction was performed by two independent researchers (L.R., L.S.), with discrepancies discussed for consensus or settled by a third assessor (N.L.).

Transparency indicators

For articles describing trial results, we also performed manual data extraction and applied previously validated algorithms for transparency features [18] to extract information on code sharing, data sharing, trial registration, disclosures of conflicts of interest, and disclosures of funding. For registered trials, we extracted whether the registration was prospective, retrospective or before completion.

Risk of Bias assessment

To assess the risk of bias of the included studies we utilized the revised Cochrane Risk-of-Bias Tool for Randomized Trials (RoB 2) [19], through the ROBoto2 webapp [20]. ROBOTO2 is an open-source, web-based platform for large language model (LLM)-assisted risk of bias assessment of clinical trials. For graphical visualization, we used the Risk-of-bias VISualization (robvis) webapp [21].

Data synthesis

The results were described narratively and through tables and graphs, where appropriate. Descriptive statistics were presented for the characteristics of the eligible studies. Trial characteristics and main outcomes were reported as counts, percentages for categorical variables, and medians with interquartile ranges (IQRs) for continuous variables.

As post hoc synthesis, we classified each study based on the combination of number of personalized/individualized/precision intervention(s) and the way of tailoring it, and the number of different doses. For example, a study with personalized feedback was classified as a single intervention, with no dose involved, and tailored differently for each participant.

Statistical analysis

Exploratory analyses were conducted to assess whether there were differences in the characteristics and features of articles depending on whether they were (1) protocols or results; (2) used the term "personaliz(s)ed", "individualiz(s)ed", or "precision"; and (3) had favorable conclusions for the personalized intervention (among those that had results rather than just protocols). We used Pearson"s χ^2 or Fisher"s exact tests to assess whether each category differed between type of study, term used and conclusions of the abstract. For associations between ordinal categorical variables with more than two levels, we applied a linear-by-linear association test (Mantel test for trend) to account for the ordered structure of the data. For continuous variables, the Kruskal–Wallis test or the Mann–Whitney U test was applied, as appropriate.

Agreement between Rtransparent automated extraction and human extraction for transparency-related fields was assessed using Cohen"s kappa coefficient. For discordant cases, discrepancies were reviewed to determine the correct classification.

A 2-sided $P < .005$ was considered statistically significant, and $P < .05$ was considered suggestive [22]. All analyses were performed using R version 4.4.0 (2024-04-24) for Windows.

Results

A total of 435 records were identified and screened by title and abstract. Of these, 313 articles were assessed for full-text eligibility, and 262 were finally included in the review. Fifty-five articles were excluded after full-text assessment for the following reasons: ineligible intervention either non-medical condition (n=8) or non-personalized intervention (n = 3); article not possible to retrieve (n = 10); unsuitable study design including secondary analysis (n=11), non-randomized design (n=10), sub-study of an RCT (n=4), economic evaluation (n=1); non-English language (n = 2); and retracted article (n = 2). Figure 1 shows the PRISMA flowchart.

Characteristics of the included trials

Among the 262 included trials (Table 1), the term "personalized" was used most frequently (n=129; 49.2%), followed by "individualized" (n=120; 45.8%) while "precision" was uncommon (n=13; 5.0%). The majority were reports of trial results but the corpus included also 28 protocols. Geographic location was highly diverse. Most trials had two arms (84.4%), while 12.2% and 3.1% had three and four arms, respectively, while one had 6 arms. The median sample size was 136 (IQR 63.3–302.3). Studies were largely classified in group 1a (85.9%) and group 1b (8.0%). Among 218 RCTs that compared personalized interventions versus non-personalized controls, the conclusion of the Abstract was favorable to the personalized/individualized/precision intervention in 156 (70.6%), mixed in 36 (16.3%), and unfavorable in only 29 (13.1%).

Characteristics of the intervention

As shown in Table 2, more than two-thirds of the interventions were therapeutic rather than preventive. Intervention types varied widely, with the most common being medication-based interventions (26%), behavioral approaches (24%), and digital interventions (13.0%). Within the experimental arm, differences most frequently pertained to the dosage of therapy (51.9%). A total of 573 individual features were identified as the main categories on which personalization was based. Within this vast variety of features, lifestyle and behavioral features (37.4%), classification

of the disease (28.6%), and psychological features (27.1%), were the most common, while genetic and omics features were uncommon. Most studies employed a single intervention (67.6%), and most studies tailored the intervention(s) to each patient (82.8%). In those with dosage variations, the dosage most often differed for each patient (73.2%). Hence, the most common combinations were one intervention with both dosage and tailoring different for each patient (34.0%), and one intervention, with tailoring different for each patient and no dose involved (29.8%) (Table 3).

Association analyses

The association analysis revealed almost no differences in applications and features among studies employing different terms (Table 2), having different abstract conclusions (Supplementary Table 1) or studies published versus protocols (Supplementary Table 2). The few significant patterns observed were that digital and behavioral interventions almost always were termed "personalized" while medication interventions were termed "individualized", studies with interventions tailored on lifestyle and behavioral employed more commonly "personalized" while those tailored on anthropometrics features were more frequently "individualized". Finally, studies on weight outcomes were largely termed "individualized" (Supplementary Table 3).

Transparency among trials

Statements on code sharing were rare, with only 1 study (0.4%) providing open code. Similarly, open data availability was rare (13 studies, 5.1%). On the other hand, presence of statements on funding source (93.4%) and on conflict of interest (93.4%) were present in almost all of the studies. 202 studies (78.3%) were registered in a trial repository, but of those only 155 (59.1% of the total) were registered prospectively.

Comparison between the Rtransparent automated extraction and human assessment revealed moderate agreement for funding declarations ($\kappa = 0.59$) and conflict of interest declarations ($\kappa = 0.51$), and lower agreement for study registration ($\kappa = 0.39$). Agreement was more substantial for data availability ($\kappa = 0.63$). Open code availability was proposed in 2 cases by Rtransparent, and 2 cases by manual extraction (with none overlapping) ($\kappa = -0.01$) (Table 4).

Risk of bias assessment

The results of the risk of bias evaluation are reported in Figure 2 and in Supplementary File 2. Overall, 162 studies (68.6%) were assessed as being at high risk of overall bias, 32 (13.6%) raised some concerns, and 42 (17.8%) were judged to be at low risk. The domains most frequently contributing to a high risk of bias were bias in selection of the reported result (53.4%), bias in measurement of the outcome (35.2%), and bias arising from the randomization process (33.5%) (Supplementary Table 4).

In exploratory analysis, no association was found between conclusion of the abstract and overall RoB. Favorable conclusions were seen in 106 (65.4%) of the RCTs at high risk of bias, in 19 (59.4%) of the RCTs with some concerns, and in 31 (73.8%) of the RCTs with low risk of bias (Supplementary Table 5). The respective rate of unfavorable conclusions were seen in 13.0%, 12.5%, and 9.5%, respectively.

Discussion

Our study found extensive heterogeneity in the features and characteristics of the interventions labeled as "personalized", "individualized", or, far less commonly, "precision". Nevertheless, despite frequent conceptual distinctions in the literature, these labels were applied largely interchangeably across trials. Evaluation of these studies showed that most of them had some favorable results for the experimental intervention(s), very few had transparency in terms of data and code sharing, and most suffered from high risk of bias. Most interventions described with these terms relied on non-genomic features, such as lifestyle behaviors, psychological characteristics, disease classification, or functional capacity. Genetic and omics-based information has been often portrayed as the defining hallmark of precision medicine [23–25]. Precision oncology [26,27] has a strong focus on targeted pharmacological therapies based on single genetic mutations [28]. However, in our study, these applications were comparatively rare. This may reflect a substantial disconnect between dominant narratives in precision medicine discourse and the realities of clinical trial implementation. An alternative explanation is that precision oncology trials do not use that terminology in the titles of their articles and thus they were not included in our eligible RCTs. Regardless, the whole discipline of personalized/individualized/precision medicine is shifting towards an integrative approach that incorporates also lifestyle, environmental exposures, education, and socioeconomic factors [29–31], rather than just molecular profiling. In our sample of RCTs, interventions labeled as personalized, individualized, or precision were more frequently behavioral in nature and primarily based on lifestyle-related characteristics. These contextual elements are fundamental while personalizing interventions [32,33].

In our included interventions, personalization most often involved adjusting intervention content, intensity, or dosage based on readily observable or self-reported individual characteristics. Moreover, the predominance of single interventions tailored differently for each participant suggests that personalization in RCTs is frequently operationalized as flexible delivery rather than

as fundamentally distinct therapeutic strategies [34]. In this sense, personalized interventions are often heterogeneous in their application but not in their underlying structure, design, or conceptual framework. More simply, the intervention remains essentially the same but is adapted to individual characteristics [34,35].

This approach is particularly relevant in studies involving rehabilitation [36] or physical exercise interventions [37]. Moving beyond rigid programs with identical targets for all participants may help overcome a common limitation of such interventions, namely the failure to achieve predefined goals [38]. In practice, objectives are often expressed as fixed thresholds, such as a minimum number of steps per day or minutes of physical activity. However, for individuals with different baseline characteristics, abilities, or health conditions, personalized targets may lead to greater adherence, improved health outcomes, or enhanced preventive effects, ultimately resulting in greater overall benefit [39,40]. While such tailoring may be clinically sensible, its methodological implications are important. When interventions vary extensively across participants, interpretability, reproducibility, and causal attribution become more challenging, particularly if tailoring algorithms are insufficiently described [41].

Among the published RCTs in our study, the conclusions of the Abstract almost always had some favorable elements (either exclusively favorable or mixed) for personalized interventions. Very few published trials reached only unfavorable conclusions. This extremely high rate of success seems spurious. It therefore raises concerns about possible biases that may have distorted, selected, and/or exaggerated the research findings of these studies. The threat of this literature being unreliable is increased given that the majority of studies were already judged to be at high risk of bias based on RoB2 assessments. Moreover, additional biases may exist beyond those captured by the RoB2 tool.

To safeguard this research literature and enhance trust in its findings, transparency would be most welcome. However, we also documented that critical transparency practices were also very limited, especially with regard to sharing of data and code. The very low rate of data sharing and the

almost non-existent sharing of code places these trials at lower transparency ratings, compared with the average biomedical literature and other trials [18]. However, other studies of high-profile trials also show very low rates of data and code sharing (e.g. 4.4% and 4.5%, respectively in Siena et al. [42]). This lack of transparency is especially problematic in studies where intervention tailoring relies on decision rules, algorithms, or adaptive components that are central to the intervention itself [41]. Enhanced sharing is currently feasible for RCTs in general [43] and should be promoted also for these specific trials.

Careful pre-registration with availability of detailed protocols with extensively documented statistical analysis plans would also enhance trust in this literature. In our eligible RCTs, we documented that three-quarters were registered, with less than 60% were registered before the trial started. While we did not examine in depth the registration entries, it is typical for registry entries of RCTs to give very limited information on details of the proposed analysis, and adherence to registration guidance is low [44–46]. Therefore, this poor documentation may leave many researcher degrees of freedom on what they report and what to conclude about the trial.

We observed no substantive differences in study characteristics based on whether authors used the term "personalized", "individualized", or "precision". This supports the interpretation that these terms function largely as rhetorical labels [47,48], rather than as indicators of distinct methodological or practical approaches, and that in clinical research they are often used interchangeably [9]. In this respect, their use in practice diverges from theoretical definitions, particularly for personalized and precision medicine, which are frequently treated as distinct concepts in the literature and by experts [49,50].

Limitations

This study has some limitations. First, our search was restricted to RCTs that included the relevant terms in the title, potentially excluding studies that implemented individualized approaches without explicitly labeling them as such. However, this choice is consistent with our objective of examining how these terms are publicly and prominently used in the scientific literature. Second,

only a limited number of RCTs used the term "precision", which restricts the generalizability of findings related to this label.. Third, we focused on the three key terms denoting personalization that may apply to any field. Some additional terms are more specific to specific fields and interventions. It would be too convoluted to unearth every single trial that has personalization features. For example, some oncology trials with personalization features may use a variety of terms such as "targeted" therapy, "master" protocols, "umbrella" and "basket" designs, "enrichment" or "adaptive"[51].

Conclusions

Contemporary RCTs that include interventions labeled as "personalized", "individualized", or "precision" encompass a wide and heterogeneous set, and most rely on lifestyle and behavioral approaches. In practice, these terms are used largely interchangeably, reflecting a lack of clear distinction in their use in clinical research and practice. The fact that most of these trials reach favorable conclusions and have low transparency, and high risk of bias suggests that there is substantial room for improving the rigor and reliability of this important line of clinical and translational investigation. Our evaluated sample of 262 RCTs provides a recent baseline against which future progress in the field can be probed.

Declarations

Acknowledgements: The authors would like to sincerely thank Dr. Salvatore Di Grande and Dr. Alessio Perilli for their support during the early phases of the study. The work of Luigi Russo has been supported by the European Network Staff Exchange for Integrating Precision Health in the Healthcare Systems project (Marie Skłodowska-Curie Research and Innovation Staff Exchange no. 823995).

Ethical approval: Not required (evaluation of published literature).

Contributors: LR: Conceptualization, Methodology, Validation, Formal analysis, Investigation, Data Curation, Writing - Original Draft, Writing - Review & Editing, Visualization. NL: Validation, Formal analysis, Data Curation, Writing - Original Draft, Writing - Review & Editing, Visualization. LS: Formal analysis, Investigation, Writing - Review & Editing. RP: Conceptualization, Methodology, Writing - Original Draft, Writing - Review & Editing, Supervision. SB: Conceptualization, Methodology, Validation, Resources, Writing - Original Draft, Writing - Review & Editing, Supervision, funding acquisition. JPAI: Conceptualization, Methodology, Validation, Formal analysis, Writing - Original Draft, Writing - Review & Editing, Visualization, Supervision. JPAI and SB are the guarantors of the study.

Data availability statement and code sharing: The full list of the included RCTs in contained in Supplementary File 1. The dataset used for the analysis will be available on Open Science Framework at this link: https://osf.io/cgy4k/overview.

Funding: None declared.

Competing interests: None declared.

Transparency: The lead authors (the manuscript''s guarantors) affirm that the manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

Figure 1. PRISMA flow diagram of the study search and selection process.

Figure 2. Summary of the Risk of bias with RoB-2 tool of the included RCTs.

Table 1. Characteristics of the eligible RCTs.

Variable		N (%)	N (%)	N (%)	N (%)	p
		Overall	**Individualized**	**Personalized**	**Precision**	
		262 (100%)	120 (45.8%)	129 (49.2%)	13 (5.0%)	
Year	2020	78 (29.8)	35 (29.2)	40 (31.0)	3 (23.1)	0.517
	2021	82 (31.3)	37 (30.8)	43 (33.3)	2 (15.4)	
	2022	102 (38.9)	48 (40.0)	46 (35.7)	8 (61.5)	
Type of article	Trial	235 (89.7)	110 (91.7)	116 (89.9)	9 (69.2)	0.048
	Protocol	27 (10.3%)	10 (8.3%)	13 (10.1%)	4 (31.8%)	
Country	Europe	86 (32.8%)	35 (29.2%)	47 (36.4%)	4 (30.8%)	**<0.001**
	Germany	14 (5.3)	9 (7.5)	5 (3.9)	0 (0.0)	
	UK	13 (5.0)	2 (1.7)	11 (8.5)	0 (0.0)	
	France	12 (4.6)	8 (6.7)	3 (2.3)	1 (7.7)	
	Europe others	47 (17.9)	16 (13.3)	28 (21.7)	3 (23.1)	
	USA	58 (22.1)	13 (10.8)	39 (30.2)	6 (46.2)	
	China	42 (16.0)	27 (22.5)	12 (9.3)	3 (23.1)	
	Australia	19 (7.3)	9 (7.5)	10 (7.8)	0 (0.0)	
	Canada	12 (4.6)	5 (4.2)	7 (5.4)	0 (0.0)	
	World others	45 (17.2)	31 (25.8)	14 (10.9)	0 (0.0)	
Medical specialty	Cardiology	32 (12.2)	19 (15.8)	10 (7.8)	3 (23.1)	0.017
	Psychiatry	29 (11.1)	6 (5.0)	22 (17.1)	1 (7.7)	
	Oncology	28 (10.7)	8 (6.7)	16 (12.4)	4 (30.8)	
	Endocrinology	27 (10.3)	12 (10.0)	15 (11.6)	0 (0.0)	
	Obstetrics and gynecology	16 (6.1)	9 (7.5)	5 (3.9)	2 (15.4)	
	Orthopedics	16 (6.1)	9 (7.5)	7 (5.4)	0 (0.0)	
	Infectious Diseases	14 (5.3)	5 (4.2)	8 (6.2)	1 (7.7)	
	Respiratory medicine	13 (5.0)	7 (5.8)	6 (4.7)	0 (0.0)	
	Internal medicine	13 (5.0)	5 (4.2)	8 (6.2)	0 (0.0)	
	Others	72 (27.5%)	40 (33.3%)	30 (23.3%)	2 (15.4%)	
Number of trial arms	Two arms trial	221 (84.4)	101 (84.2)	109 (84.5)	11 (84.6)	0.787
	Three arms trial	32 (12.2)	15 (12.5)	16 (12.4)	1 (7.7)	
	Four arms trial	8 (3.1)	4 (3.3)	3 (2.3)	1 (7.7)	
	Six arms trial	1 (0.4)	0 (0.0)	1 (0.8)	0 (0.0)	
Sample size	Median (IQR)	136.0 (63.3, 302.3)	114.5 (62.8, 229.3)	150.0 (60.0, 355.0)	200.0 (150.0, 493.0)	0.045
Study group	1a	225 (85.9)	106 (88.3)	109 (84.5)	10 (76.9)	
	1b	21 (8.0)	5 (4.2)	14 (10.9)	2 (15.4)	0.138
	2	16 (6.1)	9 (7.5)	6 (4.7)	1 (7.7)	
Conclusions of	Favorable for	156 (70.6)	71 (69.6)	79 (71.8)	6 (66.7)	0.581

the Abstract [a]	the personalized intervention					
	Mixed conclusions	36 (16.3)	15 (14.7)	20 (18.2)	1 (11.1)	
	Unfavorable for the personalized intervention	29 (13.1)	16 (15.7)	11 (10.0)	2 (22.2)	

In bold p<0.005. [a]=N=221 since the conclusions were not assessed for protocols and group 2.

Table 2. Characteristics of the personalized, individualized, or precision intervention in eligible trials

Variable		Overall N (%)	Individualized N (%)	Personalized N (%)	Precision N (%)	p-value
Type of application	Therapy	186 (71.0)	94 (78.3)	85 (65.9)	7 (53.8)	0.069
	Primary prevention	74 (28.2)	25 (20.8)	43 (33.3)	6 (46.2)	
	Secondary prevention	2 (0.8)	1 (0.8)	1 (0.8)	0 (0.0)	
Type of intervention [a]	Behavioral	87 (33.2%)	27 (22.5%)	56 (43.4%)	4 (30.8%)	**0.002**
	Digital	77 (29.4%)	17 (14.2%)	58 (45.0%)	2 (15.4%)	**<0.001**
	Medication	72 (27.5%)	41 (34.2%)	24 (18.6%)	7 (53.8%)	**0.002**
	Psychological	57 (21.8%)	19 (15.8%)	37 (28.7%)	1 (7.7%)	0.024
	Physical exercises	44 (16.8%)	26 (21.7%)	17 (13.2%)	1 (7.7%)	0.161
	Dietary	41 (15.6%)	17 (14.2%)	24 (18.6%)	0 (0.0%)	0.181
	Educational	33 (12.6%)	13 (10.8%)	19 (14.7%)	1 (7.7%)	0.659
	Rehabilitation	22 (8.4%)	10 (8.3%)	10 (7.8%)	2 (15.4%)	0.514
	Surgery	17 (6.5%)	12 (10.0%)	5 (3.9%)	0 (0.0%)	0.132
	Anesthesia	16 (6.1%)	13 (10.8%)	3 (2.3%)	0 (0.0%)	0.015
	Omics	6 (2.3%)	0 (0.0%)	5 (3.9%)	1 (7.7%)	0.034
	Physiotherapy	4 (1.5%)	3 (2.5%)	1 (0.8%)	0 (0.0%)	0.474
	Radiotherapy	4 (1.5%)	3 (2.5%)	1 (0.8%)	0 (0.0%)	0.474
Difference within the intervention arm [a]	Dosage of therapy	136 (51.9)	71 (59.2)	61 (47.3)	4 (30.8)	0.055
	Schedule of therapy	75 (28.6)	44 (36.7)	29 (22.5)	2 (15.4)	0.027
	Drug used	32 (12.2)	16 (13.3)	12 (9.3)	4 (30.8)	0.069
	Testing	18 (6.9)	6 (5.0)	9 (7.0)	3 (23.1)	0.064
	Anatomical target	14 (5.3)	6 (5.0)	8 (6.2)	0 (0.0)	0.897
	Follow-up/visits	14 (5.3)	8 (6.7)	5 (3.9)	1 (7.7)	0.427
	Other	47 (17.9)	16 (13.3)	28 (21.7)	3 (23.1)	0.188
Control arm	Other treatment	110 (42.0)	50 (41.7)	53 (41.1)	7 (53.8)	0.246
	Standard of care	68 (26.0)	28 (23.3)	35 (27.1)	5 (38.5)	
	Usual care	34 (13.0)	17 (14.2)	17 (13.2)	0 (0.0)	
	No intervention	21 (8.0)	9 (7.5)	12 (9.3)	0 (0.0)	
	Placebo	18 (6.9)	13 (10.8)	4 (3.1)	1 (7.7)	
	Waitlist	11 (4.2)	3 (2.5)	8 (6.2)	0 (0.0)	
Main category of individual features [a]	Lifestyle and behavioral	98 (37.4)	34 (28.3)	61 (47.3)	3 (23.1)	**0.005**
	Disease classification	75 (28.6)	30 (25.0)	41 (31.8)	4 (30.8)	0.472
	Psychological	71 (27.1)	27 (22.5)	43 (33.3)	1 (7.7)	0.045
	Vital signs	56 (21.4)	35 (29.2)	19 (14.7)	2 (15.4)	0.019
	Physical or functional capacity	55 (21.0)	32 (26.7)	21 (16.3)	2 (15.4)	0.124
	Anthropometrics	47 (17.9)	32 (26.7)	15 (11.6)	0 (0.0)	**0.002**
	Test on biological material	42 (16.0)	20 (16.7)	19 (14.7)	3 (23.1)	0.604
	Imaging tools	29 (11.1)	16 (13.3)	10 (7.8)	3 (23.1)	0.117
	Demographics	24 (9.2)	8 (6.7)	13 (10.1)	3 (23.1)	0.111
	Complex scores (specify)	22 (8.4)	13 (10.8)	9 (7.0)	0 (0.0)	0.405
	Socioeconomic	14 (5.3)	3 (2.5)	8 (6.2)	3 (23.1)	0.013

	Genetic	13 (5.0)	2 (1.7)	8 (6.2)	3 (23.1)	0.006
	Omics	8 (3.1)	1 (0.8)	7 (5.4)	0 (0.0)	0.133
	Other (specify)	19 (7.3)	11 (9.2)	7 (5.4)	1 (7.7)	0.455
Subcategories [a]	Patient preferences	41 (15.6)	16 (13.3)	24 (18.6)	1 (7.7)	0.455
	Weight	41 (15.6)	30 (25.0)	11 (8.5)	0 (0.0)	**<0.001**
	Blood	35 (13.4)	17 (14.2)	16 (12.4)	2 (15.4)	0.785
	Motivation	25 (9.5)	10 (8.3)	15 (11.6)	0 (0.0)	0.421
	Blood pressure	17 (6.5)	12 (10.0)	5 (3.9)	0 (0.0)	0.132
	Education	14 (5.3)	3 (2.5)	8 (6.2)	3 (23.1)	0.013
	Pulse rate	12 (4.6)	7 (5.8)	4 (3.1)	1 (7.7)	0.315
	Respiration rate	8 (3.1)	5 (4.2)	2 (1.6)	1 (7.7)	0.174
	BMI	7 (2.7)	3 (2.5)	4 (3.1)	0 (0.0)	1.000
	Genetic testing	6 (2.3)	1 (0.8)	3 (2.3)	2 (15.4)	0.026
	Lifetime risk	5 (1.9)	1 (0.8)	3 (2.3)	1 (7.7)	0.196
	Age	3 (1.1)	1 (0.8)	2 (1.6)	0 (0.0)	1.000
	Biopsy	2 (0.8)	0 (0.0)	1 (0.8)	1 (7.7)	0.097
	Metabolomics	2 (0.8)	1 (0.8)	1 (0.8)	0 (0.0)	1.000
	Temperature	2 (0.8)	1 (0.8)	1 (0.8)	0 (0.0)	1.000
	Genetic risk score (GRS)	1 (0.4)	0 (0.0)	1 (0.8)	0 (0.0)	1.000
	Microbiome	1 (0.4)	0 (0.0)	1 (0.8)	0 (0.0)	1.000
	Nutrigenomics	1 (0.4)	0 (0.0)	1 (0.8)	0 (0.0)	1.000
	Proteomics	1 (0.4)	0 (0.0)	1 (0.8)	0 (0.0)	1.000
	Urine	1 (0.4)	0 (0.0)	1 (0.8)	0 (0.0)	1.000
	Other	2 (0.8)	2 (1.7)	0 (0.0)	0 (0.0)	0.306
Difference in dosage	1	4 (1.5)	2 (1.7)	1 (0.8)	1 (7.7)	0.015
	2	4 (1.5)	2 (1.7)	1 (0.8)	1 (7.7)	
	3	3 (1.1)	1 (0.8)	2 (1.6)	0 (0.0)	
	4	4 (1.5)	2 (1.7)	2 (1.6)	0 (0.0)	
	>4	25 (9.5)	16 (13.3)	6 (4.7)	3 (23.1)	
	For each patient	109 (41.6)	55 (45.8)	52 (40.3)	2 (15.4)	
	NA	113 (43.1)	42 (35.0)	65 (50.4)	6 (46.2)	
Number of personalized intervention	1	177 (67.6)	78 (65.0)	93 (72.1)	6 (46.2)	0.015
	2	42 (16.0)	15 (12.5)	23 (17.8)	4 (30.8)	
	3	13 (5.0)	6 (5.0)	7 (5.4)	0 (0.0)	
	4	3 (1.1)	2 (1.7)	1 (0.8)	0 (0.0)	
	>4	27 (10.3)	19 (15.8)	5 (3.9)	3 (23.1)	
Difference in tailoring	1	1 (0.4)	1 (0.8)	0 (0.0)	0 (0.0)	0.014
	2	5 (1.9)	2 (1.7)	2 (1.6)	1 (7.7)	
	3	2 (0.8)	2 (1.7)	0 (0.0)	0 (0.0)	
	4	1 (0.4)	0 (0.0)	1 (0.8)	0 (0.0)	
	>4	21 (8.0)	12 (10.0)	6 (4.7)	3 (23.1)	
	For each patient	217 (82.8)	94 (78.3)	116 (89.9)	7 (53.8)	
	NA	15 (5.7)	9 (7.5)	4 (3.1)	2 (15.4)	

[a] = Same studies employed multiple measurements. NA=Not Applicable. In bold p<0.005.

Table 3. Combination of number of intervention(s), number of dosage(s) and tailoring of the intervention(s).

Combination in the intervention			Term used			Conclusion of the abstract			Type of study	
Number of interventions	Tailoring	Dosage	Individualized (N (%))	Personalized (N (%))	Precision (N (%))	Favorable (N (%))	Mixed (N (%))	Unfavorable (N (%))	Protocol (N (%))	Trial (N (%))
One	Different for each	Different for each	44 (36.7)	43 (33.3)	2 (15.4)	58 (37.2)	9 (25.0)	11 (37.9)	9 (33.3)	80 (34.0)
One	Different for each	No dose	29 (24.2)	47 (36.4)	2 (15.4)	46 (29.5)	14 (38.9)	10 (34.5)	8 (29.6)	70 (29.8)
One	NA	Fixed	2 (1.7)	1 (0.8)	1 (7.7)	3 (1.9)	0 (0.0)	1 (3.4)	0 (0.0)	4 (1.7)
One	Multiple (>4)	Multiple (>4)	0 (0.0)	1 (0.8)	1 (7.7)	0 (0.0)	1 (2.8)	0 (0.0)	1 (3.7)	1 (0.4)
One	Four	Four	0 (0.0)	1 (0.8)	0 (0.0)	1 (0.6)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.4)
One	Two	Multiple (>4)	1 (0.8)	0 (0.0)	0 (0.0)	1 (0.6)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.4)
One	Same for each	No dose	1 (0.8)	0 (0.0)	0 (0.0)	1 (0.6)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.4)
One	NA	Multiple (>4)	1 (0.8)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (3.4)	0 (0.0)	1 (0.4)
One	NA	Two	1 (0.8)	0 (0.0)	0 (0.0)	1 (0.6)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.4)
Two	NA	Fixed	1 (0.8)	1 (0.8)	1 (7.7)	2 (1.3)	0 (0.0)	0 (0.0)	1 (3.7)	2 (0.9)
Two	Different for each	No dose	5 (4.2)	10 (7.8)	1 (7.7)	7 (4.5)	2 (5.6)	1 (3.4)	1 (3.7)	15 (6.4)
Two	Different for each	Different for each	6 (5.0)	7 (5.4)	0 (0.0)	6 (3.8)	1 (2.8)	0 (0.0)	3 (11.1)	10 (4.3)
Two	Multiple combinations	Multiple combinations	1 (0.8)	3 (2.3)	1 (7.7)	2 (1.3)	1 (2.8)	0 (0.0)	1 (3.7)	4 (1.7)
Two	Two	No dose	0 (0.0)	2 (1.6)	1 (7.7)	3 (1.9)	0 (0.0)	0 (0.0)	0 (0.0)	3 (1.3)
Two	Two	Multiple (>4)	1 (0.8)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.4)
Three	Different for each	Different for each	3 (2.5)	1 (0.8)	0 (0.0)	4 (2.6)	0 (0.0)	0 (0.0)	0 (0.0)	4 (1.7)
Three	NA	Fixed	1 (0.8)	2 (1.6)	0 (0.0)	3 (1.9)	0 (0.0)	0 (0.0)	0 (0.0)	3 (1.3)
Three	Different for each	No dose	0 (0.0)	3 (2.3)	0 (0.0)	0 (0.0)	2 (5.6)	0 (0.0)	0 (0.0)	3 (1.3)
Three	Three	No dose	2 (1.7)	0 (0.0)	0 (0.0)	2 (1.3)	0 (0.0)	0 (0.0)	0 (0.0)	2 (0.9)
Three	Multiple combinations	Multiple combinations	1 (0.8)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.4)
Four	NA	Fixed	1 (0.8)	0 (0.0)	0 (0.0)	1 (0.6)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.4)
Four	Different for each	Different for each	1 (0.8)	0 (0.0)	0 (0.0)	0 (0.0)	1 (2.8)	0 (0.0)	0 (0.0)	1 (0.4)
Multiple (>4)	Multiple (>4)	Multiple (>4)	12 (10.0)	2 (1.6)	1 (7.7)	7 (4.5)	3 (8.3)	5 (17.2)	0 (0.0)	15 (6.4)

Multiple (>4)	Different for each	No dose	5 (4.1)	3 (2.3)	2 (15.4)	8 (5.1)	1 (2.8)	0 (0.0)	1 (3.7)	9 (3.8)
Multiple (>4)	Different for each	Different for each	1 (0.8)	1 (0.8)	0 (0.0)	0 (0.0)	1 (2.8)	0 (0.0)	1 (3.7)	1 (0.4)
Multiple (>4)	Different for each	Multiple (>4)	0 (0.0)	1 (0.8)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (3.7)	0 (0.0)

Table 4. Comparison of transparency-related patterns across studies assessed by a human reviewer and the Rtransparent algorithm.

Variable	N (%)	Kappa	Conflicts in assessment	Accurate extraction	
				Human	Rtransparent
Declaration of Funding	244 (93.1)	0.541	21 (8.0%)	9 (42.9%)	12 (57.1%)
Declaration of Conflict of interest	245 (93.5)	0.508	11 (4.2%)	2 (18.2%)	9 (81.8%)
Registration on a trial repository	204 (77.9)	0.370	44 (16.8%)	41 (93.2%)	3 (6.8%)
Before completion	*25 (12.3)*				
Not known	*2 (1.0)*				
Retrospectively	*22 (10.8)*				
Prospectively	*155 (76.0)*				
Data availability		0.640	46 (17.6%)	29 (63.0%)	17 (37.0%)
Data fully available	*13 (5.0)*	-	-	-	-
Data not available	*171 (65.3)*	-	-	-	-
Data available upon request	*78 (29.8)*	-	-	-	-
Code availability		-0.008	4 (1.5%)	3 (75.0%)	1 (25.0%)
Code available	*1 (0.4)*	-	-	-	-
Code not available	*257 (98.1)*	-	-	-	-
Code available upon request	*4 (1.5)*	-	-	-	-

References

[1]   National Human Genome Research Institute (NIH). Personalized-Medicine 2025. https://www.genome.gov/genetics-glossary/Personalized-Medicine (accessed December 17, 2025).

[2]   European Commission. European Council Conclusion on personalised medicine for patients (2015/C 421/03). 2015.

[3]   Rituraj, Pal RS, Wahlang J, Pal Y, Chaitanya M, Saxena S. Precision oncology: transforming cancer care through personalized medicine. Medical Oncology 2025;42:246. https://doi.org/10.1007/s12032-025-02817-y.

[4]   Ahmad F. Optimizing Treatment: The Role of Pharmacology, Genomics, and AI in Improving Patient Outcomes. Drug Dev Res 2025;86. https://doi.org/10.1002/ddr.70093.

[5]   Leopold JA, Loscalzo J. Emerging Role of Precision Medicine in Cardiovascular Disease. Circ Res 2018;122:1302–15. https://doi.org/10.1161/CIRCRESAHA.117.310782.

[6]   Franks PW, Cefalu WT, Dennis J, Florez JC, Mathieu C, Morton RW, et al. Precision medicine for cardiometabolic disease: a framework for clinical translation. Lancet Diabetes Endocrinol 2023;11:822–35. https://doi.org/10.1016/S2213-8587(23)00165-1.

[7]   Konerding U, Redaèlli M, Ackermann K, Altin S, Appelbaum S, Biallas B, et al. A pragmatic randomised controlled trial referring to a Personalised Self-management SUPport Programme (P-SUP) for persons enrolled in a disease management programme for type 2 diabetes mellitus and/or for coronary heart disease. Trials 2021;22:659. https://doi.org/10.1186/s13063-021-05636-4.

[8]   McCreedy EM, Gutman R, Baier R, Rudolph JL, Thomas KS, Dvorchak F, et al. Measuring the effects of a personalized music intervention on agitated behaviors among nursing home residents with dementia: design features for cluster-randomized adaptive trial. Trials 2021;22:681. https://doi.org/10.1186/s13063-021-05620-y.

[9]   Delpierre C, Lefèvre T. Precision and personalized medicine: What their current definition says and silences about the model of health they promote. Implication for the development of personalized health. Frontiers in Sociology 2023;8. https://doi.org/10.3389/fsoc.2023.1112159.

[10]   Goetz LH, Schork NJ. Personalized medicine: motivation, challenges, and progress. Fertil Steril 2018;109:952–63. https://doi.org/10.1016/j.fertnstert.2018.05.006.

[11]   Mentis A-FA, Pantelidi K, Dardiotis E, Hadjigeorgiou GM, Petinaki E. Precision Medicine and Global Health: The Good, the Bad, and the Ugly. Front Med (Lausanne) 2018;5. https://doi.org/10.3389/fmed.2018.00067.

[12]   Fournier V, Prebet T, Dormal A, Brunel M, Cremer R, Schiaratura L. Definition of Personalized Medicine and Targeted Therapies: Does Medical Familiarity Matter? J Pers Med 2021;11:26. https://doi.org/10.3390/jpm11010026.

[13]   Delpierre C, Lefèvre T. Precision and personalized medicine: What their current definition says and silences about the model of health they promote. Implication for the development of personalized health. Frontiers in Sociology 2023;8. https://doi.org/10.3389/fsoc.2023.1112159.

[14]   Salari K, Watkins H, Ashley EA. Personalized medicine: hope or hype? Eur Heart J 2012;33:1564–70. https://doi.org/10.1093/eurheartj/ehs112.

[15]   Chan IS, Ginsburg GS. Personalized Medicine: Progress and Promise. Annu Rev Genomics Hum Genet 2011;12:217–44. https://doi.org/10.1146/annurev-genom-082410-101446.

[16]   Luigi Russo, Salvatore Di Grande, Alessio Perilli, Roberta Pastorino, Stefania Boccia, John P.A. Ioannidis. Use Of The Words "Personalized", "Individualized" And "Precision" In Randomized Controlled Trials: Definitions And Common Characteristics. 1017605/OSFIO/CGY4K 2022. https://osf.io/cgy4k/overview (accessed December 17, 2025).

[17] European Commission. Directive 2005/36/EC of the European Parliament and of the Council of 7 September 2005 on the recognition of professional qualifications (Text with EEA relevance). 2005. https://eur-lex.europa.eu/legal-content/EN/ALL/?uri=CELEX:32005L0036 (accessed December 30, 2025).

[18] Serghiou S, Contopoulos-Ioannidis DG, Boyack KW, Riedel N, Wallach JD, Ioannidis JPA. Assessment of transparency indicators across the biomedical literature: How open is open? PLoS Biol 2021;19:e3001107. https://doi.org/10.1371/journal.pbio.3001107.

[19] Sterne JAC, Savović J, Page MJ, Elbers RG, Blencowe NS, Boutron I, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. BMJ 2019:l4898. https://doi.org/10.1136/bmj.l4898.

[20] Anthony Hevia, Sanjana Chintalapati, Veronica Ka Wai Lai, Thanh Tam Nguyen, Wai-Tat Wong, Terry Klassen, et al. ROBOTO2: An interactive system and dataset for LLM-assisted clinical trial risk of bias assessment (pre-print). ArXiv 2025.

[21] McGuinness LA, Higgins JPT. Risk‐of‐bias VISualization (robvis): An R package and Shiny web app for visualizing risk‐of‐bias assessments. Res Synth Methods 2021;12:55–61. https://doi.org/10.1002/jrsm.1411.

[22] Benjamin DJ, Berger JO, Johannesson M, Nosek BA, Wagenmakers E-J, Berk R, et al. Redefine statistical significance. Nat Hum Behav 2017;2:6–10. https://doi.org/10.1038/s41562-017-0189-z.

[23] Mani S, Lalani SR, Pammi M. Genomics and multiomics in the age of precision medicine. Pediatr Res 2025;97:1399–410. https://doi.org/10.1038/s41390-025-04021-0.

[24] Zhao J, Feng Q, Wei W-Q. Integration of Omics and Phenotypic Data for Precision Medicine, 2022, p. 19–35. https://doi.org/10.1007/978-1-0716-2265-0_2.

[25] Ginsburg GS, Willard HF. Genomic and personalized medicine: foundations and applications. Translational Research 2009;154:277–87. https://doi.org/10.1016/j.trsl.2009.09.005.

[26] Fujiwara Y, Kato S, Kurzrock R. Evolution of Precision Oncology, Personalized Medicine, and Molecular Tumor Boards. Surg Oncol Clin N Am 2024;33:197–216. https://doi.org/10.1016/j.soc.2023.12.004.

[27] Bode AM, Dong Z. Precision oncology- the future of personalized cancer medicine? NPJ Precis Oncol 2017;1:2. https://doi.org/10.1038/s41698-017-0010-5.

[28] Russo L, Siena LM, Farina S, Pastorino R, Boccia S, Ioannidis JPA. High-impact trials with genetic and -omics information focus on cancer mutations, are industry-funded, and less transparent. J Clin Epidemiol 2025;180:111676. https://doi.org/10.1016/j.jclinepi.2025.111676.

[29] Yurkovich JT, Evans SJ, Rappaport N, Boore JL, Lovejoy JC, Price ND, et al. The transition from genomics to phenomics in personalized population health. Nat Rev Genet 2024;25:286–302. https://doi.org/10.1038/s41576-023-00674-x.

[30] Ramaswami R, Bayer R, Galea S. Precision Medicine from a Public Health Perspective. Annu Rev Public Health 2018;39:153–68. https://doi.org/10.1146/annurev-publhealth-040617-014158.

[31] Sarigiannis D, Karakitsios S, Anesti O, Stem A, Valvi D, Sumner SCJ, et al. Advancing translational exposomics: bridging genome, exposome and personalized medicine. Hum Genomics 2025;19:48. https://doi.org/10.1186/s40246-025-00761-6.

[32] Krause KR, Ravaud P, Tran V-T. Integrating personomics into precision medicine. J Clin Epidemiol 2025;184:111859. https://doi.org/10.1016/j.jclinepi.2025.111859.

[33] Domenach P, Krause KR, Malmartel A, Ravaud P, Tran V-T. Identifying psychosocial and contextual markers considered by physicians to personalize care. BMC Med 2024;22:415. https://doi.org/10.1186/s12916-024-03616-4.

[34] Davidson KW, Silverstein M, Cheung K, Paluch RA, Epstein LH. Experimental Designs to Optimize Treatments for Individuals. JAMA Pediatr 2021;175:404. https://doi.org/10.1001/jamapediatrics.2020.5801.

[35] Lee KM, Turner RM, Thwaites GE, Walker AS, White IR. The Personalised Randomized Controlled Trial: Evaluation of a new trial design. Stat Med 2023;42:1156–70. https://doi.org/10.1002/sim.9663.

[36] French MA, Roemmich RT, Daley K, Beier M, Penttinen S, Raghavan P, et al. Precision Rehabilitation: Optimizing Function, Adding Value to Health Care. Arch Phys Med Rehabil 2022;103:1233–9. https://doi.org/10.1016/j.apmr.2022.01.154.

[37] Strohacker K, Sudeck G, Keegan R, Ibrahim AH, Beaumont CT. Contextualising flexible nonlinear periodization as a person-adaptive behavioral model for exercise maintenance. Health Psychol Rev 2024;18:285–98. https://doi.org/10.1080/17437199.2023.2233592.

[38] Goddard SG, Dossetor J, Barry S, Lawrence A, Stevens CJ, Swann C. "It Took Away the Trauma of Failing": A Mixed Methods Feasibility Trial of an Open Goals Physical Activity Program. Res Q Exerc Sport 2025;96:389–400. https://doi.org/10.1080/02701367.2024.2412661.

[39] Xu X, Zhang G, Xia Y, Xie H, Ding Z, Wang H, et al. Influencing Factors and Implementation Pathways of Adherence Behavior in Intelligent Personalized Exercise Prescription: Qualitative Study. JMIR Mhealth Uhealth 2024;12:e59610. https://doi.org/10.2196/59610.

[40] Adams MA, Todd M, Angadi SS, Hurley JC, Stecher C, Berardi V, et al. Adaptive Goals and Reinforcement Timing to Increase Physical Activity in Adults: A Factorial Randomized Trial. Am J Prev Med 2022;62:e57–68. https://doi.org/10.1016/j.amepre.2021.09.014.

[41] Malmartel A, Ravaud P, Ghosn L, Tran V-T. A classification of methods used to personalize participative interventions revealed inadequate reporting in trial protocols. J Clin Epidemiol 2021;133:80–93. https://doi.org/10.1016/j.jclinepi.2021.01.004.

[42] Siena LM, Papamanolis L, Siebert MJ, Bellomo RK, Ioannidis JPA. Industry Involvement and Transparency in the Most Cited Clinical Trials, 2019-2022. JAMA Netw Open 2023;6:e2343425. https://doi.org/10.1001/jamanetworkopen.2023.43425.

[43] Naudet F, Siebert M, Pellen C, Gaba J, Axfors C, Cristea I, et al. Medical journal requirements for clinical trial data sharing: Ripe for improvement. PLoS Med 2021;18:e1003844. https://doi.org/10.1371/journal.pmed.1003844.

[44] Shang B-H, Yang F-H, Lin Y, Bialka S, Rensburg DCJ van, Tonelli AR, et al. Discrepancies between pre-specified and reported primary outcomes: A cross-sectional analysis of randomized controlled trials in gastroenterology and hepatology journals. PLoS One 2024;19:e0305027. https://doi.org/10.1371/journal.pone.0305027.

[45] Al-Durra M, Nolan RP, Seto E, Cafazzo JA. Prospective registration and reporting of trial number in randomised clinical trials: global cross sectional study of the adoption of ICMJE and Declaration of Helsinki recommendations. BMJ 2020:m982. https://doi.org/10.1136/bmj.m982.

[46] El Boghdadly K, Wiles MD, Atton S, Bailey CR. Adherence to guidance on registration of randomised controlled trials published in *Anaesthesia*. Anaesthesia 2018;73:556–63. https://doi.org/10.1111/anae.14103.

[47] Juengst E, McGowan ML, Fishman JR, Settersten RA. From "Personalized" to "Precision" Medicine: The Ethical and Social Implications of Rhetorical Reform in Genomic Medicine. Hastings Center Report 2016;46:21–33. https://doi.org/10.1002/hast.614.

[48] March R, Schott C. Personalized/Precision Medicine/Personalised Healthcare: The Art of Giving Different Names to the Same Thing? Per Med 2017;14:463–6. https://doi.org/10.2217/pme-2017-0073.

[49] De Grandis G, Halgunset V. Conceptual and terminological confusion around personalised medicine: a coping strategy. BMC Med Ethics 2016;17:43. https://doi.org/10.1186/s12910-016-0122-4.

[50] Tran A, Klossner Q, Crain T, Prasad V. Shifting, overlapping and expanding use of "precision oncology" terminology: a retrospective literature analysis. BMJ Open 2020;10:e036357. https://doi.org/10.1136/bmjopen-2019-036357.

[51] Janiaud P, Serghiou S, Ioannidis JPA. New clinical trial designs in the era of precision medicine: An overview of definitions, strengths, weaknesses, and current use in oncology. Cancer Treat Rev 2019;73:20–30. https://doi.org/10.1016/j.ctrv.2018.12.003.