Integrative Multimodal Subtyping, Risk of Incident Mild Cognitive Impairment, and Differential Cardiometabolic Treatment Effects: A Prospective Cohort Study in the All of Us Research Program

1   **Yinjun Zhao[1], PhD**, **Karen Marder, MD[2], MPH**, **Yuanjia Wang, PhD[1]***

1. Department of Biostatistics, Columbia University Irving Medical Center, New York, NY, USA, 10032

2. Department of Neurology, Columbia University Irving Medical Center, New York, NY, USA, 10032

*Corresponding Author:

Yuanjia Wang, PhD

Email: yw2016@cumc.columbia.edu

Abstract

Background: Cognitively unimpaired (CU) adults vary substantially in their risk of developing mild cognitive impairment (MCI), yet most subtyping approaches focus on downstream neurobiological or cognitive markers rather than upstream, modifiable risk factors. We aimed to identify clinically meaningful subgroups of CU adults defined by integrated comorbid, behavioral, and social risk profiles, and to evaluate heterogeneity in both incident MCI risk and cardiometabolic treatment effects.

Methods: We conducted a prospective cohort study of 121,322 CU adults aged ≥ 50 years from the All of Us Research Program. Baseline comorbidities, lifestyle behaviors, and social determinants of health were jointly modeled using the Bayesian Mixed Integrative Data Subtyping framework, which integrates binary and continuous modalities via modality-specific likelihoods and shared latent constructs. Subtype-specific risk of incident MCI was assessed using multivariable Cox proportional hazards models adjusting for demographics and baseline medication use. A double/debiased machine learning interactive regression model with inverse probability of censoring weights to mitigate bias from informative censoring was implemented to estimate the average treatment effects of antihypertensive agents, Glucagon-Like Peptide (GLP) receptor agonists, and non–GLP antidiabetic medications on time to MCI.

Results: Four distinct subtypes were identified: I low-risk healthy aging, II

behavioral/social vulnerability, III cardiometabolic–depressive multimorbidity, and IV

mixed social-medical vulnerability profiles. Compared with Subtype I, Subtype III

demonstrated the highest risk of incident MCI (HR: 3.69, 95% CI: 3.14–4.33), followed

by Subtype IV and Subtype II. In treatment effect analyses, antihypertensive use was

associated with a modest prolongation of MCI-free survival overall (time ratio:1.04, 95%

CI: 1.03–1.06), with the largest benefit observed in Subtype III (time ratio: 1.14, 95% CI:

1.09–1.19). Non–GLP antidiabetic therapies were similarly associated with modest

overall delay, with significant benefits in Subtypes I and III. GLP-class therapies were

not associated with overall delay but showed a significant association in Subtype III.

Conclusions: Integrative subtyping based on comorbid, behavioral, and social risk

factors reveals clinically meaningful heterogeneity in both cognitive risk and treatment

response. Aligning dementia prevention strategies with dominant vulnerability pathways

may enhance the effectiveness and equity of population-level precision prevention.

Key Words: multi-modal integrative subtyping; differential cognitive risk; treatment

effect heterogeneity; All of Us Research Program

Background

Mild cognitive impairment (MCI) is a common intermediate stage between normal aging

and dementia, affecting approximately 15–20% of U.S. adults aged 60 years or older(1).

Because disease-modifying therapies for dementia remain limited, with only two

monoclonal antibody treatments currently approved and in clinical use after the

withdrawal of aducanumab from the market(2), prevention efforts increasingly focus on

identifying cognitively unimpaired (CU) individuals at elevated risk for future cognitive

decline. In this context, accurate risk stratification among CU adults is central to

targeted prevention, trial enrichment, and efficient allocation of preventive resources.

Cognitive impairment and dementia are influenced by a broad range of modifiable risk

factors, including cardiometabolic disease, depression, sensory impairment, social

isolation, and socioeconomic disadvantage(2,3). A growing body of data-driven

research has demonstrated substantial heterogeneity among CU individuals with

respect to subsequent risk of MCI and dementia. Most existing subtyping studies in CU

populations have concentrated on neurobiological and cognitive modalities, including

structural magnetic resonance imaging, such as cortical thickness and regional brain

volume; amyloid and tau positron emission tomography; fluid biomarkers, including

cerebrospinal fluid and blood-based biomarkers; and detailed neuropsychological

profiles(4–7). Using unsupervised clustering, latent class models, and disease-

progression frameworks, these studies have shown that individuals who are clinically

normal at baseline may nonetheless occupy distinct biomarker- or cognition-defined

subgroups with markedly different longitudinal trajectories and rates of conversion.

However, by focusing on downstream pathology, these approaches offer limited insight

into upstream, modifiable drivers of risk and have restricted utility for population-level

prevention.

In contrast, epidemiologic research has consistently implicated comorbid disease

burden, lifestyle behaviors, and social determinants of health (SDOH) as central

contributors to cognitive aging. Multimorbidity, particularly cardiometabolic and vascular

conditions, has been robustly associated with elevated risk of incident MCI and

dementia in large population-based cohorts(8–10). Lifestyle factors such as smoking,

physical inactivity, and obesity further accelerate cognitive decline and account for a substantial proportion of potentially preventable dementia cases(11). Socioeconomic disadvantage, neighborhood deprivation, and psychosocial stress have likewise been linked to increased dementia incidence and faster cognitive aging(2,12). Despite this strong evidence base, most studies treat these factors as covariates or examine them in isolation, emphasizing population-average associations and obscuring the co-occurring vulnerability profiles and distinct risk pathways present in real-world populations(2,13,14).

The All of Us Research Program provides a unique opportunity to address this gap. As a national precision-medicine initiative designed to reflect the diversity of the U.S. population, All of Us integrates electronic health records with detailed lifestyle and SDOH survey data, enabling large-scale evaluation of how biological, behavioral, and social factors jointly shape cognitive risk(15). However, these data are high-dimensional, correlated, and measured on mixed scales, posing challenges for conventional clustering and feature-based multimodal methods that are not designed to jointly model heterogeneous data types. To address these challenges, we apply a Bayesian integrative subtyping framework (MINDS) that jointly models binary and continuous variables and identifies latent constructs capturing shared variation across multimodal domains(16). By integrating comorbidity burden, lifestyle behaviors, and SDOH indicators into a unified latent space, MINDS enables discovery of interpretable subtypes that reflect real-world vulnerability profiles rather than single-modality patterns. In this study, we sought to (1) identify clinically meaningful subtypes among cognitively unimpaired adults using integrated modeling of comorbidities, lifestyle factors, and

SDOH; (2) evaluate heterogeneity in incident MCI risk across these subtypes; and (3)

assess whether commonly used cardiometabolic therapies antihypertensive, GLP-1

receptor agonist , and non–GLP antidiabetic medications demonstrate differential

associations with time to MCI across vulnerability profiles. We focused on these

medications because hypertension and type 2 diabetes are established upstream

cardiometabolic risk factors for cognitive decline and dementia(2,13,17,18). These

therapies have shown potential to delay cognitive impairment in both randomized trials

and large meta-analyses(14,19–21). By explicitly modeling heterogeneity in both

baseline risk and treatment responsiveness, this work advances a population-level

framework for precision dementia prevention and contributes to understanding how

upstream, modifiable risk factors jointly shape cognitive aging.

Methods

Integrative subtyping framework: MINDS

To identify data-driven subtypes that capture heterogeneity across comorbidities,

lifestyle behaviors, and social determinants of health, we employed the Mixed

INtegrative Data Subtyping (MINDS) framework, a Bayesian hierarchical model for joint

analysis of mixed-type data(16). MINDS was originally developed to integrate

multimodal clinical, cognitive, and neuroimaging data for psychiatric subtyping and is

designed to perform simultaneous dimension reduction and clustering while preserving

shared structure across heterogeneous modalities.

At its core, MINDS models each data modality through a modality-specific likelihood, i.e.,

Bernoulli for binary indicators and Gaussian for continuous measures, linked by shared

cluster-specific latent centers and subject-specific latent construct. Individuals are

assumed to belong to one of the latent clusters, each characterized by distinct mean

profiles in the latent construct space, while allowing subject-specific deviations. This

formulation enables coherent probabilistic integration of diverse data types without

requiring ad hoc transformations or two-stage clustering procedures. Computationally,

MINDS leverages Pólya–Gamma data augmentation to yield conditionally conjugate

posterior distributions for binary outcomes, enabling efficient Gibbs sampling for

posterior inference(22). The number of subtypes is selected using a Bayesian

information criterion (IC) that balances model fit and parsimony, as described in the

original methodological work(16).

Cohort for subtyping

We applied MINDS to identify data-driven subtypes among participants in the All*of*Us

Research Program by jointly integrating comorbidities, SDOH, and lifestyle factors. The

Curated Data Repository Version 8, released by the All*of*Us Research Program and

containing participant data through October 1, 2023, served as the source for all

analyses(23). We began with all eligible participants aged 50 years or older and linked

their electronic health records (EHR), SDOH survey responses, and lifestyle survey

data. Participants missing an entire modality at baseline or with documented cognitive

function finding at baseline were excluded to reduce potential bias arising from

systematically different health or survey-response profiles. Participants with any

recorded cognitive function finding were defined as the SNOMED CT parent concept

"Cognitive function finding" (OMOP concept id 4162723) and all its descendant

concepts, identified in the OMOP CONDITION domain. This definition captures

clinician-documented cognitive abnormalities (e.g., memory, attention, or thought

findings). The final analytic cohort consisted of 121,322 individuals with complete

multimodal data and follow-ups to assess incident MCI. Figure 1 depicts the derivation

of this cohort.

Comorbidity status at baseline including stroke, type 2 diabetes, heart failure, hearing

loss, depression, hypertension, and brain injury was obtained from EHR data. Lifestyle

behaviors were extracted from the lifestyle survey. Among current drinkers, we

constructed the AUDIT–C score (range, 0–12) using three items assessing (1) drinking

frequency, (2) typical quantity consumed per drinking occasion, and (3) frequency of

binge drinking (defined as consuming 6 or more drinks on a single occasion). Each item

included five ordered response options scored from 0 (lower consumption) to 4 (higher

consumption). Alcohol misuse was defined as an AUDIT–C score of 3 or higher for

women and 4 or higher for men(24). Current smoker status was obtained from the

corresponding lifestyle survey question. SDOH measures were obtained from validated

instruments included in the SDOH survey. These measures encompassed four

conceptual domains: social and community context, economic stability, neighborhood

and built environment, and health and healthcare(25). Constructs such as social

cohesion, social support, loneliness, perceived discrimination, perceived stress, and

daily spiritual experiences were assessed through established multi-item scales.

Economic stability was evaluated using indicators of food insecurity, housing instability,

and housing quality. Neighborhood characteristics, including physical and social

disorder, walk ability, and crime, were captured using standardized environmental

assessment tools, and perceived discrimination in healthcare settings was assessed

using a dedicated subscale. Higher values of the SDOH measures reflected worse

situation, consistent with the direction of comorbidity and lifestyle measures. Detailed

descriptions of each construct and measurement instrument are provided in **Error!**

Reference source not found.. Collectively, these multidimensional measures served

as inputs for the subtyping analysis.

MCI association analysis

To investigate the prognostic relevance of the identified subtypes, we conducted an

exploratory survival analysis to examine the association between subtypes and the

subsequent risk of developing MCI. Time-to-event was defined as the interval from

cohort entry to the first documented diagnosis of MCI or censoring, whichever occurred

earlier. Kaplan–Meier curves were generated to provide an initial descriptive

comparison of incident MCI across clusters. We then fit Cox proportional hazards

models to estimate hazard ratios (HRs) and 95% confidence intervals (CIs) for MCI

associated with each subtype, controlling for demographics and antihypertensive, non-

GLP antidiabetic and GLP class medication use at baseline. The medication use was

ascertained from electronic health records based on prescriptions recorded within one

year before the baseline assessment. Participants with evidence of use of any

medication in a given class during this one-year window before baseline were classified

as users of that medication class. Details of the agents that were included in the

medication class are in **Error! Reference source not found.**.

Treatment effect analysis

We estimated the causal effects of antihypertensives, non–GLP antidiabetic, and GLP-class medications use on time to incident MCI using a double/debiased machine learning (DoubleML) interactive regression model (IRM)(26). DoubleML is a causal inference approach designed for observational studies with complex, high-dimensional confounding. The method leverages flexible machine-learning algorithms to model both treatment assignment and outcome processes ("double") and combines them in a way that reduces bias from model misspecification ("debiased"). It ensures valid causal inference when either the treatment model or the outcome model is correctly specified, increasing robustness in observational data analyses. Medication use was treated as a binary exposure, and confounding variables included age, sex, race/ethnicity, educational attainment, and baseline cardiometabolic comorbidities.

The primary estimand was the average treatment effect (ATE), defined as the expected difference in outcomes had all individuals received versus not received the treatment. To assess heterogeneity in treatment response, we additionally estimated group average treatment effects (GATEs), which quantify treatment effects within each data-driven subtype identified by the integrative subtyping model. The outcome was defined as the logarithm of observed follow-up time to MCI, such that treatment effects can be interpreted on a multiplicative time scale. Exponentiating the estimated effects yields a geometric mean time ratio, representing the relative prolongation (or shortening) of MCI-free survival associated with medication use.

To account for right censoring, we incorporated inverse probability of censoring weights (IPCW). Specifically, we estimated the probability of remaining uncensored over time

213   conditional on treatment status and covariates, and weighted complete observations by

214   the inverse of this probability. This procedure reduces bias due to informative censoring

215   under standard assumptions(27). A fixed cutoff time of 48 months was used to define

216   the at-risk set, including individuals who either experienced the event before this time or

217   remained under observation beyond it.

Results

Subtyping results

We evaluated models with the number of clusters ranging from $k = 3$ to $k = 9$ using

IC(28) to determine the optimal cluster structure. As shown in Supple. **Error! Reference**

source not found.(a), the IC reached its minimum at $k = 4$, indicating that the four-

cluster model achieved the best balance between model fit and parsimony. To assess

convergence and model stability, we examined posterior log-likelihood trace plot in

Supple. **Error! Reference source not found.**(b), which demonstrated stable

convergence and consistent mixing of the Markov chain Monte Carlo (MCMC) samples

after the burn-in period. These results suggest that the four-cluster solution provides a

well-fitted and robust model for integrating the multimodal data.

As shown in Figure 2, the four subtypes demonstrated distinct and coherent patterns

across comorbidity burden, lifestyle risk behaviors, and social determinants of health.

Subtype I reflected a broadly healthy profile, characterized by minimal chronic disease,

low rates of smoking and alcohol misuse, and consistently favorable social and

environmental conditions, suggesting a Low-Risk Healthy Aging group. Subtype II

showed markedly elevated smoking and alcohol misuse alongside the highest levels of

social and economic adversity, including neighborhood disorder, discrimination, social

isolation, and housing instability, indicating a Behavioral-Risk and Socially Vulnerable subgroup. Subtype III exhibited the greatest medical burden, with pronounced hypertension, depression, hearing loss, and cardiometabolic disease, paired with moderate psychosocial stressors, aligning with a Cardiometabolic–Depressive Multimorbidity subtype. Finally, Subtype IV demonstrated intermediate levels of comorbidity and moderately elevated SDOH disadvantage, representing a Mixed-Risk Mild Vulnerability group. Together, these patterns highlight the multidimensional heterogeneity in health, lifestyle, and social context within the population. The characteristics of the four types are summarized in Table 3, which are also reflected in the summary statistics by the four subtypes in Supple. **Error! Reference source not found.**.

The radar plots provide a visual comparison of subtype differences across SDOH, lifestyle, and comorbidity measures in another way. The first plot (Figure 3) illustrates variation in social determinants of health across the four subtypes, highlighting pronounced contrasts in constructs such as discrimination in healthcare, social support, social cohesion, perceived stress, loneliness, neighborhood disorder, and material hardship (e.g., food insecurity, housing instability). Subtypes II and IV show elevated levels on multiple adverse SDOH indicators, whereas Subtype I generally reflects more favorable social and environmental conditions. The second plot (Figure 3) summarizes subtype differences in lifestyle risk factors and chronic disease burden, including smoking, alcohol misuse, and major comorbidities such as hypertension, diabetes, heart failure, stroke, and depression. Subtype III shows elevated levels on comorbidities. Together, these plots highlight distinct multidimensional profiles across subtypes,

demonstrating how combinations of social context, behavior, and health conditions

jointly characterize the heterogeneity captured by the clustering model.

The three latent constructs that MINDS estimated reflect distinct but complementary

domains of vulnerability. As shown in Figure 4, LC1 is best characterized as

Neighborhood Disorder and Safety Risk, defined by high levels of neighborhood

physical and social disorder and low perceptions of crime safety, indicating exposure to

adverse environmental conditions. LC2 represents a pattern of High-Burden Chronic

Multimorbidity, marked by strong associations with stroke, type 2 diabetes, heart failure,

hypertension, hearing loss, brain injury, and depression. LC3 reflects Psychosocial

Stress and Social Vulnerability, driven by elevated food insecurity, poor housing quality,

loneliness, discrimination (every day and within health-care settings), perceived stress,

and low social support and cohesion. See Supple. **Error! Reference source not found.**

for the exact values of the loading matrix.

Supple. Figure A.2 illustrates the distribution of the three latent constructs (LC1–LC3)

across the four subtypes, highlighting clear and interpretable subtype-specific patterns.

Subtype I shows consistently low to moderate expression of LC1 and minimal

contributions from LC2 and LC3, reflecting a comparatively low burden of adverse social

and health factors. Subtype II demonstrates a stronger LC1 profile, driven by elevated

social and psychological stressors such as loneliness, discrimination, and perceived

stress. In contrast, Subtype III is distinguished by a dominant LC2 pattern, with high

scores on multiple comorbidities including hypertension, type 2 diabetes, hearing loss,

depression, and other chronic conditions. Subtype IV exhibits the strongest expression

of LC3, characterized by markedly higher levels of neighborhood physical/social

disorder and crime safety concerns. Together, these distinct profiles demonstrate that

the subtypes are differentiated not only in magnitude but also in the composition of their

latent constructs, capturing qualitatively different combinations of social, environmental,

and health-related vulnerabilities.

Subtype's association with MCI

The models were first examined unadjusted (Supple. **Error! Reference source not**

found.) and then adjusted for relevant demographic, clinical, and medication-use

covariates(Table 2) to explore whether the observed associations persisted after

accounting for potential confounders.

The four data-driven subtypes demonstrated clear heterogeneity in their risk of

developing MCI. Using Subtype I as the reference, individuals in Subtype III exhibited

the highest risk, with HR 3.69 (95% CI: 3.14–4.33), indicating more than a threefold

increase in the hazard of MCI. Subtype IV also showed significantly elevated risk (HR =

1.87, 95% CI: 1.48–2.35), whereas Subtype II demonstrated a more modest but still

significant increase (HR = 1.43, 95% CI: 1.07–1.91). These subtype differences were

further reflected in the Kaplan–Meier survival curves in Figure 5, where Subtype III

displayed the steepest decline in MCI-free survival, followed by Subtype IV and Subtype

II, while Subtype I maintained the most favorable survival trajectory. Together, these

findings indicate that Subtype III represents a particularly vulnerable group with

accelerated cognitive decline, whereas Subtype I appears comparatively resilient, with

Subtypes II and IV occupying intermediate risk profiles.

Causal treatment effect

Figure 6 displays the estimated exponential ATEs of antihypertensive, non–GLP

antidiabetic, and GLP-class medications on time to incident MCI in the overall cohort

and across four data-driven subtypes. Antihypertensive use led a modestly longer time

to MCI overall (time ratio = 1.04; 95% CI, 1.03–1.06), corresponding to an

approximately 4% increase in geometric-mean follow-up time(months). Subtype III

showed a markedly stronger association (time ratio = 1.14; 95% CI, 1.09–1.19; ~14%

longer time), while Subtype IV demonstrated a smaller but statistically significant

association (time ratio = 1.04; 95% CI, 1.00–1.08). No significant associations were

observed for Subtypes I or II.

For non–GLP antidiabetic medications, the overall association was similarly modest but

statistically significant (time ratio = 1.04; 95% CI, 1.02–1.05; ~4% longer time). At the

subtype level, significant associations were observed in Subtype I (time ratio = 1.03; 95%

CI, 1.01–1.05; ~3% longer time) and Subtype III (time ratio = 1.05; 95% CI, 1.01–1.09;

~5% longer time), whereas Subtypes II and IV showed no evidence of benefit. GLP-

class medications were not associated with time to MCI overall; however, Subtype III

demonstrated a significant association (time ratio = 1.08; 95% CI, 1.02–1.16; ~8%

longer time). Overall, these results indicate pronounced heterogeneity in medication-

associated delay of MCI onset, with Subtype III consistently exhibiting the largest

estimated benefit across medication classes.

Discussion

In this large and diverse cohort from the All of Us Research Program, we identified four

data-driven subtypes of cognitively unimpaired adults using integrative modeling of

comorbidities, lifestyle behaviors, and SDOH. These subtypes captured distinct

multidimensional vulnerability profiles that were associated with heterogeneous risk of

incident MCI and have differential estimated effect of antihypertensive and antidiabetic

medication use. By integrating correlated biological, behavioral, and social risk factors

into coherent latent profiles, this study provides a population-level framework for

understanding heterogeneity in cognitive aging and offers insight into why

cardiometabolic interventions have yielded modest and inconsistent average effects in

prior dementia-prevention studies(2,10,29).

The four identified subtypes reflected distinct configurations of medical burden,

behavioral risk, and social disadvantage. Subtype I represented a low-risk healthy aging

profile, characterized by minimal chronic disease, low smoking and alcohol misuse, and

favorable social conditions. Subtype II was marked by prominent behavioral risk and

psychosocial adversity, but relatively limited cardiometabolic burden. Subtype III

exhibited concentrated cardiometabolic and depressive multimorbidity, particularly

hypertension, diabetes, hearing loss, and depression, with comparatively less social

disadvantage. Subtype IV combined moderate medical comorbidity with pronounced

social and environmental adversity, including neighborhood disorder, food insecurity,

and housing instability.

These patterns align with prior epidemiologic studies of comorbidity and social

disadvantage associated with adverse cognitive outcomes(8,12,30). Unlike studies that

model comorbidities or SDOH as covariates or examine single domains in

isolation(2,31–33), our integrative approach treats these factors as defining dimensions

of heterogeneity, yielding subtypes that more closely reflect real-world co-occurring risk

profiles.

Consistent with epidemiologic evidence, the subtypes showed marked differences in

risk of incident MCI. Subtype III exhibited the highest hazard, aligning with evidence that

cardiometabolic and vascular disease burden strongly predicts cognitive impairment

and dementia(8,32,34). Subtype IV also had substantially elevated risk, supporting

links between chronic social and environmental disadvantage and accelerated cognitive

aging (2,33,35). In contrast, Subtype II showed a more modest increase in risk, while

Subtype I maintained the most favorable MCI-free survival trajectory.

Notably, the relatively modest MCI risk observed in Subtype II, despite substantial

behavioral and psychosocial adversity, may reflect delayed manifestation of stress-

related effects, unmeasured intermediate conditions, or attenuation due to competing

risks, differential healthcare engagement, or under-ascertainment of cognitive

diagnoses(12,30,35).

We observed substantial heterogeneity in the estimated effect of cardiometabolic

medication use. Subtype III consistently exhibited the largest estimated effect by

antihypertensive, non–GLP antidiabetic, and GLP-class medications, suggesting that

pharmacologic modification of cardiometabolic risk factors may be most impactful

among individuals with substantial underlying comorbidity burden. This pattern is

concordant with randomized evidence demonstrating that intensive blood pressure

control reduces incident MCI among individuals at elevated vascular risk(14) and

metformin improves memory performance(36). In contrast, Subtypes II and IV showed

limited or selective effects by pharmacologic therapy, suggesting that when cognitive

risk is driven primarily by behavioral or structural adversity, medication alone may be

insufficient to meaningfully alter trajectories. Subtype I showed modest estimated

effects by non–GLP antidiabetic medications despite low baseline MCI risk, raising the

possibility that some cardiometabolic therapies may influence cognitive aging through

mechanisms beyond prevention of vascular events, such as effects on insulin signaling,

inflammation, or cerebral metabolism(21,37). This interpretation aligns with metformin

clinical trials in cognitive aging, including Metformin in Mild Cognitive Impairment and

TAME, which target insulin resistance and inflammation as upstream mechanisms of

neurodegeneration (36,38).

Importantly, these findings should be interpreted as estimated causal effects carefully.

Although we applied a double/debiased machine learning causal framework with

inverse probability of censoring weights to mitigate confounding and selection bias,

residual confounding related to disease severity, medication indication, adherence,

healthcare access, and treatment duration cannot be excluded. Failure to account for

underlying vulnerability profiles may contribute to the modest and inconsistent average

effects reported in prior observational and interventional studies(10,21).

The latent constructs identified by the integrative model highlight distinct yet interacting

pathways linking subtype profiles to cognitive risk and inform targeted prevention

strategies. The construct dominated by cardiometabolic and depressive multimorbidity

likely reflects vascular, metabolic, and inflammatory mechanisms contributing to

neurodegeneration(34), suggesting that individuals in Subtype III may benefit most from

intensified cardiometabolic and neurological disease management. In contrast, the

construct characterized by psychosocial stress, discrimination, loneliness, and material

hardship may act through chronic stress–related pathways, including hypothalamic–

pituitary–adrenal axis dysregulation and reduced cognitive reserve(2), indicating that

Subtype II may respond best to psychosocial and community-based interventions. The

construct reflecting neighborhood disorder and safety concerns likely affects cognition

indirectly by limiting physical activity, social engagement, and access to health-

promoting resources(35), underscoring the need for structural interventions in Subtype

IV. These pathways likely interact over the life course, reinforcing the value of

multidimensional risk stratification for mechanism-informed and targeted dementia

prevention.

Together, these findings underscore that dementia prevention strategies should be

aligned with dominant vulnerability pathways rather than applied uniformly across

heterogeneous populations. Integrative subtyping frameworks embedded within large-

scale health systems can identify individuals most likely to benefit from pharmacologic

modification of upstream cardiometabolic risk factors, while distinguishing subgroups for

whom behavioral or structural interventions addressing psychosocial and environmental

risk may be more relevant. By explicitly modeling heterogeneity in both baseline

vulnerability and treatment responsiveness, this approach supports subgroup-enriched

pragmatic trials and advances a more mechanism-informed, targeted, and equitable

framework for precision prevention in cognitive aging.

Importantly, this population-level perspective is complementary to biomarker-based

approaches. Whereas biomarkers capture downstream, disease-specific biological

processes, integrative subtyping focuses on upstream vulnerability relevant to early risk

stratification and prevention, together providing a more comprehensive foundation for

dementia prevention across the life course(2).

Several limitations warrant consideration. First, the analysis focused on all-cause MCI,

and information on etiologic MCI subtypes and severity was unavailable, as

standardized cognitive test scores or continuous measures of cognitive performance

were not incorporated. Second, cognitive, neuroimaging, and molecular biomarkers

were not included; accordingly, the identified subtypes should be interpreted as

population-level vulnerability profiles rather than direct proxies for underlying

neuropathology(7). Third, medication exposure was defined using baseline prescription

records and did not capture dose, adherence, or changes over follow-up. Fourth,

subtypes were derived from baseline measures and therefore do not capture

longitudinal changes that may influence MCI risk or treatment responsiveness. Also,

although the All of Us cohort enhances generalizability through its size and diversity(15),

healthy volunteer bias, differential healthcare access, and diagnostic ascertainment may

affect absolute risk estimates. Last, subtype definitions are model-dependent, and

alternative modeling choices or the inclusion of additional modalities could yield different

subgroup structures. Independent validation in external cohorts such as the UK Biobank

is therefore needed to assess the robustness, reproducibility, and generalizability of the

identified subtypes.

Conclusion

In summary, this study demonstrates that integrative, data-driven subtyping based on

comorbidities, lifestyle behaviors, and social determinants of health can uncover

clinically and socially meaningful heterogeneity in cognitive aging trajectories. The

identified subtypes differed substantially in risk of incident MCI and in estimated

associations with cardiometabolic therapies, highlighting the importance of aligning

prevention strategies with dominant vulnerability pathways. By explicitly recognizing that

cognitive risk emerges from interacting biological mechanisms, modifiable health

behaviors, and structural determinants, this multidimensional framework provides a

foundation for more targeted, equitable, and effective approaches to dementia

prevention.

446   List of abbreviations

Abbreviation	Definition
ATE	Average Treatment Effect
AUDIT-C	Alcohol Use Disorders Identification Test–Consumption
CI	Confidence Interval
CU	Cognitively Unimpaired
DoubleML	Double/Debiased Machine Learning
EHR	Electronic Health Record
GATE	Group Average Treatment Effect
GLP	Glucagon-Like Peptide
HR	Hazard Ratio
IC	Information Criterion
IRM	Interactive Regression Model
LC	Latent Construct
MCI	Mild Cognitive Impairment
MCMC	Markov Chain Monte Carlo
MINDS	Mixed Integrative Data Subtyping
OMOP	Observational Medical Outcomes Partnership
PANES	Physical Activity Neighborhood Environment Scale
SDOH	Social Determinants of Health
SNOMED CT	Systematized Nomenclature of Medicine Clinical Terms

Declarations

Ethics approval and consent to participate

Non applicable.

Consent for publication

Non applicable.

Availability of data and materials

The data that support the findings of this study are available from the National Institutes of Health's All of Us Research Program upon reasonable request and with permission of the All of Us Research Program.

Competing interests

Non applicable.

Funding

This work was supported by the National Institute of Mental Health [MH123487], National Institute of Neurological Disorders and Stroke [NS073671], and National Institute of Health [TL1TR001875].

Authors' contributions

YZ conducted the analyses and drafted the manuscript. KM contributed to clinical interpretation and critical revisions. YW conceived and supervised the study and revised the manuscript. All authors approved the final version.

Acknowledgements

We gratefully acknowledge All of Us participants for their contributions, without whom this research would not have been possible. We also thank the National Institutes of

469   Health's All of Us Research Program for making available the participant data examined

470   in this study.

Reference

1. Petersen RC. Mild Cognitive Impairment. Continuum. 2016 Apr;22(2):404–18. doi:10.1212/CON.0000000000000313

2. Livingston G, Huntley J, Sommerlad A, Ames D, Ballard C, Banerjee S, et al. Dementia prevention, intervention, and care: 2020 report of the Lancet Commission. The Lancet. 2020 Aug 8;396(10248):413–46. doi:10.1016/S0140-6736(20)30367-6 PubMed PMID: 32738937.

3. Kivimäki M, Singh-Manoux A, Pentti J, Sabia S, Nyberg ST, Alfredsson L, et al. Physical inactivity, cardiometabolic disease, and risk of dementia: an individual-participant meta-analysis. BMJ. 2019 Apr 17;365:l1495. doi:10.1136/bmj.l1495 PubMed PMID: 30995986; PubMed Central PMCID: PMC6468884.

4. Estarellas M, Oxtoby NP, Schott JM, Alexander DC, Young AL, for the Alzheimer's Disease Neuroimaging Initiative. Multimodal subtypes identified in Alzheimer's Disease Neuroimaging Initiative participants by missing-data-enabled subtype and stage inference. Brain Commun. 2024 Aug 1;6(4):fcae219. doi:10.1093/braincomms/fcae219

5. Capogna E, Sørensen Ø, Watne LO, Roe J, Strømstad M, Idland AV, et al. Subtypes of brain change in aging and their associations with cognition and Alzheimer's disease biomarkers. Neurobiol Aging. 2025 Mar;147:124–40. doi:10.1016/j.neurobiolaging.2024.12.009 PubMed PMID: 39740372.

6. Dadar M, Shafiee N, Collins L, Camicioli R, Duchesne S. Subtyping mild cognitive impairment based on imaging and CSF biomarker levels. Alzheimer's & Dementia. 2021 Dec;17(S4):e054129. doi:10.1002/alz.054129

7. Jack Jr CR, Bennett DA, Blennow K, Carrillo MC, Dunn B, Haeberlein SB, et al. NIA☐AA research framework: toward a biological definition of Alzheimer's disease. Alzheimer's & dementia. 2018;14(4):535–62. doi:DOI: 10.1016/j.jalz.2018.02.018

8. Kivimäki M, Singh-Manoux A, Pentti J, Sabia S, Nyberg ST, Alfredsson L, et al. Physical inactivity, cardiometabolic disease, and risk of dementia: an individual-participant meta-analysis. bmj. 2019;365.

9. Katabathula S, Davis PB, Xu R. Comorbidity☐driven multi☐modal subtype analysis in mild cognitive impairment of Alzheimer's disease. Alzheimer's & Dementia. 2023 Apr;19(4):1428–39. doi:10.1002/alz.12792

10. Calvin CM, Conroy MC, Moore SF, Kuźma E, Littlejohns TJ. Association of multimorbidity, disease clusters, and modification by genetic factors with risk of dementia. JAMA network open. 2022;5(9):e2232124–e2232124.

11. Barnes DE, Yaffe K. The projected effect of risk factor reduction on Alzheimer's disease prevalence. The Lancet Neurology. 2011;10(9):819–28.

12. Wang K, Fang Y, Zheng R, Zhao X, Wang S, Lu J, et al. Associations of socioeconomic
status and healthy lifestyle with incident dementia and cognitive decline: two prospective
cohort studies. eClinicalMedicine. 2024 Oct 1;76. doi:10.1016/j.eclinm.2024.102831
PubMed PMID: 39290638.

13. Ou YN, Tan CC, Shen XN, Xu W, Hou XH, Dong Q, et al. Blood Pressure and Risks of
Cognitive Impairment and Dementia. Hypertension. 2020 Jul;76(1):217–25.
doi:10.1161/HYPERTENSIONAHA.120.14993

14. The SPRINT MIND Investigators for the SPRINT Research Group. Effect of Intensive vs
Standard Blood Pressure Control on Probable Dementia: A Randomized Clinical Trial.
JAMA. 2019 Feb 12;321(6):553–61. doi:10.1001/jama.2018.21442

15. Investigators A of URP. The "All of Us" research program. New England Journal of
Medicine. 2019;381(7):668–76.

16. Zhao Y, Wang Y, Liu Y. An Integrative Approach for Subtyping Mental Disorders Using
Multimodal Data [Internet]. medRxiv; 2025 [cited 2026 Jan 29]. p. 2025.05.27.25328416.
Available from: https://www.medrxiv.org/content/10.1101/2025.05.27.25328416v2
doi:10.1101/2025.05.27.25328416

17. Chatterjee S, Peters SAE, Woodward M, Mejia Arango S, Batty GD, Beckett N, et al. Type 2
Diabetes as a Risk Factor for Dementia in Women Compared With Men: A Pooled Analysis
of 2.3 Million People Comprising More Than 100,000 Cases of Dementia. Diabetes Care.
2016 Feb;39(2):300–7. doi:10.2337/dc15-1588 PubMed PMID: 26681727; PubMed Central
PMCID: PMC4722942.

18. McGrath ER, Beiser AS, DeCarli C, Plourde KL, Vasan RS, Greenberg SM, et al. Blood
pressure from mid- to late life and risk of incident dementia. Neurology. 2017 Dec
12;89(24):2447–54. doi:10.1212/WNL.0000000000004741 PubMed PMID: 29117954;
PubMed Central PMCID: PMC5729797.

19. Gudala K, Bansal D, Schifano F, Bhansali A. Diabetes mellitus and risk of dementia: A
meta□analysis of prospective observational studies. J Diabetes Investig. 2013 Nov
27;4(6):640–50. doi:10.1111/jdi.12087 PubMed PMID: 24843720; PubMed Central PMCID:
PMC4020261.

20. He J, Zhao C, Zhong S, Ouyang N, Sun G, Qiao L, et al. Blood pressure reduction and all-
cause dementia in people with uncontrolled hypertension: an open-label, blinded-endpoint,
cluster-randomized trial. Nat Med. 2025 Jun;31(6):2054–61. doi:10.1038/s41591-025-
03616-8

21. Pierret AC, Mizuno Y, Saunders P, Lim E, De Giorgi R, Howes OD, et al. Glucagon-like
peptide 1 receptor agonists and mental health: a systematic review and meta-analysis. JAMA
psychiatry. 2025.

22. Polson NG, Scott JG, Windle J. Bayesian Inference for Logistic Models Using Pólya–
Gamma Latent Variables. Journal of the American Statistical Association. 2013
Dec;108(504):1339–49. doi:10.1080/01621459.2013.829001

23. All of Us Research Program. Curated Data Repository (CDR) Version 8 Release Notes
[Internet]. All of Us Research Program; 2024 [cited 2026 Jan 29]. Available from:
https://support.researchallofus.org/hc/en-us/articles/30294451486356-Curated-Data-
Repository-CDR-version-8-Release-Notes

24. Bradley KA, DeBenedetti AF, Volk RJ, Williams EC, Frank D, Kivlahan DR. AUDIT–C as
a brief screen for alcohol misuse in primary care. Alcoholism: Clinical and Experimental
Research. 2007;31(7):1208–17.

25. Koleck TA, Dreisbach C, Zhang C, Grayson S, Lor M, Deng Z, et al. User guide for social
determinants of health survey data in the All of Us Research Program. Journal of the
American Medical Informatics Association. 2024;31(12):3032–41.

26. Chernozhukov V, Chetverikov D, Demirer M, Duflo E, Hansen C, Newey W, et al.
Double/debiased machine learning for treatment and structural parameters. Oxford
University Press Oxford, UK; 2018.

27. Robins JM, Finkelstein DM. Correcting for Noncompliance and Dependent Censoring in an
AIDS Clinical Trial with Inverse Probability of Censoring Weighted (IPCW) Log–Rank
Tests. Biometrics. 2000 Sep;56(3):779–88. doi:10.1111/j.0006-341X.2000.00779.x

28. Spiegelhalter DJ, Best NG, Carlin BP, Linde A. The Deviance Information Criterion: 12
Years on. J R Stat Soc Ser B Stat Methodol. 2014 Jun 1;76(3):485–93.
doi:10.1111/rssb.12062

29. Richard E, Moll van Charante EP, Hoevenaar-Blom MP, Coley N, Barbera M, van der Groep
A, et al. Healthy ageing through internet counselling in the elderly (HATICE): a
multinational, randomised controlled trial. Lancet Digit Health. 2019 Dec;1(8):e424–34.
doi:10.1016/S2589-7500(19)30153-0 PubMed PMID: 33323224.

30. Petersen JD, Wehberg S, Packness A, Svensson NH, Hyldig N, Raunsgaard S, et al.
Association of Socioeconomic Status With Dementia Diagnosis Among Older Adults in
Denmark. JAMA Netw Open. 2021 May 18;4(5):e2110432.
doi:10.1001/jamanetworkopen.2021.10432

31. Guan Y, Ebrahimzadeh SA, Cheng C hsin, Chen W, Leung T, Bigornia S, et al. Association
of Diabetes and Hypertension With Brain Structural Integrity and Cognition in the Boston
Puerto Rican Health Study Cohort. Neurology. 2022 Apr 12;98(15):e1534–44.
doi:10.1212/WNL.0000000000200120

32. Ma H, Mu X, Jin Y, Luo Y, Wu M, Han Z. Multimorbidity, lifestyle, and cognitive function:
A cross-cultural study on the role of diabetes, cardiovascular disease, cancer, and chronic
respiratory diseases. Journal of Affective Disorders. 2024 Oct 1;362:560–8.
doi:10.1016/j.jad.2024.07.053

33. Yang ZG, Sun X, Han X, Wang X, Wang L. Relationship between social determinants of
health and cognitive performance in an older American population: a cross-sectional
NHANES study. BMC Geriatr. 2025 Jan 10;25(1):25. doi:10.1186/s12877-024-05672-0
PubMed PMID: 39794733; PubMed Central PMCID: PMC11720803.

34. Gottesman RF, Albert MS, Alonso A, Coker LH, Coresh J, Davis SM, et al. Associations
between midlife vascular risk factors and 25-year incident dementia in the Atherosclerosis
Risk in Communities (ARIC) cohort. JAMA neurology. 2017;74(10):1246–54.

35. Becerril A, Pfoh ER, Hashmi AZ, Mourany L, Gunzler DD, Berg KA, et al. Racial, ethnic
and neighborhood socioeconomic differences in incidence of dementia: A regional
retrospective cohort study. Journal of the American Geriatrics Society. 2023;71(8):2406–18.
doi:10.1111/jgs.18322

36. Luchsinger JA, Perez T, Chang H, Mehta P, Steffener J, Pradabhan G, et al. Metformin in
Amnestic Mild Cognitive Impairment: results of a pilot randomized placebo controlled
clinical trial. J Alzheimers Dis. 2016;51(2):501–14. doi:10.3233/JAD-150493 PubMed
PMID: 26890736; PubMed Central PMCID: PMC5079271.

37. Yu X, He H, Wen J, Xu X, Ruan Z, Hu R, et al. Diabetes-related cognitive impairment:
mechanisms, symptoms, and treatments. Open Medicine. 2025;20(1):20241091.

38. Barzilai N, Crandall JP, Kritchevsky SB, Espeland MA. Metformin as a Tool to Target
Aging. Cell Metab. 2016 Jun 14;23(6):1060–5. doi:10.1016/j.cmet.2016.05.011 PubMed
PMID: 27304507; PubMed Central PMCID: PMC5943638.

39. Gower JC. A general coefficient of similarity and some of its properties. Biometrics.
1971;857–71.

40. Jin Y, Liang J, Hong C, Liang R, Luo Y. Cardiometabolic multimorbidity, lifestyle
behaviours, and cognitive function: a multicohort study. The lancet Healthy longevity.
2023;4(6):e265–73.

41. Pierret ACS, Mizuno Y, Saunders P, Lim E, De Giorgi R, Howes OD, et al. Glucagon-Like
Peptide 1 Receptor Agonists and Mental Health: A Systematic Review and Meta-Analysis.
JAMA Psychiatry. 2025 Jul 1;82(7):643–53. doi:10.1001/jamapsychiatry.2025.0679
PubMed PMID: 40366681; PubMed Central PMCID: PMC12079569.

Legends

612 Table 1: Characteristics of the four data-driven subtypes based on comorbidity, lifestyle

613 behaviors, and social determinants of health.

628   *Table 3: Characteristics of the four data-driven subtypes based on comorbidity, lifestyle*
629   *behaviors, and social determinants of health.*

Subtype	Key Features	Name
I	Lowest comorbidity burden; low smoking and alcohol misuse; favorable social and environmental conditions.	Low-Risk Healthy Aging
II	Intermediate chronic disease burden; moderate smoking/alcohol levels; moderate disadvantage across social determinants of health.	Mixed-Risk Mild Vulnerability
III	Greatest medical burden with pronounced hypertension, depression, hearing loss, and cardiometabolic disease; moderate psychosocial stress levels.	Cardiometabolic– Depressive Multimorbidity
IV	Highest social and economic adversity including neighborhood disorder, discrimination, social isolation, and housing instability; moderate chronic disease.	Social Vulnerability

631   *Table 4: Cox Regression Results on time to MCI by subtype, demographics, and medication use.*

Characteristic	N	HR	95% CI	p-value
Subtype				**<0.001**
I	66,422	—	—	
II	9,039	1.11	0.53, 2.32	
III	30,070	4.71	3.40, 6.52	
IV	15,791	1.96	1.17, 3.30	
Age (years)	121,322	1.85	1.69, 2.03	**<0.001**
Gender				**<0.001**
Female	74,270	—	—	
Male	47,052	1.37	1.16, 1.61	
Race				0.9
Non-White	15,914	—	—	
White	105,408	1.02	0.77, 1.35	
Ethnicity				0.2
Non-Hispanic	110,172	—	—	
Hispanic	11,150	1.22	0.88, 1.69	
Educational attainment				0.5
Below college	46,794	—	—	
College & above	74,528	0.94	0.79, 1.12	
Antihypertensive use				**<0.001**
No	80,649	—	—	
Yes	40,673	3.23	2.35, 4.42	
Non-GLP antidiabetic use				0.5
No	108,002	—	—	
Yes	13,320	1.13	0.77, 1.65	
GLP-1 class use				0.077
No	116,665	—	—	
Yes	4,657	2.16	1.01, 4.66	
Subtype × Antihypertensive use	121,322			**<0.001**
II * Yes	2,605	1.25	0.51, 3.07	
III * Yes	14,025	0.31	0.21, 0.46	
IV * Yes	5,713	0.87	0.47, 1.62	
Subtype × Non-GLP antidiabetic use	121,322			>0.9
II * Yes	842	1.19	0.45, 3.10	
III * Yes	4,576	0.94	0.57, 1.54	
IV * Yes	2,071	1.13	0.58, 2.18	
Subtype × GLP class use	121,322			0.7
II * Yes	239	0.44	0.05, 3.79	
III * Yes	2,418	0.71	0.30, 1.67	
IV * Yes	928	0.58	0.19, 1.72	

634
635 *Figure 7: Flowchart of participant selection and exclusion criteria in the All of Us cohort.*

636
637

638
639 *Figure 8: Heatmaps of binary and continuous measures across subtypes by the MINDS method,*
640 *integrating lifestyle factors, comorbidity and social determinants of health measures.*

Figure 9: Radar plot illustrating differences in SDOH and comorbidity profiles across the four identified latent subtypes (I–IV).

Figure 10: Heatmap displaying the magnitude of standardized loadings for each observed measure across the three latent constructs (LC1–LC3).

652
653 *Figure 11: Kaplan–Meier survival curves depicting the probability of remaining free from MCI*
654 *across Subtypes I–IV.*

Treatment	Subtype	Medication Use(%)	Exponential ATE (95%CI)	
Antihypertensives				
	Overall (N = 37,577)	20,535 (55%)		1.04 (1.03, 1.06)
	I (N = 19,715)	8,430 (43%)		1.00 (0.99, 1.02)
	II (N = 2,926)	1,224 (42%)		0.99 (0.95, 1.03)
	III (N = 9,725)	7,906 (81%)		1.14 (1.09, 1.19)
	IV (N = 5,211)	2,975 (57%)		1.04 (1.00, 1.08)
Non-GLP Antidiabetics				
	Overall (N = 37,577)	5,377 (14%)		1.04 (1.02, 1.05)
	I (N = 19,715)	2,373 (12%)		1.03 (1.01, 1.05)
	II (N = 2,926)	357 (12%)		1.01 (0.96, 1.06)
	III (N = 9,725)	1,766 (18%)		1.05 (1.01, 1.09)
	IV (N = 5,211)	881 (17%)		1.04 (0.99, 1.09)
GLP class				
	Overall (N = 37,577)	1,710 (4.6%)		1.02 (0.98, 1.05)
	I (N = 19,715)	366 (1.9%)		0.98 (0.94, 1.01)
	II (N = 2,926)	84 (2.9%)		1.01 (0.91, 1.12)
	III (N = 9,725)	900 (9.3%)		1.08 (1.02, 1.16)
	IV (N = 5,211)	350 (6.9%)		0.98 (0.89, 1.07)

656 *Figure 12: Estimated exponential average treatment effects for antihypertensive, non-GLP*
657 *antidiabetic and GLP class medications across the four data-driven subtypes.*

661