# Mediating Effects of Healthy Lifestyle Factors on Associations between Mental Health and Functional Outcomes in Early Adolescence

Jason Smucny, Ph.D.[1]
Tyler A. Lesh, Ph.D.[1]
Tara A. Niendam, Ph.D.[1]
Nicole R. Karcher, Ph.D.[2]

[1]Department of Psychiatry and Behavioral Sciences, University of California, Davis
[2]Department of Psychiatry, Washington University St. Louis

Correspondence:

Nicole Karcher
nkarcher@wustl.edu

**Abstract**

**Importance:** Although mental health and healthy lifestyle interventions are associated with functional outcomes in adolescence, the extent to which specific healthy lifestyle factors explain relationships between mental health and outcome is unclear. It is also unclear the extent to which environmental adversity alters the influence of these associations.

**Objective:** To examine the mediating effects of healthy lifestyle factors on the relationships between mental health at ages 10-11 and two functional outcomes measured 2-3 years later (academic functioning and social problems). Secondarily, to examine the moderating effect of environmental risk factors on mediation strength.

**Design, Setting, and Participants:** This cohort study used data from 3 waves of follow-up data from the Adolescent Brain Cognitive Development (ABCD) Study (ages 10-11, 11-12, and 12-13). Data included 6754 adolescents for the academic functioning outcome and 7550 adolescents for the social problems outcome.

**Main Outcomes and Measures:** The mediating effects of four healthy lifestyle measures (sleep quality, screen time, physical activity and Mediterranean diet) on the relationships between four mental health measures (depression, anxiety, psychotic-like experience (PLE) distress, and total problems) and two subsequent functional outcomes (academic functioning and social problems) were examined. Secondary analyses

included neighborhood safety, family conflict, school environment, and family financial adversity as moderators.

**Results:** Sleep quality was the strongest mediating factor predicting academic functioning outcome, mediating 18.5%, 36.3%, 8.3%, and 3.4% of the relationships between depression, anxiety, PLE distress and total problems, respectively. Screen time was the second strongest mediating factor. For the social problems outcome, sleep quality was the only factor that showed > 3% mediation (19.6% - 23.3% mediation). Moderation analyses found the mediating effects of sleep and screen time on academic functioning decreased as financial adversity increased. Conversely, mediating effects of sleep quality on social problems increased with worsening family conflict, financial adversity, and school environment.

**Conclusion and Relevance:** These results suggest that healthy lifestyle factors, in particular sleep quality, may partially explain the associations between mental health and functioning in adolescents. The results also provide evidence that the effect of lifestyle is modulated by environment factors, including financial adversity, family conflict, and school environment. These results may have important implications for future intervention studies.

**Introduction**

Adolescence represents a critical developmental window during which mental health concerns, including depression, anxiety, and psychotic-like experiences (PLEs), often emerge [1-3]. Globally, approximately one in seven 10- to 19-year-olds experience a mental disorder, and depression and anxiety are leading causes of illness and disability among young people [4,5]. Understanding both the functional correlates of early mental health concerns and the targetable mechanisms underlying their development is of great importance for both public health and developmental science. The current paper directly addresses this need by examining potentially targetable mechanisms linked to the emergence of early mental health concerns.

Mental health difficulties experienced in adolescence often have a number of downstream consequences, including impact on everyday functioning such as academic achievement and peer/social functioning [6,7]. Mood and anxiety disorders have been associated with school absenteeism, lower grades, and withdrawal from peer relationships [8-12]. Likewise, psychotic-like experiences (PLEs) may disrupt social engagement, peer support, and academic performance [13-15]. Failure to examine functioning alongside symptomatology risks overlooking the real-world consequences of adolescent mental health concerns and misses opportunities to intervene early and prevent these potential downstream impacts.

Given the relationship between early mental health concerns and everyday functioning, elucidating mechanisms underlying these associations becomes imperative. Among readily modifiable mechanisms, lifestyle factors, including healthy sleep, physical activity, diet quality, and screentime hold promise as leverageable targets in

prevention and intervention. Research in "lifestyle psychiatry" has underscored that physical activity, sleep, and diet contribute meaningfully to the onset, maintenance, and alleviation of mental disorders [16]. For example, higher levels of exercise have been linked to reduced depressive and anxiety symptoms in youth, and emerging evidence also points to benefits for social and academic functioning [17]. Sleep disruptions, including but not limited to insufficient duration, poor quality, or irregular timing, have been associated with increased internalizing symptoms and poorer school and interpersonal functioning in adolescents [18,19]. Diet quality, especially adherence to a Mediterranean diet, has been connected to lower levels of depressive and anxiety symptoms in children and adolescents [20,21]. Screen time presents another critical lifestyle parameter: extensive screen exposure, especially via social media and passive screen use, has been consistently tied to worse mental health outcomes in adolescents, and may further impair social and academic engagement through sleep displacement, reduced activity, and diminished in-person peer interaction [22,23]. Consistent with this, research using the Adolescent Brain Cognitive Development (ABCD) study has found evidence that increased screen time [24,25], greater sleep disturbances [26,27], and lower physical activity [28] are associated with greater endorsement of mental health concerns. The current manuscript aims to examine the untested model that these lifestyle behaviors act as mediators that link adolescent mental health (e.g., depression, anxiety, and PLEs) to everyday academic and social functioning.

It is also likely that these lifestyle mechanisms (i.e., sleep, physical activity, diet, and screen time) do not operate uniformly across all youth but rather are shaped by environmental context. For instance, environmental risk factors including financial

adversity (e.g., socioeconomic strain), neighborhood and school conditions, and/or familial conflict may moderate the strength of the pathways linking lifestyle behaviors to mental health and functioning. Adolescents growing up in economically disadvantaged conditions [29-32] or high-conflict homes [33,34] may face compounding risk, such that the same amount of sleep disruption or elevated screen time may confer greater detriment in the presence of environmental risk factors, or lifestyle buffers such as exercise may be undermined by environmental risk factors. The possibility of moderated mediation, whereby the environment (e.g., family conflict, disadvantage) influences the degree to which lifestyle factors mediate associations between symptoms with functioning, is an unexplored yet potentially compelling framework for understanding heterogeneity in adolescent risk for functioning decline.

In the present study, we leveraged data from the longitudinal, multi-site ABCD Study to test a conceptual model whereby adolescent depression, anxiety, and PLEs predict academic and social functioning, with lifestyle behaviors (screen time, sleep quality, physical activity, and diet) acting as mediators partially linking these associations. Although mediating effects of many interventions could be examined, we focused on these four lifestyle factors not only because of their established links to functioning but also because they can be addressed in outpatient therapeutic settings. Follow-up analyses explored the possibility of moderated mediation, exploring whether links between lifestyle factors and functioning were moderated by environmental risk factors (i.e., financial adversity, neighborhood safety, school quality, and familial conflict). This work aims to provide novel evidence identifying actionable targets for

early intervention and to delineate under which conditions healthy lifestyle factors are most impactful.

**Methods**

*Participants*

ABCD data used in this report came from release 5.1 (DOI: 10.15154/z563-zd24). DOIs can be found at https://nda.nih.gov/abcd/abcd-annual-releases.html. This release includes baseline data (ABCD wave 1, ages 9-10), 1-year follow-up data (wave 2, ages 10-11), 2-year follow-up data (wave 3, ages 11-12) and 3-year follow-up data (wave 4, ages 12-13). The present study primarily used 1-year follow-up data as clinical predictors, 2-year follow-up data as mediators (except for Mediterranean diet, which was only available at 1-year follow-up), and 3-year follow-up data as the outcome measure in mediation models. Potential participants were excluded from ABCD study participation for the following reasons: child not fluent in English, major neurological disorder, gestational age less than 28 weeks or birthweight less than 1,200 grams, history of traumatic brain injury, or has a current diagnosis of schizophrenia, autism spectrum disorder (moderate, severe), mental retardation/intellectual disability, or alcohol/substance use disorder. For families with siblings, only one sibling from the family was included (based on alphabetical order from the Global Unique Identifier).

*Mental Health Predictors*

The following mental health measures were included as predictors in mediation models: Child Behavior Checklist (CBCL) [6] Depression T-score, CBCL Anxiety T-score,

CBCL Total Problems T-score (a composite of all CBCL ratings), and Prodromal Questionnaire-Brief Child Version (PQ-BC) PLEs distress score sum [7]. Depression and anxiety scores were taken from the DSM-5-oriented affective problems and anxiety problems CBCL scales. All predictive clinical measures were taken from 1-year follow-up data.

*Healthy Lifestyle Mediators*

The following "healthy lifestyle" measures were included as potential mediators in analyses: diet, physical activity, screen time, and sleep quality. "Diet" was captured by the parent-reported sum of the Mediterranean-DASH Intervention for Neurodegenerative Delay (MIND) diet questionnaire (see Supplement for more details) [19]. "Physical activity" was captured by the child's answer to the question: "During the past 7 days, on how many days were you physically active for a total of at least 60 minutes per day?" (from 0 to 7). "Screen time" was captured by parent-reported screen time (up to 15 minute increments) on weekends or weekdays. Weekend and weekday screen time was examined separately because they may reflect different aspects of time allocation (i.e., recreational time allocation vs. educational ("occupational") time allocation). "Sleep quality" was captured by taking the caregiver-reported sum of the Sleep Disturbance Scale for Children (SDSC) [18] (see Supplement). Screen time and sleep quality measures were reverse coded such that greater values on all scales corresponded to greater adherence towards following healthy lifestyle habits. Physical activity, screen time, and sleep quality were taken from 2-year follow-up data. Diet information was taken from 1-year follow-up data (as it was not included as part of 2-

year follow-up data; it should be noted that year-to-year diets are unlikely to substantially change between ages 10-12 [35]).

*Functional Outcomes*

Two functional outcomes were examined: academic functioning during follow-up years 2-3, and social problems at 3-year follow-up. Parents were asked to quantify academic functioning at the 3-year follow-up visit by reporting the child's average grades during the preceding year. Academic functioning was quantified as the parental report of school grades converted to a 1-12 scale, with 1 corresponding to "A+" or 97-100%, 2 with "A" or 93-97%, 3 with "A-" or 90-92%, etc. down to 12 with "F" or < 65%. These scores were reverse coded such that higher scores were associated with higher grades. Children whose parents refused to report grades were excluded from any analyses with grades. Social problems were quantified by CBCL Social Problems T-score which includes questions regarding their getting along with other children, getting teased, feeling lonely, and acting too young for their age [36].

*Environmental Adversity Moderators*

The following environmental adversity moderators were examined: neighborhood safety, family conflict, school environment, and family financial adversity. Neighborhood safety was captured by the parental report ABCD Neighborhood Crime and Safety Survey [37] (lower scores = safer neighborhood), family conflict by the ABCD Parent Family Environment Scale-Family Conflict Subscale (modified from PhenX) [38] (higher scores = more conflict), school environment by the youth report School Risk and

Protective Factors School Environment subscale (modified from PhenX) [39] (higher

scores = better environment), and family financial adversity by the summed

endorsement of 7 parent-reported questions of financial difficulties experienced during

the past 12 months from a demographic questionnaire [40] (higher scores = more

adversity). More detailed information on these instruments is provided in the

Supplement.

*Mediation Analyses*

Mediation analyses were performed using the PROCESS v.5.0 toolbox

(processmacro.org) in SPSS v.30 (IBM). The primary analyses included eight mediation

models; each model included one of the mental health measures (i.e., either

depression, anxiety, PLE distress, or total problems) as the independent variable,

age/sex/site as covariates, the four healthy lifestyle factors (diet, physical activity,

weekend screen time, and sleep quality) as simultaneous (parallel) mediators, and one

of two functional outcome measures (academic functioning and social problems) as the

dependent variable. Total problems (CBCL Total Problems score) was not included in

models with social problems as the outcome due to overlap with CBCL Social Problems

score.  A set of secondary analyses was performed similarly except it included NIH

Cognition Toolbox Composite [41] raw score (an IQ measure) as a covariate as previous

research indicates grades [42] and social functioning [43] may be influenced by IQ. A third

set of analyses examined weekday screen time instead of weekend screen time.

Independent variables were taken from 1-year follow-up data, mediators from 2-year

follow-up (except for diet, which was only available at 1-year follow-up), and outcome

measures from 3-year follow-up (i.e., academic functioning since 2-year follow-up, and

CBCL Social Problems score at 3-year follow-up). Significance was determined by

calculating the 95% confidence interval (CI) of the indirect effect based on 5000 bias-

corrected bootstrapped samples; an indirect effect in which the corresponding 95% CI

was entirely positive or negative implies that $p < .05$. Percent mediation was also

calculated using the indirect effect for each individual mediator as well as across all

mediators combined by the equation *indirect effect ÷ total effect* × 100. Significant

indirect effects were only considered to be of interest if the total effect (between the

independent and dependent variable, including the mediator) was significant ($p < .05$).

*Moderated Mediation Effects*

Effects of environmental moderators were also examined using PROCESS v.5.0.

A diagram of the analysis strategy for environmental moderation effects is shown in

Figure 1. Briefly, these moderated mediation models examined the degree to which

each moderator (e.g., family financial adversity) affected the mediation of associations

between mental health with outcome by affecting the relationships between lifestyle

factors and outcome (i.e., the mediation model "b" path). We examined moderation of

the "b" path (i.e., the path linking lifestyle factors to outcome), as the goal of this study

was to examine everyday lifestyle mediators (e.g., screen time habits) that can be

altered independent of a clinical provider. Concordantly, we were particularly interested

to examine how the beneficial effects of these mediators might be influenced by living

conditions. Similar to the primary (non-moderated) mediation analyses, significant

moderated mediation was determined by calculating the 95% confidence interval (CI) of the index of moderated mediation based on 5000 bias-corrected bootstrapped samples.

**Results**

*Sample*

The sample average age at two-year follow-up was 12 years, and the sample was 53% male. The analyses with academic functioning as the outcome included 6754 participants (after excluding siblings and children with missing data) and the analyses with social problems as the outcome included 7550 participants. Additional, clinical, behavioral, environmental, and functional information for children included in the study are presented in Table 1.

*Mediation Analyses: Mental Health Predicting School Grades*

First, we ran a set of mediation analyses in which a mental health measure (depression, anxiety, distressing PLEs, and CBCL total problems) was the independent variable, four healthy lifestyle factors (sleep quality, weekend screen time, diet, and physical activity) were mediators, sex/age/site were covariates, and academic functioning was the outcome. When entered simultaneously, these lifestyle factors combined significantly mediated associations between all four mental health variables with academic functioning (see Figure 2 for mediation models and Supplementary Table 1 for detailed statistics). The relationship between anxiety and academic functioning showed the highest mediation from these lifestyle mediators (48%), followed by depression (30%), distressing PLEs (18%), and finally total problems (10%).

Furthermore, in all four models, sleep showed evidence of being the most important mediator, showing 36.3% (of 48% total) mediation for anxiety, 18.5% (of 30% total) mediation for depression, 8.3% (of 18% total) mediation for distressing PLEs, and 4.6% (of 10% total) mediation for total problems (Figure 2, Supplementary Table 1). Screen time was the second-most important mediator, although its contributions were often noticeably less than sleep quality (e.g., 36.3% and 18.5% mediation by sleep quality vs. 5.0% and 6.3% mediation by screen time when anxiety and depression were the independent variables, respectively). Exercise also showed significant mediation, although its effects were even smaller than for screen time (e.g., 2.8% and 4.0% mediation for models predicting anxiety and depression, respectively). Diet did not show significant mediation for depression, distressing PLEs, or total problems.

A highly similar pattern of findings was observed when including cognition as a covariate (Supplementary Figure 1, Supplementary Table 2). When weekday screen time was used instead of weekend screen time, the mediating effects of screen time were consistently smaller (e.g., 3.7% (weekday) vs. 6.3% (weekend) for depression), although it remained statistically significant for all four psychopathological domains tested (Supplementary Figure 2, Supplementary Table 3).

*Mediation Analyses: Mental Health Predicting Social Functioning*

Next, we ran a set of mediation analyses in which a mental health measure was the independent variable, the four healthy lifestyle factors were mediators, sex/age/site were covariates, and social problems (CBCL Social Problems score) was the outcome. As was the case for academic functioning, the four lifestyle mediators combined to show

significant mediation in all of the psychopathological models (depression, anxiety, and distressing PLEs; Figure 3, Supplementary Table 4). The relationship between distressing PLEs and social problems showed the highest mediation from these lifestyle mediators (27%), followed by depression (21%) and anxiety (21%). Strikingly, amongst the lifestyle mediators, only sleep quality was a robust, consistently significant mediator for all models examined. More specifically, for depression sleep showed 19.9% (of 21% total) mediation, for anxiety sleep showed 19.6% (of 21% total) mediation, and for PLE distress sleep showed 23.3% (of 27% total) mediation. Thus, almost all of the healthy lifestyle mediation effects were driven by sleep quality for all three mental health domains examined.

A highly similar pattern of findings was observed when including cognition as a covariate (Supplementary Figure 3, Supplementary Table 5). When weekday screen time was used instead of weekend screen time, the mediating effects of screen time became consistently smaller and non-significant for all psychopathological domains tested (Supplementary Figure 4, Supplementary Table 6).

*Environmental Moderation Effects*

Supplementary Table 7 depicts how environmental factors moderate associations between lifestyle mediators with academic functioning, as well as moderator effects on percent mediation. As can be seen in Table 2, as financial adversity increased, the mediating role of sleep quality and/or weekend screen time (reverse-scored) on associations between mental health and academic functioning decreased.

Supplementary Table 8 depicts how environmental factors moderate associations between lifestyle mediators with social problems was the outcome. As can be seen in Table 3, as family conflict and financial adversity increased and school environment worsened, the mediating effects of sleep quality on associations between mental health and social problems increased.

**Discussion**

The present study examined the degree to which potentially modifiable lifestyle behaviors, including sleep quality, physical activity, diet quality, and screen time, serve as mechanisms linking adolescent mental health symptoms to academic functioning and social problems, as well as whether these associations were moderated in the context of environmental adversity. Using data from the ABCD Study, we found evidence that healthy lifestyle factors, especially sleep, mediated associations between depression, anxiety, and PLEs with both academic and social functioning. We also found evidence that financial adversity modified associations between lifestyle factors with both academic and social problems, and that school environment and family conflict additionally modified associations with social problems. These findings clarify modifiable pathways that may contribute to functional decline in adolescence and identify contextual conditions under which lifestyle-focused interventions may be less effective.

**Lifestyle Mediators Linking Symptoms to Academic Functioning**

Consistent with predictions, healthy lifestyle factors jointly mediated the associations between symptoms and academic functioning across all mental health domains examined. For academic functioning, mediation was strongest for models using anxiety as the predictor (48% of the total effect), followed by depression, PLEs, and total mental health concerns. This pattern may suggest that academic functioning is particularly sensitive to internalizing processes (e.g., cognitive rumination, worrying), domains that may be especially intertwined with lifestyle behaviors [9,44,45]. Sleep quality emerged as the dominant mediator across all analyses (i.e., explaining the majority of the total mediated effect), with screen time providing additional contributions. These findings point to sleep quality as potentially a key mechanism through which early symptoms translate into reduced academic performance, consistent with evidence linking sleep irregularities to daytime fatigue, attention problems, and poorer executive functioning [9,11,44,45]. Additionally, screen time showed a moderate influence while exercise and diet showed less robust mediation effects, with their small but significant contributions in some models highlighting the value of a multi-behavioral approach to supporting academic outcomes [16,17,20,21,46]. Notably, weekend screen time showed a qualitatively stronger mediating effect than weekend screen time. One possible reason for the discrepancy is the dynamic range for weekday screen time is lower (as children are in school, and as evidenced by the lower mean weekday screen time (Table 1)), reducing its ability to mediate functional outcomes. Overall, these results suggest that interventions that improve sleep quality and/or (to a lesser extent) restrict screen time may reduce the deleterious effects of mental health problems on academic functioning.

**Lifestyle Mediators Linking Symptoms to Social Problems**

When social problems were the outcome, a similar overall mediation pattern emerged as compared to academic functioning, although the strength of the mediation differed. Mediation was strongest for PLE distress (27%), consistent with some psychosis spectrum research [13-15], followed by depression and anxiety (both ~21%) [8,12]. Unlike the academic functioning models where multiple healthy lifestyle factors contributed, sleep quality was the only consistently significant mediator of social functioning across all domains. This may indicate that sleep plays a uniquely central role in how symptoms translate into impairments in social functioning. This is potentially consistent with previous research linking mental health concerns, including social anhedonia and irritability, with social problems [47,48]. In contrast, screen time, diet, and physical activity contributed minimally to social problems pathways, suggesting these behaviors exert a smaller influence on associations between mental health with social problems. The contrast between academic functioning and social problems mediation patterns underscores that healthy lifestyle factors may operate differently across functional domains, with sleep exerting a more domain-general influence, and other lifestyle factors (e.g., screentime) playing more domain-specific roles. As with academic functioning, these results also suggest that interventions that improve sleep quality may help mitigate the effects of poor mental health on social outcomes.

**Moderated Mediation by Environmental Adversity**

Environmental moderation analyses revealed that the pathways described above were not uniform across youth. For example, for academic functioning models, financial

adversity was a significant moderator in that higher financial adversity weakened associations between sleep quality and the academic outcome in these models. This may suggest that, in the context of financial strain, lifestyle improvements may confer smaller academic functioning benefits.  This may be because economic stressors may more directly affect academic functioning (e.g., through resource limitations, unstable housing), overshadowing lifestyle-related influences [30,31], although this hypothesis requires further investigation.

For social problems, the pattern of moderated mediation differed from that observed for models examining academic functioning. Whereas financial adversity weakened lifestyle-related mediation for academic functioning, the opposite was true for social problems: as family conflict and financial adversity increased and school environment worsened, the mediating effects of sleep quality on associations between mental health and social problems strengthened. One possibility is that, under high-conflict or high-adversity conditions, sleep disruptions may exacerbate emotional reactivity, irritability, or social withdrawal, thereby magnifying their downstream impact on social functioning [8,34]. Taken together, these findings indicate that healthy lifestyle mechanisms operate differently across functional domains, with sleep quality becoming a stronger mediator of social functioning under higher adversity, in contrast to the diminished mediation observed for academic functioning.

**Implications for Prevention and Intervention**

This study identifies sleep quality and to a lesser extent screen time as potentially tractable intervention targets for improving outcomes among adolescents

experiencing mental health concerns. Sleep interventions, such as school-based sleep hygiene programs to improve sleep quality, may meaningfully improve academic and social functioning [46,49]. Yet the moderating effects of financial adversity and family conflict suggest that lifestyle-focused interventions should be tailored to the youth's environmental context. For youth experiencing economic hardship or high family conflict, sleep-based supports may need to be paired with additional structural supports (e.g., resource assistance, family-based therapies) to maximize impact.

**Limitations**

Despite the strengths of the present study, several limitations warrant consideration. First, although the present work focused on modifiable lifestyle variables, it is likely that other lifestyle factors may influence both mental health as well as social and academic functioning. Second, even though mediation effects were statistically robust, several of the effect sizes were modest and may not generalize uniformly across developmental periods. Third, while sleep problems are a diagnostic feature of several mental health conditions (e.g., mood disorders), notably, sleep disturbances partially mediated associations with a range of mental health symptoms, suggesting sleep may function as a separatable and potentially modifiable factor, rather than intrinsic to experiencing specific symptoms such as depression. Finally, although mediation models incorporated independent variables, mediators, and outcomes in chronological order (except for diet, as this information was only available for 1-year follow-up (and would be unlikely to change by 2-year follow-up [35])), this does not necessarily imply that these relationships were causal. Future studies using more sophisticated structural equation

models incorporating data from multiple timepoints are necessary to examine the longitudinal relationships between these factors in more comprehensively, and intervention studies are necessary to firmly establish cause-and-effect relationships. This may necessitate using data from future ABCD studies, as later releases will include more timepoints (including data during middle and late adolescence, when mental issues become increasingly prevalent [50]).

## Conclusions

Findings from this large, multi-site study provide novel evidence that lifestyle factors, especially sleep quality, are key factors linking adolescent mental health symptom domains to academic and social functioning. These pathways are sensitive to contextual conditions, with financial adversity and family conflict moderating the impact lifestyle-related mediation. Together, these results underscore the importance of integrating lifestyle-based and family-informed approaches to support adolescents experiencing early mental health concerns.

## Acknowledgements

Dr. Smucny is supported by grant K01-MH125906 from the National Institutes of Mental Health. Dr. Karcher is supported by grant R01-MH139880.

Data used in the preparation of this article were obtained from the Adolescent Brain Cognitive Development[SM] (ABCD) Study (https://abcdstudy.org) release 5.1 (DOI: 10.15154/z563-zd24), held in the NIMH Data Archive (NDA). This is a multisite, longitudinal study designed to recruit more than 10,000 children ages 9-10 and follow

them over 10 years into early adulthood. The ABCD Study® is supported by the National Institutes of Health and additional federal partners under award numbers U01DA041048, U01DA050989, U01DA051016, U01DA041022, U01DA051018, U01DA051037, U01DA050987, U01DA041174, U01DA041106, U01DA041117, U01DA041028, U01DA041134, U01DA050988, U01DA051039, U01DA041156, U01DA041025, U01DA041120, U01DA051038, U01DA041148, U01DA041093, U01DA041089, U24DA041123, U24DA041147. A full list of supporters is available at https://abcdstudy.org/federal-partners.html. A listing of participating sites and a complete listing of the study investigators can be found at https://abcdstudy.org/consortium_members/. ABCD consortium investigators designed and implemented the study and/or provided data but did not necessarily participate in the analysis or writing of this report. This manuscript reflects the views of the authors and may not reflect the opinions or views of the NIH or ABCD consortium investigators.

# References

1.     Barch DM, Albaugh MD, Avenevoli S, et al. Demographic, physical and mental health assessments in the adolescent brain and cognitive development study: Rationale and description. *Dev Cogn Neurosci*. Aug 2018;32:55-66. doi:10.1016/j.dcn.2017.10.010

2.     Galvan A. Adolescence, brain maturation and mental health. *Nat Neurosci*. Mar 29 2017;20(4):503-504. doi:10.1038/nn.4530

3.     Olfson M, Druss BG, Marcus SC. Trends in Mental Health Care among Children and Adolescents. *N Engl J Med*. Sep 10 2015;373(11):1079. doi:10.1056/NEJMc1507642

4.     Carvajal L, Requejo JH, Irwin CE. The Measurement of Mental Health Problems Among Adolescents and Young Adults Throughout the World. *J Adolesc Health*. Sep 2021;69(3):361-362. doi:10.1016/j.jadohealth.2021.06.009

5.     Remschmidt H, Belfer M. Mental health care for children and adolescents worldwide: a review. *World Psychiatry*. Oct 2005;4(3):147-53.

6.     Achenbach TM, Rescorla LA. *Manual for the ASEBA School-Age Forms & Profiles*. University of Vermont, Research Center for Children, Youth and Families; 2001.

7.     Karcher NR, Barch DM, Avenevoli S, et al. Assessment of the Prodromal Questionnaire-Brief Child Version for Measurement of Self-reported Psychoticlike Experiences in Childhood. *JAMA Psychiatry*. Aug 1 2018;75(8):853-861. doi:10.1001/jamapsychiatry.2018.1334

8.     Achterbergh L, Pitman A, Birken M, Pearce E, Sno H, Johnson S. The experience of loneliness among young people with depression: a qualitative meta-synthesis of the literature. *BMC Psychiatry*. Aug 24 2020;20(1):415. doi:10.1186/s12888-020-02818-3

9.     Dalforno RW, Wengert HI, Kim LP, Jacobsen KH. Anxiety and school absenteeism without permission among adolescents in 69 low- and middle-income countries. *Dialogues Health*. Dec 2022;1:100046. doi:10.1016/j.dialog.2022.100046

10.    Finning K, Ukoumunne OC, Ford T, et al. The association between child and adolescent depression and poor attendance at school: A systematic review and meta-analysis. *J Affect Disord*. Feb 15 2019;245:928-938. doi:10.1016/j.jad.2018.11.055

11.    Kearney CA, Dupont R, Fensken M, Gonzálvez C. School attendance problems and absenteeism as early warning signals: review and implications for health-based protocols and school-based practices. Review. *Frontiers in Education*. 2023-August-30 2023;Volume 8 - 2023doi:10.3389/feduc.2023.1253595

12.     Kupferberg A, Hasler G. The social cost of depression: Investigating the impact of impaired social emotion regulation, social cognition, and interpersonal behavior on social functioning. *Journal of Affective Disorders Reports*. 2023/12/01/ 2023;14:100631. doi:https://doi.org/10.1016/j.jadr.2023.100631

13.     El Bouhaddani S, van Domburgh L, Schaefer B, Doreleijers TAH, Veling W. Peer status in relation to psychotic experiences and psychosocial problems in adolescents: a longitudinal school-based study. *Eur Child Adolesc Psychiatry*. Jun 2018;27(6):701-710. doi:10.1007/s00787-017-1063-2

14.     Huckle C, Lemmel F, Johnson S. Experiences of friendships of young people with first-episode psychosis: A qualitative study. *PLoS One*. 2021;16(7):e0255469. doi:10.1371/journal.pone.0255469

15.     Steenkamp LR, Bolhuis K, Blanken LME, et al. Psychotic experiences and future school performance in childhood: a population-based cohort study. *J Child Psychol Psychiatry*. Mar 2021;62(3):357-365. doi:10.1111/jcpp.13281

16.     Firth J, Solmi M, Wootton RE, et al. A meta-review of "lifestyle psychiatry": the role of exercise, smoking, diet and sleep in the prevention and treatment of mental disorders. *World Psychiatry*. Oct 2020;19(3):360-380. doi:10.1002/wps.20773

17.     Ruiz-Ranz E, Asin-Izquierdo I. Physical activity, exercise, and mental health of healthy adolescents: A review of the last 5 years. *Sports Med Health Sci*. May 2025;7(3):161-172. doi:10.1016/j.smhs.2024.10.003

18.     Bruni O, Ottaviano S, Guidetti V, et al. The Sleep Disturbance Scale for Children (SDSC). Construction and validation of an instrument to evaluate sleep disturbances in childhood and adolescence. *J Sleep Res*. Dec 1996;5(4):251-61. doi:10.1111/j.1365-2869.1996.00251.x

19.     Nagata JM, Bashir A, Weinstein S, et al. Social epidemiology of the Mediterranean-dietary approaches to stop hypertension intervention for neurodegenerative delay (MIND) diet among early adolescents: the Adolescent Brain Cognitive Development Study. *Pediatr Res*. Jul 2024;96(1):230-236. doi:10.1038/s41390-023-02959-7

20.     Camprodon-Boadas P, Gil-Dominguez A, De la Serna E, Sugranyes G, Lazaro I, Baeza I. Mediterranean Diet and Mental Health in Children and Adolescents: A Systematic Review. *Nutr Rev*. Feb 1 2025;83(2):e343-e355. doi:10.1093/nutrit/nuae053

21.     O'Neil A, Quirk SE, Housden S, et al. Relationship between diet and mental health in children and adolescents: a systematic review. *Am J Public Health*. Oct 2014;104(10):e31-42. doi:10.2105/AJPH.2014.302110

22.     Santos RMS, Mendes CG, Sen Bressani GY, et al. The associations between screen time and mental health in adolescents: a systematic review. *BMC Psychology*. 2023/04/20 2023;11(1):127. doi:10.1186/s40359-023-01166-7

23.     Zablotsky B, Ng AE, Black LI, et al. Associations Between Screen Time Use and Health Outcomes Among US Teenagers. *Prev Chronic Dis*. Jul 10 2025;22:E38. doi:10.5888/pcd22.240537

24.     Nagata JM, Al-Shoaibi AAA, Leong AW, et al. Screen time and mental health: a prospective analysis of the Adolescent Brain Cognitive Development (ABCD) Study. *BMC Public Health*. Oct 7 2024;24(1):2686. doi:10.1186/s12889-024-20102-x

25.     Zink J, O'Connor SG, Blachman-Demner DR, Wolff-Hughes DL, Berrigan D. Examining the Bidirectional Associations Between Sleep Duration, Screen Time, and Internalizing Symptoms in the ABCD Study. *J Adolesc Health*. Mar 2024;74(3):496-503. doi:10.1016/j.jadohealth.2023.09.001

26.     Goldstone A, Javitz HS, Claudatos SA, et al. Sleep Disturbance Predicts Depression Symptoms in Early Adolescence: Initial Findings From the Adolescent Brain Cognitive Development Study. *J Adolesc Health*. May 2020;66(5):567-574. doi:10.1016/j.jadohealth.2019.12.005

27.     Reeve S, Bell V. Sleep disorders predict the 1-year onset, persistence, but not remission of psychotic experiences in preadolescence: a longitudinal analysis of the ABCD cohort data. *Eur Child Adolesc Psychiatry*. Sep 2023;32(9):1609-1619. doi:10.1007/s00787-022-01966-z

28.     Damme KSF, Vargas TG, Walther S, Shankman SA, Mittal VA. Physical and mental health in adolescence: novel insights from a transdiagnostic examination of FitBit data in the ABCD study. *Transl Psychiatry*. Feb 3 2024;14(1):75. doi:10.1038/s41398-024-02794-2

29.     Aliyas Z, Mahmoudian A, Cloutier M-S. Association Between Socio-Emotional Health, Physical Activity and Screen Time Among Children. *American Journal of Health Education*. 2025/07/04 2025;56(4):366-375. doi:10.1080/19325037.2024.2396602

30.     Gautam N, Dessie G, Rahman MM, Khanam R. Socioeconomic status and health behavior in children and adolescents: a systematic literature review. *Front Public Health*. 2023;11:1228632. doi:10.3389/fpubh.2023.1228632

31.     Gautam N, Rahman MM, Khanam R. Socioeconomic inequalities in health behaviors in children and adolescents: evidence from an Australian cohort. *BMC Public Health*. Jan 24 2025;25(1):314. doi:10.1186/s12889-025-21472-6

32.     Steare T, Evans-Lacko S, Araya M, et al. Economic inequalities in adolescents' internalising symptoms: longitudinal evidence from eight countries. *Lancet Psychiatry*. Nov 2024;11(11):890-898. doi:10.1016/S2215-0366(24)00255-4

33.     Al-Shoaibi AAA, Zamora G, Chu J, et al. Family conflict and less parental monitoring were associated with greater screen time in early adolescence. *Acta Paediatr*. Nov 2024;113(11):2452-2458. doi:10.1111/apa.17349

34. Cummings EM, Koss KJ, Davies PT. Prospective relations between family conflict and adolescent maladjustment: security in the family system as a mediating process. *J Abnorm Child Psychol*. Apr 2015;43(3):503-15. doi:10.1007/s10802-014-9926-1

35. da Costa MP, Severo M, Araujo J, Vilela S. Longitudinal tracking of diet quality from childhood to adolescence: The Interplay of individual and sociodemographic factors. *Appetite*. May 1 2024;196:107279. doi:10.1016/j.appet.2024.107279

36. Achenbach TM, Ruffle TM. The Child Behavior Checklist and related forms for assessing behavioral/emotional problems and competencies. *Pediatr Rev*. Aug 2000;21(8):265-71. doi:10.1542/pir.21-8-265

37. Echeverria SE, Diez-Roux AV, Link BG. Reliability of self-reported neighborhood characteristics. *J Urban Health*. Dec 2004;81(4):682-701. doi:10.1093/jurban/jth151

38. Moos RH. *Family Environment Scale Manual: Development, Applications, Research*. Consult. Psychol. Press; 1994.

39. Zucker RA, Gonzalez R, Feldstein Ewing SW, et al. Assessment of culture and environment in the Adolescent Brain and Cognitive Development Study: Rationale, description of measures, and early data. *Dev Cogn Neurosci*. Aug 2018;32:107-120. doi:10.1016/j.dcn.2018.03.004

40. Karcher NR, Loewy RL, Savill M, et al. Replication of Associations With Psychotic-Like Experiences in Middle Childhood From the Adolescent Brain Cognitive Development (ABCD) Study. *Schizophr Bull Open*. Jan 2020;1(1):sgaa009. doi:10.1093/schizbullopen/sgaa009

41. Luciana M, Bjork JM, Nagel BJ, et al. Adolescent neurocognitive development and impacts of substance use: Overview of the adolescent brain cognitive development (ABCD) baseline neurocognition battery. *Dev Cogn Neurosci*. Aug 2018;32:67-79. doi:10.1016/j.dcn.2018.02.006

42. Roth B, Becker N, Romeyke S, Schäfer S, Domnick F, Spinath FM. Intelligence and school grades: A meta-analysis. *Intelligence*. 2015/11/01/ 2015;53:118-137. doi:https://doi.org/10.1016/j.intell.2015.09.002

43. Almat NS, Aliya MS, Zhanna UT, Gulmira DS. The Relationship Between Social Intelligence And IQ: A Psychometric Analysis. *The Open Psychology Journal*. 2023/01/01/ 2023;16doi:https://doi.org/10.2174/18743501-v16-e230120-2022-78

44. Fergusson DM, Woodward LJ. Mental health, educational, and social role outcomes of adolescents with depression. *Arch Gen Psychiatry*. Mar 2002;59(3):225-31. doi:10.1001/archpsyc.59.3.225

45.     Finning K, Ukoumunne OC, Ford T, et al. Review: The association between anxiety and poor attendance at school - a systematic review. *Child Adolesc Ment Health*. Sep 2019;24(3):205-216. doi:10.1111/camh.12322

46.     Green JG, McLaughlin KA, Alegria M, et al. School mental health resources and adolescent mental health service use. *J Am Acad Child Adolesc Psychiatry*. May 2013;52(5):501-10. doi:10.1016/j.jaac.2013.03.002

47.     Qiu J, Morales-Munoz I. Associations between Sleep and Mental Health in Adolescents: Results from the UK Millennium Cohort Study. *Int J Environ Res Public Health*. Feb 7 2022;19(3)doi:10.3390/ijerph19031868

48.     Woodfield M, Butler NG, Tsappis M. Impact of sleep and mental health in adolescence: an overview. *Curr Opin Pediatr*. Aug 1 2024;36(4):375-381. doi:10.1097/MOP.0000000000001358

49.     Fenwick-Smith A, Dahlberg EE, Thompson SC. Systematic review of resilience-enhancing, universal, primary school-based mental health promotion programs. *BMC Psychol*. Jul 5 2018;6(1):30. doi:10.1186/s40359-018-0242-3

50.     *Substance Abuse and Mental Health Services Administration (2013). Behavioral Health, United States, 2012. HHS Publication No. (SMA) 13-4797. Rockville, MD: Substance Abuse and Mental Health Services Administration*.

**Table 1.** Demographic, clinical, behavioral, environmental, and functional information for all participants included in analyses. Values are from parental report data unless otherwise specified (parental report data was used unless unavailable). Numbers in parentheses represent the standard deviation. *A grades score of 3 corresponds to a GPA of 3.5-3.69 (A-) and a score of 4 to a GPA of 3.30-3.49 (B+). CBCL = Child Behavior Checklist, NIH = National Institutes of Health, PQ-BC = Prodromal Questionnaire-Brief Child Version.

| Analysis Dependent Variable | Grades in School, Years 2-3 | CBCL Social Problems Score, Year 3 |
|---|---|---|
| *N* | 6754 | 7550 |
| **Demographic Covariates** | | |
| Age in Years at Year 2 | 11.99 (.66) | 12.00 (.66) |
| Sex *n* M/F/Intersex | 3578/3174/2 | 3988/3560/2 |
| NIH Cognition Toolbox Total Composite Raw Score, Baseline | 87.11 (8.89) | 86.79 (9.01) |
| **Clinical, Independent (Predictor) Variables** | | |
| CBCL Depression T-Score, Year 1 | 53.87 (5.97) | 53.97 (6.08) |
| CBCL Anxiety T-Score, Year 1 | 53.55 (6.10) | 53.63 (6.25) |
| PQ-BC Distress Score, Year 1 | 4.40 (9.05) | 4.48 (9.09) |
| CBCL Total Problems T-Score, Year 1 | 45.51 (10.95) | 45.65 (11.14) |
| **Healthy Lifestyle Mediators** | | |
| Sleep Disturbances Score, Year 2 | 36.43 (8.01) | 36.48 (8.06) |
| Mediterranean Diet Score, Year 1 | 8.08 (2.39) | 8.07 (2.42) |
| Youth Reported Days Physically Active in the Last Week, Year 2 Questionnaire | 3.80 (2.15) | 3.78 (2.16) |
| Weekend Screen Time Hours, Year 2 | 4.63 (3.07) | 4.69 (3.13) |
| Weekday Screen Time Hours, Year 2 | 3.30 (3.08) | 3.38 (3.15) |
| **Environmental Moderators** | | |
| Neighborhood Crime Rating, Year 2 | 3.88 (.93) | 3.87 (.94) |
| Family Conflict Score, Year 2 | 2.36 (1.94) | 2.38 (1.96) |
| Youth Reported School Environment Score, Year 2 | 19.63 (2.76) | 19.63 (2.78) |
| Family Financial Adversity Score, Baseline | .40 (1.01) | .42 (1.05) |
| **Functional, Dependent (Outcome) Variables** | | |
| Grades in School*, Years 2-3 | 3.73 (2.31) | – |
| CBCL Social Problems Score, Year 3 | – | 52.65 (4.90) |

**Table 2.** Changes in %mediation dependent on moderator level for significant moderators for models predicting academic functioning between years 2-3. See Table 1 for mean (SD) of moderator variables. *Because mean - 1 SD < 0, mediation effects at this level were calculated when the moderator = 0.

| Predictor Mediator | Moderator | Moderator Score | Indirect Effect (95% CI) | %Mediation |
|---|---|---|---|---|
| **Depression** | | | | |
| **Sleep Quality** | Family Financial Adversity | Mean - 1 SD* | -.0138 (-.0193 – -.0084) | 20.2% |
| | | Mean | -.0120 (-.0172 – -.0070) | 17.6% |
| | | Mean + 1 SD | -.0076 (-.0139 – -.0015) | 11.1% |
| **Weekend Screen Time (Reverse-Scored)** | Family Financial Adversity | Mean - 1 SD* | -.0045 (-.0063 – -.0030) | 6.6% |
| | | Mean | -.0040 (-.0056 – -.0027) | 5.9% |
| | | Mean + 1 SD | -.0027 (-.0043 – -.0015) | 4.0% |
| **Anxiety** | | | | |
| **Sleep Quality** | Family Financial Adversity | Mean - 1 SD* | -.0153 (-.0199 – -.0112) | 38.0% |
| | | Mean | -.0138 (-.0181 – -.0101) | 34.3% |
| | | Mean + 1 SD | -.0102 (-.0153 – -.0055) | 25.4% |
| **Weekend Screen Time (Reverse-Scored)** | Family Financial Adversity | Mean - 1 SD* | -.0021 (-.0035 – -.0010) | 5.2% |
| | | Mean | -.0019 (-.0031 – -.0009) | 4.7% |
| | | Mean + 1 SD | -.0013 (-.0025 – -.0006) | 3.2% |
| **PLE Distress** | | | | |
| **Sleep Quality** | Family Financial Adversity | Mean - 1 SD* | -.0032 (-.0047 – -.0022) | 8.6% |
| | | Mean | -.0029 (-.0043 – -.0020) | 7.8% |
| | | Mean + 1 SD | -.0022 (-.0035 – -.0013) | 5.9% |
| **Weekend Screen Time (Reverse-Scored)** | Family Financial Adversity | Mean - 1 SD* | -.0023 (-.0035 – -.0015) | 6.2% |
| | | Mean | -.0021 (-.0032 – -.0013) | 5.6% |
| | | Mean + 1 SD | -.0015 (-.0025 – -.0008) | 4.0% |
| **Total Problems** | | | | |
| **Weekend Screen Time (Reverse-Scored)** | Family Financial Adversity | Mean - 1 SD* | -.0027 (-.0038 – -.0019) | 4.0% |
| | | Mean | -.0024 (-.0034 – -.0017) | 3.5% |
| | | Mean + 1 SD | -.0016 (-.0026 – -.0009) | 2.3% |

**Table 3.** Changes in %mediation dependent on moderator level for significant moderators for models predicting social problems (CBCL Social Problems Score) at year 3. See Table 1 for mean (SD) of moderator variables. *Because mean - 1 SD < 0, mediation effects at this level were calculated when the moderator = 0.

| Predictor<br>Mediator | Moderator | Moderator Score | Indirect Effect B (95% CI) | %Mediation |
|---|---|---|---|---|
| **Depression** | | | | |
| **Sleep Quality** | Family Conflict | Mean - 1 SD | .0455 (.0318 – .0603) | 13.1% |
| | | Mean | .0595 (.0484 – .0716) | 17.1% |
| | | Mean + 1 SD | .0735 (.0593 – .0890) | 21.2% |
| **Sleep Quality** | School Environment | Mean - 1 SD | .0777 (.0632 – .0933) | 22.4% |
| | | Mean | .0667 (.0549 – .0795) | 19.2% |
| | | Mean + 1 SD | .0557 (.0408 – .0719) | 16.1% |
| **Sleep Quality** | Family Financial Adversity | Mean - 1 SD* | .0590 (.0463 – .0717) | 17.0% |
| | | Mean | .0639 (.0523 – .0761) | 18.4% |
| | | Mean + 1 SD | .0762 (.0617 – .0928) | 22.0% |
| **Anxiety** | | | | |
| **Sleep Quality** | Family Conflict | Mean - 1 SD | .0443 (.0332 – .0554) | 13.8% |
| | | Mean | .0547 (.0457 – .0648) | 17.0% |
| | | Mean + 1 SD | .0651 (.0539 – .0777) | 20.3% |
| **Sleep Quality** | School Environment | Mean - 1 SD | .0690 (.0576 – .0816) | 21.5% |
| | | Mean | .0613 (.0519 – .0719) | 19.1% |
| | | Mean + 1 SD | .0536 (.0419 – .0673) | 16.7% |
| **Sleep Quality** | Family Financial Adversity | Mean - 1 SD* | .0543 (.0451 – .0650) | 16.9% |
| | | Mean | .0584 (.0494 – .0685) | 18.2% |
| | | Mean + 1 SD | .0683 (.0564 – .0821) | 21.3% |
| **PLE Distress** | | | | |
| **Sleep Quality** | Family Conflict | Mean - 1 SD* | .0139 (.0010 – .0188) | 17.2% |
| | | Mean | .0165 (.0121 – .0215) | 20.4% |
| | | Mean + 1 SD | .0191 (.0140 – .0251) | 23.6% |
| **Sleep Quality** | School Environment | Mean - 1 SD* | .0202 (.0151 – .0265) | 25.0% |
| | | Mean | .0184 (.0137 – .0240) | 22.8% |
| | | Mean + 1 SD | .0167 (.0120 – .0223) | 20.7% |
| **Sleep Quality** | Family Financial Adversity | Mean - 1 SD* | .0172 (.0126 – .0224) | 21.3% |
| | | Mean | .0179 (.0133 – .0233) | 22.2% |
| | | Mean + 1 SD | .0196 (.0146 – .0257) | 24.3% |

**Figure Legends**

**Figure 1.** Diagram illustrating moderation analysis procedure. The effect of each individual moderator (e.g., family financial adversity) was tested on the relationships between all mediators (e.g., sleep quality) and each outcome (e.g., academic functioning). In this fashion, moderated mediation models examined the degree to which each moderator affected the mediation of associations between mental health with outcome through lifestyle factors.

**Figure 2.** Results of mediation models predicting academic functioning (school grades between 2-3 year follow-up) while including age (2-year follow-up), sex, and site as covariates. More detailed results are provided in Supplementary Table 1. PLE = Psychotic-like Experiences.

**Figure 3.** Results of mediation models predicting social problems (3 year CBCL Social Problems score) while including age (2-year follow-up), sex, and site as covariates. More detailed results are provided in Supplementary Table 4. PLE = Psychotic-like Experiences.

**Clinical Predictor**

**Outcome: Social Problems**

*p < .05

**Depression**

Depression (Year 1)

-.5845 (.0137)*  Sleep Quality (Year 2)  -.1180 (.0069)*
Indirect Effect: .0690 (.0063)* %Mediation: 19.9%

-.0260 (.0041)*  Physical Activity (Year 2)  -.0762 (.0235)*
Indirect Effect: .0020 (.0007)* %Mediation: .6%

-.0659 (.0045)*  Diet (Year 1)  -.0225 (.0212)
Indirect Effect: .0015 (.0015) %Mediation: .4% (n.s.)

-.0528 (.0059)*  Weekend Screen Time (Year 2)  -.0253 (.0165)
Indirect Effect: .0013 (.0010) %Mediation: .4% (n.s.)

Social Problems (Year 3)

Total Effect: .3470 (.0084)*
Direct Effect: .2732 (.0092)*
Sum Indirect Effects: .0738 (.0064)*
Sum %Mediation: 21%

**Anxiety**

Anxiety (Year 1)

-.4595 (.0139)*  Sleep Quality (Year 2)  -.1370 (.0067)*
Indirect Effect: .0630 (.0051)* %Mediation: 19.6%

-.0379 (.0044)*  Diet (Year 1)  -.0617 (.0211)*
Indirect Effect: .0023 (.0009)* %Mediation: .7%

-.0210 (.0040)*  Physical Activity (Year 2)  -.0798 (.0235)*
Indirect Effect: .0017 (.0006)* %Mediation: .5%

-.0263 (.0057)*  Weekend Screen Time (Year 2)  -.0420 (.0165)*
Indirect Effect: .0011 (.0006) %Mediation: .3%

Social Problems (Year 3)

Total Effect: .3213 (.0082)*
Direct Effect: .2532 (.0086)*
Sum Indirect Effects: .0681 (.0052)*
Sum %Mediation: 21%

**PLE Distress**

PLE Distress (Year 1)

-.0947 (.0102)*  Sleep Quality (Year 2)  -.1989 (.0066)*
Indirect Effect: .0188 (.0027)* %Mediation: 23.3%

-.0174 (.0027)*  Physical Activity (Year 2)  -.0981 (.0247)*
Indirect Effect: .0017 (.0005)* %Mediation: 2.1%

-.0281 (.0039)*  Weekend Screen Time (Year 2)  -.0282 (.0173)
Indirect Effect: .0008 (.0006) %Mediation: 1.0% (n.s.)

-.0029 (.0031)  Diet (Year 1)  -.1042 (.0221)*
Indirect Effect: .0003 (.0004) %Mediation: .4% (n.s.)

Social Problems (Year 3)

Total Effect: .0808 (.0061)*
Direct Effect: .0592 (.0058)*
Sum Indirect Effects: .0216 (.0028)*
Sum %Mediation: 27%

Mediation Strength (%)