Patterns and Clinical Outcomes of Physical Activity and Sedentary Behavior
Across 20 Million Days of Wearable Monitoring in U.S. Adults

Arash A. Nargesi, MD, MPH[1,2,3]*, Valentina D'Souza, MSc[1]*, Tal Shnitzer, PhD[1]*, Andri
Kadaifciu[1], Aurora Cremer[1], Sean Joseph Jurgens, MD[4], Nakia Mack, MSc, MPH[5], Ryan Lupi[5],
Romuladus Azuine, DrPH, MPH[5], Geoffrey S. Ginsburg, MD, PhD[5], Chris Lunt[5], Christopher D.
Anderson, MD, MSc[6,7], Sam Friedman, PhD[1], Patrick T. Ellinor, MD, PhD[4,8], Mahnaz Maddah,
PhD[1]
8
1. Data Sciences Platform, Broad Institute of MIT and Harvard, Cambridge, Massachusetts, USA
2. Division of Cardiovascular Medicine, Brigham and Women's Hospital, Boston, Massachusetts, USA
3. Clinical Informatics, Mass General Brigham, Somerville, Massachusetts, USA
4. Cardiovascular Disease Initiative, Broad Institute of MIT and Harvard, Cambridge, Massachusetts, USA
5. All of Us Research Program, National Institutes of Health, Bethesda, MD, USA
6. McCance Center for Brain Health, Massachusetts General Hospital, Boston, USA.
7. Department of Neurology, Brigham and Women's Hospital, Boston, USA.
8. Cardiovascular Research Center, Massachusetts General Hospital, Boston, Massachusetts, USA
17
18 *Authors contributed equally to this project
19
Corresponding Author:
Mahnaz Maddah, PhD
Director, Machine Learning for Health
Broad Institute of MIT and Harvard
415 Main Street, Cambridge, MA 02142
Email: maddah@broadinstitute.org
Phone: (617) 714-7000

27
28 **Short Title:** Physical Activity, Sedentary Behavior, and Clinical Outcomes

29 **Keywords:** Physical activity; step count; sedentary behavior; wearables; digital health.
30
Conflict of Interest Disclosures: None

Funding/Support: This work is supported by a grant from the National Institutes of Health, *All*
of Us Program (3OT2OD035404-01S3D). Dr. Nargesi is supported by a US National Institutes
of Health National Heart, Lung, and Blood Institute training grant (T32HL007604) and has
consulted for Novo Nordisk unrelated to the presented work. Dr. Anderson has received
sponsored research support from Bayer AG, and has consulted for MPM BioImpact and
ApoPharma, unrelated to the presented work. Dr. Ellinor is supported by the NIH
(1R01HL092577, K24HL105780), AHA (18SFRN34110082), Foundation Leducq (14CVD01),
and by MAESTRIA (965286). Dr. Maddah is supported by grants from the National Institutes of
Health (3OT2OD035404-01S3, 1R01NS134597 and 1UG3HG014379-01) and from the
American Heart Association (961045).
42
43 **Acknowledgment:** We gratefully acknowledge *All of Us* participants for their contributions,
44 without whom this research would not have been possible. We also thank the National Institutes
45 of Health's *All of Us* program for making available the participant data used in this study.

Abstract

Background: Physical activity is a key modifiable determinant of health, yet current guidelines primarily emphasize moderate-to-vigorous physical activity and provide limited guidance on other actionable dimensions such as daily step count and sedentary time. The growing availability of wearables enables high-resolution measurement of these metrics and assessment of their associations with clinical outcomes.

Methods: Minute-level wearable data from adult participants of *All of Us* between 2015-2023 were used to calculate step count and sedentary time. Patterns of step count and sedentary behavior were assessed across temporal scales, U.S. states, and sociodemographic groups. Phenome-wide association (PheWAS) and dose-response analyses were conducted to evaluate associations with incident disease.

Results: Among 50,300 participants (68% female, 71% White, median age 55 years), 19,845,612 person-days of wearable monitoring over a median follow-up of 13 months were included in this analysis. Step counts peaked at midday with weekly highs on Saturdays and seasonal highs in May, while sedentary time was highest in midafternoon and winter. Higher step counts and lower sedentary time were observed among males, individuals with higher income and education, and residents of the Northeast, upper Midwest, and West Coast. In PheWAS (n=31,446), higher step count and lower sedentary time were associated with lower risk of multiple diseases, including obesity, cardiometabolic risk factors, and mood and anxiety disorders. Dose response analyses demonstrated heterogeneous relationships across disease groups, with cardiovascular benefits plateauing beyond 9,000-10,000 steps/day and below 600-640 minutes/day of sedentary time.

70   **Conclusions:** In this nationwide cohort of U.S. adults, wearable-derived physical

71   activity and sedentary behavior showed distinct temporal and geographic patterns with

72   significant disease-specific associations with clinical outcomes. These findings support

73   the use of longitudinal high-resolution wearable data for advancing personalized

74   guidelines of physical activity and sedentary behavior.

Introduction

Physical activity is widely recognized as one of the most effective and modifiable determinants of health[1]. Clinical guidelines from the World Health Organization (WHO) and American Heart Association (AHA) recommend adults to engage in at least 150 minutes of moderate or 75 minutes of vigorous aerobic activity per week[2,3]. The Physical Activity Guidelines for Americans also recommend reducing sedentary behavior as a key strategy for preventing chronic diseases[4]. Despite the robust evidence supporting the role of physical activity in human health, current guidelines mostly focus on moderate and vigorous physical activity (MVPA), while target levels for other dimensions of physical activity, such as daily step count and sedentary time have not been clearly established.

Physical activity encompasses diverse combinations of movement intensity and duration, and its patterns among individuals are highly variable and yet poorly characterized. A key barrier to developing universally applicable guidelines has been the limited scope and short follow-up of previous studies in this domain. Large-scale studies, including community cohorts such as the UK biobank[1,5] and nationally representative surveys such as National Health and Nutrition Examination Survey (NHANES)[6], were limited to 1-2 weeks of monitoring. Nevertheless, the growing availability of wearable technology has significantly enhanced the feasibility and accuracy of tracking physical activity at scale. Among the contemporary cohorts, the *All of Us* (AoU) program from the US National Institutes of Health (NIH) provides a unique opportunity to access longitudinal wearable data collected from a broad sample of individuals across the US. The long-term follow-up and linkage of wearable data with

electronic health records (EHRs) facilitate the assessment of longitudinal patterns of

physical activity[7]. Furthermore, understanding the exact relationship of step count and

sedentary time with future risk of disease is essential to inform personalized and

actionable guidance on physical activity and complement existing recommendations

based on MVPA alone.

In this study, we used minute-level wearable data from adult participants in the

AoU Program to characterize temporal, geographic, and sociodemographic patterns of

daily step count and sedentary time across the US. We further evaluated their

associations with incident disease using phenome-wide and dose–response analyses.

By linking high-resolution wearable data with longitudinal clinical outcomes, this study

provides evidence relevant to the use of wearable-derived physical activity metrics in

population health and preventive medicine. In addition to describing population-level

activity patterns, we aimed to identify disease-specific step count and sedentary time

thresholds associated with a lower risk of incident disease.

Methods

Study Design and Participants

This was a prospective cohort study using longitudinal wearable data and EHRs from

the National Institutes of Health AoU Program[8,9]. Details of study protocols have been

previously published[10,11]. We used registered and controlled tier datasets (Version 8, ID

numbers R2024Q3R3 and C2024Q3R4) which include data from participants enrolled

between 2015 and 2023. Adults aged 18 years or older with available wearable data

were eligible for inclusion. The analytic cohort was derived as shown in the CONSORT

flow diagram (**Figure 1**). Demographic and survey data were collected at enrollment,

and EHRs were linked from participating health systems. All participants provided

informed consent at enrollment, including consent to share wearable data. This study

was conducted in accordance with the Strengthening the Reporting of Observational

Studies in Epidemiology (STROBE) reporting guideline.

Wearable Data

Wearable data in the AoU Program were collected using Fitbit devices. Participants

were enrolled through 2 pathways: (1) Bring Your Own Device (BYOD), in which

participants linked a personal Fitbit account to the All of Us participant portal, and (2)

the Wearable Enhancing All of Us Research (WEAR) study, in which Fitbit devices

(Charge 4 or Versa 3) were provided to participants at no cost. Details of enrollment

procedures, device distribution, and data collection have been reported previously[12,13].

Fitbit devices continuously measure movement and physiologic signals, including

accelerometer-based activity and photoplethysmography-derived heart rate, which are

processed using proprietary algorithms to generate activity metrics. AoU provides

second- or minute-level step counts and daily summaries of physical activity across

intensity zones to the researchers, including total daily minutes spent at sedentary, light,

moderate, vigorous, and peak activity zones.

Data Preprocessing

We used minute-level step count and heart rate to assess the patterns of physical

activity and sedentary behavior in the study population. The raw minute-level data are

noisy and require substantial preprocessing to ensure adequate quality for downstream

analysis. We converted step count and heart rate data into numerical minute-level

vectors, identified and excluded non-wear times, removed outliers based on

physiological ranges of heart rate and step count, and excluded records with high

missingness. Fitbit trackers do not provide data on sedentary behavior at minute-level

scale. To comprehensively assess the patterns of physical activity, we estimated

minute-level sedentary times from the accelerometer and heart rate data. Procedural

methods, quality control, and calculations of sedentary time are described in

Supplement (**Method description**).

Study Outcomes and Covariates

We assessed temporal and geographic patterns of step count and sedentary time,

evaluated their associations with the incidence of clinical phenotypes in a PheWAS

analysis, and conducted dose-response analyses to identify potential thresholds of step

count and sedentary time associated with lower risk of incident disease.

Sociodemographic data, including age, sex, self-reported race, education,

income, employment, and the state of residence were collected at enrollment. Clinical

characteristics were derived from the EHRs using standardized coding systems.

Corresponding codes for clinical phenotypes were grouped using phecodeX, a

phenotyping tool that adds granularity and improves the hierarchical structure of under-

represented phenotypes[14]. Using phecodeX definitions, we performed a PheWAS

analysis to examine the association of step count and sedentary time with incident

cardiovascular, metabolic, musculoskeletal, gastrointestinal, respiratory, neoplastic,

neurological, and mental diseases.

We identified incident events as the first appearance of disease diagnosis codes

in the EHRs after applying a blanking period of 30 days from the last wearable

monitoring. All wearable data over a maximum time horizon of 3 years preceding the

incident event were included in the analysis. The 3-year limit was applied to minimize

the influence of long-term behavioral changes that could weaken the temporal

relevance between wearable-derived predictors and disease outcomes. In all

prospective analyses, data from the control group were restricted to wearable

monitoring between 2015-2022 to temporally align with the incident group. All data

beyond the incidence date or the last clinical encounter were censored.

Statistical Analysis

*a. Descriptive Analyses***:** Temporal patterns of daily step count and sedentary time were

examined across multiple time scales, including within-day, weekly, monthly, and yearly

intervals, as well as over an extended follow-up period of up to 5 years. For each

participant, average daily profiles were first computed for each metric and subsequently

aggregated to generate population-level estimates, with equal weighting assigned to all

participants. In sensitivity analyses, we examined daily and weekly trends of step count

across sociodemographic subgroups and within each enrollment pathway (BYOD vs.

WEAR). As sedentary time is calculated in minutes, its daily distribution represents the

proportion of participants engaged in such activities at each minute point in time. To

ensure consistency and minimize the confounding effect of early dropouts, yearly

analyses were restricted to individuals with available data for at least 12 contiguous

months and multi-year analyses to participants with complete five-year follow-ups. For

participants with non-contiguous periods of monitoring and gaps exceeding 3 months,

only the longest continuous data were used in yearly and multi-year analyses. To assess

geographical variations of physical activity, each metric was aggregated at the state level

and visualized in choropleth maps. Similar to our analysis of temporal patterns, equal

weights were given to all participants within each state.

b. PheWAS: We also examined the association of step count and sedentary time with

incident disease conditions using phenotypic codes described above and identified the

top five associations in each disease category based on the numerical p-values. These

analyses were restricted to individuals with available EHRs (n=31,446). Each physical

activity metric was tested in multivariable Cox regression models adjusted for age, sex,

and self-reported race. Hazard ratios per one standard deviation change in each metric

are presented against adjusted p-values in volcano plots after applying Bonferroni

correction for multiple comparisons.

c. Dose–Response Analyses: We assessed cumulative risk of incident disease across

eight disease categories against varying levels of step count and sedentary time. All

disease phenotypes under the same category were grouped together with equal

weights given to each condition. Segmented regression was used to test for non-linear

associations between levels of daily step count and sedentary time with the incidence of

any phenotype within each disease category after adjusting for age, sex, and self-

reported race. For categories with a statistically significant non-linear association, the

elbow of the dose-response curve (inflection point) was subsequently identified.

Categorical variables are presented in frequency and percentage. Continuous

variables are presented in mean and standard deviation or median and interquartile

range, as appropriate. Analysis of variance (ANOVA) and Kruskal–Wallis tests were

used to assess group differences, and post-hoc pairwise comparisons were performed

with Bonferroni correction as appropriate. Bonferroni correction was also applied to

PheWAS and dose-response analysis for multiple comparisons. All statistical tests are

2-sided with a pre-adjusted level of significance of 0.05.

Computational and statistical analyses were performed using Python 3.9[15] and

on the secure cloud-based platform of AoU Researcher Workbench[10]. All codes are

available at https://github.com/broadinstitute/ml4h_aou.

Results

In the AoU, wearable data were available for 59,018 participants, with a total of

41,721,100 person-days of monitoring over a median follow-up of 18 months. Data from

50,300 participants, including 19,845,612 person-days of monitoring over a median

follow-up of 13 months, passed the quality control and were included in our final

analysis (CONSORT diagrams in **Figure 1** and **Supplement, Figure S1**).

The study population was predominantly female (n= 34,258, 68%) and White

(n=35,784, 71%), with a majority holding a college or advanced degree (n=31,645, 63%,

Table S1). The median age of study participants was 55 (IQR 40-68) years, 26,723

(53%) were employed at the time of enrollment with a median income between $100k-

200k. The average daily step count and sedentary time were 7,503 (95% CI 7,471-

7,535) steps/day, and 591 (95% CI 590-592) min/day, respectively. Characteristics of

participants with available EHRs and those in each enrollment pathway (BYOD vs

WEAR) are presented in **Supplement** (**Tables S2-S3**). Adherence to wearable

monitoring was higher among participants in the WEAR than those in the BYOD

enrollment pathway (average wear time 1397 min/day in WEAR vs. 1393 min/day in

BYOD, p-value < 0.001, **Supplement, Figure S2**). Geographic distribution of AoU

participants and those with available wearable data are presented in **Supplement**

(**Figure S3**). The study period included the COVID-19 pandemic, which was associated

with a decrease in physical activity during the lockdown and post-lockdown periods

between 2020-2022 (**Supplement, Figure S4**), similar to previous reports from

wearable studies in the AoU[16]. The impact of quality control measures on the

distribution of physical activity is represented in **Supplement** (**Figures S5**).

Temporal and Geographic Patterns of Physical Activity

We assessed the patterns of step count and sedentary time across various time

intervals. Minute-level step count showed a bimodal distribution with a peak around

12:00 pm and spikes at whole hour timestamps (**Figure 2A**). Aside from the expected

increase in sedentary time during sleep hours, daily sedentary behavior showed a

downward trend through the morning and upward trend in early afternoon with peak

sedentary time at 2:57 pm.

Participants recorded the highest daily average step count (7,674 [95%CI:7,638-

7,710] steps/day) and lowest sedentary time (mean 567, 95% CI 566-568 min/day) on

Saturdays, whereas Sundays exhibited the opposite pattern. Despite numerically

significant differences in day-to-day variations of step count and sedentary time over

different days of the month (p-value < 0.001), no consistent pattern emerged

(**Supplement, Figure S6)**. However, both step count and sedentary behavior displayed

significant seasonal variations (**Figure 2A-B)**. Step count peaked in May (mean 8,237,

95% CI 8,187-8,287 step/day) and trended down in fall toward its lowest level in

January (mean 7,379, 95% CI 7,333-7,425 per day). Conversely, sedentary time

reached its highest level in February (mean 598, 95% CI 597-600 min/day), followed by

a marked decline in the spring, reaching its minimum of 583 (95% CI 582-585) min/day

in July. Among individuals with at least 5 years of contiguous wearable monitoring, step

count trended down and sedentary time trended up in each subsequent year of follow-

up (p-value < 0.001, **Supplement, Figure S6**).

We also examined geographic variations in step count and sedentary behavior

across the US states (**Figure 2C and 2D**). Average daily step count was relatively

higher in participants living in the Northeast, upper Midwest, and West Coast and lower

in those residing in central Southern states. Step counts were highest in the District of

Columbia, Hawaii, and New York, and lowest in West Virginia, Oklahoma, and

Arkansas. Patterns of sedentary time showed an inverse trend with highest levels

recorded in participants living in South Dakota, Kentucky, Wyoming, and Oklahoma,

and lowest values observed in residents of Mississippi, Hawaii and Louisiana. Seasonal

variations of step count and sedentary time across US states are presented in

Supplement (**Figure S7**).

Physical Activity Across Sociodemographic Subgroups

Step count demonstrated unique patterns across sociodemographic subgroups (**Figure**

3). Participants aged 18-44 and 45-65 years showed similar bimodal distributions, with

peak activity in late morning and late afternoon hours. In contrast, those who were 65

years of age or older had peak step counts in the early morning with a gradual decline

through the rest of the day. Males consistently recorded higher step counts than

females (daily average 8,396 [95% CI 8,334-8,458] vs 7,092 [95% CI 7,056-7,129]

steps/day, p-value < 0.001). Asian and White participants had higher step counts than

Black participants (daily average for Asians 8,502 [95% CI 8,360-8,645], Whites 7,433

[95% CI 7,395-7,471], and Blacks 7,170 [95% CI 7,053-7,287], p-value < 0.001). In

addition, Asian participants showed a unique pattern of activity with peak step count in

the evening (**Figure 3**).

Step count patterns also varied by educational attainment and income.

Individuals with college or advanced degrees recorded the highest step counts (mean

daily step count 7,842 [95% CI 7,802-7,881] steps/day). A similar gradient was

observed by income, as individuals with a household income greater than $200k had

the highest step counts (mean 8,874 [95% CI 8,773-8,976] steps/day), followed by

those in the $100k–200k and $50k–100k groups, with the lowest step counts observed

in participants earning less than $50k. Among employment categories, retired

participants had their peak step count in the early morning, while students and

employed individuals showed bimodal patterns with peak step counts in the evening.

Weekly patterns of step count across demographic groups are presented in

Supplement (**Figure S8**). Temporal patterns of step count across demographic groups

within BYOD and WEAR enrollment pathways are also presented in **Supplement**

(**Figures S9-S12**).

PheWAS and Dose-Response Analysis

We performed a PheWAS analysis to assess the relationship between step count and

sedentary time with incident clinical phenotypes across eight disease categories

(**Figure 4**). In this prospective analysis, the median follow up time was 16 months for

incident cases and 29 months for controls. Higher step counts were significantly

associated with lower incidence of multiple conditions across the disease spectrum,

including obesity (HR = 0.60 [95% CI 0.56-0.64]), sleep apnea (HR = 0.64 [0.60-0.68]),

abnormalities of breathing (HR = 0.72 [0.68-0.77]), hypotension (HR = 0.49 [0.43-0.56]),

gastroesophageal reflux (HR = 0.72 [0.67-0.76]), mood disorders (HR = 0.72 [0.67-

0.77]) and myalgia (HR = 0.71 [0.65-0.77]). In contrast, longer sedentary times were

associated with higher incidence of morbid obesity (HR = 1.85 [1.72-2.00]), sleep apnea

(HR = 1.41 [1.33-1.50]), essential hypertension (HR = 1.27 [1.21-1.34]), hypoxemia (HR

= 1.59 [1.38-1.83]), fatty liver disease (HR = 1.31 [1.20-1.44]) and dysthymic disorders

(HR = 1.46 [1.25-1.70]). Detailed results of our phenome-wide association analysis are

presented in **Supplement** (**Tables S4-S5**).

To better understand the nature of the relationships between physical activity and

sedentary behavior with clinical outcomes, we performed a dose-response analysis

between daily step count and sedentary time with cumulative risk of incident disease

across eight disease groups. Increasing numbers of daily steps and shorter duration of

sedentary time were associated with lower incidence of all eight disease categories (p

for trend < 0.001 for all, **Figure 5**). The inverse association between step count and

disease groups plateaued beyond approximately 11,000 to 12,000 steps/day for

neurologic disorders (p-value <0.001) and 9,000 to 10,000 steps/day for cardiovascular

diseases (p-value<0.001). Similarly, the association between daily sedentary time and

neurologic, cardiovascular, and metabolic diseases reached a plateau below 600 to 640

min/day of sedentary behavior. For all other disease groups, we did not find a

statistically significant risk threshold for either step count or sedentary time

(**Supplement Table S6-S7**). Dose-response associations between deciles of these two

metrics and incident disease categories are presented in **Supplement** (**Figure S13**).

Discussion

Leveraging data from over 19 million person-days of wearable monitoring in broad

populations, this study represents one of the largest longitudinal analyses

of physical activity with minute-level resolution and multi-year follow-up. Our findings

provide a detailed picture of temporal and geographic patterns of physical activity and

sedentary behavior among US adults, including their variations across

sociodemographic groups. Our PheWAS analysis complements prior studies on the

association of step count and sedentary behavior with clinical outcomes. Moreover, we

identify daily thresholds for step count and sedentary time that are associated with the

lowest observed risk of disease across the spectrum of clinical conditions. Collectively,

these findings provide a comprehensive picture of physical activity and sedentary

behavior across the US and can potentially inform actionable recommendations in

future guidelines to improve population health.

We observed substantial temporal variations in both physical activity and

sedentary behavior. Step count generally followed a bimodal daily distribution, peaking

at late morning, followed by lower activities in mid-afternoon hours before reaching a

second peak in the evening. We also observed higher levels of physical activity on

Saturdays and a seasonal trend with greater activity during spring and summer.

Temporal patterns of physical activity are known to correlate with distinct risk profiles of

obesity and metabolic syndrome[17], highlighting the importance of characterizing these

patterns at the population level. Our observations are also consistent with previous

studies on environmental and behavioral factors, such as daylight availability, work

schedules, and weather conditions, which can significantly influence the patterns of

physical activity[18].

We also observed substantial variations in physical activity across demographic

subgroups which highlight the influence of structural and societal factors on health-

related behaviors. Our findings support prior reports on the association of demographic

features with physical activity in other populations[19,20], while offering a comprehensive

characterization of such patterns among US adults. Geographic variations of physical

activity similarly reflect a complex interaction of individual characteristics, built

environments, and broader societal factors that shape opportunities for physical activity

across the US. Notably, we observed differential seasonal patterns between northern

and southern states, highlighting the significant role of environmental factors on physical

activity. The breadth of data collected in the AoU, including longitudinal surveys of

social determinants of health and linkage to geospatial environmental exposures,

empowers future research to investigate the multifaceted interactions between

individual characteristics, environmental conditions, and health behaviors at scale.

The significance of these observations is underscored by the association of

physical activity and sedentary behavior with a broad range of clinical outcomes. In our

analysis, cardiometabolic disorders such as obesity and hypertension, and psychiatric

diseases such as mood and anxiety disorders consistently stood out as correlates of

inactivity and sedentary behavior. Consistent associations between step count and

sedentary time with incident disease, together with findings from our dose-response

analyses reinforce the existing literature that associates physical inactivity with poor

health outcomes[21-24]. Furthermore, our analysis demonstrates the heterogeneous

relationship of varying levels of step counts and sedentary time with clinical outcomes.

In many disease groups, these relationships appear to be linear, with no limit to the

benefits of higher step counts and lower sedentary times. In contrast, the cardiovascular

and neurologic benefits appear to plateau beyond certain thresholds of daily step counts

and sedentary times. Across disease categories, graded associations between physical

activity metrics and incident disease revealed distinct nonlinear patterns, enabling

identification of step count and sedentary time thresholds associated with lower disease

risk. These findings suggest disease-specific relationships between step counts and

sedentary time with clinical outcomes, with both linear and threshold-dependent

patterns that can inform personalized guidance for different populations in future.

At a broader scale, our study demonstrates the value of wearables to

characterize dynamic patterns of physical activity in real-world settings[25]. The richness

of wearable data extends beyond self-reported estimates of physical activity or

aggregate measures of daily exercise. Importantly, current guidelines do not universally

recognize wearable technology as a tool for measurement and promotion of physical

activity[2,3]. Wearable monitoring can inform a data-driven approach to understand

behavior-disease relationships when linked with EHRs in large-scale cohorts such as

AoU[26]. Although integration of wearable data into clinical workflows may not be

straightforward[27], a wearable-centered approach has the potential to facilitate the

design of tailored interventions for promotion of physical activity in target populations.

Because step count and sedentary time are routinely captured by consumer wearables,

these metrics offer intuitive and monitorable targets that may complement existing

physical activity counseling in clinical and public health settings.

Limitations

First, the study sampling is prone to selection bias as it is more likely for health-

conscious individuals who use wearables to be enrolled through the BYOD pathway.

The AoU attempts to mitigate this bias by enrolling a community-based cohort through

the WEAR study. However, the selection process remains non-random, and the study

sample is not representative of the broader US population[28]. Second, characterization

of clinical phenotypes relies on EHRs which are inherently subject to incomplete and

variable adjudication at local sites, particularly as data are primarily collected for clinical

and not research indications. Third, participating health systems follow varying

measurement and documentation standards, introducing heterogeneity in clinical

phenotypes that may affect the consistency of covariate or outcome ascertainment.

Fourth, wearable data in this study were derived exclusively from Fitbit devices;

therefore, the observed patterns and associations may not generalize to users of other

vendors with different sensors or algorithms. Finally, this study is observational in nature

and despite our attempts for prospective investigation in PheWAS and dose-response

analysis, causal inferences between patterns of physical activity and clinical outcomes

cannot be definitively established. Despite these limitations, the large sample size,

longitudinal follow-up, and high-resolution wearable data provide robust evidence of

associations between physical activity and incident disease.

Conclusions

This study provides a large-scale, high-resolution characterization of physical activity

and sedentary behavior among U.S. adults in the All of Us Program. By analyzing more

than 19 million person-days of wearable monitoring, we identified temporal, geographic,

and sociodemographic patterns of physical activity and demonstrated associations with

incident disease across multiple clinical domains, including disease-specific nonlinear

relationships. In particular, cardiovascular risk reduction plateaued at approximately

9,000 to 10,000 steps per day, while higher disease risk was observed beyond

approximately 600 to 640 minutes per day of sedentary time. These findings underscore

the value of integrating granular wearable data with electronic health records to inform

more actionable physical activity guidance and population-level disease prevention

strategies.

References

References

1.   Khurshid, S., *et al*. Wearable accelerometer-derived physical activity and incident
disease. *NPJ Digit Med* **5**, 131 (2022).
2.   Arnett, D.K., *et al*. 2019 ACC/AHA Guideline on the Primary Prevention of
Cardiovascular Disease: A Report of the American College of Cardiology/American
Heart Association Task Force on Clinical Practice Guidelines. *Circulation* **140**,
e596-e646 (2019).
3.   Organization, W.H. Global Recommendations on Physical Activity for Health. *World*
Health Organization (2010).
4.   Piercy, K.L., *et al*. The Physical Activity Guidelines for Americans. *JAMA* **320**, 2020-
2028 (2018).
5.   Strain, T., *et al*. Wearable-device-measured physical activity and future health risk.
Nat Med **26**, 1385-1391 (2020).
6.   Saint-Maurice, P.F., *et al*. Association of Daily Step Count and Step Intensity With
Mortality Among US Adults. *JAMA* **323**, 1151-1160 (2020).
7.   Zheng, N.S., *et al*. PheMap: a multi-resource knowledge base for high-throughput
phenotyping within electronic health records. *J Am Med Inform Assoc* **27**, 1675-
1687 (2020).
8.   Bianchi, D.W., *et al*. The All of Us Research Program is an opportunity to enhance
the diversity of US biomedical research. *Nat Med* **30**, 330-333 (2024).
9.   Khajouei, E., *et al*. Phenome-wide association of APOE alleles in the All of Us
Research Program. *EBioMedicine* **117**, 105768 (2025).
10.   All of Us Research Program, I., *et al*. The "All of Us" Research Program. *N Engl J Med*
381, 668-676 (2019).
11.   Mayo, K.R., *et al*. The All of Us Data and Research Center: Creating a Secure,
Scalable, and Sustainable Ecosystem for Biomedical Research. *Annu Rev Biomed*
Data Sci **6**, 443-464 (2023).
12.   Master, H., *et al*. How Fitbit data are being made available to registered researchers
in All of Us Research Program. *Pac Symp Biocomput* **28**, 19-30 (2023).
13.   Lederer, L., *et al*. Considerations while using Fitbit Data in the All of Us Research
Program. *All of Us Research Program* (2022).
14.   Shuey, M.M., *et al*. Next-generation phenotyping: introducing phecodeX for
enhanced discovery research in medical phenomics. *Bioinformatics* **39**(2023).
15.   Van Rossum, G. & Drake, F.L. Python 3 Reference Manual. *CreateSpace*, 242
(2009).
16.   Desine, S., *et al*. Daily Step Counts Before and After the COVID-19 Pandemic
Among All of Us Research Participants. *JAMA Netw Open* **6**, e233526 (2023).
17.   Aqeel, M., *et al*. Temporal physical activity patterns are associated with obesity in
U.S. adults. *Prev Med* **148**, 106538 (2021).
18.   Turrisi, T.B., *et al*. Seasons, weather, and device-measured movement behaviors: a
scoping review from 2006 to 2020. *Int J Behav Nutr Phys Act* **18**, 24 (2021).

19. Strain, T., *et al.* National, regional, and global trends in insufficient physical activity
among adults from 2000 to 2022: a pooled analysis of 507 population-based
surveys with 5.7 million participants. *Lancet Glob Health* **12**, e1232-e1243 (2024).
20. Armstrong, S., *et al.* Association of Physical Activity With Income, Race/Ethnicity,
and Sex Among Adolescents and Young Adults in the United States: Findings From
the National Health and Nutrition Examination Survey, 2007-2016. *JAMA Pediatr*
172, 732-740 (2018).
21. Stamatakis, E., *et al.* Association of wearable device-measured vigorous
intermittent lifestyle physical activity with mortality. *Nat Med* **28**, 2521-2529 (2022).
22. Raichlen, D.A., *et al.* Sedentary Behavior and Incident Dementia Among Older
Adults. *JAMA* **330**, 934-940 (2023).
23. Pearce, M., *et al.* Association Between Physical Activity and Risk of Depression: A
Systematic Review and Meta-analysis. *JAMA Psychiatry* **79**, 550-559 (2022).
24. Master, H., *et al.* Association of step counts over time with the risk of chronic
disease in the All of Us Research Program. *Nat Med* **28**, 2301-2308 (2022).
25. Khurshid, S., Al-Alusi, M.A., Churchill, T.W., Guseh, J.S. & Ellinor, P.T.
Accelerometer-Derived "Weekend Warrior" Physical Activity and Incident
Cardiovascular Disease. *JAMA* **330**, 247-252 (2023).
26. Lee, I.M., Keadle, S.K. & Matthews, C.E. Fitness Trackers to Guide Advice on Activity
Prescription. *JAMA* **330**, 1733-1734 (2023).
27. Ginsburg, G.S., Picard, R.W. & Friend, S.H. Key Issues as Wearable Digital Health
Technologies Enter Clinical Care. *N Engl J Med* **390**, 1118-1127 (2024).
28. Zeng, C., *et al.* Comparison of phenomic profiles in the All of Us Research Program
against the US general population and the UK Biobank. *J Am Med Inform Assoc* **31**,
846-854 (2024).

Figure 1. Study Population. The CONSORT diagram illustrates the inclusion and exclusion of participants from the All of Us Research Program. Wearable data, including step counts and heart rates, that passed quality control were included in the final analysis. Abbreviations: EHR, Electronic Health Records.

Figure 2. Temporal and Geographic Variations in Step Counts and Sedentary Times. (A) and (B) represent minute-level times series and aggregated distributions of step counts and sedentary times across daily, weekly, and yearly intervals. (C) and (D) illustrate geographic variations in daily average step counts and sedentary times across all U.S. states and the District of Columbia. Color scales represent the distribution of each metric across states, with yellow indicating higher and blue indicating lower values.

A Steps Trend

Daily   Weekly   Yearly

C Step Count Map

B Sedentary Trend

Daily   Weekly   Yearly

D Sedentary Map

Figure 3. Step Counts Across Sociodemographic Subgroups. Minute-level time series of step counts shown across sociodemographic subgroups of age, sex, self-reported race, education, income, and employment status. Lines represent minute-level average step counts with 95% confidence intervals.

Figure 4. Association of Step Count and Sedentary Time with Incident Disease.
Associations between daily step counts (A) and sedentary times (B) with incident
phenotypes are shown across eight major disease categories. Each point represents
an individual phenotype, plotted by the hazard ratio (x-axis, per one standard deviation
change in each metric) and the corresponding –log10(p-values) on the y-axis. Vertical
lines denote hazard ratio of 1.0 and horizontal lines indicate the Bonferroni-corrected
significance threshold. All models are adjusted for age, sex, and self-reported race.
Phenotypes are grouped and color-coded by disease category and the top 5
phenotypes by significance are listed where available. Among 31,661 participants with
wearable data and electronic health records, the last clinical encounter or
demographic covariates could not be adjudicated in 215 participants who were
excluded from this analysis (n=31,446).

A

Association of Step Counts with Incident Clinical Phenotypes (n=31,446)

Top Phenotypes

Metabolic
1. Overweight and obesity
2. Obesity
3. Morbid obesity
4. Type 2 diabetes
5. Diabetes mellitus

Neurologic
1. Sleep apnea
2. Obstructive sleep apnea
3. Sleep disorders
4. Polyneuropathies
5. Disorders of autonomic nervous system

Respiratory
1. Abnormalities of breathing
2. Dyspnea [Shortness of breath]
3. Respiratory failure
4. Snoring*
5. Hypoxemia

Cardiovascular
1. Hypotension
2. Essential hypertension
3. Hypertension
4. Orthostatic hypotension
5. Heart failure

Gastrointestinal
1. Gastro-esophageal reflux disease [GERD]
2. Fatty liver disease (FLD)
3. Chronic liver disease
4. Symptoms involving digestive system
5. Aphagia and dysphagia

Mental
1. Mood [affective] disorders
2. Major depressive disorder
3. Anxiety and anxiety disorders
4. Generalized anxiety disorder
5. Mental disorder, not otherwise specified

Muscloskeletal
1. Myalgia
2. Low back pain
3. Other symptoms and disorders of the soft tissue
4. Disorder of ligament
5. Spinal stenosis

B

Association of Sedentary Time with Incident Clinical Phenotypes (n=31,446)

Top Phenotypes

Metabolic
1. Morbid obesity
2. Obesity
3. Overweight and obesity
4. Type 2 diabetes
5. Diabetes mellitus

Neurologic
1. Sleep apnea
2. Obstructive sleep apnea
3. Sleep disorders
4. Polyneuropathies
5. Hypersomnia

Cardiovascular
1. Essential hypertension
2. Hypertension
3. Heart failure
4. Diastolic heart failure
5. Hypotension

Respiratory
1. Hypoxemia
2. Asphyxia and hypoxemia
3. Respiratory failure
4. Snoring*
5. Abnormalities of breathing

Gastrointestinal
1. Fatty liver disease (FLD)
2. Chronic liver disease
3. Gallstones [Cholelithiasis]

Mental
1. Dysthymic disorder
2. Posttraumatic stress disorder
3. Mood [affective] disorders

Figure 5. Dose-Response Analysis of Daily Step Counts and Sedentary Time. The figures show cumulative incidence rate of eight major disease categories across varying levels of daily step counts (A) and sedentary time (B). Histograms at the bottom of each panel indicate the number of participants contributing data to each exposure bin.

A   **Step Counts**   **B**   **Sedentary Time**

Group

- Neurologic
- Cardiovascular
- Metabolic
- Mental
- Gastrointestinal
- Muscloskeletal
- Cancer
- Respiratory