Unmet palliative care needs in England and Wales: population-based estimates and future projections (2025-2050)

Authors:

Therese Johansson[1], https://orcid.org/0000-0001-6101-5925

Katherine E Sleeman[1], https://orcid.org/0000-0002-9777-4373

Anne Finucance[2], https://orcid.org/0000-0002-3056-059X

Joanna M Davies[1], https://orcid.org/0000-0002-6375-0023

Lorna K Fraser[1], https://orcid.org/0000-0002-1360-4191

Irene J Higginson[1], https://orcid.org/0000-0002-3687-1313

Melanie Diggle[1], https://orcid.org/0000-0002-4137-1946

Fliss E M Murtagh[3], https://orcid.org/0000-0003-1289-3726

Anna E Bone[1]*, https://orcid.org/0000-0002-8800-9581

*Corresponding author: Anna Bone, anna.bone@kcl.ac.uk

Affiliations:

[1] Cicely Saunders Institute of Palliative Care, Policy & Rehabilitation, Florence Nightingale Faculty of Nursing, Midwifery & Palliative Care, King's College London, London, UK

[2] Clinical Psychology, School of Health in Social Science, University of Edinburgh, Edinburgh, UK

[3] Wolfson Palliative Care Research Centre, Hull York Medical School, University of Hull, Hull, UK

Word count: 4422

Tables: 3

Figures: 2

Supplement files: 9

Abstract

Introduction: With global populations ageing, demand for palliative care is increasing. Population-level assessments of unmet palliative care needs are essential for strategic planning, yet rigorous methods to estimate unmet needs are lacking. This study aimed to develop methods and estimate current and future population-level prevalence of unmet palliative care needs among adults in England and Wales.

Methods: Secondary analyses of data from a nationally representative post-bereavement survey in England and Wales in 2022 (n=1,194). Unmet needs in the survey sample were estimated using two methods: (1) reported unresolved symptoms and concerns using Integrated Palliative care Outcome Scale scores, cutoff ≥34/68; and (2) reported insufficient care provision from general practitioners. These methods were combined to further provide a conservative estimate (1 and 2) and a broad estimate (1 or 2). We examined associations with unmet needs using modified Poisson regression. Age-, gender- and nation-specific sample estimates were applied to mortality data for 2022 and projections from the Office for National Statistics to calculate population-level estimates and prevalence from 2025 to 2050.

Results: In 2022, 247,993 (46%) adult decedents in England and 17,209 (49%) in Wales had unmet palliative care needs using method 1; 244,612 (46%) and 15,280 (43%), respectively, using method 2. According to conversative and broad estimates, 32% and 61% could have unmet needs in England, and 29% and 62% in Wales. By 2050, prevalence of unmet needs are projected to rise by 21−26% in England and 14−19% in Wales depending on estimate used, with the largest absolute increase among those aged ≥85 years.

Conclusions: Unmet palliative care needs are high in England and Wales and projected to increase by 2050, regardless of method. We contrast methods based on unresolved symptoms and concerns or insufficient care provision, or both, to inform the planning and evaluating of equitable care.

Keywords: Palliative Care; Health Services Needs and Demand; Terminal Care; Mortality; Forecasting

Key Messages

What is already known on this topic

- Although understanding population-level unmet palliative care needs is critical for effective service planning, robust and standardised methods to estimate these needs remain limited.

What this study adds

- We use two methods to provide four estimates of population-level prevalence of unmet palliative care needs and discuss their strengths and limitations.

- Regardless of estimate, prevalence of unmet palliative care needs is high, ranging 32-61% in England and in 29-62% in Wales; the number of people with unmet needs is anticipated to increase by 21–26% in England and 14–19% in Wales by 2050.

How this study might affect research, practice or policy

- This advancement in methods to estimate unmet palliative care needs can inform the development and evaluation of population-level strategies to improve end-of-life care.

- Our population-level estimates do not account for multiple long-term conditions which are rising and will likely increase the complexity of needs.

- There needs to be more investment in primary and community-based services to ensure high-quality symptom management and support for people and their families towards the end of life.

Introduction

Worldwide, mortality patterns are changing as populations age and health-related suffering of non-communicable diseases continues to rise. Consequently, more people are likely to benefit from palliative care.[1-4] Palliative care is a person-centred approach to care for people with life-limiting illness and their families, focusing on management of symptoms, alleviation of distress and improvement of quality of life. It can be delivered by specialist palliative care services, such as hospices, but is also commonly provided in primary and community care setting by generalists such as General Practitioners (GPs), community and district nurses, care home providers and hospital teams.[5] Despite this, many people do not receive the care and support they need towards the end of life, which can have significant implications for their health, wellbeing, and comfort. Understanding and addressing palliative care needs and the extent to which these are met in a population is therefore essential for effective and equitable healthcare planning and service delivery.[6 7]

A large body of work exists assessing palliative care needs in populations, often based on prevalence of diseases that are amenable to palliative, using administrative data such as death certificates.[8-10] Some population-level palliative care needs assessments have also considered symptom burden[11] and service utilisation,[12 13] and others have included subjective measures of needs.[14] However, very little research has attempted to quantify the extent or nature of unmet palliative care needs at the population level. Empirical evidence on how many people experience unmet palliative care needs, and characteristics of affected groups remains limited. In the literature, unmet needs are typically understood as the gap between the care required and the care received,[15] though definitions and measurement approaches vary widely and few studies are based on representative samples or include populations with non-cancer diagnoses.[15-17]

This study builds on earlier phases of the DUECare project, including a scoping review that identified three key domains for defining and measuring unmet needs: (1) symptoms and concerns; (2) access to services; and (3) sufficiency of service provision to address symptoms.[16] Stakeholder workshops using nominal group technique further refined priorities for measurement, highlighting the importance of capturing both symptom-related and service-related dimensions of unmet palliative care needs.[18] Through this work we developed a definition of unmet palliative care needs as occurring 'when a person with life-limiting illness has symptoms, psychosocial concerns, or care requirements that are not adequately addressed through available services, with inability to access or receive person-centred care'.[18]

The aim of this study was to develop methods to estimate the prevalence of unmet palliative care needs among adults in England and Wales, examine associated socio-demographic and clinical factors, and project baseline and future population-level prevalence of adults with unmet palliative care needs to 2050.

Materials and Methods

Study design

Secondary analysis of a nationally representative mortality follow-back survey data sent to next of kin of people who died in 2022 in England and Wales,[19] and of routinely available national death

registry data and mortality projections.[20][21] We report the study in line with Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) recommendations.[22]

Study data

Better End of Life survey data

As part of the Better End of Life study, a modified version of the QUALYCARE survey [23] was conducted with adults who had registered the death of a family member six to ten months prior. A nationally representative sample of decedents were identified using stratified sampling applied to the ONS' mortality data. Deaths were eligible if they had causes of death considered to be amenable to palliative care (ICD-10 codes sampled available in Supplement Table 1). The sampling process has been described in detail elsewhere.[19] To ensure adequate power for regression analyses with 25 predictors, we applied a conservative rule of 20 observations per variable, indicating a minimum sample size of 550. Data were collected between May and November 2023 (n=1,194), for deaths occurring between August and December 2022. Details of the sampling frame composition and characteristics of decedents for whom responses were received are published on https://doi.org/10.6084/m9.figshare.31100068.

Proxy reports from bereaved family members offer a valuable lens for understanding unmet needs in the final weeks or months of life.[23][24] While not without limitations, these accounts provide the best available population-based insights into symptom burden, care experiences, and perceived gaps in support. The Better End of Life survey collected data from bereaved family members on decedent socio-demographic and clinical characteristics (e.g., age, gender, financial circumstances, and cause of death). The survey also contained validated measures, including the Integrated Palliative care Outcome Scale (IPOS – proxy version), a 17-item measure assessing multidimensional symptoms and concerns across three sub-scales (*physical symptoms* (10 items), *emotional symptoms* (4 items), and *communication and practical issues* (3 items).[25] Each IPOS item relates to how a symptom or concern affected the person who died during their last week of life, rated using a 5-point Likert scale from 0 ('not at all') to 4 ('overwhelmingly' or 'always'). Scores are summed up, with possible total scores ranging from 0 to 68. The survey also included questions about which health care services were received and whether enough help was received from these services.

National Mortality Registry Data

We extracted publicly availably mortality data for England and Wales from the Office for National Statistics (ONS)[20] for 2022 (same year as deaths sampled in the Better End of Life survey), for adults aged 20 years and above (available age categories in the mortality data by cause of death precluded extraction for ages 18-19).

Official National mortality projections

Publicly available mortality projections for England and Wales for the years 2025 to 2050 were obtained from the ONS.[21] We used the 'principal projections', which provides mid-year annual mortality projections by age and gender.

Analysis

Estimates from Better End of Life survey sample

To estimate the proportion of decedents experiencing unmet needs in the survey sample, we developed and applied two estimation methods informed by the approaches to measuring unmet palliative care needs identified in our scoping review and stakeholder consultation workshops: (1) unresolved symptoms and concerns; (2) insufficient care provision to address symptoms and concerns.[16 18] The estimation methods were combined in different ways, generating four sample estimates, which are summarised in Box 1.

Estimate 1: Unresolved symptoms and concerns in the last week of life

Estimate 1 was based on reported unresolved symptoms and concerns, by using summed IPOS scores (scores of 0-4 across 17 items). A cut-off at the 50th percentile of the maximum possible total score range (≥34 out of 68) was applied to indicate unmet needs. A total score of ≥34 equates to being 'moderately' affected by all items or 'overwhelmingly/always' affected for at least nine items. Observations with eight or more IPOS items missing were excluded from the analytic sample and examined in a sensitivity analysis. For observations with one to seven IPOS items missing, we used mean scores from the non-missing items to impute missing values.

Estimate 2: Insufficient care provision in the last 3 months of life

For estimate 2, unmet palliative care needs were indicated by reported *insufficient care provision*. In this study, we focus on primary palliative care, which is palliative care provided by care professionals working in primary and community-based care services.[26 27] Primary palliative care should be available to everyone with advanced illness, and involves palliative care delivery integrated with chronic disease management.[28] We did not focus on specialist palliative care provision, because it was not possible in our dataset to distinguish between those who did not need specialist palliative care and those who needed it but did not receive it. Estimate 2 was based on the survey question "*In the last three months before they died, overall, do you feel they got as much help as needed from GPs?*". Responses of 'no', 'sometimes', or 'did not receive care' were coded as having unmet needs (i.e., not receiving enough help or not receiving care at all); 'yes, most of the time' was coded as having needs met (i.e., receiving enough help). Given that the survey sampled causes of deaths considered amenable to palliative care (i.e., expected deaths), we assumed decedents would have required at least some primary care in their final months of life.

Estimate 3: Conservative estimate

We calculated a conservative estimate based on the proportion of decedents who had unmet palliative care needs identified by both estimation methods 1 and 2 (i.e., those with a high burden of symptoms and concerns who also did not receive enough, or any, help from GPs).

Estimate 4: Broad estimate

A broad estimate was calculated based on decedents identified by either estimation method 1 or 2 (i.e., those with either a high symptom burden or those who did not receive enough help from GPs).

For each sample estimate, we calculated the 95% confidence intervals based on proportions using the Wilson score method.[29] For each estimate we excluded observations with information about decedent gender and/or age missing.

Regression analysis

We report frequences and proportions for categorical variables and means with standard deviations for continuous variables. Associations between unmet need and socio-demographic or clinical characteristics were examined using modified Poisson regression with robust error variance.[30] Separate multivariable models were fitted for binary outcomes (unmet needs vs no unmet needs) derived using the main two estimation methods, adjusting for gender, age, ethnicity, financial circumstances, nation, cause of death, and comorbidities. We report unadjusted and adjusted prevalence ratios (PR) with 95% confidence intervals. Analyses were performed in Stata 18.0.[31]

Box 1. *Overview of how estimates of unmet palliative care needs are derived in this study.*

1. Unresolved symptoms and concerns	An individual is identified as having unmet palliative care needs if they have an IPOS total score ≥34 (≥50% of maximum possible score).
2. Insufficient care provision	An individual is identified as having unmet palliative care needs if it is reported that they did not receive enough, or any, help from GPs.
3. Conservative estimate (1 and 2 combined)	An individual is identified as having unmet palliative care needs if they have an IPOS total score ≥34 **and** did not receive enough, or any, help from GPs.
4. Broad estimate (either 1 or 2)	An individual is identified as having unmet palliative care needs if they either have an IPOS total score ≥34 **or** did not receive enough, or any, help from GPs.

Population-level estimates

To estimate unmet palliative care needs among all adult deaths (age $\geq$20) in England and Wales, we first identified the population considered 'at risk' in the ONS mortality data for 2022[21] based on the main expected causes of death considered amenable to palliative care that were sampled in the Better End of Life survey (Supplement table 1). We assumed that those dying from other causes of death (including sudden deaths and accidents) did not have unmet palliative care needs. We grouped these deaths by gender (male and female) and age (20-64, 65-84, ≥85 years) for each nation separately. For each of the four sample estimates, we then applied age-, gender- and nation-specific proportions to the 'at risk' deaths in England and Wales to derive population-level estimates of unmet palliative care needs. We report these as numbers and proportions of decedents with unmet palliative care needs out of all adult deaths in each nation. In addition to the point estimates, we derived upper and lower population-level estimates by applying the upper and lower limits of the 95% confidence intervals of each sample estimate to the population data.

Future projections

We estimated future trends in unmet palliative care needs for adults in England and Wales over a 25-year period (2025–2050), using ONS national mortality projections.[21] Age-, gender- and nation-specific population-level estimates were applied to projected mortality data for all adult deaths,

using age groups 20-64, 65-84, ≥85 years. We assumed that the distribution of causes of death and the proportion of unmet palliative care needs remained constant throughout the projection period.

Patient and public involvement

This study is part of the Marie Curie-funded DUECare project (MCCRP-23-02), which has had input from our patient and public involvement (PPI) group throughout. Our operationalisation of methods to estimate unmet palliative care needs was informed by discussions with our PPI representatives and built on findings from stakeholder workshops conducted in an earlier phase of the project.[18]

Ethics

The study was conducted in accordance with the ethical standards set out in the Declaration of Helsinki and its amendments.[32] Ethical approval for the Better End of Life survey was received from King's College London Research Ethics Committee (HR/DP-21/22-24690). The analysis of routinely collected aggregated and anonymised data that are publicly available does not require ethical approval.

Results

Sample estimates

The mean age of decedents in the Better End of Life survey sample (n=1,194) was 81.5 years; 51.9% were female, 97.4% were White. There was a similar distribution of decedents across England (n=619, 51.8%) and Wales (n=575, 48.2%). The most common single cause of death was cancer (28.5%), followed by heart disease (15.3%) and dementia (15.3%). Full sample characteristics are presented in Table 1. Twenty-one observations missing information about decedent gender and/or age were removed from the sample estimations, along with observations with missing outcome for either method 1 (n=42) or method 2 (n=46). Overall, IPOS data was imputed for 316 observations with one to seven missing IPOS items.

For estimation method 1 (*unresolved symptoms and concerns*), the mean total IPOS score was 36.1 (SD 12.2, ranging 0 to 66.9). Overall, 696 of 1135 (61.3%) decedents had a score ≥34, indicating unmet palliative care needs (see Supplement file 2 for details about score distribution, IPOS data completeness, and descriptives of the excluded group with ≥8 missing IPOS items).

Using estimation method 2 (*insufficient care provision*), 655 of 1130 (58.0%) decedents were reported to not have received enough help or no help at all from GPs in the last three months of life (response distribution provided in Supplement Table 3).

After combining estimation methods 1 and 2 (see Supplement Table 4), 429 of 1104 decedents (38.9%) were identified as having have unmet palliative care needs using the conservative estimate (both method 1 and 2). The broad estimate (either method 1 or 2) identified that 886 of 1104 (80.3%) had unmet palliative care needs. Prevalence of unmet palliative care needs by decedent characteristics are reported for estimate 1 and 2 in Table 1 (Supplement Table 5 shows prevalence by decedent characteristics using estimates 3 and 4). Age-, gender- and nation-specific proportions for each sample estimate are provided in Supplement Table 6.

Table 1. Sociodemographic and clinical characteristics of decedents in the Better End of Life survey sample (n=1,194).

	Full sample (n=1194)	Estimate 1: Unresolved symptoms and concerns			Estimate 2: Insufficient care provision		
		No unmet needs (n=447)	Unmet needs (n=705)	Missing (n=42)	No unmet needs (n=482)	Unmet needs (n=666)	Missing (n=46)
	n (col %)	n (row %)			n (row %)		
Gender							
Male	571 (47.8%)	209 (36.6%)	334 (58.5%)	28 (4.9%)	235 (41.2%)	311 (54.5%)	25 (4.4%)
Female	615 (51.5%)	236 (38.4%)	365 (59.4%)	14 (2.3%)	245 (39.8%)	349 (56.8%)	21 (3.4%)
Missing	8 (0.7%)	2 (25.0%)	6 (75.0%)	-	2 (25.0%)	6 (75.0%)	-
Age							
18-64	109 (9.1%)	34 (31.2%)	71 (65.1%)	4 (3.7%)	38 (34.9%)	64 (58.7%)	7 (6.4%)
65-84	529 (44.3%)	160 (30.3%)	349 (66.0%)	20 (3.8%)	196 (37.1%)	313 (59.2%)	20 (3.8%)
≥85	543 (45.5%)	247 (45.5%)	279 (51.4%)	17 (3.1%)	243 (44.8%)	281 (51.8%)	19 (3.5%)
Missing	13 (1.1%)	6 (46.2%)	6 (46.1%)	1 (7.7%)	5 (38.5%)	8 (61.5%)	-
Ethnicity							
White	1163 (97.4%)	439 (37.8%)	683 (58.7%)	41 (3.5%)	472 (40.6%)	647 (55.6%)	44 (3.8%)
Non-white	27 (2.3%)	7 (25.9%)	20 (74.1%)	-	10 (37.0%)	15 (55.6%)	2 (7.4%)
Missing	4 (0.3%)	1 (25.0%)	2 (50.0%)	1 (25.0%)	-	4 (100.0%)	-
Financial circumstances							
Living comfortably	545 (45.6%)	244 (44.8%)	279 (51.2%)	22 (4.0%)	234 (42.9%)	287 (52.7%)	24 (4.4%)
Doing alright	398 (33.3%)	138 (34.7%)	252 (63.3%)	8 (2.0%)	158 (39.7%)	227 (57.0%)	13 (3.3%)
Just about getting by or finding it difficult	231 (19.4%)	61 (26.4%)	164 (71.0%)	6 (2.6%)	85 (36.8%)	140 (60.6%)	6 (2.6%)
Missing	20 (1.7%)	4 (20.0%)	10 (50.0%)	6 (30.0%)	5 (25.0%)	12 (60.0%)	3 (15.0%)
Nation							
England	619 (51.8%)	238 (38.5%)	361 (58.3%)	20 (3.2%)	233 (37.6%)	364 (58.8%)	22 (3.6%)
Wales	575 (48.2%)	209 (36.4%)	344 (59.8%)	22 (3.8%)	249 (43.3%)	302 (52.5%)	24 (4.2%)
Cause of death							
Cancer	340 (28.5%)	97 (28.5%)	240 (70.6%)	3 (0.9%)	139 (40.9%)	193 (56.8%)	8 (2.4%)
Heart disease	172 (14.4%)	82 (47.7%)	85 (49.4%)	5 (2.9%)	77 (44.8%)	93 (54.1%)	2 (1.2%)
Stroke	40 (3.4%)	22 (55.0%)	17 (42.5%)	1 (2.5%)	13 (32.5%)	26 (65.0%)	1 (2.5%)

Lung disease	59 (4.9%)	26 (44.1%)	32 (54.2%)	1 (1.7%)	23 (39.0%)	34 (57.6%)	2 (3.4%)
Liver disease	19 (1.6%)	4 (21.1%)	13 (68.4%)	2 (10.5%)	5 (26.3%)	12 (63.2%)	2 (10.5%)
Kidney disease	8 (0.7%)	4 (50.0%)	4 (50.0%)	-	4 (50.0%)	3 (37.5%)	1 (12.5%)
Neurological disease	21 (1.8%)	9 (42.9%)	9 (42.9%)	3 (14.3%)	12 (57.1%)	7 (33.3%)	2 (9.5%)
Dementia	177 (14.8%)	70 (39.6%)	97 (54.8%)	10 (5.7%)	83 (46.9%)	81 (45.8%)	13 (7.3%)
Other*	342 (28.6%)	132 (38.6%)	201 (58.8%)	9 (2.6%)	123 (36.0%)	209 (61.1%)	10 (2.9%)
Missing	16 (1.3%)	1 (6.3%)	7 (43.8%)	8 (50.0%)	3 (18.8%)	8 (50.0%)	5 (31.3%)
Comorbidities							
None	415 (34.8%)	188 (45.3%)	205 (49.4%)	22 (5.3%)	167 (40.2%)	221 (53.3%)	27 (6.5%)
1 condition	392 (32.8%)	146 (37.2%)	231 (58.9%)	15 (3.8%)	154 (39.3%)	223 (56.9%)	15 (3.8%)
2 conditions	240 (20.1%)	77 (32.1%)	159 (66.3%)	4 (1.7%)	94 (39.2%)	142 (59.2%)	4 (1.67%)
3 conditions or more	147 (12.3%)	36 (24.5%)	110 (74.8%)	1 (0.7%)	67 (45.6%)	80 (54.4%)	-

* Including observations with two or more reported causes of death.

Associations between decedent characteristics and unmet palliative care needs

Estimation method 1: Unresolved symptoms and concerns

In the unadjusted model, there was higher prevalence of unmet needs among decedents aged 18-64 and 65–84 compared to those aged 85 years or older. This association slightly attenuated in the adjusted model and only remained statistically significant for those aged 65–84. A gradient between unmet palliative care needs and financial circumstance was observed in the unadjusted model, with higher prevalence among decedents who were *"just about getting by or finding it difficult"* or *"doing alright"* compared to those *"living comfortably"*. This gradient remained significant in the adjusted model. Prevalence of unmet needs was higher among decedents who died from cancer compared to those who died from other conditions in both the unadjusted model and adjusted model. A gradient was also observed between unmet needs and comorbidities, with prevalence of unmet needs increasing with higher numbers of comorbidities. This association was slightly strengthened after adjustment.

Estimation method 2: Insufficient care provision

In the unadjusted model, decedents aged 65–84 had higher risk of unmet palliative care needs compared with those aged 85 years or older, and this association slightly strengthened after adjustment. Compared to those *"living comfortably"*, unmet needs were more prevalent among decedents who were *"just about getting by or finding it difficult"*. However, this association was attenuated in the adjusted model. Prevalence of unmet needs was lower among decedents in Wales compared to those in England, though this association attenuated and was no longer statistically significant in the adjusted model. Compared to those who died from other conditions, those who

died from dementia had lower prevalence of unmet needs, and this association remained significant after adjustment.

The PRs for the decedent characteristics in each adjusted model are illustrated in Figure 1. Results from both unadjusted and adjusted models are presented in Supplementary table 7.

[Figure 1 about here]

Population-level estimates

In 2022, there were 536,311 adult deaths (age 20+) in England and 35,511 in Wales. Of those, 404,995 (75.5%) and 26,996 (76.0%) decedents, respectively, died from one of the main causes of death considered amenable to palliative care (ICD-10 codes listed in Supplement table 1). The four population-level point estimates of unmet palliative care needs, with upper and lower estimates based on the 95% confidence intervals of sample estimates, are reported separately for England and Wales in Table 2. Age-, gender- and nation-specific population-level estimates of unmet needs are reported in Supplement table 8.

Future projections (2025-2050)

Applying age- and gender-specific estimates to national mortality projections indicates that the number of adults in England and Wales with unmet palliative care needs will rise markedly over the next 25 years. By 2050, the annual number of adults with unmet palliative care needs is projected to increase by 21–26% in England and by 14–19% in Wales compared with 2025, depending on the estimation method used (see Figure 2). Depending on the estimate used, it is projected that the number of people with unmet palliative care needs will increase by 39,000–78,700 in England and by 1,400–4,300 in Wales between 2025 and 2050. The greatest absolute increase in prevalence is expected among adults aged 85 years or older (see Supplement table 9).

Table 2. Estimated numbers and proportions of adult decedents (≥20) in England and Wales with unmet palliative care needs in 2022 using the four estimates.

	England (N=536,311)		Wales (N=35,511)	
Estimate	Estimated *n* (%)	Lower–Upper estimate* (%)	Estimated *n* (%)	Lower–Upper estimate* (%)
1. Symptoms and concerns	247,993 (46.2)	210,506–285,479 (39.3–53.2)	17,209 (48.5)	14,576–19,841 (41.0–55.9)
2. Insufficient care provision	244,612 (45.6)	205,812–283,412 (38.4–52.8)	15,280 (43.0)	12,575–17,986 (35.4–50.6)
3. Conservative estimate (1 **and** 2 combined)	169,537 (31.6)	129,959–209,116 (24.2–39.0)	10,246 (28.9)	7,536– 12,956 (21.2–36.5)
4. Broad estimate (either	325,113	293,657–356,568	22,109	20,063–24,156

| | 1 or 2) | (60.6) | (54.8–66.5) | (62.3) | (56.5–68.0) |

* Lower and upper estimates for population prevalence derived from applying the upper and lower 95% confidence interval limits for each sample estimate to the population data.

[Figure 2 about here]

Discussion

To our knowledge, this is the first study to develop robust empirical estimates of unmet palliative care needs and apply these to all adult decedents at a nation-level. We present two estimation methods and four estimates of unmet palliative care needs, developed from existing literature and with stakeholder engagement, accounting for both objective and subjective aspects of the construct. We show that prevalence of unmet palliative care needs is high, irrespective of estimate used: between a third and two-thirds of all deaths in England and Wales in 2022 are estimated to have unmet palliative care needs. The risk of having unmet palliative care needs was higher among those aged 65-84, people in poorer financial circumstances, and those who had a higher number of comorbidities. Future unmet palliative care needs are projected to increase markedly, by 21–26% in England and 14–19% in Wales by 2050, with the largest absolute rise among adults aged 85 years or older.

Our findings suggest that even in the UK, which has a well-developed system of palliative care, provision for people approaching the end of life is falling short. Workforce planning and resource allocation must account for the projected rise in unmet palliative care needs, especially among the oldest age groups, where mortality will increase most rapidly, and in primary care services. This will be important if the ambitions of 'ageing in place' and shifting from acute to community-based care outlined in health care policy recommendations in the UK[33 34] and internationally[35-37] are to be achieved. GPs and community nurses already provide a large part of the care dying people receive in community settings,[38] and patients and families often expect that their GP will be involved in provision of end-of-life care.[39] The prevalence of unmet needs using estimation method 2, which concerns primary palliative care (specifically GP provision) was high in this study (45.6% in England and 43.0% in Wales), and the ongoing shift to more community-based care will increase pressures on primary and community-based care services further. A previous comparative study of 16 countries also identified private homes and nursing homes as the settings where need for palliative care provision is the highest.[40] Integration of palliative care into primary care is also highlighted by the World Health Organization as a crucial step to improve access to palliative and end-of-life care globally.[41]

Our study makes several important contributions to the evidence base on unmet palliative care needs. We address an evidence gap concerning the lack of operational clarity in measuring unmet palliative care needs identified in recent reviews.[15 16 42] We advance the field by developing two estimation methods to generate four clearly operationalised estimates to quantify unmet palliative care needs, and demonstrate how these compare within one nationally representative sample of decedents who died from causes of death amenable to palliative care. These methods for estimation offer potential for use in other nations with comparable health systems.

While no estimation method is perfect, person-centred outcome measures completed by patients are arguably best current practice for assessing unmet palliative care needs. Service access measures are often used as an indicator of unmet palliative care needs at the population level[43] but are limited as they do not capture whether the services accessed were effective in addressing patients' symptoms and concerns. For example, in 2011 it was estimated that of all annual deaths in England with palliative care needs, 20-25% were not receiving specialist palliative care and thus had unmet needs.[17] These estimations, however, did not regard of subjective experiences of unmet palliative care needs. Routine national collection of validated person-centred outcome measures that identify the nature of a patient's symptoms and concerns and the extent to which care has resolved or improved them may therefore provide the most comprehensive measurement of unmet palliative care needs (aligning with the conservative estimate in this study). Such measures would provide important opportunity to monitor care quality and improve service delivery.[44 45] Despite the availability of numerous validated tools designed to assess unmet palliative care needs,[42 46] their implementation into clinical practice remains limited. However, development of a prototype national Palliative Care Outcomes Registry,[47] together with evidence from other countries[48 49] suggest that routine use of person-centred outcome measures is feasible. Inclusion of self-report measures of unmet palliative care needs in nationally representative surveys is another possible means of assessment.[50 51]

Several socio-demographic and clinical factors were associated with the prevalence of unmet palliative care needs across both estimation method 1 (*unresolved symptoms and concerns*) and 2 (*insufficient care provision*). Patterns were broadly consistent between methods, except for comorbidities where method 1 showed a clear gradient of higher prevalence of unmet needs and with higher number of comorbidities, whereas method 2 did not. This divergence likely reflects limitations with capturing the relationship using method 2 rather than actual lower prevalence among people with multiple conditions, given previous research findings of high levels of healthcare consumption and unmet needs for primary care for this group.[52-54] A gradient was also observed between unmet needs and financial hardship for both methods, though only statistically significant for method 1, which aligns with existing evidence that people with lower socioeconomic position experience poorer healthcare outcomes and greater symptom burden towards the end of life.[55 56]

For both estimation methods, unmet needs were more prevalent among decedents aged 65-84 than those aged 85 or older. This could in part be due to reduced insight about GP care provision among survey respondents reporting for relatives in the oldest age group, many of whom lived in care homes. This explanation might also relate to the lower prevalence of unmet needs among those who died from dementia using method 2, which diverges from research identifying barriers to care access for people with dementia[57]. No association between ethnicity and the risk of unmet palliative care needs was observed in this study, likely due to the small number of non-White British decedents in the Better End of Life survey sample and the use of binary ethnicity categories. The proportion of white decedents in our sample reflects national demographic patterns for older populations in England and Wales,[58] however the aggregation of non-white ethnicities is likely to mask important differences between ethnic minority groups. In future surveys, attempts to oversample in areas of high ethnic diversity could help to provide more data to explore differences in unmet need by ethnicity.

Strengths and limitations

A key strength of this study is the application of multiple approaches to estimate unmet palliative care needs. Our two estimation methods operationalise different aspects of unmet palliative care needs, adopting a needs-based rather than diagnosis-based perspective that is more closely aligned with the palliative care approach. The estimation methods were based on existing measurement approaches identified in our scoping review and informed by stakeholder-prioritised elements of unmet needs. We derived our sample estimates from the Better End of Life survey, the first nationally representative post-bereavement survey in England and Wales since VOICES in 2015.[59]

Some limitations to this study should be noted. First, the Better End of Life survey was not designed to directly measure *unmet* palliative care needs. The sample estimates applied here should be tested in other studies, to compare findings and address limitations identified for this study. Survey data comprise proxy-reports by bereaved family members; although non-family respondents are included, decedents without family are less well represented in these estimates. Furthermore, respondents who were not closely involved in their relative's care might have less insight into their needs and whether these were met or not. IPOS data on unresolved symptoms and concerns used in estimation method 1 relate to the last week of life; while this does not capture the extent to which needs were addressed earlier in the illness trajectory, it remains a proxy for good prior managements. Real-time recording of symptoms and concerns over time would allow for more comprehensive understanding of individual needs for palliative and end-of-life care and the extent to which these are met. For estimation method 2, we only focused on primary palliative care (i.e., receiving enough help from GPs in the last 3 months of life). A method for estimating unmet needs for specialist palliative care services specifically is an important area for future research.

There are also limitations for the population-level estimates. We only consider unmet palliative care needs at the end of life among people who died; unmet palliative care needs earlier in the illness trajectory are not reflected in these estimates. We may be missing unmet needs among other causes of death that are non-sudden but not included in the list of ICD-10 codes. It should be noted that for the projections we assumed that the distribution of causes of death would remain constant over time and did not account for rising of prevalence of major disease groups such as cancer and dementia. Projections also do not account for multiple-long term conditions, which is common, especially in older age groups,[60 61] and will continue to grow, likely increasing the complexity of needs. Importantly, we assumed in the projections that service delivery will remain stable. In the UK, hospices are increasingly struggling with funding, with a rise in closures, which will likely have knock-on effects on care access. Taken together, these considerations mean that population-level estimates and future projections of unmet palliative care needs might be underestimated.

Conclusion

This study has developed empirical methods for estimating unmet palliative care needs at a population-level. We show that prevalence of unmet needs is high regardless of estimate used and projected to increase by 2050. Our study contrasts estimation methods based on unresolved symptoms and concerns and insufficient care provision for planning and evaluating of equitable care at the end of life depending on which type of metric is available at population-level.

Figures

Figure 1. Forest plot of adjusted prevalence ratios (PR) with 95% confidence intervals for associations between decedent characteristics and unmet palliative care needs using estimation methods 1 and 2.

Notes: Prevalence ratios adjusted for all covariates in the figure.

Figure 2. Projected prevalence of unmet palliative care needs among adult decedents (≥20) in England and Wales (2025–2050), based on levels in 2022 (assuming no major policy or service changes).

Notes: The shaded area for each plot illustrates upper and lower population-level estimates applied to mortality projections

Author declarations

Funding statement:

This work was supported by Marie Curie (grant MCCRP-23-02). The Better End of Life Survey was funded by the Marie Curie grant MCSON-20-102. Marie Curie supports research into palliative and end of life care to improve the care that is provided to people affected by any terminal illness. For more information visit http://www.mariecurie.org.uk/.

K.E.S. is the Laing Galazka Chair in palliative care at King's College London, funded through an endowment from Cicely Saunders International and the Kirby Laing Foundation. AF is funded by a Marie Curie Senior Research Fellowship (MCRFS-20-101). I.J.H. is an UK National Institute for Health and Care Research (NIHR) Senior Investigator (Emeritus), F.E.M.M is an NIHR Senior Investigator, and L.K.F is an NIHR Research Professor. The views expressed in this article are those of the authors and not necessarily those of the UK NIHR, or the UK Department of Health and Social Care.

Competing interests statement:

The authors declare no competing interests with respect to the research, authorship, or publication of this article.

Data sharing statement:

All data for the population-level estimates and projections are publicly available on the ONS website (2022 mortality registry data: https://www.nomisweb.co.uk/datasets/mortsa; national mortality projections: https://www.ons.gov.uk/peoplepopulationandcommunity/populationandmigration/populationprojections/datasets/2014basednationalpopulationprojectionstableofcontents).

Contributors:

Contributors TJ, AEB and FEMM accept full responsibility for the finished work and the conduct of the study, analysed the data and controlled the decision to publish. AEB, FEMM, KES, AF, TJ, JMD, LF and IJH contributed to the conception and design of the study. TJ and KES acquired and curated the Better End of Life Survey data. TJ and AEB performed the statistical analyses and drafted the manuscript. TJ, AEB, FEMM, AF, KES, JMD, IJH, LF and MD contributed to analysis and interpretation of the data and critically reviewed the manuscript for important intellectual content. TJ, AEB and FEMM are guarantors for the overall content. The corresponding author (AB) attests that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted.

The authors would like to acknowledge Joel Bates for his contributions to data visualisation, and Margaret Ogden, David Chandler, and Pam Smith from our Public and Patient Involvement (PPI) group for critical input to refining estimation methods and interpretation of findings.

References

1. Bone AE, Gomes B, Etkind SN, et al. What is the impact of population ageing on the future provision of end-of-life care? Population-based projections of place of death. *Palliative medicine* 2018;32(2):329-36. doi: 10.1177/0269216317734435
2. Etkind SN, Bone AE, Gomes B, et al. How many people will need palliative care in 2040? Past trends, future projections and implications for services. *BMC Med* 2017;15(1):102. doi: 10.1186/s12916-017-0860-2 [published Online First: 2017/05/19]
3. Kingston A, Robinson L, Booth H, et al. Projections of multi-morbidity in the older population in England to 2035: estimates from the Population Ageing and Care Simulation (PACSim) model. *Age Ageing* 2018;47(3):374-80. doi: 10.1093/ageing/afx201
4. Murtagh FE, Bausewein C, Verne J, et al. How many people need palliative care? A study developing and comparing methods for population-based estimates. *Palliative medicine* 2014;28(1):49-58.
5. Dixon J, King D, Matosevic T, et al. Equity in the provision of palliative care in the UK: review of evidence: Personal Social Services Research Unit, London School of Economics and Political Science, 2015.
6. Fantoni ER, Wynne N, Finucane AM. Estimates of population-level palliative care need in the UK: a descriptive analysis of mortality data before and during the COVID-19 pandemic. *Bmc Palliat Care* 2024;23(1):271. doi: 10.1186/s12904-024-01574-5 [published Online First: 20241125]
7. Kitreerawutiwong N, Kitreerawutiwong K, Keeratisiroj O, et al. Methods used to identify the prevalence of palliative care needs: An integrative review. *Palliat Support Care* 2023:1-16. doi: 10.1017/S1478951523000494 [published Online First: 20230522]
8. Jeba J, Taylor C, O'Donnell V. Projecting palliative and end-of-life care needs in Central Lancashire up to 2040: an integrated palliative care and public health approach. *Public Health* 2021;195:145-51. doi: 10.1016/j.puhe.2021.04.018 [published Online First: 20210608]
9. Rosenwax L, Mcnamara B, Blackmore A, et al. Estimating the size of a potential palliative care population. *Palliative medicine* 2005;19(7):556-62.
10. Scholten N, Günther AL, Pfaff H, et al. The size of the population potentially in need of palliative care in Germany-an estimation based on death registration data. *BMC palliative care* 2016;15:1-7.
11. Higginson IJ, Hart S, Koffman J, et al. Needs assessments in palliative care: an appraisal of definitions and approaches used. *Journal of pain and symptom management* 2007;33(5):500-05.
12. McNamara B, Rosenwax LK, Holman CD. A method for defining and estimating the palliative care population. *J Pain Symptom Manage* 2006;32(1):5-12. doi: 10.1016/j.jpainsymman.2005.12.018
13. Slotina E, Ditscheid B, Meissner F, et al. A refined ICD-10 diagnoses-based approach for retrospective analysis of potential palliative care need and coverage in claims data of deceased. *SAGE Open Med* 2024;12:20503121241269599. doi: 10.1177/20503121241269599 [published Online First: 20240813]
14. Gómez-Batiste X, Martínez-Muñoz M, Blay C, et al. Prevalence and characteristics of patients with advanced chronic conditions in need of palliative care in the general population: a cross-sectional study. *Palliative medicine* 2014;28(4):302-11.
15. Wang T, Molassiotis A, Chung BPM, et al. Unmet care needs of advanced cancer patients and their informal caregivers: a systematic review. *Bmc Palliat Care* 2018;17(1):96. doi: 10.1186/s12904-018-0346-9 [published Online First: 20180723]
16. Johansson T, Diggle M, Finucance A, et al. Defining and measuring unmet palliative care needs among people with life-limiting illness: a scoping review of international evidence. *medRxiv* 2025:2025.07. 07.25330993.

17. Hughes-Hallett T, Craft A, Davies C, et al. Palliative care funding review. Independent report for the Secretary of State for Health, 2011.
18. Bone AE, Diggle M, Johansson T, et al. Coproducing a conceptual understanding of unmet palliative care needs: stakeholder workshops using modified nominal group technique. *Bmc Palliat Care* 2025 doi: 10.1186/s12904-025-01971-4 [published Online First: 20251223]
19. Johansson T, Pask S, Goodrich J, et al. Time to care: Findings from a nationally representative survey of experiences at the end of life in England and Wales. London, UK: King's College London, Cicely Saunders Institute; Hull York Medical School at the University of Hull; and University of Cambridge, 2024.
20. Office for National Statistics. Mortality statistics-underlying cause, sex and age. https://www.nomisweb.co.uk/datasets/mortsa (accessed 10 October 2025), 2024.
21. Office for National Statistics. National population projections: Projected deaths. https://www.ons.gov.uk/peoplepopulationandcommunity/populationandmigration/populationprojections/datasets/2014basednationalpopulationprojectionstableofcontents (accessed 17 July 2025), 2025.
22. Vandenbroucke JP, Von Elm E, Altman DG, et al. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): explanation and elaboration. *International journal of surgery* 2014;12(12):1500-24.
23. Gomes B, McCrone P, Hall S, et al. Variations in the quality and costs of end-of-life care, preferences and palliative outcomes for cancer patients by place of death: the QUALYCARE study. *BMC Cancer* 2010;10:1-10.
24. Addington-Hall J, McPherson C. After-death interviews with surrogates/bereaved family members: some issues of validity. *Journal of pain and symptom management* 2001;22(3):784-90.
25. Murtagh FE, Ramsenthaler C, Firth A, et al. A brief, patient-and proxy-reported outcome measure in advanced illness: Validity, reliability and responsiveness of the Integrated Palliative care Outcome Scale (IPOS). *Palliative medicine* 2019;33(8):1045-57.
26. Munday D, Boyd K, Jeba J, et al. Defining primary palliative care for universal health coverage. *Lancet* 2019;394(10199):621-22. doi: 10.1016/S0140-6736(19)31830-6
27. Murray SA, Pastrana T. Primary palliative care: Onwards and upwards! *Palliat Med* 2024;38(8):762-65. doi: 10.1177/02692163241271049 [published Online First: 20240909]
28. Murray SA, Firth A, Schneider N, et al. Promoting palliative care in the community: production of the primary palliative care toolkit by the European Association of Palliative Care Taskforce in primary palliative care. *Palliat Med* 2015;29(2):101-11. doi: 10.1177/0269216314545006 [published Online First: 2014/11/15]
29. Newcombe RG. Two-sided confidence intervals for the single proportion: comparison of seven methods. *Statistics in medicine* 1998;17(8):857-72.
30. Zou G. A modified poisson regression approach to prospective studies with binary data. *American journal of epidemiology* 2004;159(7):702-06.
31. StataCorp. Stata Statistical Software: Release 18. . College Station, TX: StataCorp LLC, 2023.
32. World Medical Association. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA* 2013;310(20):2191-94.
33. NHS England. Fit for the future: 10 year health plan for England. NHS England, 2025.
34. Welsh Government. A healthier Wales: long term plan for health and social care, 2018.
35. Policy RCoN, Department I. Moving care to the community: an international perspective: Royal College of Nursing 2013.
36. Organization WH. Integrated care for older people (ICOPE): guidance for person-centred assessment and pathways in primary care: World Health Organization 2025.
37. Assembly WH. Strengthening of palliative care as a component of comprehensive care throughout the life course. *Resolution of May 24, 2014 at the 67th session of the World Health Assembly (WHA67 19)* 2014

38. Mitchell GK, Senior HE, Johnson CE, et al. Systematic review of general practice end-of-life symptom control. *BMJ Support Palliat Care* 2018;8(4):411-20. doi: 10.1136/bmjspcare-2017-001374 [published Online First: 20180120]

39. Carmont SA, Mitchell G, Senior H, et al. Systematic review of the effectiveness, barriers and facilitators to general practitioner engagement with specialist secondary services in integrated palliative care. *BMJ Support Palliat Care* 2018;8(4):385-99. doi: 10.1136/bmjspcare-2016-001125 [published Online First: 20170214]

40. Morin L, Aubry R, Frova L, et al. Estimating the need for palliative care at the population level: A cross-national study in 12 countries. *Palliative medicine* 2017;31(6):526-36. doi: 10.1177/0269216316671280 [published Online First: 2016/09/30]

41. World Health Organization. Integrating palliative care and symptom relief into primary health care: a WHO guide for planners, implementers and managers. 2018

42. Schmidt EB, Blum D, Domeisen Benedetti F, et al. Tools for guiding interventions to address patient-perceived multidimensional unmet healthcare needs in palliative care: Systematic literature review. *BMJ Supportive and Palliative Care* 2023;13(e1):E1-E9. doi: https://dx.doi.org/10.1136/bmjspcare-2020-002495

43. Franks PJ, Salisbury C, Bosanquet N, et al. The level of need for palliative care: a systematic review of the literature. *Palliative medicine* 2000;14(2):93-104.

44. Currow DC, Allingham S, Yates P, et al. Improving national hospice/palliative care service symptom outcomes systematically through point-of-care data collection, structured feedback and benchmarking. *Supportive Care in Cancer* 2015;23(2):307-15.

45. Etkind SN, Daveson BA, Kwok W, et al. Capture, transfer, and feedback of patient-centered outcomes data in palliative care populations: Does it make a difference? A systematic review. *Journal of pain and symptom management* 2015;49(3):611-24. doi: https://dx.doi.org/10.1016/j.jpainsymman.2014.07.010

46. Shunmugasundaram C, Rutherford C, Butow PN, et al. Content comparison of unmet needs self-report measures used in patients with head and neck cancer: A systematic review. *Psycho-Oncology* 2019;28(12):2295-306. doi: 10.1002/pon.5257

47. RESOLVE. Resolving symptoms in cancer: Wolfson Palliative Care Research Centre, University of Hull; 2025 [Available from: https://www.hyms.ac.uk/research/research-centres-and-groups/wolfson/projects/resolve accessed 19 December 2025.

48. Palliative Care Outcomes Collaboration (PCOC). University of Wollongong; 2020 [Available from: https://www.uow.edu.au/australasian-health-outcomes-consortium/pcoc/ accessed 19 December 2025.

49. Martinsson L, Heedman PA, Lundstrom S, et al. Improved data validity in the Swedish Register of Palliative Care. *PLoS One* 2017;12(10):e0186804. doi: 10.1371/journal.pone.0186804 [published Online First: 20171019]

50. Kowal P, Corso B, Anindya K, et al. Prevalence of unmet health care need in older adults in 83 countries: measuring progressing towards universal health coverage in the context of global population ageing. *Popul Health Metr* 2023;21(1):15. doi: 10.1186/s12963-023-00308-8 [published Online First: 20230915]

51. Rosenberg M, Kowal P, Rahman MM, et al. Better data on unmet healthcare need can strengthen global monitoring of universal health coverage. *BMJ* 2023;382:e075476. doi: 10.1136/bmj-2023-075476 [published Online First: 20230905]

52. Cassell A, Edwards D, Harshfield A, et al. The epidemiology of multimorbidity in primary care: a retrospective cohort study. *British Journal of General Practice* 2018;68(669):e245-e51.

53. Damarell RA, Morgan DD, Tieman JJ. General practitioner strategies for managing patients with multimorbidity: a systematic review and thematic synthesis of qualitative research. *BMC Fam Pract* 2020;21(1):131. doi: 10.1186/s12875-020-01197-8 [published Online First: 20200701]

54. Rolewicz L, Keeble E, Paddison C, et al. Are the needs of people with multiple long-term conditions being met? Evidence from the 2018 General Practice Patient Survey. *BMJ open* 2020;10(11):e041569. doi: 10.1136/bmjopen-2020-041569 [published Online First: 20201114]

55. Bowers SP, Chin M, O'Riordan M, et al. The end of life experiences of people living with socio-economic deprivation in the developed world: an integrative review. *Bmc Palliat Care* 2022;21(1):193. doi: 10.1186/s12904-022-01080-6 [published Online First: 20221105]

56. Davies JM, Sleeman KE, Leniz J, et al. Socioeconomic position and use of healthcare in the last year of life: A systematic review and meta-analysis. *PLoS Med* 2019;16(4):e1002782. doi: 10.1371/journal.pmed.1002782 [published Online First: 20190423]

57. Williamson LE, Sleeman KE, Evans CJ. Exploring access to community care and emergency department use among people with dementia: A qualitative interview study with people with dementia, and current and bereaved caregivers. *Int J Geriatr Psychiatry* 2023;38(7):e5966. doi: 10.1002/gps.5966

58. Office for National Statistics (ONS). Profile of the older population living in England and Wales in 2021 and changes since 2011, 2023.

59. Office for National Statistics (ONS). National Survey of Bereaved People (VOICES) England, 2015: Quality of care delivered in the last 3 months of life for adults who died in England., 2016.

60. Barnett K, Mercer SW, Norbury M, et al. Epidemiology of multimorbidity and implications for health care, research, and medical education: a cross-sectional study. *The Lancet* 2012;380(9836):37-43.

61. Brameld K, Spilsbury K, Rosenwax L, et al. Issues using linkage of hospital records and death certificate data to determine the size of a potential palliative care population. *Palliat Med* 2017;31(6):537-43. doi: 10.1177/0269216316673550 [published Online First: 20161024]

Adjusted ORs for estimation methods 1 and 2

Gender (ref male)	Female
Age (ref <45)	45-74
	>=74
Ethnicity (ref white)	Asian
	Black
	Other
Financial circumstances (ref living comfortably)	Doing alright
	Just about getting by
	Finding it difficult
Nation (ref England)	Wales
	Scotland
	Northern Ireland
Cause of death (ref cancer)	Cancer
	Heart disease
	Dementia
	Lung disease
Comorbidities (ref none)	1 condition
	2 conditions
	3 conditions

OR < 1 indicates worse for PC > 1 indicates higher risk

● Estimation method 1: unresolved symptoms and services
▲ Estimation method 2: insufficient care provision

England

Number of deaths

- Estimate 4: +23.7%
- Estimate 2: +25.6%
- Estimate 1: +21.2%
- Estimate 3: +22.5%

Wales

Number of deaths

- Estimate 4: +19.3%
- Estimate 1: +18.1%
- Estimate 2: +17.3%
- Estimate 3: +13.8%

Estimate 1: Unresolved symptoms and concerns
Estimate 2: Insufficient care provision
Estimate 3: Estimates 1 and 2 combined
Estimate 4: Either estimate 1 or 2