Title

Long-read metagenomics and methylation-based binning allow the description of the emerging high-risk antibiotic resistance genes and their hidden hosts in complex communities

Short title

Antibiotic resistance gene carriers revealed by a methylation-based approach

Melina Markkanen[1,2]*, Heidi Putkuri[3], Dovydas Kičiatovas[2,3], Ville Mustonen[2,3], Marko Virta[1,2], Antti Karkman[1,2]*

[1]Department of Microbiology, University of Helsinki, Helsinki, Finland
[2]Multidisciplinary Center of Excellence in Antimicrobial Resistance Research, University of Helsinki, Helsinki, Finland
[3]Department of Computer Science, Department of Organismal and Evolutionary Biology, University of Helsinki, Helsinki, Finland

*Corresponding authors: melina.markkanen@helsinki.fi, antti.karkman@helsinki.fi

Abstract

Antibiotic resistance genes (ARGs) circulating among clinically relevant bacteria pose serious challenges to public health. Given the ancient and environmental bacterial origins of ARGs, a better understanding of the carriers of ARGs beyond the clinically most relevant species is urgently needed for more farsighted resistance monitoring and intervention measures. While the risks of emerging ARGs from environmental sources have been recognized, the identification bottlenecks stem from the limitations of shotgun metagenomic sequencing and bioinformatic methods. Here, we used long-read metagenomic sequencing and bacteria-specific methylation profiles to re-establish the links between established (well-described) or latent (absent in databases) ARGs and their bacterial and genetic contexts in wastewater. The base modification data produced by PacBio SMRT sequencing was analyzed by an in-house pipeline utilizing position weight matrices and UMAP visualizations. The approach was validated by a synthetic community with known bacterial composition. Our analysis revealed several previously unreported ARGs and their hosts with varying risk levels defined by their potential as emerging public health threats. For instance, *Arcobacter*, as one of the prevalent taxa in influent wastewater, was shown to carry a latent beta-lactamase gene with high predicted mobility potential. Of the other emerging beta-lactamases, we provided a real-life example of ongoing p*dif* module-mediated genetic reshuffling of the *bla*MCA gene occurring at least within *Acinetobacter* hosts in our samples. Additionally, we identified *Simplicispira*, Phycisphaerae, and environmental groups of the Bacteroidales order as the carriers of established, clinically important ARGs. These findings support the intermediate host roles of strictly environmental bacteria for the further dissemination of mobilized ARGs, highlighting the importance of exploring the uncultivated, or non-pathogenic, carriers of ARGs for the early detection of newly arising ARGs and mobility mechanisms.

Key words

Antibiotic resistance, latent antibiotic resistance genes, DNA methylation, shotgun metagenomic sequencing, wastewater, uncultivated species, environmental bacteria, gene mobility, integron gene cassettes, p*dif*-modules, IS6100 transposase

Introduction

The introduction of antibiotics into therapeutic practice represents a pivotal milestone in the advancement of medicine (1). However, the simultaneous increase in emergence and circulation of antibiotic resistance genes (ARGs) has severely diminished the efficacy of the antibiotic drugs available in human and veterinary medicine, challenging the entire health care system. According to the most recent estimates, deaths of 4.71 million people alone in 2021 were associated with bacterial resistance, whereas for the year 2050, the estimate stands at 8.2 million (2). Although the ARGs are ancient components of some bacterial genomes owing to the "producer hypothesis" (3–5), their rapid increase and spread across bacterial lineages are driven by the gene mobilization events occurring between bacteria, which is fueled by the selection pressures arising from the clinical use of antibiotics (6). ARG mobilization is enabled by horizontally transferred mobile genetic elements (MGEs) and their adjacent genes. Sequential ARG mobilization events, starting from the source organism and proceeding further toward the putative opportunistic pathogen hosts, may include several steps of so-called intermediate carriers of ARGs (7, 8).

The resistance features carried by the most common opportunistic pathogens are relatively well described owing to their immediate clinical relevance, but also due to practical reasons, as many of these bacteria are cultivable. However, given the complexity of the resistance gene dynamics – including non-pathogenic bacteria – and rapidly evolving resistance situation, a better understanding of the ARG mobility mechanisms and ARG carriers beyond the cultivable clinical strains is urgently needed (7). The study of environmental resistomes enables the identification of both established and latent ARGs – that is, those that are well-characterized in databases and those that are largely unexplored – regardless of the cultivability of their host bacteria (9).

For instance, both established and predicted versatile beta-lactamase genes are highly frequent among environmental bacteria, especially within wastewater communities (5, 9, 10). Indeed, wastewater has been identified as one of the key environments regarding the presence of latent ARGs (9) with elevated risk of becoming clinically problematic if mobilized into relevant opportunistic pathogen hosts (8, 9). This is not difficult to envision due to the high diversity of bacteria present in wastewater originating from both the human body and various environmental sources, striving to persist despite a range of environmental stressors (11). Newly emerging ARGs are expected to arise from the largely unknown repertoire of bacteria inhabiting external environments and commonly referred to as environmental bacteria (8, 12), whose description and detection are still limited in contrast to cultivable human health-related species. The transition from a non-mobile context in environmental bacteria to clinically relevant ARGs circulating among human pathogens contains multiple gene decontextualization and host transition events by

intermediate hosts (7, 8). Unfortunately, considering the prevailing conditions of the post-
antibiotic era, this transfer may be irreversible (13). Several environmental origin bacteria and
intermediate hosts (or recent hosts) of the current clinical ARGs have been resolved (3, 5, 6, 14–
16). Although due to detection difficulties (7), this is typically done retrospectively. Knowledge
about the intermediate ARG carriers is particularly important, as they may supply the foundational
genetic material for the emergence of clinically relevant resistant bacteria (17).
99

Ranking ARGs in environmental resistomes by their impact on human health is inherently difficult
and remains debatable (12, 18). The risk-ranking schemes that have given prominence to mobile
ARGs that are already present in pathogens (18, 19) have been challenged by more farsighted
frameworks, which define novel resistance determinants arising from environmental microbiomes,
especially with the implications of gene mobility, as the most critical contributors to the current
rapidly developing resistance problem. These schemes aim for early capture of emerging resistance
threats by assessing the likelihood of ARG transmission from environmental species to clinically
relevant bacteria and the severity of such an event (12), using ARG mobility as the proxy for this
(7).
109

However, the inability of shotgun metagenomic sequencing (20) and assembly (21) in linking the
ARGs, especially those encoded by MGEs, to their host bacteria has substantially limited our
understanding of the non-pathogenic or environmental carriers of ARGs and has hindered the risk
assessment (7, 12). Even though metagenomic assembly with long-reads may allow simultaneous
identification of ARGs and their near genetic context, their hosts remain unresolved, especially in
highly complex communities, such as influent wastewater (7). The current metagenomic binning
algorithms, relying on sequence composition and coverage, tend to overlook the ARGs residing
within MGEs, for instance, by the inability to connect plasmids to their carrier bacteria due to
differential coverage (22, 23). This remains one of the most significant challenges in the entire
field of microbial metagenomics and antibiotic resistance research, as plasmids carry vital
adaptation-increasing and evolution-driving features (23, 24). Additionally, binning algorithms are
confused by the presence of multiple strains of one species, biased toward the high-abundance taxa
while ignoring the genomes of rare species, which the ARG-carrying bacteria commonly are (9,
25). Consequently, these biases leave us with persistent uncertainties regarding the role of
environmental lineages on the ARG carriage and spread.
125

Bacterial DNA methylation signals are base modifications mediated by target sequence motif-
specific methyltransferase enzymes, often acting as part of the restriction-modification (R-M)
systems (23). As the main epigenetic modification mechanism of bacterial genomes, methylation
has roles in various regulatory functions, gene expression (26, 27), and genome defense against
phages (28) by enabling the distinction between own and foreign DNA (27). Due to the unique set
of methylation-mediating enzymes and their unique target motifs, all sequence material of a
bacterial strain (chromosomes and plasmids) is expected to show strain-specific methylation
profiles (23), allowing their bioinformatic reconnection into bins of connected contigs (14, 23). So
far, the bacterial methylation detection with the latest technology by PacBio and Nanopore,
allowing direct methylation detection without pretreatment (27), has been focused on isolates,
while fewer analysis tools and workflows are available for metagenomic datasets, except those
with low bacterial complexity (23, 29–32). For instance, the computational demands required for
a comprehensive analysis of highly complex wastewater microbial communities with the

MultiMotifMaker algorithm (33) extend beyond what is practically achievable. Tools similar to
Nanomotif (22), which is specifically designed for Nanopore data, are greatly needed to take
advantage of the kinetic data produced by PacBio SMRT sequencing.

Here, we present a pipeline for summarizing the PacBio base modification data using position
weight matrices (PWMs) and Uniform Manifold Approximation and Projection (UMAP) to cluster
metagenomic contigs based on their methylation profiles to generate bins of connected contigs (or
"genome bins" hereafter) and thereby reveal the ARG-carrying bacteria present in wastewater. A
synthetic community with known bacterial composition and available sequence data was used to
validate our methodology. In contrast to individual metagenomic contigs, the ability to form
genome bins, even in incomplete form, provides sufficient genomic context to support predictions
of ARG mobility and host species assignment. We focused on carriers of both established and
latent ARGs detected in influent, effluent, and dried sludge to expand our understanding of the
(hidden) environmental bacterial diversity responsible for the carriage and spread of ARG, as well
as the hosts of uncharacterized ARGs showing potential for transmission to clinically relevant
bacteria.

Materials and Methods

Sample collection and DNA extraction

Detailed sample descriptions of the wastewater metagenomes (PRJEB106782, PRJEB83306)
investigated here are provided in our previous work (14). Briefly, composite samples were
collected from influent (INF1, INF2, INF3), effluent (EFF1, EFF2, EFF3), and dried sludge (SLU1,
SLU2, SLU3) at one of the two wastewater treatment plants responsible for processing wastewater
in the Helsinki metropolitan area in Finland during three days in 2019 (18.3.2019, 20.3.2019,
26.3.2019). For wastewater samples, after filtration through a 0.22 µm membrane, the DNA was
extracted from the filter using the Qiagen DNeasy PowerWater kit with bead beating.

Synthetic community data (PRJEB85727) of the study by Partanen and colleagues (34) and
associated genome sequence data (PRJNA1047486) by Hogle and colleagues (35) were utilized as
the validation data for the correctly reconnected metagenomic contigs. A comprehensive
description of the synthetic communities is provided in the work by Partanen (34) and Hogle and
colleagues (35) and **Supplementary Methods**.

Sequencing and preparatory analyses

Library preparation and PacBio Single Molecule Real-Time (SMRT) sequencing for metagenomic
long-reads of wastewater communities were performed at the Institute of Biotechnology,
University of Helsinki. Highly accurate long reads (HiFi reads) of metagenomic community
samples with kinetic tags were generated from the subreads using PacBio BAM toolkit programs
(https://github.com/PacificBiosciences/pbtk) to allow methylation detection in downstream
analyses. Instead, HiFi reads without the tags were used for metagenomic assembly. Each of the
synthetic and wastewater community samples (five and nine samples, respectively) was assembled

separately using hifiasm-meta (v.0.18.0) (36). Detailed Snakemake (v7.17.1) (37) workflows are
found in https://github.com/melinamarkkanen/Methylation.

Primary data analyses of the metagenomic contigs

HiFi reads with attached kinetic tags were aligned to assembled metagenomic contigs using the
PacBio program pbmm2 (v1.10) (https://github.com/PacificBiosciences/pbmm2/). Comparison of
the observed interpulse duration (IPD) values of each sequence position from aligned reads to
those expected for unmodified sequences allowed the detection of base modifications by the
ipdSummary program (v.3.0) (38). As a result, statistically significant comparisons are gathered
as rows into the resulting General Feature Format (GFF) output files for all contigs separately.
Base modifications are identified as either of types m6A or m4C or of unknown type, referred to
as 'modified base'. Along with the contig name and modification type, the position of the
methylated base and its flanking region of 20 bp upstream and downstream is recorded for the
following steps.

Methylation data conversion to Position Weight Matrices (PWM)

The obtained contig-wise summaries of the detected methylated bases were converted into Position
Weight Matrices (PWM) to be used as unique signatures for further analysis. For that, in-house
Python scripts were used to extract the flanking sequences separately for each modification type
and sample (5 and 9 synthetic and wastewater community samples, respectively) and then convert
those data into PWMs. In the PWM files, each row represented one nucleotide (A, T, G, or C) and
a column indicated each position surrounding the methylated base [−20, 20] (more detailed
description of the constructed PWMs is provided in the Supplementary Methods). To discover the
balance between noise and an adequate amount of data (given the high complexity of wastewater
samples, especially influent) for downstream methylation analysis, the minimum thresholds for
detected instances of modified bases per contig were explored with the data of known bacterial
composition (**Table S1**). Of the tested values (20, 50, 100, 200, and 500), 50 was selected as it
provided the highest model accuracy values for the Random Forest classifier (**Table S2**).

Random Forest classifier

To test our methylation-based approach and the high-dimensional data presentation using PWMs
described above, a Random Forest model was trained and implemented using Scikit-learn (v1.4.2)
(39) Python library on synthetic community data with *a priori* known taxonomical composition
(**Supplementary Methods**). Random Forest model was chosen due to its ability to generate
feature importance scores, offering insights into the relative contribution of individual methylation
types, positions, and bases to the prediction of the origin of bacterial genomes of metagenomic
contigs. Model performance was evaluated using a confusion matrix, which summarizes the
number of correct and incorrect predictions. Overall accuracy was calculated as the proportion of
correctly classified instances to the total number of instances.

231
Visualization of PWMs with the synthetic community data

UMAP (Uniform Manifold Approximation and Projection) (40) was applied to visualize and
reduce the complexity of the high-dimensional data by projecting it into two dimensions. By the
non-linear dimensionality reduction technique of UMAP, known for its ability to preserve both
local and global structures in data (40), this approach allowed us to explore the clusters represented
by methylation profiles. In addition to visual inspection of the clustering of species-wise colored
contigs in UMAP, contigs clustering together were combined into (sample-wise) bins of connected
contigs to confirm their completeness and putative redundancy, which were evaluated by
CheckM2 (v 1.0.1) (41). GTDB-Tk (v2.4.1, database release 220) (42) was used to confirm the
taxonomical identity of the genome bins. We also investigated whether contigs within the same
clusters exhibit similar nucleotide information content by generating sequence logos showing the
nucleotide preferences across the context sequences at each position (43), which, in our case,
illustrates the methylation target motif sequences (**Fig. S4**).

Identification of established and latent ARG-carrying genome bins in wastewater

Established and latent ARGs present in the contigs were predicted using ResFinder (v 4.7.0) (44)
and fARGene (Fragmented Antibiotic Resistance Gene iENntifiEr) (v0.1) (45), respectively. By
color-coding the contigs with ARGs in the UMAP visualization, we directed the focus on the
formation of genome bins with ARGs. The need for manual processing in methylation-based
binning restricted the volume of data that could be examined. This was especially highlighted
regarding the latent ARGs detected by fARGene, with which we decided to focus on the putative
beta-lactamases due to their higher likelihood of serving the antibiotic resistance function and thus
being mobile in contrast to chromosomal housekeeping genes that are incorrectly assigned as
putative resistance genes (10). The clustering contigs were combined similarly to the synthetic
community data, and the quality and the taxonomical identity of the resulting genome bins were
evaluated similarly as described for the synthetic community above.

Analysis of ARG mobility potential and genetic context

Five bins of connected contigs with established or latent ARGs were investigated in more detail
as example cases representing newly resolved ARG-host linkages. These genome bins were
determined using evidence from the initial exploratory phase: The established or latent ARG was
detected in multiple different contigs and potentially hosted by different species, indicated by its
location in separate clusters of the UMAP visualization (**Fig. S8**).

The following genetic context analysis was performed for all five examined bins of connected
contigs with ARGs: In addition to the ARG present in the binned contigs, hits to the same ARGs
were queried among all wastewater data using BLAST (46). ARG flanking regions (5 or 10 kb up-
and downstream of the ARG) of the ARG of the matching sequences (including both those binned
using methylation and those that remained unbinned due to scarce or absent methylation data) were
extracted using an in-house script applying SeqKit tools (v 2.5.1) (47). Additional mobility

predictions for the entire contigs or extracted flanking regions, respectively, were performed using
geNomad (v1.8.0) (48) and pdifFinder (49). Bakta (v1.11.0) (50) was used for annotating the genes
in the flanking regions. To examine the diversity within the genetic contexts and to reduce the
number of sequences for visualization purposes, the genetic contexts were dereplicated to exclude
identical or similar genetic contexts by vsearch (v2.22.1) (51). After removing the redundancy of
the contexts, the ARGs in those that remained were extracted for phylogenetic tree analysis. For
that, gene alignment was done using MAFFT (v7.505) (52), for which the tree was computed by
RAxML (command raxmlHPC-PTHREADS) (v8.2.12) (53) and visualized with Interactive Tree
of Life (iTOL) (54).
286
The flanking sequences were visualized using pyGenomeViz (v1.6.1) (55) using a BLAST (46)
query for highlighting the aligned nucleotide sequences between the compared sequences. The
insertion sequences located near the investigated ARGs were examined in more detail by the
ISfinder (56) online server and the *attC* sites by the HattCI program (57). Protein structures of the
Arcobacter encoded latent beta-lactamase genes were predicted using AlphaFold (58), and similar
structures were queried by FoldSeek (59).
293
294
295 **Taxonomical composition**
296
297 Relative abundance of genus-level taxa in both synthetic and wastewater communities was
298 analyzed using Sylph (sylph (v 0.8.1) and sylph-tax (v 1.5.1)), with metagenomic reads as the input
299 (60). For the wastewater data, genera with relative abundance below 0.5 % were left out of the
300 visualization.
301
302
303 **Further details**
304
305 All other figures that were not specifically mentioned above were drawn using Plotly (61) and
306 RStudio (2024.12.1+563.pro5) (62) by applying packages phyloseq (v1.16.2) (63), ggplot2
307 (v3.5.0) (64), and vegan (v2.6-10) (65). Comprehensive descriptions of the materials and methods
308 can be found in the **Supplementary Methods**.
309
310
311 **Data availability**
312
The subread sequence data from the wastewater metagenomes are available in the European
Nucleotide Archive (ENA) under accession number PRJEB106782, along with the corresponding
HiFi reads, which can be found under accession number PRJEB83306. The other sequencing data
used in this study were accessed via the following accession numbers: synthetic community
metagenomes PRJEB85727 (34) and the whole genome sequence (WGS) data of the synthetic
communities PRJNA1047486 (35) (**Table S1**). The genome bin sequences are available in
https://doi.org/10.5281/zenodo.18239683. The analysis scripts are shared in the publicly available
GitHub repository at https://github.com/melinamarkkanen/Methylation.
321
322

Results

Position Weight Matrix (PWM) -based method for linking contigs by their shared methylation signatures

In the absence of suitable tools for methylome analysis of PacBio SMRT sequencing data from complex microbial community samples, such as wastewater, a new workflow was established here (**Fig. 1**). In our approach, the per-contig base modifications were converted into position weight matrices (PWMs), allowing their clustering according to their origin bacterial genome by Uniform Manifold Approximation and Projection (UMAP) to form bins of connected contigs, relying on the assumption that methylation profiles are unique among bacterial strains. We validated our workflow with a synthetic community of known bacterial composition, of which whole genome sequence (WGS) data were available for half of the community members (**Table S1**, **Fig. S1**). Moreover, preliminary analyses with the validation data enabled the exploration of the most appropriate signal-to-noise ratio regarding the amount of the input methylation data (>50 base modifications per contig) for the downstream Random Forest analysis (**Table S2**).

Fig. 1. Schematic representation of the key steps in the workflow. Before the analysis of the wastewater samples, the workflow (until the formation of bins of connected contigs step) was applied on a synthetic community with known bacterial composition as a method validation procedure. Figure created with BioRender.com.

The validation results confirmed that, at least for most community members, the species identity of a metagenomic contig can be predicted based on its methylation profile presented by PWMs (**Fig. S2**, **Table S2**). This was confirmed by the Random Forest confusion matrix, which summarized true and false predictions and provided the basis for calculating test accuracy. The model showed strong performance, with a test accuracy value of 0.88 at the species level (**Table S2**). For the remaining cases, prediction accuracy was low due to misclassifications across many different species, rather than repeated specific incorrect predictions, suggesting rather

heterogeneous and inconclusive methylation profiles instead of substantially overlapping profiles between multiple species (**Fig. S2, Table S2**). This was reflected in the UMAP visualization as a cloud of contigs that were not unique to any single species (**Fig. S3A**). However, as this miscellaneous group of contigs was clearly separate from the more defined, distinct clusters, the risk of false positives coming from this group is diminished (loss of data rather than false positives). In some but not all cases, a higher amount of input methylation data or the contig length seemed to be descriptive for the more distinct species-wise clusters (**Fig. S3B**).

Metagenomic contigs from synthetic communities were clustered based on their methylation profiles, aiming to recover contig relationships lost during shotgun sequencing. To test whether the contig clusters correspond to the original community member genomes, completion and redundancy values, computed by CheckM2 (v 1.0.1) (41), of some selected species metagenome-assembled genomes (MAGs) or bins of connected contigs were examined (**Table S3**). Fully complete MAGs containing both chromosomal and plasmid sequences were resolved for multiple species and samples (**Table S3**). This was also seen in the sequence logos of chromosomal and plasmid contigs showing ± 20 bases surrounding the modified base, differentially sized according to their information content (**Fig. S4**). However, in some cases, only partial genome bins could be curated due to too few methylation information observed as the absence of contigs (exclusion by initial filtering by >50 base modifications/contig) or located apart from the main cluster due to incomplete methylation profile (**Fig. 3A**, **Table S3**). Altogether, we concluded that our approach could reconnect metagenomic contigs into species-wise bins of connected contigs to allow the establishment of ARG-host linkages and description of previously unknown contributors to resistance carriage in wastewater.

General overview of the wastewater bins of connected contigs with ARGs

The miscellaneous clusters of metagenomic contigs in UMAP visualizations of wastewater samples were excluded to decrease noise and obtain a more detailed view of distinctive clusters for the manual bin curation step (**Fig. S7**). Aiming to re-establish the links between resistance genes and their wider contexts, we focused on clusters containing contigs with established or latent ARGs (**Fig. S8**). After quality assessment, 35 genome bins with established (**Table 1**) and 53 with latent (**Table 1**) ARGs were described. The average completion and redundancy values were 55.40 and 2.17, respectively. Of all genome bins, 31 were obtained from influent, 15 from effluent, and 42 from dried sludge (**Table 1**). However, due to the non-quantitative nature of the approach and the included manual steps, the resulting bins of connected contigs cannot be considered a quantitative measure of ARG-encoding bacteria in the samples. On average, the genome bins were composed of 12 contigs (range from 2 to 44) (**Table 1**). In addition, the GC-content of the genome bin corresponded to that of the bin's assigned taxa and their closest matching reference genome by GTDB-tk (**Table 1**).

Given the pivotal role of plasmid-encoded ARGs for the antibiotic resistance problem and the difficulties related to their characterization within whole-community data, we were particularly interested in exploring the plasmid sequences that were linked to their host genome bins by the methylation analysis. We were not able to link any ARG-encoding plasmids to their wider, genome bin context (**Table 1**). However, we could find linkages between non-ARG-carrying plasmids and

their host genome bins in several cases (**Table S4**). For instance, two to five plasmids were linked
to *Acinetobacter* sp. in three different genome bins from influent (bins INF1-C3f, INF1-C6f, and
INF2-C2f) (**Table S4**). Furthermore, single plasmids were identified, for example, for
Aquabacterium_A sp. (bin EFF2-C1f) and *Aeromonas media* (bin INF1-C4) (**Table S4**). Querying
these contigs against the NCBI database with BLAST resulted in matches with plasmid sequences
(data not shown), confirming our ability to connect chromosomal and plasmid sequences.
405
*bla*OXA genes carrying *Aeromonas* sp. were most commonly identified as established ARG
carriers in influent, while Bacteroidales order bacteria carrying *erm*(F) genes were descriptive for
dried sludge. The fewer genome bins with established ARGs obtained from effluent were mainly
Burkholderia order bacteria carrying various established ARGs. Regarding latent ARGs, beta-
lactamases of classes A, B 1-3, C, and D2, most notably of the latter, were identified in more
diverse hosts, including bacteria belonging to 16 different orders such as Burkholderiales,
Bacteroidales, Pseudomonadales, Chitinophagales, Campylobacterales, and Christensenellales, to
name the most common groups. In general, the latent ARG host species included more members
of the uncultivated bacterial lineages compared to the identified carriers of established ARGs. We
next focused on a couple of the most interesting newly identified carriers of established and latent
ARGs (**Table 1**).
417
418
419 **A subset of the latent class D beta-lactamase variants encoded by *Arcobacter* exhibited high**
420 **mobility potential**
421
Arcobacter species of the Campylobacterota phylum were one of the major bacterial groups
detected in influent wastewater (**Fig. S6**). Bins of connected contigs (n = 6, 3, 7, 16) assigned as
Arcobacter spp. and hosting latent class D beta-lactamase genes were recovered from each of the
influent samples (bins INF1-C5f, INF2-C7f, INF3-C4f, and INF3-C7f, **Table 1**). The genome bins
showed completeness and redundancy values ranging from 21.16 to 31.69 and 0.03 to 0.8,
respectively, and they were assigned as *Arcobacter* sp., *Arcobacter* sp., *Arcobacter sp017995775*,
and *Arcobacter cryaerophilus A,* respectively (**Table 1**). (The previously distinct genera
Arcobacter and *Aliarcobacter* are referred here as a single genus *Arcobacter* according to the latest
update on nomenclature and taxonomical classification (66).) Identical or similar ARG sequences
were also found among unbinned contigs of influent wastewater, and these were included in the
gene tree and the genetic context analysis, along with the reference sequence *bla*OXA-464,
showing the closest similarity of the established beta-lactamase genes (**Fig. 2**, **Table S5**).
434
The class D beta-lactamases carried by each of the four genome bins formed distinct branches of
the tree (**Fig. 2**). Moreover, they primarily differed from those of the reference strains. Of these,
A. butzleri carried a gene identical to the variant *bla*OXA-464, while the beta-lactamases of *A.*
defluvii, *A. cryaerophilus*, and *A. articola* showed more dissimilarities to any other variants (**Fig.**
2). On the contrary, the beta-lactamase encoded by *A. suis* showed the closest similarity to that in
one of our genome bins (**Fig. 2**). Despite differences in the beta-lactamase sequences, their
immediate flanking region showed substantial conservation across most hosts and branches of the
phylogenetic tree (**Fig. 2**, **Table 1**). The conserved region consisted of genes *rhl*E and aldose 1-
epimerase (mutarotase), encoding core functions related to RNA (67) and carbohydrate

metabolism (68), respectively, further supporting the non-mobile context of the beta-lactamase
gene among these sequences (**Fig. 2**).

As a notable exception to the sequences described above, the branch represented by the INF3-C7f
genome bin assigned as *A. cryarophilus*, harbored identical beta-lactamase genes but lacked the
conserved core region descriptive for all other branches (**Fig. 2**). Instead, the beta-lactamase gene,
together with peptide methionine sulfoxide reductase gene *msrA* involved in oxidative stress
protection (69) (not to be mixed with the established ARG *msr*(A)) were flanked by transposase
genes on both sides (**Fig. 2**). Additionally, IS21 family transposases were detected in some of these
contexts within this branch, but not among the other branches (**Fig. 2**). While no similar context
sequences of the entire module were detected in public sequence repositories, matching
transposase genes were found in various Campylobacterales such as *Sulfurospirillum* and
Arcobacter species. Furthermore, all except the bin INF3-C7f-linked sequences of the branch were
predicted as plasmid-borne, implying higher mobility potential of these sequences, unlike most of
the conserved contexts (**Fig. 2**, **Table S5**).

The amino-acid sequence of this beta-lactamase showed low sequence similarity with *bla*OXA-
464 (72 % 182/254 AA) and the best matches for its predicted protein structure were the beta-
lactamases present in *Sulfurospirillum* sp. hDNRA2 (AF-A0AAU8L8S4-F1, e-value 8.27e-33;
sequence identity 50.9%) by AlphaFold Database clustered to 50% sequence identity and
Enterobacter cloacae (9h15_A_1 e-value 1.33e-29; sequence identity 43.8%) by the Protein Data
Bank.

Altogether, these findings suggested that this particular class D beta-lactamase represented a
putative novel mobile ARG carried and possibly horizontally transferred at least among
Arcobacter and other Campylobacterales.

471
472 **Fig. 2.** Gene phylogeny tree of the latent class D beta-lactamase gene (shown in red) found in the
473 bins of connected contigs of *Arcobacter* species and unbinned sequences from influent wastewater.
474 Flanking regions of 5 kb upstream and downstream of the class D beta-lactamase were included
475 in the genetic context visualization. Reference sequences of class D beta-lactamase and their
476 contexts were embedded in the analysis.

477
478

479 **Class C beta-lactamase genes (*bla*MCA) are reshuffled by p*dif* modules across at least**
480 **plasmids and chromosomes of *Acinetobacter* species**

481
One class C beta-lactamase gene-carrying, high-quality (99.97 completeness and 3.43 redundancy)
Acinetobacter sp. genome bin INF1-C3f was characterized through the methylation analysis (**Fig.**
3A, **Table 1**, **Table S6**). This beta-lactamase was predicted by the fARGene tool, which excludes
the established ARGs from further analysis steps (45), suggesting that the gene is previously
undescribed. However, it was identified in one database, the AMRFinderPlus of the NCBI applied
by Bakta (50), where it appeared under the name *bla*MCA. We thus refer to the gene by the name
'*bla*MCA' from hereon. Matching genes were found in unbinned contigs from influent samples
and were predominantly predicted to be plasmid-derived (**Fig. 3A**, **Table S6**).

490
One major branch of the *bla*MCA gene tree was formed by the reference sequence *Acinetobacter*
sp. (CP032274.1) and unbinned 12 predicted plasmid sequences from influent wastewater
possessing identical ARG sequences but highly diverse genetic contexts (**Fig. 3A**). Of the two
other main tree branches, the *bla*MCA gene carried by *Acinetobacter* sp. genome bin INF1-C3f
showed 92 % (1122/1215 bp) sequence similarity with both references (MRSN15084 and
CP032274.1), while the beta-lactamases carried by the references (MRSN15084 and CP032274.1),

that were most distantly related within the tree, showed 89 % (1081/1216 bp) sequence similarity
with each other (**Table 1**, **Fig. 3A**).

The non-plasmid context, as well as the conserved gene arrangement across the sequences of the
branch that was represented by genome bin INF1-C3f *Acinetobacter* sp. (**Table 1**), suggested a
chromosomal context for *bla*MCA in these sequences (**Fig. 3A**, **Table S6**). This was contrary to
what was observed for the remaining sequences, where the flanking regions are variable despite
the identical ARG sequences (**Fig. 3A**), indicating recent decontextualization events. Nevertheless,
common to all sequences were the presence of p*dif* sites (**Fig. 3**, **Table S6**, **Table S7**). More
specifically, in most cases, these C|D and D|C-oriented site-specific recombination sites were
detected immediately bordering the *bla*MCA gene, forming a *bla*MCA-p*dif* module, which was,
in a few cases, present multiple times along the investigated sequence region (**Fig. 3A**).

In three instances, different IS3 or IS4 family transposase genes co-occurred with *bla*MCA in the
same p*dif* modules suggesting their possible role in the transmission or expression of *bla*MCA
gene (**Fig. 3B**). For the INF1-C3f *Acinetobacter* sp. genome bin and its cluster companions, genes
encoding proteins of unknown function were located upstream *bla*MCA of (**Fig. 3**). Overall, the
studied regions contained a high number of p*dif* sites proximal to not only *bla*MCA, but also other
genes (**Fig. 3**). For instance, p*dif* modules encoding *brnT* and toxin-antitoxin system genes,
referring to plasmid maintenance functions, with or without integrase catalytic domain containing
protein encoding gene were repeatedly observed in multiple sequences (**Fig. 3A**). Altogether, the
high frequency of p*dif* modules likely contributes to the extensive gene content variation seen
across the studied contexts suggesting active shuffling of *bla*MCA not only among plasmid but
also chromosomal contexts at least among *Acinetobacter* species (**Fig. 3**).

Fig. 3. A. The phylogeny tree of class C beta-lactamase genes (also known as *bla*MCA) (shown
in red) in the methylation-based bins of connected contigs and unbinned contigs with matching
genes. *Acinetobacter baumannii* (MRSN15084) and *Acinetobacter* sp. (CP032274.1) were
included as references. The annotations of the target gene flanking sequences are shown for 5 kb
upstream and downstream. **B.** More detailed visualization shows the gene arrangement of the p*dif*-
modules containing IS3 or IS4 family transposases alongside the *bla*MCA gene. Bin INF1-C3f
and *Acinetobacter* sp. (CP032274.1) sequences are included in the figure for comparison.

530
531

Identification of *Simplicispira* sp. as a newly described host for the integron-encoded ESBL
gene *bla*OXA-129

The established extended-spectrum beta-lactamase (ESBL) phenotype encoding gene *bla*OXA-
129 was identified in two methylation-based genome bins assigned as *Simplicispira sp017988165*
(prev. *Giesbergeria sp017988165*) from two separate influent samples (bins INF1-C3, INF3-C5)
(**Fig. 4**, **Table 1**, **Table S8**). The bins of connected contigs were composed of 20 and 18 contigs
and showed 31.41 and 38.56 completion and 3.4 and 1.35 redundancy, respectively (**Table 1**).
Further exploration of matching genes in the full data revealed two more occurrences of this gene
among the unbinned contigs from influent samples (**Fig. 4**, **Table S8**). The ARGs in the genome
bins were 99.90-100 % and in the unbinned contigs 99.75-100 % identical (802-804/804 bp) to the
*bla*OXA-129 genes in the reference sequences of clinical class 1 integron encoded genes carried
by *E. hormaechei* (FJWZ01000025, the reference gene in the ResFinder database) and of
Pseudomonas aeruginosa (CP031449.2).

The flanking region of the *bla*OXA-129 displayed high similarity between the two *Simplicispira*
sp. genome bins but differed more from the reference sequences (**Fig. 4**). However, a common
feature among all but one of the sequences was the presence of traits indicating integron gene
cassette context of the *bla*OXA-129 gene (**Fig. 4**). These included the presence of a gene annotated
as integron integrase upstream of *bla*OXA-129, detection of aminoglycoside resistance gene *aadA*
(*aadA1* in *P. aeruginosa* and *aadA5* in *Simplicispira* sp.) downstream of *bla*OXA-129 and finally
numerous *attC* sites flanking the genes which were numerous especially in the *Simplicispira* sp.
sequences (**Fig. 4**). The integrases of *Simplicispira* sp. were dissimilar to those in *E. hormaechei*
and *P. aeruginosa* which in turn resembled each other (**Fig. 4**). However, according to BLAST
search against public sequence databases, similar integron integrase genes to those in *Simplicispira*
sp. are found in multiple other bacterial species such as *Klebsiella* sp. and *Acidovorax* sp. (data not
shown). Another notable difference between the context regions of the clinical *E. hormaechei* and
of *Pseudomonas aeruginosa* integron cassettes and those in *Simplicispira* sp. was the presence or
absence of IS6 family IS6100 transposases, respectively (**Fig. 4**). Particularly, the arrangement of
the two identical IS6100 genes flanking entire the integron gene cassette in composite transposon
orientation in *Pseudomonas aeruginosa* may have a key role in mobilizing the cassette (**Fig. 4**).
While one copy of IS6100 is observed also in *E. hormaechei*, none are found in the cassettes of
this study (**Fig. 4**).

Fig. 4. A comparative analysis of genetic features of the *bla*OXA-129 gene (shown in red) hosts identified in sequences of this study and previously described isolates of *Enterobacter hormaechei* (FJWZ01000025.1) and *Pseudomonas aeruginosa* (CP031449.2). The annotations of the target gene flanking sequences are shown for 10 kb upstream and downstream.

Class 1 integron cassette encoded *sul1* was hosted by scarcely studied bacteria of the order UBA1845 (class Phycisphaerae)

Sulfonamide resistance gene variant *sul1-9* was detected within clinical class 1 integron gene cassette in two high-quality genome bins (bins SLU2-C4, SLU3-C1) from two separate sludge samples (**Table 1**, **Table S9**, **Fig. 5**). These bins constituting of 12 and 4 contigs showed high completion and redundancy values of 90.86 and 97 and 3.74 and 4.14 respectively and were assigned as *JAUCQB01* sp. of UBA1845 order within Phycisphaerae class and Planctomycetota phylum (**Table 1**, **Table S9**). Gene *sul1-9* differs from its most similar variant *sul1-2* by an 11 bp long tail and one nucleotide substitution.

Exploring the *sul1-9* contexts of the sequences of this study and of the previously reported hosts of *Escherichia coli* (CP026207.1) and *Salmonella enterica* (AY963803.6, the reference gene in the ResFinder database) revealed that common to all these sequences was the presence of the *intI1* gene, located adjacent to the *sul1-9* (**Fig. 5**, **Table S9**). Moreover, similarly to the *bla*OXA-129 encoding integron gene cassettes, IS6 family IS6100 transposase was detected in all investigated sequences including both reference sequences from public databases as well as the newly described *sul1-9* hosts of *JAUCQB01* sp. and unbinned contig characterized in this study (**Fig. 5**). In contrast to the *E. coli* and *S. enterica* reference strains, no additional ARGs were present in the integron cassettes of this study (**Fig. 5**). The partially shared sequence between the *JAUCQB01* sp. genome bin sequences and the unbinned contig described in this study and extending the core structure of the integron supports the wide occurrence and hence mobility of this *sul1-9* cassette with IS6100

594   in different contexts (**Fig. 5**). However, unlike for the reference strains, no *attC* sites were reported
595   for the sequences of this study (**Fig. 5**).

596
597   **Fig. 5.** The phylogeny tree of *sul1* genes (shown in red) in the methylation-based bins of connected
598   contigs and unbinned contigs. Reference sequences of *Escherichia coli* (CP026207.1), *Klebsiella*
599   sp. (CP055486.1), and *Pseudomonas aeruginosa* (CP021775.1) were included in the comparative
600   genetic analysis of the *sul1-9* flanking region of 10 kb up- and downstream length.

601
602

Gene *erm*(F) is carried by multiple different environmental Bacteroidales species in the
sludge

The acquired macrolide–lincosamide–streptogramin B (MLSB) resistance gene *erm*(F) was the
most common established resistance gene present among the ARG-encoding genome bins and was
also detected in numerous unbinned contigs (**Table 1**, **Fig. 6**, **Table S10**). The 13 visualized *erm*(F)
encoding bins of connected contigs were primarily derived from sludge samples, but one influent
derived *erm*(F) was also reported (**Table 1**, **Fig. 6**). The completeness and redundancy values of
the *erm*(F) genome bins ranged from 17.49 to 94.92 and 0.02 to 3.84 (average completion 58.51
and redundancy 2.11), and the number of included contigs per genome bin was on average 11
(**Table 1**). Altogether, these genome bins represented 7 different bacterial families within the
Bacteroidales order (**Table 1**). Of these, only the one genome bin represented the human gut-
associated *Bacteroidetes* sp. (*Bacteroidaceae* family) *erm*(F) host, and this bin was recovered from
the influent. Instead, the remaining bins of connected contigs from sludge represented the so-called
environmental lineages of Bacteroidales, including families *Dysgonomonadaceae,*
Paludibacteraceae, Tenuifilaceae, Prolixibacteraceae, VadinHA17, and *ML635J-15* (**Fig. 6**,
Table S10). These findings align with the observed Bacteroidota phylum dynamics throughout the
wastewater treatment process (**Fig. S6**, **Fig. S9**): While the *Bacteroidaceae* family was clearly
dominant in the influent, the taxa profile was substantially different in sludge (**Fig. S5**, **Fig. S6**),
where multiple other Bacteroidales families were present in high relative abundances, particularly
the *erm*(F) hosting families in the sludge (**Fig. 6**, **Fig. S6**, **Fig. S9**).

624
Phylogenetic analysis revealed that the identified *erm*(F) genes were highly similar despite the
variable genetic context indicative of horizontal gene transfer (HGT) (**Fig. 6**). Most of the genes
shared 99-100 % sequence identity with the reference *erm*(F) gene (*Bacteroides fragilis*
(CP054002.1 and M17808.1, reference gene in ResFinder database) while for the two most
dissimilar genes the similarity to reference gene was lower (95% and 92%) (**Fig. 6**). High plasticity
among the *erm*(F) flanking regions was emphasized by the observation of clearly distinct context
sequences, even when originating from genome bins of the same species (*Seramator thermalis*
bins SLU1-C5, SLU1-C2, SLU3-C3) (**Fig. 6**, **Table S10**). Moreover, in some cases, MGEs such
as transposases of IS30 and IS987 families are detected near *erm*(F), supporting the rearrangement
of genes. However, no CTnDOT transposases, typically explaining the transfer of *erm*(F) among
human gut-associated *Bacteroidales* (70), were detected. In the *Bacteroides* sp. genome bin (bin
INF3-C2), the *erm*(F) seems to be part of an integron gene cassette along with *aadS* and *bla*OXA
ARGs (**Fig. 6**). Altogether, both the high diversity of *erm*(F) hosts and contexts suggested that
erm(F) is horizontally moving between various, not only human-gut-associated *Bacteroidales*
hosts.
640

641
Fig. 6. Gene phylogeny tree of the *erm*(F) gene (shown in red) found in genome bins of different
Bacteroidales order bacteria and unbinned sequences from wastewater. The *erm*(F) genes encoded
by *Bacteroides fragilis* (M17808.1) and *B. fragilis* (CP054002.1) and their flanking regions were
included as references. Flanking regions of 5 kb upstream and downstream of the *erm*(F) were
included in the genetic context visualization. Family-level taxonomical identity of the genome bins
is shown in bolded text.
648
649
650
651

Discussion

Although the novel resistance determinants arising from environmental microbiomes, especially with the implications of gene mobility, have been assigned as critical contributors to the current rapidly developing resistance problem, methodological limitations have impaired our ability to recognize them (7, 12). While a wide variety of ARGs are being detected in diverse environments and microbiomes, the crucial information regarding their genetic and bacterial contexts is often missing (12, 18). Instead, in many metagenomic studies, the highest-abundance taxa are often reported separately, although those bacteria rarely represent the ARG carriers in those communities (9, 25). In this study, by applying metagenomic long-read sequencing and bacteria-specific methylation signals, we were able to identify the host bacteria of the emerging resistance threats whose characterization has been delayed by technical limitations and prioritized clinical settings.

The latent class D beta-lactamase gene that was carried by *Arcobacter cryaerophilus* showed potential as a newly emerging, uncharacterized ARG that could be transferred among *Arcobacter* or related species primarily due to the composite structure of the flanking transposase genes, similar to what was reported before with other ARGs (71–73). The IS21 family transposase situated in proximity may also have a role in the mobilization (74). Although the studied beta-lactamase did not match any of the existing proteins in the databases, the best matches included proteins encoded by *Sulfurospirillum,* which, similarly to *Arcobacter,* represents the Campylobacterales order. Moreover, the observation of similar transposase genes but not their adjacent genes, both in *Sulfurospirillum* and *Arcobacter* sequences of the public sequence databases, could suggest that transmission of genes facilitated by these composite transposases occurs at least between closely related species. These observations may also point to *Sulfurospirillum* as the ancestral host of the beta-lactamase described here before its transfer to the species *Arcobacter cryaerophilus.*

According to our data, the remaining class D beta-lactamase genes represented likely chromosomal, non-mobile variants of this gene, indicated by the conserved context across different *Arcobacter* species, including *A. butzleri*, *A. defluvii*, *A. articola*, *A. sp017995775, A. suis,* and unidentified species of *Arcobacter*. Of these genes, only the one hosted by *A. butzleri* showed similarity to any previously established ARGs, namely the *bla*OXA-464 gene, reported previously in wastewater (75) (Dekic Rozman *et al.* manuscript titled "Emerging Food- and Waterborne Pathogen *Arcobacter* in Wastewater: Diversity and Antibiotic Resistance" under review).

Although *Arcobacteraceae* are among the most dominant families in influent wastewater (76, 77), a substantial diversity within this bacterial group has remained uncharacterized (Dekic Rozman *et al.* manuscript under review). At the same time, indications of the contribution of *Arcobacter* species to the ARG load in wastewater have been reported, yet the extent and underlying mechanisms remain poorly understood (76, 77). The lack of standardized cultivation and phenotyping protocols (Dekic Rozman *et al.* manuscript under review) and the genome plasticity, challenging the genome-resolved metagenomics for *Arcobacter* species (78), have delayed the description of genotypic and phenotypic resistance features of *Arcobacter* species. Therefore, the description of the emerging mobile class D beta-lactamase here elucidates the less-understood genetic mechanisms underlying resistance of *Arcobacter* species. In addition to the human clinical

698 cases and wastewater, where they act as anoxic denitrifiers (79), *Arcobacter* have been isolated
699 from food and animals, for which they are usually asymptomatic, driving their unnoticed
700 dissemination (80).
701
The widespread distribution of *Arcobacteria* spp. across both external environments and human
activity-linked sources highlights the significance of the *Arcobacter* species serving as the
mediators of the established and emerging ARGs between non-pathogenic and pathogenic species.
Given that *A. cryaerophilus* is one of the *Arcobacter* species most commonly isolated in clinical
cases (80), the mobile beta-lactamase gene described in this study within *A. cryaerophilus* host
could cautiously be interpreted as an example of the ARG decontextualization from its original
genomic context and further into a pathogenetic background by MGEs. Given the limited evidence
presented in this study, we cautiously propose that the potential past host in this case may have
represented a species of genus *Sulfurospirillum*, which are taxonomically and functionally related
to *Arcobacter* (81).
712
Our description of the carriage and mobility of the class C beta-lactamase gene *bla*MCA provides
another instance of emerging resistance risk that is not yet widely recognized. To our best
knowledge, this gene has been referred to only once before (49). Its absence in most of the widely
used ARG databases, such as ResFinder (44) and CARD (82), and therefore its scarce
representation in the literature would suggest that *bla*MCA is considered rather an intrinsic gene
than an acquired ARG. However, based on the active reshuffling of *bla*MCA-encoding p*dif*-
modules across diverse genomic contexts demonstrated by our results, we argue that *bla*MCA may
show potential as an emerging clinically relevant ARG. While the XerC/XerD (*dif*) binding site-
specific recombination is central in chromosomal dimer resolution among all bacterial lineages,
p*dif*-module trafficking, which occurs primarily among *Acinetobacter* plasmids (83), allows the
rapid acquisition of adaptation-improving genes by site-specific homologous recombination (84).
Despite its ancient origin, this mechanism is thought to have gained additional functionality as an
ARG mobilizing mechanism along the post-antibiotic era (83). The full understanding of the
significance of the p*dif* sites as mobile genetic elements has been impeded by the difficulty in
identifying them (49) as well as the detection limitations of the current tools focusing on well-
established ARG-modules and pathogens (49, 85), which has restricted the discovery of emerging
latent ARGs and their non-pathogenic hosts possibly harboring novel p*dif* modules.
730
Building on previous knowledge on *bla*MCA that is based on observations on publicly available
plasmid sequences, which are likely biased towards opportunistic pathogens (49), the findings
presented here add a real-world example of the p*dif*-module rearrangement at a microbial
community level, beyond the boundaries of genetic context or taxonomical lineages. This was
exemplified here by the detection of *bla*MCA-p*dif* modules within both likely chromosomal
Acinetobacter sp. host and plasmid-sequence contexts. Chromosome-integrated p*dif*-modules
encoding ARGs have been reported before (86), and on the other hand, the so-called additional
chromosomal *dif* sites, observed in *Acinetobacter* spp. before (84), may act as reservoirs for
adaptation-improving genes transferred to the plasmid p*dif*-modules.
740
In addition to the p*dif* modules consisting of *bla*MCA formed by the inverted p*dif*-sites (C|D – D|C
orientation) flanking *bla*MCA in *Acinetobacter,* reported also earlier (49), we also found modules
in which *bla*MCA was coupled with IS elements. The presence of complete sets of ORFs of either

IS3 or IS4 family transposases located up- or downstream of *bla*MCA within the same p*dif* module
could be an indication for further mobility and transcription of the *bla*MCA gene by an IS-encoded
promoter (87). Similar observations of IS elements within other kinds of ARG-p*dif*-modules have
been made before (88), but not in association with the *bla*MCA.

While particularly common in *Acinetobacter* plasmids (83), identical p*dif*-modules have been
detected across genera (49), confirming that p*dif*-module-mediated spread of ARGs across
bacterial lineages does occur, and environments like wastewater could host conditions enabling
such cross-species transmission. Unfortunately, we failed to assign taxonomic identity to any other
than *Acinetobacter* sp. sequences hosting *bla*MCA. Even though the resistance phenotype of this
beta-lactamase gene remains to be confirmed, the frequent rearrangement of *bla*MCA p*dif*-
modules suggests that it confers an adaptive advantage to its recipient under selective conditions,
such as those caused by the presence of antibiotic residues in the influent wastewater. Overall, our
description of the *bla*MCA reshuffling at least among *Acinetobacter* chromosomes and plasmids,
supported by previous findings (49), adds to the extensive list of antibiotic resistance features
typical for this genus, which comprises members across ecological niches from those flagged as
the top priority pathogens regarding multidrug resistance by WHO (89) to the non-pathogenic
environmental species.

The risk potential of environmental microbiome origin ARGs is defined by the likelihood and
consequences of their transmission to human pathogens (7). As both hosts of the latent ARGs
described above, *Arcobacter* and *Acinetobacter*, represent genera that include both opportunistic
pathogens and environmental lineages, the frequency of HGT and likelihood of ARG transmission
across pathogenic and non-pathogenic strains is less rare than is considered for ARGs emerging
from strictly environmental bacteria (12, 15). This is particularly concerning, given that only one
successful transmission to a clinically important strain is required for the emergence of a novel
clinically important resistance threat (12).

In addition to the novel, predicted ARGs, our methylation-based binning approach allowed the
identification of previously uncharacterized environmental hosts for functionally confirmed ARGs,
shedding light on the hidden contributors to total resistance burden. To the best of our knowledge,
Simplicispira sp017988165 (prev. *Giesbergeria sp017988165*) described here as the carrier of the
integron-encoded *bla*OXA-129 is the first non-pathogenic host identified for this gene.
Simplicispira sp. are important denitrifiers under both aerobic and anoxic conditions in wastewater
(79, 90, 91). *bla*OXA-129 was first described in integron gene cassettes of clinical isolates of *P.*
aeruginosa collected in 2006-07 in China, showing carbapenem-sensitive ESBL phenotype (92)
and in other opportunistic pathogens (*Enterobacter* sp., *P. aeruginosa*, *K. pneumoniae*, *S. enterica*
(93) and *C. resistens* (CP125947.1)) ever since.

Although the integrase encoding genes in *Simplicispira* and the previously described pathogens
differ, the integron gene cassette context with numerous detected *attC* sites supports its possibility
of acting as a source of ARGs to be translocated into other contexts. However, integrons are not
mobile themselves (94). While two integron gene cassette flanking IS6100 transposases (72) were
present in most but not all *bla*OXA-129 sequences reported before in opportunistic pathogens (93,
95), no such genetic elements required for the mobilization of the integron gene cassette were
observed in *Simplicispira*. Altogether, our findings broaden the current knowledge about the host

range and mobility of this gene and point to the potential of *Simplicispira* sp. acting as an
environmental intermediate host of clinically relevant ARGs in wastewater.
792
Interestingly, according to our results, it appeared that the same IS6100 transposase (72), which
has also previously been strongly associated with ARG mobilization (96), would have also been
involved in the putative mobility of the class 1 integron-encoded *sul1-9* gene within the strictly
environmental class Phycisphaerae bacteria (species *JAUCQB01* sp.) described here. However,
detection of only one copy of the IS6100 gene within the *sul1-9* flanking region of both
environmental Phycisphaerae and opportunistic pathogenic reference hosts *E. coli* and *S. enterica*
may not yet warrant the transfer of its nearby genes (72).
800
Considering the available evidence, no ARGs have been described in this *JAUCQB01* sp. or within
its higher taxonomic rank, order UBA1845 bacteria before. Although Phycisphaerae are being
detected in various environments, the understanding of their functional characteristics and
ecological roles is held back by the lack of cultivable reference strains, as is the case with order
UBA1845 (97). Only recently, Lenferink and colleagues have discovered that the UBA1845
bacteria, mostly residing in engineered environments under anoxic conditions, such as wastewater,
thrive in the presence of diverse sugar and complex carbon substrates such as extracellular
polymeric substances extracted by primary producers (97). The presence of *sul1-9* as the only
ARG in these cassettes of UBA1845 might point to an early evolutionary stage of these integron
cassettes, as more ARGs are expected to accumulate in time under certain conditions (94). Our
observations could be further explained by their chromosomal context, unlike the reference
opportunistic pathogens' plasmid- or genomic island-encoded (98) integron cassettes containing
multiple consecutive ARGs.
814
The two example cases confirm that even the clinical integron-encoded ARG cassettes are well
disseminated to taxa lacking any human health association, challenging the previous conception
of integrons encoding ARG cassettes being disseminated mainly among opportunistic pathogens
(94). Furthermore, these findings reveal insights into the so-called intermediate hosts and
mobilizers of ARGs fostered by the wastewater treatment process before their putative irreversible
transfer to human health-associated species (7, 8).
821
Similarly, our analysis revealed the host dynamics of macrolide-lincosamide-streptogramin
resistance phenotype encoding (99) *erm*(F) gene throughout the wastewater treatment. While the
identified *erm*(F) carrier in influent represented gut-associated Bacteroidales, no human-associated
taxa but numerous other members of this order were distinguishable in the outlet sludge. The
particularly well-distinctive value of methylation profiles in the detection of Bacteroidales has also
been noted before, although in considerably less complex gut microbiomes (23). The *erm*(F) hosts
in the sludge comprised 7 different families and 10 genera representing environmental groups of
Bacteroidales. These species, such as those belonging to families *Dysgonomonadaceae*,
Paludibacteraceae, and *Tenuifilaceae,* are less studied but common members of the bioreactor
systems, specifically wastewater sludge (100–102). They are characterized by their metabolically
versatile functions, for instance, denitrification and the capability for fermenting a wide range of
mono- and disaccharides (79, 100–102).
834

The expanding host range and the highly variable genetic contexts in the sludge point to the horizontal mobility of the *erm*(F) among different bacteria, as noted before (103). However, as the host range revealed here and in previous work (15, 103) suggests, certain HGT barriers seem to restrict the transmission to Bacteroidota. Although various MGEs are detected among the gut-unrelated Bacteroidales hosts, no single specific mobility mechanism responsible for the horizontal mobility of *erm*(F) could be identified here or by previous work (103). While the CTnDOT (Conjugative Transposon from strain *Bacteroidetes thetaiotaomicron* DOT) conjugative transposons are the central carriers of *erm*(F) among gut-associated Bacteroidales species (70), some other mobility mechanism could explain the presence of identical or highly similar *erm*(F) gene sequences within highly variable genetic contexts among the different environmental Bacteroidales hosts, even within species, as described here for *Seramator thermalis*. Similarly, invertible promoter-regulated ARGs located in integrative conjugative elements (ICEs) are highly enriched within Bacteroidales, particularly among the gut-associated species (70, 99, 104). Although previous work has reported a lower prevalence of these mobile invertible promoter-associated ARGs among environmental Bacteroidota lineages (105), wastewater was not included among the environments explored in this study by Jiang and colleagues (105), leaving their occurrence among wastewater Bacteroidales families unexplored.

It may also be that the acquisition and later chromosomal integration of *erm*(F) by diverse Bacteroidales species has occurred in the distant evolutionary past, and the selective conditions present in sludge promote the growth of *erm*(F) carrying strains. Altogether our results regarding the observed increase in the abundance of *erm*(F) hosting taxa in the sludge supported the view that the high abundance of this gene, particularly through its enrichment in the sludge, noted also by previous studies (75, 103, 106, 107), is not due to the proliferation of one species by the treatment process but rather multiple taxa that are enriched under the conditions present in the sludge. Furthermore, the geographical differences reported for the *erm*(F) prevalence in sludge (103) might reflect the differences in wastewater treatment systems, mirrored as the Bacteroidales composition differences and their interactions facilitating the *erm*(F) carriage and spread. Bacteria of the *Bacteroides fragilis* group have been recognized as resistance gene reservoirs before (99). However, our results suggest that this characteristic is not exclusive to the gut-associated Bacteroidales but is likewise observed in environmental lineages of Bacteroidales, which might serve as environmental reservoirs of *erm*(F) for further transmission to the clinically relevant species.

Taken together, by linking ARGs to their broader genetic contexts and hosts at the microbial community level using bacteria-specific methylation signals, this study addresses one of the key challenges in the field hindering the environmental resistome risk assessment (7, 12). Another recent approach to this task is high-throughput single-cell metagenomic sequencing, which uses individually barcoded cells for sequencing (108). While promising, this approach has certain limitations, including laborious sample preparation, dependency on specific reagents and equipment, and possible bias against gram-negative bacteria (108). A strictly computational strategy for coupling taxonomy, ARGs, and their genetic context draws upon alignment-free methods with CARD k-mers (109). However, the main weakness of this method is its reliance on established ARGs in a certain database and the assumption that these ARGs occur in expected context sequences, and especially among pathogens, while latent ARGs and any ARGs within new contexts are missed. In contrast, by the characterization of newly described ARG hosts and

881 contexts for both latent and established ARGs without the dependency of prior knowledge, we
882 show that the approach presented in our study mitigates this limitation.
883
The methylation-based binning here enabled the description of several putative emerging ARGs
and their hosts and confirmed the intermediate host roles of strictly environmental and
understudied taxa. Even though the obtained genome bins represented predominantly partial
genomes rather than fully complete MAGs, we were able to assign the taxonomic identity and set
the ARG into its wider genetic context to evaluate its mobility potential. One limitation of our
approach, however, was the loss of data, as not all contigs displayed distinct methylation profiles
or no modifications at all, which led to fewer data available for downstream analyses, namely the
genome bin curation. Technical factors such as the lower sensitivity for detecting certain base
modifications (110, 111) or simply lower coverage, as in the fewer mapping reads, and their
attached modification data of certain strains (23) may have affected this. Another technical
bottleneck arose from the manual work required for refining the genome bins of clustered contigs.
895
Importantly, the data-loss limitation did not introduce false positives but rather gaps in the dataset,
and the potential biases introduced by our approach would be different from those of more
traditional binning algorithms. For instance, our findings regarding the multiple *erm*(F) *Seramator*
thermalis hosts showing diverse methylation profiles and genetic contexts indicate the concurrent
presence of multiple strains of this species and their distinct identification by methylation analysis
in complex communities. Previous observations on strain- (111, 112) or serovar- (113) level
differences in methylation profiles supported our findings. This variation is explained by the
transcriptional regulation roles of epigenetic variation mediated by methylation (27, 111, 112) as
well as more frequent HGT of restriction modification systems between closely related taxa than
more distant lineages (114). Additionally, our approach circumvented the common bias towards
the high-abundance taxa that is descriptive for traditional binning methods (9, 25). This is of high
importance when aiming to understand the ARG dynamics often mediated by rare taxa (9, 25)
within highly complex environmental microbiomes.
909
Finally, despite considerable research efforts (23, 115), the task of assigning the host-plasmid
linkages in metagenomic data has remained without a definitive resolution, particularly in complex
communities. Given that the use of methylation profiles ignores the variation in coverage and
sequence composition, which currently constrains the correct binning of sequences rich in MGEs,
holds great potential for this (22, 23). Building on the valuable work carried out on simpler
communities (23), our study demonstrates that methylation-based binning can be used to resolve
plasmid origins also in highly complex communities, although with critical limitations. Although
we could assign the bacterial hosts for multiple plasmids, unfortunately, none of them represented
ARG-encoding plasmids, which are more likely to be those categorized as conjugative or
mobilizable plasmids (116). Despite their multi-host history, they are considered to acquire the
methylation motif profile of their current host rapidly (22, 23), owing to the evolutionary arms
race between plasmids and host restriction-modification systems (116, 117). However, the
simultaneous carriage of similar plasmids by different host bacteria may confuse the methylation
profile detection by our approach, as reads coming from different strains and hence with mixed
methylation signatures are aligned to the same sequence. Given that these most frequently
horizontally moving broad-host range plasmids are also the most important drivers of ARG
carriage and spread (118), this highlights a critical limitation of our method, in the context of data

with the level of complexity examined here. Alternatively, erroneous assembly (21) might explain
the poor detectability of these plasmids and their methylation-resolved origins because their
mosaic architecture, shaped by embedded MGEs, is prone to fragmentation into smaller contigs
(118). This, in turn, would lead to a decreased likelihood of capturing unique methylation profiles
for the plasmid sequences to match with those of their host genomes (23).
932
933
934 **Conclusions**
935
Our work sheds light on the previously hidden intermediate hosts and mobility mechanisms of
established and newly emerging resistance threats in highly diverse wastewater samples.
Furthermore, these results highlight the importance of the culture-independent investigations of
ARGs and their carriers. Further investigations are needed to examine the broader occurrence of
the latent beta-lactamases and, most importantly, to verify the resistance phenotypes of these
ARGs, as their description here relied solely on computational predictions.
942
943
944 **Acknowledgements and study funding**
945

This work was supported by the Research Council of Finland funding for the Multidisciplinary
Center of Excellence in Antimicrobial Resistance Research (V.M. 364234, 346128, M.V. 364231,
346125). M.M. received funding from the MBDP doctoral program at the University of Helsinki.
We would like to acknowledge CSC, IT Center for Science, Finland, for providing the
computational resources for the study, and the DNA Sequencing and Genomics Laboratory
(supported by HiLIFE and Biocenter Finland funding), Institute of Biotechnology, University of
Helsinki, for sequencing. Open access was funded by the Helsinki University Library.

953
954
955 **Author contributions**
956
Conceptualization, A.K., M.M., V.M. Methodology — H.P. developed the methylation-detection
algorithm under the supervision of V.M., and D.K. — M.M. performed the application of the
methylation-detection algorithm, curation of bins of connected contigs, comparative genomics
investigation, and all associated bioinformatic analyses under the supervision of A.K. Data
acquisition, A.K. Writing — original draft, M.M. Writing — review and editing, all authors.
Supervision, A.K., M.V., and V.M. Funding acquisition, A.K., M.V., and V.M.
963
964
965 **Table 1.** Methylation-based bins of connected contigs encoding latent and established ARGs.
966

Bin	C[1]	R[2]	Species	Order	Number of contigs	GC % (exp.[3])	Established ARG	Latent ARG
INF1-C3	31.41	3.4	*Simplicispira sp0179881165*[4]	Burkholderiales	20	61 (61.5)	*blaOXA-129_1; aadA5_1*	
INF1-C4	36.71	4.24	*Aeromonas media*	Enterobacterales	25	61 (61)	*blaOXA-427_1; blaOXA-427_1*	
INF1-C1f	65.09	3.04	*Comamonas denitrificans_A*	Burkholderiales	31	61 (60.5)		beta_lactamase_d_2
INF1-C2f	87.47	0.03	*Chryseobacterium* sp.	Flavobacteriales	9	34 (34.5)		beta_lactamase_a; beta_lactamase_b_3

INF1-C3f	99.97	3.43	*Acinetobacter* sp.	Pseudomonadales	13	38 (38.5)		beta_lactamase_a; beta_lactamase_d_2; beta_lactamase_c
INF1-C4f	20.82	0.41	*JAFGUR01 sp017988045*	Fusobacteriales	9	31 (29)		beta_lactamase_d_2
INF1-C5f	27.03	0.8	*Arcobacter* sp.	Campylobacterales	6	28 (27.5)		beta_lactamase_d_2
INF1-C6f	53.85	1.31	*Acinetobacter* sp.	Pseudomonadales	9	38 (38.5)		beta_lactamase_d_2
INF2-C3	66.95	7	*Acinetobacter bohemicus*	Pseudomonadales	12	39 (39.5)	*blaOXA-296_1*	
INF2-C4	33.93	2.06	*Aeromonas media*	Enterobacterales	8	61 (61)	*mcr-3.6_1*	
INF2-C5	22.78	1.56	*Aeromonas* sp.	Enterobacterales	32	59 (61)	*blaOXA-427_1*	
INF2-C6	24.01	3.08	*Aeromonas media*	Enterobacterales	13	62 (61)	*blaOXA-427_1*	
INF2-C1f	27.46	1.97	*JAFGUR01 sp017988045*	Fusobacteriales	28	29 (29)		beta_lactamase_b_3
INF2-C2f	99.83	2.34	*Acinetobacter* sp.	Pseudomonadales	17	38 (38.5)		beta_lactamase_d_2
INF2-C3f	22.46	0.07	*Acinetobacter* sp.	Pseudomonadales	14	61 (61)		beta_lactamase_d_2
INF2-C4f	45.82	3.38	*Acinetobacter* sp.	Pseudomonadales	6	38 (38.5)		beta_lactamase_d_2
INF2-C5f	60.19	0.93	*Acinetobacter celticus*	Pseudomonadales	5	39 (39.5)		beta_lactamase_c
INF2-C6f	34.31	0.17	*Alistipes putredinis*	Bacteroidales	3	54 (53.5)		beta_lactamase_a
INF2-C7f	21.16	0.03	*Arcobacter* sp.	Campylobacterales	3	28 (27.5)		beta_lactamase_d_2
INF3-C1	19.42	0	*Aeromonas media*	Enterobacterales	8	62 (61)	*blaOXA-427_1*	
INF3-C2	29.96	1.1	*Bacteroides* sp.	Bacteroidales	13	44 (47.5)	*erm(F)_3*	
INF3-C3	40.91	0.06	*Aeromonas media*	Enterobacterales	13	61 (61)	*mph(A)_2; blaOXA-427_1; aph(6)-Id_1*	
INF3-C4	22.65	2.05	*JABFXX01* sp.	Haliangiales	21	59 (59)	*ere(A)_2*	
INF3-C5	38.56	1.35	*Simplicispira sp017988165[4]*	Burkholderiales	18	61 (61.5)	*blaOXA-129_1; aadA5_1*	
INF3-C1f	99.97	0.75	*Aquabacterium_A* sp.	Burkholderiales	8	68 (67.5)		beta_lactamase_c
INF3-C2f	46.2	3.58	*Methylotenera* sp.	Burkholderiales	3	42 (44)		beta_lactamase_b_1_2
INF3-C3f	83.88	0.03	*Ferruginibacter* sp.	Chitinophagales	2	38		beta_lactamase_d_2
INF3-C4f	31.69	0.13	*Arcobacter sp017995775*	Campylobacterales	7	27 (27.5)		beta_lactamase_d_2
INF3-C5f	49.34	4.67	*Comamonas denitrificans_A*	Burkholderiales	37	60 (61)		beta_lactamase_d_2
INF3-C6f	50.23	2.32	*Zoogloea sp016720685*	Burkholderiales	21	64 (64.5)		beta_lactamase_d_2
INF3-C7f	27.71	0.21	*Arcobacter cryaerophilus_A*	Campylobacterales	16	28 (27.5)		beta_lactamase_d_2
EFF1-C1	19.53	5	*Acinetobacter* sp.	Pseudomonadales	17	44 (38.5)	*blaOXA-296*	
EFF1-C1f	92.88	0.16	*Aquabacterium_A* sp.	Burkholderiales	7	68 (67.5)		beta_lactamase_c
EFF1-C2f	42.63	1.99	*Aquella sp003538525*	Burkholderiales	14	37 (38)		tet_enzyme; beta_lactamase_a
EFF2-C1	77.76	0.37	*Fluviibacter phosphoraccumulans*	Burkholderiales	9	55 (54)	*ant(3″)-Ia_1*	
EFF2-C1f	93.2	0.5	*Aquabacterium_A* sp.	Burkholderiales	5	68 (67.5)		beta_lactamase_c
EFF2-C2f	37.21	1.59	*Pleomorphomonas* sp.	Rhizobiales	8	65 (64.5)		beta_lactamase_a; aminoglycoside_c; aminoglycoside_h; aminoglycoside_i
EFF2-C3f	41.82	0.11	*Ideonella* sp.	Burkholderiales	4	67 (66.5)		beta_lactamase_a
EFF2-C4f	77.76	6.04	*Rhodoferax_C* sp.	Burkholderiales	5	62 (63)		beta_lactamase_a
EFF2-C5f	47.17	0.05	*ELB16-189* sp.	Cytophagales	6	40		beta_lactamase_b_1_2
EFF2-C6f	58.76	0	*Ferruginibacter* sp.	Chitinophagales	2	38		beta_lactamase_b_3
EFF3-C1	65.35	2.46	*Rhodoferax_C* sp.	Burkholderiales	7	60 (61.5)	*dfrB7_1*	
EFF3-C1f	88.64	2.3	*Ferruginibacter* sp.	Chitinophagales	6	38		beta_lactamase_d_2; beta_lactamase_b_3
EFF3-C2f	44.96	0.04	*Flavobacterium* sp.	Flavobacteriales	4	32		beta_lactamase_d_2; beta_lactamase_b_1_2
EFF3-C3f	50.05	0.38	*JADJSS01 sp.*	Chitinophagales	6	35		beta_lactamase_b_1_2; beta_lactamase_a

EFF3-C4f	90.39	1.5	JACMQJ01	UBA7694	2	29 (29)		beta_lactamase_d_1
SLU1-C1	75.61	6.14	*Seramator thermalis*	Bacteroidales	5	62 (42)	*tet(W)_5*	
SLU1-C2	56.2	1.74	*Seramator thermalis*	Bacteroidales	8	41 (42)	*erm(F)_3*	
SLU1-C3	25.87	1.92	*UBA1064 sp002316235*	Bacteroidales	26	45 (45)	*erm(F)_3*	
SLU1-C5	34.44	0.02	*Seramator thermalis*	Bacteroidales	14	42 (42)	*erm(F)_3*	
SLU1-C6	89.54	3.18	*DUVQ01 sp012840965*	DTJE01	5	51 (50.5)	*sul4_1*	
SLU1-C7	19	0	*Seramator thermalis*	Bacteroidales	9	42 (42)	*erm(F)_3*	
SLU1-C8	24.06	3.95	*LD21 sp029564935*	Bacteroidales	18	39 (42)	*erm(F)_3*	
SLU1-C1f	33.8	0.95	*JAAYJM01* sp.	Kapaibacteriales	10	32		beta_lactamase_d_2
SLU1-C2f	37.6	0.09	*Afipia sp017308595*	Rhizobiales	2	60 (60)		beta_lactamase_d_2
SLU1-C3f	65.42	0.3	*LD21* sp.	Bacteroidales	12	44		beta_lactamase_b_1_2
SLU1-C4f	73.95	0.15	*UBA1413* sp.	Bacteroidales	2	46		erm_f
SLU1-C5f	71.21	3.11	*JAFGBI01* sp.	Polyangiales	7	73		mph
SLU2-C1	69.04	3.84	*Bact-19 sp002412425*	Bacteroidales	14	32 (30.5)	*erm(F)_3*	
SLU2-C2	29.47	1.45	*MVYY01 sp012517575*	Bacteroidales	10	37 (37.5)	*erm(F)_3*	
SLU2-C3	94.92	3.25	*DUQS01 sp012837685*	Bacteroidales	9	40 (40.5)	*erm(F)_3*	
SLU2-C4	90.86	3.74	*JAUCQB01 sp.*	UBA1845	12	62 (62.5)	*sul1_9*	
SLU2-C5	76.97	1.31	Aristaeellaceae	Christensenellales	4	60	*tet(W)*	
SLU2-C6	38.2	5.05	*LD21 sp029564935*	Bacteroidales	44	41 (42)	*erm(F)_3*	
SLU2-C8	45.14	0.49	*UBA6115* sp.	Christensenellales	8	52 (54.5)	*erm(A)*	
SLU2-C9	41.41	0.17	*Petrimonas sp002069095*	Bacteroidales	5	39 (40)	*erm(F)_3*	
SLU2-C10	17.7	0.83	*SR-FBR-E99 sp009881065*	Bacteroidales	20	49 (49.5)	*erm(F)_3*	
SLU2-C1f	96.37	4.78	*PLanc-01 sp012744475*	Sedimentisphaerales	23	62 (62.5)		beta_lactamase_b_1_2
SLU2-C2f	79.16	7.5	*UBA5266 sp002069895*	Bacteroidales	23	35 (35)		beta_lactamase_b_1_2
SLU2-C3f	58.02	7.16	*LD21* sp.	Bacteroidales	9	40		beta_lactamase_b_3
SLU2-C4f	82.9	1.06	*Neoroseomonas* sp.	Acetobacterales	10	69 (68.5)		beta_lactamase_a
SLU2-C5f	77.17	7.85	*UBA1413* sp.	Bacteroidales	11	47		erm_f
SLU2-C6f	92.79	3.58	*LD21* sp.	Bacteroidales	15	45		beta_lactamase_b_1_2
SLU2-C7f	62.08	2.75	*Pullichristensenella* sp.	Christensenellales	3	62		beta_lactamase_b_3
SLU3-C1	97	4.14	*JAUCQB01 sp.*	UBA1845	4	62 (62.5)	*sul1_9*	
SLU3-C2	77.07	3.33	*MVYY01 sp012517575*	Bacteroidales	24	38 (37.5)	*erm(F)_3*	
SLU3-C3	34.65	0.63	*Seramator thermalis*	Bacteroidales	17	41 (42)	*erm(F)_3*	
SLU3-C5	52.83	3.49	*Petrimonas* sp.	Bacteroidales	15	40 (40)	*erm(F)_3*	
SLU3-C7	17.49	0.6	*UBA1413 sp002304925*	Bacteroidales	11	48 (50)	*erm(F)_3*	
SLU3-C1f	69.32	2.45	*JAAYJM01* sp.	Kapaibacteriales	17	32		beta_lactamase_d_2
SLU3-C2f	71.63	2.06	*MWCO01 sp002070115*	Polyangiales	9	57 (57)		beta_lactamase_a
SLU3-C3f	77.46	3.37	*LD21* sp.	Bacteroidales	5	44		beta_lactamase_b_1_2
SLU3-C4f	42.73	1.06	*UBA6115* sp.	Christensenellales	8	49 (54.5)		beta_lactamase_d_2
SLU3-C5f	46.31	5.14	*UBA1413 sp002304925*	Bacteroidales	26	49 (50)		beta_lactamase_d_2; erm_f; tet_rpg
SLU3-C6f	67.29	4.46	*JAFGBI01* sp.	Polyangiales	11	73		beta_lactamase_a; mph
SLU3-C7f	92.89	0.14	Aristaeellaceae	Christensenellales	5	62		erm_f; aminoglycoside_e
SLU3-C8f	27.63	0.19	*JAKQIM01* sp.	Anaerolineales	10	58 (59)		beta_lactamase_d_2
SLU3-C9f	92.44	5.49	*PLanc-01 sp012744475*	Sedimentisphaerales	7	62 (62.5)		beta_lactamase_b_1_2

[1] C denotes genome bin completion

2 R denotes genome bin redundancy
3 GC % of the closest reference genome by GTDB-Tk
4 Previous taxonomical identity *Giesbergeria sp017988165*

References

1.   Hutchings M, Truman A, Wilkinson B. 2019. Antibiotics: past, present and future. Curr Opin Microbiol 51:72–80.

2.   Naghavi M, Vollset SE, Ikuta KS, Swetschinski LR, Gray AP, Wool EE, Robles Aguilar G, Mestrovic T, Smith G, Han C, Hsu RL, Chalek J, Araki DT, Chung E, Raggi C, Gershberg Hayoon A, Davis Weaver N, Lindstedt PA, Smith AE, Altay U, Bhattacharjee N V., Giannakis K, Fell F, McManigal B, Ekapirat N, Mendes JA, Runghien T, Srimokla O, Abdelkader A, Abd-Elsalam S, Aboagye RG, Abolhassani H, Abualruz H, Abubakar U, Abukhadijah HJ, Aburuz S, Abu-Zaid A, Achalapong S, Addo IY, Adekanmbi V, Adeyeoluwa TE, Adnani QES, Adzigbli LA, Afzal MS, Afzal S, Agodi A, Ahlstrom AJ, Ahmad A, Ahmad S, Ahmad T, Ahmadi A, Ahmed A, Ahmed H, Ahmed I, Ahmed M, Ahmed S, Ahmed SA, Akkaif MA, Al Awaidy S, Al Thaher Y, Alalalmeh SO, AlBataineh MT, Aldhaleei WA, Al-Gheethi AAS, Alhaji NB, Ali A, Ali L, Ali SS, Ali W, Allel K, Al-Marwani S, Alrawashdeh A, Altaf A, Al-Tammemi AB, Al-Tawfiq JA, Alzoubi KH, Al-Zyoud WA, Amos B, Amuasi JH, Ancuceanu R, Andrews JR, Anil A, Anuoluwa IA, Anvari S, Anyasodor AE, Apostol GLC, Arabloo J, Arafat M, Aravkin AY, Areda D, Aremu A, Artamonov AA, Ashley EA, Asika MO, Athari SS, Atout MMW, Awoke T, Azadnajafabad S, Azam JM, Aziz S, Azzam AY, Babaei M, Babin FX, Badar M, Baig AA, Bajcetic M, Baker S, Bardhan M, Barqawi HJ, Basharat Z, Basiru A, Bastard M, Basu S, Bayleyegn NS, Belete MA, Bello OO, Beloukas A, Berkley JA, Bhagavathula AS, Bhaskar S, Bhuyan SS, Bielicki JA, Briko NI, Brown CS, Browne AJ, Buonsenso D, Bustanji Y, Carvalheiro CG, Castañeda-Orjuela CA, Cenderadewi M, Chadwick J, Chakraborty S, Chandika RM, Chandy S, Chansamouth V, Chattu VK, Chaudhary AA, Ching PR, Chopra H, Chowdhury FR, Chu DT, Chutiyami M, Cruz-Martins N, da Silva AG, Dadras O, Dai X, Darcho SD, Das S, De La Hoz FP, Dekker DM, Dhama K, Diaz D, Dickson BFR, Djorie SG, Dodangeh M, Dohare S, Dokova KG, Doshi OP, Dowou RK, Dsouza HL, Dunachie SJ, Dziedzic AM, Eckmanns T, Ed-Dra A, Eftekharimehrabad A, Ekundayo TC, El Sayed I, Elhadi M, El-Huneidi W, Elias C, Ellis SJ, Elsheikh R, Elsohaby I, Eltaha C, Eshrati B, Eslami M, Eyre DW, Fadaka AO, Fagbamigbe AF, Fahim A, Fakhri-Demeshghieh A, Fasina FO, Fasina MM, Fatehizadeh A, Feasey NA, Feizkhah A, Fekadu G, Fischer F, Fitriana I, Forrest KM, Fortuna Rodrigues C, Fuller JE, Gadanya MA, Gajdács M, Gandhi AP, Garcia-Gallo EE, Garrett DO, Gautam RK, Gebregergis MW, Gebrehiwot M, Gebremeskel TG, Geffers C, Georgalis L, Ghazy RM, Golechha M, Golinelli D, Gordon M, Gulati S, Gupta R Das, Gupta S, Gupta VK, Habteyohannes AD, Haller S, Harapan H, Harrison ML, Hasaballah AI, Hasan I, Hasan RS, Hasani H, Haselbeck AH, Hasnain MS, Hassan II, Hassan S, Hassan Zadeh Tabatabaei MS, Hayat K, He J, Hegazi OE, Heidari M, Hezam K, Holla R, Holm M, Hopkins H, Hossain MM, Hosseinzadeh M, Hostiuc S, Hussein NR, Huy LD, Ibáñez-Prada ED, Ikiroma A, Ilic IM, Islam SMS, Ismail F, Ismail NE, Iwu CD, Iwu-Jaja CJ, Jafarzadeh A, Jaiteh F, Jalilzadeh Yengejeh R, Jamora RDG, Javidnia J, Jawaid T, Jenney AWJ, Jeon HJ, Jokar M, Jomehzadeh N, Joo T, Joseph N, Kamal Z, Kanmodi KK,

Kantar RS, Kapisi JA, Karaye IM, Khader YS, Khajuria H, Khalid N, Khamesipour F, Khan A, Khan MJ, Khan MT, Khanal V, Khidri FF, Khubchandani J, Khusuwan S, Kim MS, Kisa A, Korshunov VA, Krapp F, Krumkamp R, Kuddus M, Kulimbet M, Kumar D, Kumaran EAP, Kuttikkattu A, Kyu HH, Landires I, Lawal BK, Le TTT, Lederer IM, Lee M, Lee SW, Lepape A, Lerango TL, Ligade VS, Lim C, Lim SS, Limenh LW, Liu C, Liu X, Liu X, Loftus MJ, Amin HIM, Maass KL, Maharaj SB, Mahmoud MA, Maikanti-Charalampous P, Makram OM, Malhotra K, Malik AA, Mandilara GD, Marks F, Martinez-Guerra BA, Martorell M, Masoumi-Asl H, Mathioudakis AG, May J, McHugh TA, Meiring J, Meles HN, Melese A, Melese EB, Minervini G, Mohamed NS, Mohammed S, Mohan S, Mokdad AH, Monasta L, Moodi Ghalibaf AA, Moore CE, Moradi Y, Mossialos E, Mougin V, Mukoro GD, Mulita F, Muller-Pebody B, Murillo-Zamora E, Musa S, Musicha P, Musila LA, Muthupandian S, Nagarajan AJ, Naghavi P, Nainu F, Nair TS, Najmuldeen HHR, Natto ZS, Nauman J, Nayak BP, Nchanji GT, Ndishimye P, Negoi I, Negoi RI, Nejadghaderi SA, Nguyen QAP, Noman EA, Nwakanma DC, O'Brien S, Ochoa TJ, Odetokun IA, Ogundijo OA, Ojo-Akosile TR, Okeke SR, Okonji OC, Olagunju AT, Olivas-Martinez A, Olorukooba AA, Olwoch P, Onyedibe KI, Ortiz-Brizuela E, Osuolale O, Ounchanum P, Oyeyemi OT, Padukudru PAM, Paredes JL, Parikh RR, Patel J, Patil S, Pawar S, Peleg AY, Peprah P, Perdigão J, Perrone C, Petcu IR, Phommasone K, Piracha ZZ, Poddighe D, Pollard AJ, Poluru R, Ponce-De-Leon A, Puvvula J, Qamar FN, Qasim NH, Rafai CD, Raghav P, Rahbarnia L, Rahim F, Rahimi-Movaghar V, Rahman M, Rahman MA, Ramadan H, Ramasamy SK, Ramesh PS, Ramteke PW, Rana RK, Rani U, Rashidi MM, Rathish D, Rattanavong S, Rawaf S, Redwan EMM, Reyes LF, Roberts T, Robotham J V., Rosenthal VD, Ross AG, Roy N, Rudd KE, Sabet CJ, Saddik BA, Saeb MR, Saeed U, Saeedi Moghaddam S, Saengchan W, Safaei M, Saghazadeh A, Saheb Sharif-Askari N, Sahebkar A, Sahoo SS, Sahu M, Saki M, Salam N, Saleem Z, Saleh MA, Samodra YL, Samy AM, Saravanan A, Satpathy M, Schumacher AE, Sedighi M, Seekaew S, Shafie M, Shah PA, Shahid S, Shahwan MJ, Shakoor S, Shalev N, Shamim MA, Shamshirgaran MA, Shamsi A, Sharifan A, Shastry RP, Shetty M, Shittu A, Shrestha S, Siddig EE, Sideroglou T, Sifuentes-Osornio J, Silva LMLR, Simões EAF, Simpson AJH, Singh A, Singh S, Sinto R, Soliman SSM, Soraneh S, Stoesser N, Stoeva TZ, Swain CK, Szarpak L, Sree Sudha TY, Tabatabai S, Tabche C, Taha ZMA, Tan KK, Tasak N, Tat NY, Thaiprakong A, Thangaraju P, Tigoi CC, Tiwari K, Tovani-Palone MR, Tran TH, Tumurkhuu M, Turner P, Udoakang AJ, Udoh A, Ullah N, Ullah S, Vaithinathan AG, Valenti M, Vos T, Vu HTL, Waheed Y, Walker AS, Walson JL, Wangrangsimakul T, Weerakoon KG, Wertheim HFL, Williams PCM, Wolde AA, Wozniak TM, Wu F, Wu Z, Yadav MKK, Yaghoubi S, Yahaya ZS, Yarahmadi A, Yezli S, Yismaw YE, Yon DK, Yuan CW, Yusuf H, Zakham F, Zamagni G, Zhang H, Zhang ZJ, Zielińska M, Zumla A, Zyoud SHH, Zyoud SH, Hay SI, Stergachis A, Sartorius B, Cooper BS, Dolecek C, Murray CJL. 2024. Global burden of bacterial antimicrobial resistance 1990–2021: a systematic analysis with forecasts to 2050. The Lancet 404:1199–1226.

3. Ebmeyer S, Kristiansson E, Larsson DGJ. 2025. Unraveling the origins of mobile antibiotic resistance genes using random forest classification of large-scale genomic data. Environ Int 198:109374.

1062   4.   Jiang X, Ellabaan MMH, Charusanti P, Munck C, Blin K, Tong Y, Weber T, Sommer
1063   MOA, Lee SY. 2017. Dissemination of antibiotic resistance genes from antibiotic
1064   producers to pathogens. Nat Commun 8.
1065

1066   5.   Gholipour S, Lee D, Tokuriki N. 2025. Molecular Evolution and Origins of Antibiotic
1067   Resistance Genes. Biochemistry 64:2516–2528.
1068

1069   6.   Ebmeyer S, Kristiansson E, Larsson DGJ. 2021. A framework for identifying the recent
1070   origins of mobile antibiotic resistance genes. Commun Biol 4.
1071

1072   7.   Klümper U, Fang P, Li B, Xia Y, Frigon D, Hamilton KA, Quon H, Berendonk TU, de la
1073   Cruz Barron M. 2025. Towards the integration of antibiotic resistance gene mobility into
1074   environmental surveillance and risk assessment. npj Antimicrobials and Resistance 3:81.
1075

1076   8.   Larsson DGJ, Flach CF. 2022. Antibiotic resistance in the environment. Nat Rev
1077   Microbiol 20:257–269.
1078

1079   9.   Inda-Díaz JS, Lund D, Parras-Moltó M, Johnning A, Bengtsson-Palme J, Kristiansson E.
1080   2023. Latent antibiotic resistance genes are abundant, diverse, and mobile in human,
1081   animal, and environmental microbiomes. Microbiome 11.
1082

1083   10.   Mlynarcik P, Zdarska V, Kolar M. 2025. Are Putative Beta-Lactamases Posing a Potential
1084   Future Threat? Antibiotics 14:1174.
1085

1086   11.   Karkman A, Do TT, Walsh F, Virta MPJ. 2018. Antibiotic-Resistance Genes in Waste
1087   Water. Trends Microbiol 26:220–228.
1088

1089   12.   Bengtsson-Palme J, Larsson DGJ. 2015. Antibiotic resistance genes in the environment:
1090   Prioritizing risks. Nat Rev Microbiol 13:396.
1091

1092   13.   Sundqvist M. 2014. Reversibility of antibiotic resistance. Ups J Med Sci 119:142–148.
1093

1094   14.   Markkanen M, Pezzutto D, Virta M, Karkman A. 2025. Sulfonamide resistance gene *sul4*
1095   is hosted by common wastewater sludge bacteria and found in various newly described
1096   contexts and hosts. Microbiol Spectr 14.
1097

1098   15.   Parras-Moltó M, Lund D, Ebmeyer S, Larsson DGJ, Johnning A, Kristiansson E. 2025.
1099   The transfer of antibiotic resistance genes between evolutionarily distant bacteria.
1100   mSphere 10.
1101

1102   16.   Cantón R, María González-Alba J, Galán JC, Stefani S. 2012. CTX-M enzymes: origin
1103   and diffusion. Front Microbiol 3:110.
1104

1105   17.   Berglund F, Ebmeyer S, Kristiansson E, Larsson DGJ. 2023. Evidence for wastewaters as
1106   environments where mobile antibiotic resistance genes emerge. Commun Biol 6.
1107

1108 18.   Martínez JL, Coque TM, Baquero F. 2014. What is a resistance gene? Ranking risk in
1109   resistomes. Nat Rev Microbiol 13:116–123.
1110
1111 19.   Zhang A-N, Gaston JM, Dai CL, Zhao S, Poyet M, Groussin M, Yin X, Li L-G, Van
1112   Loosdrecht MCM, Topp E, Gillings MR, Hanage WP, Tiedje JM, Moniz K, Alm EJ,
1113   Zhang T. 2021. An omics-based framework for assessing the health risk of antimicrobial
1114   resistance genes. Nat Commun 4765.
1115
1116 20.   Quince C, Walker AW, Simpson JT, Loman NJ, Segata N. 2017. Shotgun metagenomics,
1117   from sampling to analysis. Nat Biotechnol 35:833–844.
1118
1119 21.   Abramova A, Karkman A, Bengtsson-Palme J. 2024. Metagenomic assemblies tend to
1120   break around antibiotic resistance genes. BMC Genomics 25:959.
1121
1122 22.   Heidelbach S, Mølvang Dall S, Støtt Bøjer J, Nissen J, der Maas van, Sereika M,
1123   Kirkegaard RH, Jensen SI, Just S, Thorlacius-Ussing O, Hose K, Dyhre Nielsen T.
1124   Nanomotif: Identification and Exploitation of DNA Methylation Motifs in Metagenomes 1
1125   using Oxford Nanopore Sequencing 2 3 https://doi.org/10.1101/2024.04.29.591623.
1126
1127 23.   Beaulaurier J, Zhu S, Deikus G, Mogno I, Zhang XS, Davis-Richardson A, Canepa R,
1128   Triplett EW, Faith JJ, Sebra R, Schadt EE, Fang G. 2018. Metagenomic binning and
1129   association of plasmids with bacterial host genomes using DNA methylation. Nat
1130   Biotechnol 36:61–69.
1131
1132 24.   Frost LS, Leplae R, Summers AO, Toussaint A. 2005. Mobile genetic elements: The
1133   agents of open source evolution. Nat Rev Microbiol 3:722–732.
1134
1135 25.   Diebold PJ, Rhee MW, Shi Q, Trung NV, Umrani F, Ahmed S, Kulkarni V, Deshpande P,
1136   Alexander M, Thi Hoa N, Christakis NA, Iqbal NT, Ali SA, Mathad JS, Brito IL. 2023.
1137   Clinically relevant antibiotic resistance genes are linked to a limited set of taxa within gut
1138   microbiome worldwide. Nat Commun 14.
1139
1140 26.   Ghosh D, Veeraraghavan B, Elangovan R, Vivekanandan P. 2020. Antibiotic resistance
1141   and epigenetics: More to it than meets the eye. Antimicrob Agents Chemother 64.
1142
1143 27.   Seong HJ, Han SW, Sul WJ. 2021. Prokaryotic DNA methylation and its functional roles.
1144   The journal of microbiology 59:242–248.
1145
1146 28.   Tuomala H, Holtel J, Markkanen M, Patpatia S, Kaansalo K, Rolland C, Bayfield OW,
1147   Ranta K, Skurnik M, Wittmann J, Kiljunen S. 2025. The Host R-M Systems Change the
1148   Host Range of Staphylococcus Phage EBHT. Microbiologyopen 14.
1149
1150 29.   Seong HJ, Roux S, Hwang CY, Sul WJ. 2022. Marine DNA methylation patterns are
1151   associated with microbial community composition and inform virus-host dynamics.
1152   Microbiome 10.
1153

30. Tourancheau A, Mead EA, Zhang XS, Fang G. 2021. Discovering multiple types of DNA methylation from bacteria and microbiome using nanopore sequencing. Nat Methods 18:491–498.

31. Wilbanks EG, Doré H, Ashby MH, Heiner C, Roberts RJ, Eisen JA. 2022. Metagenomic methylation patterns resolve bacterial genomes of unusual size and structural complexity. ISME Journal 16:1921–1931.

32. Hiraoka S, Okazaki Y, Anda M, Toyoda A, Nakano S ichi, Iwasaki W. 2019. Metaepigenomic analysis reveals the unexplored diversity of DNA methylation in an environmental prokaryotic community. Nat Commun 10.

33. Li T, Zhang X, Luo F, Wu FX, Wang J. 2020. MultiMotifMaker: A multi-thread tool for identifying DNA methylation motifs from Pacbio reads. IEEE/ACM Trans Comput Biol Bioinform 17:220–225.

34. Partanen V, Dekić Rozman S, Karkman A, Muurinen J, Hiltunen T, Virta M. 2025. Tracking horizontal gene transfer of antimicrobial resistance genes in microbial community with sequence barcodes. ISME J 5.

35. Hogle SL, Tamminen M, Hiltunen T. 2024. Complete genome sequences of 30 bacterial species from a synthetic community. Microbiol Resour Announc 13.

36. Feng X, Cheng H, Portik D, Li H. 2022. Metagenome assembly of high-fidelity long reads with hifiasm-meta. Nat Methods 19:671–674.

37. Mölder F, Jablonski KP, Letcher B, Hall MB, Tomkins-Tinch CH, Sochat V, Forster J, Lee S, Twardziok SO, Kanitz A, Wilm A, Holtgrewe M, Rahmann S, Nahnsen S, Köster J. 2021. Sustainable data analysis with Snakemake. F1000Res 10:33.

38. Pacific Biosciences. ipdsummary - a tool to detect dna base-modifications from kinetic signatures. https://www.pacb.com/support/software-downloads/. Pacific Biosciences.

39. Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, Blondel M, Prettenhofer P, Weiss R, Dubourg V, Vanderplas J, Passos A, Cournapeau D. 2011. Scikit-learn: Machine Learning in Python. Journal of Machine Learning Research 12:2825–2830.

40. McInnes L, Healy J, Melville J. 2018. UMAP: Uniform Manifold Approximation and Projection for Dimension Reduction https://doi.org/10.48550/arXiv.1802.03426.

41. Chklovski A, Parks DH, Woodcroft BJ, Tyson GW. 2023. CheckM2: a rapid, scalable and accurate tool for assessing microbial genome quality using machine learning. Nat Methods 20:1203–1212.

1199 42. Parks DH, Chuvochina M, Rinke C, Mussig AJ, Chaumeil PA, Hugenholtz P. 2022.
1200   GTDB: An ongoing census of bacterial and archaeal diversity through a phylogenetically
1201   consistent, rank normalized and complete genome-based taxonomy. Nucleic Acids Res
1202   50:D785–D794.
1203

1204 43. Schneider TD, Stephens RM. Sequence logos: a new way to display consensus sequences.
1205   Nucleic Acids Res 18:6097.
1206

44. Bortolaia V, Kaas RS, Ruppe E, Roberts MC, Schwarz S, Cattoir V, Philippon A, Allesoe
RL, Rebelo AR, Florensa AF, Fagelhauer L, Chakraborty T, Neumann B, Werner G,
Bender JK, Stingl K, Nguyen M, Coppens J, Xavier BB, Malhotra-Kumar S, Westh H,
Pinholt M, Anjum MF, Duggett NA, Kempf I, Nykäsenoja S, Olkkola S, Wieczorek K,
Amaro A, Clemente L, Mossong J, Losch S, Ragimbeau C, Lund O, Aarestrup FM. 2020.
ResFinder 4.0 for predictions of phenotypes from genotypes. J Antimicrob Chemother
75:3491–3500.
1214

1215 45. Berglund F, Österlund T, Boulund F, Marathe NP, Larsson DGJ, Kristiansson E. 2019.
1216   Identification and reconstruction of novel antibiotic resistance genes from metagenomes.
1217   Microbiome 7.
1218

1219 46. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL.
1220   2009. BLAST+: architecture and applications. BMC Bioinformatics 10:421.
1221

1222 47. Shen W, Le S, Li Y, Hu F. 2016. SeqKit: A cross-platform and ultrafast toolkit for
1223   FASTA/Q file manipulation. PLoS One 11.
1224

1225 48. Camargo AP, Roux S, Schulz F, Babinski M, Xu Y, Hu B, Chain PSG, Nayfach S,
1226   Kyrpides NC. 2023. Identification of mobile genetic elements with geNomad. Nat
1227   Biotechnol 42:1303–1312.
1228

1229 49. Shao M, Ying N, Liang Q, Ma N, Leptihn S, Yu Y, Chen H, Liu C, Hua X. 2023. Pdif-
1230   mediated antibiotic resistance genes transfer in bacteria identified by pdifFinder. Brief
1231   Bioinform 24.
1232

1233 50. Schwengers O, Jelonek L, Dieckmann MA, Beyvers S, Blom J, Goesmann A. 2021.
1234   Bakta: Rapid and standardized annotation of bacterial genomes via alignment-free
1235   sequence identification. Microb Genom 7.
1236

1237 51. Rognes T, Flouri T, Nichols B, Quince C, Mahé F. 2016. VSEARCH: A versatile open
1238   source tool for metagenomics. PeerJ 2016.
1239

1240 52. Katoh K, Misawa K, Kuma K-I, Miyata T. 2002. MAFFT: a novel method for rapid
1241   multiple sequence alignment based on fast Fourier transform. Nucleic Acids Res 30:3059–
1242   66.
1243

1244   53.   Stamatakis A. 2014. RAxML version 8: A tool for phylogenetic analysis and post-analysis
1245   of large phylogenies. Bioinformatics 30:1312–1313.
1246

1247   54.   Letunic I, Bork P. 2021. Interactive tree of life (iTOL) v5: An online tool for phylogenetic
1248   tree display and annotation. Nucleic Acids Res 49:293–296.
1249

1250   55.   Shimoyama Y. 2024. pyGenomeViz: A genome visualization python package for
1251   comparative genomics. https://github.com/moshi4/pyGenomeViz.
1252

1253   56.   Siguier P, Perochon J, Lestrade L, Mahillon J, Chandler M. 2006. ISfinder: the reference
1254   centre for bacterial insertion sequences. Nucleic Acids Res 34.
1255

1256   57.   Pereira MB, Wallroth M, Kristiansson E, Axelson-Fisk M. 2016. HattCI: Fast and
1257   Accurate attC site Identification Using Hidden Markov Models. J Comput Biol 23:891–
1258   902.
1259

58.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K,
Bates R, Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A,
Romera-Paredes B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman D, Clancy E,
Zielinski M, Steinegger M, Pacholska M, Berghammer T, Bodenstein S, Silver D, Vinyals
O, Senior AW, Kavukcuoglu K, Kohli P, Hassabis D. 2021. Highly accurate protein
structure prediction with AlphaFold. Nature 596:583–589.
1266

1267   59.   van Kempen M, Kim SS, Tumescheit C, Mirdita M, Lee J, Gilchrist CLM, Söding J,
1268   Steinegger M. 2024. Fast and accurate protein structure search with Foldseek. Nat
1269   Biotechnol 42:243–246.
1270

1271   60.   Shaw J, Yu YW. 2025. Rapid species-level metagenome profiling and containment
1272   estimation with sylph. Nat Biotechnol 43:1348–1359.
1273

1274   61.   Kruchten N, Seier A, Parmer C. 2025. An interactive, open-source, and browser-based
1275   graphing library for Python. https://github.com/plotly/plotly.py.
1276

1277   62.   RStudio Team. Inc. Boston. 2016. Integrated development for R. RStudio.
1278   http://www.rstudio.com/.
1279

1280   63.   McMurdie PJ, Holmes S. 2013. Phyloseq: An R Package for Reproducible Interactive
1281   Analysis and Graphics of Microbiome Census Data. PLoS One 8.
1282

1283   64.   Wickham H. 2016. ggplot2: Elegant graphics for data analysis.
1284   https://ggplot2.tidyverse.org.
1285

1286   65.   Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, Mcglinn D, Minchin PR,
1287   O'hara RB, Simpson GL, Solymos P, Henry M, Stevens H, Szoecs E, Wagner H. 2020.
1288   vegan: Community Ecology Package. https://cran.r-project.org/package=vegan.
1289

66. On SLW. 2025. International Committee on Systematics of Prokaryotes (ICSP) Subcommittee on the Taxonomy of Campylobacter and related bacteria: minutes of the closed meetings, 17 January and 31 January 2024. Int J Syst Evol Microbiol 75.

67. Jain C. 2008. The E. coli RhlE RNA helicase regulates the function of related RNA helicases during ribosome assembly. RNA 14:381–389.

68. Poolman B;, Royer TJ, Mainzer SE, Schmidt BF. 1990. Carbohydrate Utilization in Streptococcus thermophilus: Characterization of the Genes for Aldose 1-Epimerase (Mutarotase) and UDPglucose 4-Epimerase. J Bacteriol 172:4037–3047.

69. Romsang A, Atichartpongkul S, Trinachartvanit W, Vattanaviboon P, Mongkolsuk S. 2013. Gene expression and physiological role of Pseudomonas aeruginosa methionine sulfoxide reductases during oxidative stress. J Bacteriol 195:3299–3308.

70. Jiang X, Hall AB, Xavier RJ, Alm EJ. 2019. Comprehensive analysis of chromosomal mobile genetic elements in the gut microbiome reveals phylum-level niche-adaptive gene pools. PLoS One 14.

71. Guo Z, Qin X, Yue M, Wu L, Li N, Su J, Jiang M. 2025. IS26 carrying blaKPC−2 mediates carbapenem resistance heterogeneity in extensively drug-resistant Klebsiella pneumoniae isolated from clinical sites. Mob DNA 16.

72. Targant H, Doublet B, Aarestrup FM, Cloeckaert A, Madec JY. 2010. IS6100-mediated genetic rearrangement within the complex class 1 integron In104 of the Salmonella genomic island 1. Journal of Antimicrobial Chemotherapy 65:1543–1545.

73. Sewunet T, Razavi M, Tellapragada C, Giske CG. 2025. Silent spread of mcr-9 in ESBL-producing Enterobacteriaceae clinical isolates, Jimma, Ethiopia. PLoS One 20.

74. Schmid S, Berger B, Haas D. 1999. Target joining of duplicated Insertion Sequence IS21 is assisted by IstB protein in vitro. J Bacteriol 181:2286–2289.

75. Díaz-García C, Sánchez-Osuna M, Serra-Compte A, Karakatsanidou I, Gómez-Sánchez I, Fidalgo B, Barbuzana-Armas C, Fittipaldi M, Rosselli R, Vinyoles J, González S, Pich OQ, Espasa M, Yáñez MA. 2025. Mapping antimicrobial resistance landscape at a city scale sewage network. Science of the Total Environment 974.

76. Dekić Rozman S, Puljko A, Karkman A, Virta M, Udiković-Kolić N. 2024. Bacterial hosts of clinically significant beta-lactamase genes in Croatian wastewaters. FEMS Microbiol Ecol 100.

77. Hultman J, Tamminen M, Pärnänen K, Cairns J, Karkman A, Virta M. 2018. Host range of antibiotic resistance genes in wastewater treatment plant influent and effluent. FEMS Microbiol Ecol 94.

78. Buzzanca D, Kerkhof PJ, Alessandria V, Rantsiou K, Houf K. 2023. Arcobacteraceae comparative genome analysis demonstrates genome heterogeneity and reduction in species isolated from animals and associated with human illness. Heliyon 9.

79. Pishgar R, Dominic JA, Sheng Z, Tay JH. 2019. Denitrification performance and microbial versatility in response to different selection pressures. Bioresour Technol 281:72–83.

80. Buzzanca D, Chiarini E, Alessandria V. 2024. Arcobacteraceae: An Exploration of Antibiotic Resistance Featuring the Latest Research Updates. Antibiotics 13.

81. Heo H, Nguyen-Dinh T, Jung MY, Greening C, Yoon S. 2025. Hydrogen-dependent dissimilatory nitrate reduction to ammonium enables growth of Campylobacterota isolates. ISME Journal 19.

82. Jia B, Raphenya AR, Alcock B, Waglechner N, Guo P, Tsang KK, Lago BA, Dave BM, Pereira S, Sharma AN, Doshi S, Courtot M, Lo R, Williams LE, Frye JG, Elsayegh T, Sardar D, Westman EL, Pawlowski AC, Johnson TA, Brinkman FSL, Wright GD, McArthur AG. 2017. CARD 2017: Expansion and model-centric curation of the comprehensive antibiotic resistance database. Nucleic Acids Res 45:D566–D573.

83. Balalovski P, Grainge I. 2020. Mobilization of pdif modules in Acinetobacter: A novel mechanism for antibiotic resistance gene shuffling? Mol Microbiol 114:699–709.

84. Mindlin S, Beletsky A, Mardanov A, Petrova M. 2019. Adaptive dif modules in permafrost strains of acinetobacter lwoffiiand their distribution and abundance among present day acinetobacter strains. Front Microbiol 10.

85. Blackwell GA, Hall RM. 2017. The tet39 determinant and the msrE-mphE genes in Acinetobacter plasmids are each part of discrete modules flanked by inversely oriented pdif (XerC-XerD) sites. Antimicrob Agents Chemother 61.

86. Girlich D, Bonnin RA, Bogaerts P, De Laveleye M, Huang DT, Dortet L, Glaser P, Glupczynski Y, Naas T. 2017. Chromosomal amplification of the blaOXA-58 carbapenemase gene in a Proteus mirabilis clinical isolate. Antimicrob Agents Chemother 61.

87. Tempel S, Bedo J, Talla E. 2022. From a large-scale genomic analysis of insertion sequences to insights into their regulatory roles in prokaryotes. BMC Genomics 23.

88. Liu H, Moran RA, Chen Y, Doughty EL, Hua X, Jiang Y, Xu Q, Zhang L, Blair JMA, McNally A, Van Schaik W, Yu Y. 2021. Transferable Acinetobacter baumannii plasmid pDETAB2 encodes OXA-58 and NDM-1 and represents a new class of antibiotic resistance plasmids. Journal of Antimicrobial Chemotherapy 76:1130–1134.

1381   89.   Geneva: World Health Organization; 2024. WHO Bacterial Priority Pathogens List, 2024:
1382   Bacterial pathogens of public health importance to guide research, development and
1383   strategies to prevent and control antimicrobial resistance.
1384

1385   90.   Lu S, Ryu SH, Chung BS, Chung YR, Park W, Jeon CO. 2007. Simplicispira limi sp.
1386   nov., isolated from activated sludge. Int J Syst Evol Microbiol 57:31–34.
1387

1388   91.   Peng C, Gao Y, Fan X, Peng P, Huang H, Zhang X, Ren H. 2019. Enhanced biofilm
1389   formation and denitrification in biofilters for advanced nitrogen removal by rhamnolipid
1390   addition. Bioresour Technol 287.
1391

1392   92.   Liu W, Liu X, Liao J, Zhang Y, Liang X. 2010. Identification of bla OXA-128 and bla
1393   OXA-129, two novel OXA-type extended-spectrum-β-lactamases in Pseudomonas
1394   aeruginosa, in Hunan Province, China. J Basic Microbiol 50.
1395

1396   93.   Valiatti TB, Streling AP, Cayô R, Santos FF, Almeida MS, Tonini MAL, Gales AC. 2024.
1397   Genomic analysis of a Pseudomonas aeruginosa strain harbouring blaKPC-2 and blaOXA-
1398   129 belonging to high-risk clone ST235 in Brazil. J Glob Antimicrob Resist 37:69–71.
1399

1400   94.   Gillings MR. 2017. Class 1 integrons as invasive species. Curr Opin Microbiol 38:10–15.
1401

1402   95.   Johansson MHK, Petersen TN, Nag S, Lagermann TMR, Birkedahl LEK, Tafaj S,
1403   Bradbury S, Collignon P, Daley D, Dougnon V, Fabiyi K, Coulibaly B, Dembélé R,
1404   Magloire N, Ouindgueta IJ, Hossain ZZ, Begoum A, Donchev D, Diggle M, Turnbull LA,
1405   Lévesque S, Berlinger L, Søgaard KK, Guevara PD, Duarte C, Maikanti P, Amlerova J,
1406   Drevinek P, Tkadlec J, Dilas M, Kaasch A, Westh HT, Bachtarzi MA, Amhis W, Salazar
1407   CES, Villacis JE, Lúzon MAD, Palau DB, Duployez C, Paluche M, Asante-Sefa S, Møller
1408   M, Ip M, Mareković I, Pál-Sonnevend A, Cocuzza CE, Dambrauskiene A, Macanze A,
1409   Cossa A, Mandomando I, Nwajiobi-Princewill P, Okeke IN, Kehinde AO, Adebiyi I,
1410   Akintayo I, Popoola O, Onipede A, Blomfeldt A, Nyquist NE, Bocker K, Ussher J, Ali A,
1411   Ullah N, Khan H, Gustafson NW, Jarrar I, Al-Hamad A, Luvira V, Paveenkittiporn W,
1412   Baran I, Mwansa JCL, Sikakwa L, Yamba K, Aarestrup FM. 2025. Investigation of
1413   mobile genetic elements and their association with antibiotic resistance genes in clinical
1414   pathogens worldwide. PLoS One 20.
1415

1416   96.   Johansson MHK, Aarestrup FM, Petersen TN. 2023. Importance of mobile genetic
1417   elements for dissemination of antimicrobial resistance in metagenomic sewage samples
1418   across the world. PLoS One 18.
1419

1420   97.   Lenferink WB, van Alen TA, Jetten MSM, Op den Camp HJM, van Kessel MAHJ,
1421   Lücker S. 2024. Genomic analysis of the class Phycisphaerae reveals a versatile group of
1422   complex carbon-degrading bacteria. Antonie van Leeuwenhoek, International Journal of
1423   General and Molecular Microbiology 117.
1424

1425   98.   Hamidian M, Holt KE, Hall RM. 2015. The complete sequence of Salmonella genomic
1426   island SGI2. Journal of Antimicrobial Chemotherapy 70:617–619.

99. Yan W, Hall AB, Jiang X. 2022. Bacteroidales species in the human gut are a reservoir of antibiotic resistance genes regulated by invertible promoters. NPJ Biofilms Microbiomes 8.

100. Owusu-Agyeman I, Plaza E, Cetecioglu Z. 2022. Long-term alkaline volatile fatty acids production from waste streams: Impact of pH and dominance of Dysgonomonadaceae. Bioresour Technol 346.

101. Hirakata Y, Mei R, Morinaga K, Katayama T, Tamaki H, Meng X ying, Watari T, Yamaguchi T, Hatamoto M, Nobu MK. 2023. Identification and cultivation of anaerobic bacterial scavengers of dead cells. ISME Journal 17:2279–2289.

102. Ueki A, Akasaka H, Suzuki D, Ueki K. 2006. Paludibacter propionicigenes gen. nov., sp. nov., a novel strictly anaerobic, Gram-negative, propionate-producing bacterium isolated from plant residue in irrigated rice-field soil in Japan. Int J Syst Evol Microbiol 56:39–44.

103. Ma J, Sun H, Li B, Wu B, Zhang X, Ye L. 2024. Horizontal transfer potential of antibiotic resistance genes in wastewater treatment plants unraveled by microfluidic-based mini-metagenomics. J Hazard Mater 465.

104. Wen J, Zhang H, Chu D, Chen X, Feng J, Wang Y, Liu G, Zhang Y, Li Y, Ning K. 2024. Deep learning revealed the distribution and evolution patterns for invertible promoters across bacterial lineages. Nucleic Acids Res 52:12817–12830.

105. Jiang X, Hall AB, Arthur TD, Plichta DR, Covington CT, Poyet M, Crothers J, Moses PL, Tolonen AC, Vlamakis H, Alm EJ, Xavier RJ. 2019. Invertible promoters mediate bacterial phase variation, antibiotic resistance, and host adaptation in the gut. Science (1979) 363:181–187.

106. Hendriksen RS, Munk P, Njage P, van Bunnik B, McNally L, Lukjancenko O, Röder T, Nieuwenhuijse D, Pedersen SK, Kjeldgaard J, Kaas RS, Clausen PTLC, Vogt JK, Leekitcharoenphon P, van de Schans MGM, Zuidema T, de Roda Husman AM, Rasmussen S, Petersen B, Amid C, Cochrane G, Sicheritz-Ponten T, Schmitt H, Alvarez JRM, Aidara-Kane A, Pamp SJ, Lund O, Hald T, Woolhouse M, Koopmans MP, Vigre H, Petersen TN, Aarestrup FM. 2019. Global monitoring of antimicrobial resistance based on metagenomics analyses of urban sewage. Nat Commun 10:1124.

107. Karkman A, Johnson TA, Lyra C, Stedtfeld RD, Tamminen M, Tiedje JM, Virta M. 2016. High-throughput quantification of antibiotic resistance genes from an urban wastewater treatment plant. FEMS Microbiol Ecol 92.

108. Ling M, Szarvas J, Kurmauskaitė V, Kiseliovas V, Žilionis R, Avot B, Munk P, Aarestrup FM. 2024. High throughput single cell metagenomic sequencing with semi-permeable capsules: unraveling microbial diversity at the single-cell level in sewage and fecal microbiomes. Front Microbiol 15.

109. Wlodarski MA, Lau TTY, Alcock BP, Raphenya AR, Ta TE, Maguire F, Beiko RG, McArthur AG. 2025. CARD k-mers: Unmasking the pathogen hosts and genomic contexts of antimicrobial resistance genes in metagenomic sequences https://doi.org/10.1101/2025.09.15.676352.

110. Beaulaurier J, Schadt EE, Fang G. 2019. Deciphering bacterial epigenomes using modern sequencing technologies. Nat Rev Genet 20:157–172.

111. Ghatak S, Armstrong CM, Reed S, He Y. 2020. Comparative Methylome Analysis of Campylobacter jejuni Strain YH002 Reveals a Putative Novel Motif and Diverse Epigenetic Regulations of Virulence Genes. Front Microbiol 11.

112. Fang G, Munera D, Friedman DI, Mandlik A, Chao MC, Banerjee O, Feng Z, Losic B, Mahajan MC, Jabado OJ, Deikus G, Clark TA, Luong K, Murray IA, Davis BM, Keren-Paz A, Chess A, Roberts RJ, Korlach J, Turner SW, Kumar V, Waldor MK, Schadt EE. 2012. Genome-wide mapping of methylated adenine residues in pathogenic Escherichia coli using single-molecule real-time sequencing. Nat Biotechnol 30:1232–1239.

113. Pirone-Davies C, Hoffmann M, Roberts RJ, Muruvanda T, Timme RE, Strain E, Luo Y, Payne J, Luong K, Song Y, Tsai YC, Boitano M, Clark TA, Korlach J, Evans PS, Allard MW. 2015. Genome-wide methylation patterns in Salmonella enterica subsp. enterica Serovars. PLoS One 10.

114. Oliveira PH, Touchon M, Rocha EPC. 2014. The interplay of restriction-modification systems with mobile genetic elements and their prokaryotic hosts. Nucleic Acids Res 42:10618–10631.

115. Zorea A, Moraïs S, Pellow D, Gershoni-Yahalom O, Probst M, Nadler S, Shamir R, Rosental B, Elia N, Mizrahi I. 2026. ProFiT-SPEci-FISH: a novel approach for linking plasmids to hosts in complex microbial communities at the single-cell level. Microbiome 14.

116. Shaw LP, Rocha EPC, Maclean RC. 2023. Restriction-modification systems have shaped the evolution and distribution of plasmids across bacteria. Nucleic Acids Res 51:6806–6818.

117. Dimitriu T, Szczelkun MD, Westra ER. 2024. Various plasmid strategies limit the effect of bacterial restriction–modification systems against conjugation. Nucleic Acids Res 52:12976–12986.

118. Coluzzi C, Rocha EPC. 2025. The Spread of Antibiotic Resistance Is Driven by Plasmids Among the Fastest Evolving and of Broadest Host Range. Mol Biol Evol 42.