Title: Mental health outcomes following a psilocybin session within Oregon's state-regulated model: A naturalistic study

Authors: Amanda Gow, MS, LPC-a,[1] Emily Shih, PhD,[2] Ryan Reid, BS,[1] Jimmy J. Qian, MS,[2] Charan Mellor, MA,[1] Lynne Alison McInnes, MD,[2] Robin Carhart-Harris, PhD,[3] Jennie N. Davis, PhD[1,4*]

Affiliations:

1. Bendable Therapy, Bend, Oregon, USA

2. Osmind, San Francisco, California, USA

3. Weill Institute for Neurosciences, University of California San Francisco, San Francisco, California, USA

4. Center for Health Systems Effectiveness, Oregon Health & Science University, Portland, Oregon, USA

*Corresponding Author:

Jennie N. Davis, Bendable Therapy, PO Box 1229, Bend, OR 97709,

jdavis@bendabletherapy.org, 541-408-0790

Abstract

Background: In 2020, Oregon became the first U.S. state to establish a regulated framework

for adults to access psilocybin services using naturally-derived mushroom products. No studies

have examined mental health outcomes among individuals receiving psilocybin in this context.

Aims: To evaluate changes in self-reported symptoms of depression, anxiety, and well-being

30-days post-psilocybin session under the Oregon state-regulated model , and document

session-related adverse events and doses consumed.

Methods: This was a naturalistic study (March 2024-April 2025) among adults ≥21 years

participating in a legal psilocybin services program. Online surveys were completed pre-session,

1-day, and 30-days post-session. Primary outcomes were change in depression, anxiety, and

well-being symptoms pre-session to 30-days post-session evaluated using linear mixed-effects

models (random effect: timepoint; fixed effects: sex, concurrent psychiatric medication use, age,

session dose [total psilocybin equivalents, TPE, mg: psilocybin mg + 1.39 * psilocin mg]).

Adverse events (e.g., hallucinogen persisting perception disorder [HPPD]) were assessed at 1-

day and 30-days post-session.

Results: Participants (n=88; median age 43 years; 52% male) were predominantly Oregon

residents (53.4%), psychedelic-experienced (64.8%), and concurrently using psychiatric

medication (46.6%). All outcomes improved significantly at 30-days post-session (p<0.001),

including in sensitivity analyses stratified by concurrent psychiatric medication usage (p<0.001

all outcomes, both groups). Two participants (2.3%) reported symptoms consistent with HPPD

at 1-day post-session, but none at 30-days. Mean dose was 27.8 mg (SD 8.2) TPE.

Conclusions: Psilocybin sessions delivered under the Oregon regulatory model were

associated with clinically meaningful improvements in depression, anxiety, and well-being 30-

days post-session, supporting therapeutic effectiveness of legal psilocybin services.

Keywords: psilocybin, Oregon, depression, anxiety

Introduction

Psilocybin has emerged as a promising therapeutic intervention for mental health disorders, including depression and anxiety (Carhart-Harris et al., 2016, 2018; Dorczok et al., 2025; Goodwin, Aaronson, et al., 2023; Goodwin et al., 2022; Griffiths et al., 2016). As a naturally occurring psychoactive alkaloid found in certain species of mushrooms, once ingested, psilocybin is converted to psilocin to produce its characteristic psychedelic effects by acting on serotonin receptors (e.g., 5-HT2$_A$) (Passie et al., 2002). These neuropharmacological actions are thought to underlie the improvements in mental health symptoms observed across controlled clinical trials and naturalistic observational studies (Calnan et al., 2025; Carhart-Harris et al., 2021; Compass, 2025; Davis et al., 2021; de la Salle et al., 2024; Goodwin, Aaronson, et al., 2023; Goodwin et al., 2022; Griffiths et al., 2016; Irrmischer et al., 2025; Nygart et al., 2022; Raison et al., 2023; Ross et al., 2016; von Rotz et al., 2023).

Most psilocybin clinical trials enroll highly selective samples under tightly controlled conditions, typically excluding individuals with complex psychiatric diagnoses or active psychiatric medication use, and administering synthetic psilocybin within standardized dosing protocols (Carhart-Harris et al., 2021; Davis et al., 2021; Griffiths et al., 2016; Raison et al., 2023; Ross et al., 2016; von Rotz et al., 2023). Only one trial has used a natural mushroom formulation to examine impacts of psilocybin on alcohol use disorder, with findings highlighting the therapeutic potential of natural mushroom formulations (Jensen et al., 2025). Naturalistic studies introduce greater variability in participants' medical and psychiatric histories, session formats, and psilocybin source (often whole fruit mushrooms), yet still demonstrate improvements in depression, anxiety, and well-being (Calnan et al., 2025; de la Salle et al., 2024; Irrmischer et al., 2025; Morton et al., 2023; Nygart et al., 2022; Tap et al., 2025). However, additional research bridging controlled clinical efficacy trials and real-world effectiveness is needed to understand psilocybin's mental health benefits in diverse, community-based populations.

In 2020, Oregon voters approved Measure 109, establishing the first regulated

psilocybin services model in the United States and globally. These services are administered by

the Oregon Health Authority (OHA) through its Oregon Psilocybin Services (OPS) section

(Oregon Health Authority, n.d.) Oregon's non-medical model permits adults aged 21 years and

older to access psilocybin by consuming products derived from whole fruit mushrooms

(*Psilocybe cubensis*) at licensed service centers under the supervision of licensed facilitators

(Oregon Health Authority, 2024). The model does not require a medical diagnosis, prescription,

or clinical referral, despite mental health applications being common reasons for psilocybin use

(Dorczok et al., 2025). In 2025, statewide aggregate data released under Senate Bill 303 began

providing basic information, including demographics, psilocybin dosages, and session types.

However, these data do not include clinical outcomes or follow-up assessments, limiting insight

into the model's therapeutic effectiveness.

To address this gap, this study aimed to evaluate changes in self-reported symptoms of

depression, anxiety, and well-being following a psilocybin session under Oregon's regulated

model, and document session-related adverse events and psilocybin doses consumed.

Predictors of session dosage were additionally explored to understand potential correlates of

dosing decisions (e.g., participant characteristics, mental health history). Characterizing who

uses psilocybin services and with what outcomes is critical to understanding the benefits and

limitations of Oregon's regulated framework.

Methods

Setting and Participants

This prospective, naturalistic study (March 2024-April 2025) was integrated into an

existing Oregon legal psilocybin services program at Bendable Therapy ("Bendable"). Oregon

regulatory minimums require one facilitator-client preparation session 24-hours in advance of a

psilocybin administration session and to offer an optional integration session post-administration

(Oregon Health Authority, n.d.). Bendable enhanced the Oregon requirements by including

additional screening and support sessions **(Figure 1)**. Eligibility requirements for legal psilocbyin

services are: aged 21 years or older, no lithium use in the past 30 days (due to seizure risk

(Nayak et al., 2021)), no active psychosis, no active suicidal or homicidal ideation, not pregnant

or breastfeeding [Bendable only]) (Oregon Health Authority, 2024). During the study period, all

individuals who attended the virtual interview (Figure 1) and met eligibility criteria were invited to

the study and to complete the written informed consent. No compensation was provided to

study participants. WCG Institutional Review Board approved this study.

Procedures and measures

A pre-session screening application (completed on average 75 days prior to the

psilocybin administration session) collected demographics, medical, mental health, psychedelic

use histories, medication usage, and validated scales assessing self-reported symptoms of

depression (Patient Health Questionnaire, PHQ-9), anxiety (Generalized Anxiety Disorder,

GAD-7), and well-being (World Health Organization Well-Being Index, WHO-5) (Figure 1).

The PHQ-9 maps nine depression symptoms over the past two weeks onto the

Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, Text Revision criteria for

major depressive disorder; items are scored on a 4-point Likert scale (0="not at all" to 3="nearly

every day") (Kroenke et al., 2001)). For this study, the PHQ-9 was modified to PHQ-8 by

excluding the suicidal ideation item; scores were adjusted accordingly (Shin et al., 2019). The

GAD-7 is a 7-item self-report scale assessing generalized anxiety severity over the past two

weeks with items rated on a 4-point Likert scale (0="not at all" to 3="nearly every day") (Spitzer

et al., 2006). The WHO-5 is a 5-item measure of subjective psychological well-being where

participants rated positive mood states (e.g., feeling cheerful, relaxed, or energetic) over the

past two weeks on a 5-point scale (0="at no time" to 5="all of the time") (Topp et al., 2015).

On the administration session day, before consuming psilocybin, participants completed

the Imperial Psychedelic Predictor Scale (iPPS), a validated 8- (individual sessions) or 9-item

(group sessions) multidimensional self-report instrument designed to predict acute features of

the psychedelic experience across the domains of set, intention, and rapport; items were rated

0-100 (Angyus et al., 2024). Consumed psilocybin and psilocin doses (mg) were recorded.

A survey emailed to participants 30-days post-session (completed on average 34 days

post-session) included follow-up PHQ-8, GAD-7, and WHO-5 assessments, integration session

attendance (a post-session participant-facilitator meeting to discuss and process the psilocybin

session), and collected information on adverse events by assessing lingering unpleasant

psychological effects attributed to the psilocybin session (i.e., anxiety, existential struggle, social

disconnection, depersonalization/derealization) (Schlag et al., 2022), and assessed symptoms

of Hallucinogen Persisting Perception Disorder (HPPD), a condition of enduring, distressing

visual perceptions post-psychoactive substance intake (also assessed in a survey sent 1-day

post-session) (Figure 1). HPPD symptoms attributed to the psilocybin session and not due to

other medical or mental health conditions may be candidates for HPPD diagnosis (Zhou et al.,

2025). For participants completing two sessions within the follow-up period (e.g., weeklong

retreats), data were collected according to the first session only. Online assessments were

completed via Jotform (Jotform Inc.). **Supplemental Table 1** contains a summary of the study's

validated scales.

Participants completed individual or group facilitator-supervised psilocybin sessions at

licensed service centers following Oregon regulations (Oregon Health Authority, n.d.), and

Bendable's protocols were not altered for this study. Psilocybin was consumed as whole

mushrooms or tea made with ground whole mushrooms. Session dosing followed Bendable's

protocol of a total dose of 20-30 mg psilocybin including optional 'booster doses,' or additional

offered approximately 1-hr after the initial dose. Per Oregon rules, participants may request

lower or higher doses (up to 50 mg) (Oregon Psilocybin Services, 2025, n.d.).

Statistical analysis

Participant pre-session characteristics are presented as medians (range) or frequencies

(proportions). Primary outcomes were change in mental health status from pre-session to 30-

days post-session using the PHQ-8, GAD-7, and WHO-5 scales.

Scores for the PHQ-8 (0-24) were summed, and cut-points of 5, 10, 15, and 20 denoted

mild, moderate, moderately severe, and severe depression respectively; scores ≥10 indicated

clinically significant depression (Kroenke et al., 2001). GAD-7 scores (0-21) were summed with

cut-points of 5, 10, and 15 denoting mild, moderate, and severe anxiety respectively, with ≥10

indicating clinically significant anxiety (Spitzer et al., 2006). WHO-5 scores (0–25) were summed

with scores <13 indicating poor well-being. The PHQ-8, GAD-7, and WHO-5 demonstrated

strong internal consistency measured with Cronbach's alpha (PHQ-8: $\square$=0.87 pre-session and

30-days post-session; GAD-7: α =0.91 pre-session and α=0.89 30-days post-session; WHO-5:

α=0.96 pre-session and α=0.93 30-days post-session).

Per Oregon regulations, mushrooms were grown, processed, and potency-tested for

analyte content (psilocybin and psilocin) in Oregon licensed facilities; product packaging listed

analyte content in milligrams per gram of whole mushroom (Oregon Psilocybin Services, n.d.).

Oregon regulations define dose by psilocybin content; because psilocybin rapidly converts to its

pharmacologically active form of psilocin once ingested (Passie et al., 2002), Bendable

calculates dose as Total Psilocybin Equivalents (TPE = psilocybin mg + 1.39 × psilocin mg), a

metric assuming complete conversion of psilocybin to psilocin based on molecular weight

(MacCallum et al., 2022). We report both TPE and psilocybin dose.

We used separate linear mixed effects models using the 'lme4' package in R to evaluate

changes in outcomes across two timepoints (means, SE). Timepoint was modeled as a

categorical variable (pre-session as reference), with fixed effects of birth sex (male/female), age

(years), concurrent psychiatric medication use (yes/no), and dose (TPE, mg). Participant-level

random intercepts and slopes accounted for individual differences in pre-session symptom

levels and change over time. Prior to analyses, we evaluated covariate collinearity and excluded

the session type variable (group vs. individual) due to strong correlation with TPE. Effect sizes

30-days post session (relative to pre-session) were calculated using Cohen's *d*. This was

calculated by dividing the estimated marginal mean differences by the pooled standard

deviation, which incorporated both the residual variance and the random intercept variance from

the fitted model. Effect sizes were classified as small (0.20-0.49), moderate (0.50-0.79), large

($\geq$0.80), or very large ($\geq$1.2) (Cohen, 1992). Model fit was summarized using marginal and

condition R^2 to indicate variance explained by fixed effects and by the full model, respectively. A

post-hoc simulation-based power analysis ($\square$=0.05) indicated approximately 100% power for

large effects and 89% for small effects.

Sensitivity and exploratory analyses

In a sensitivity analysis, we repeated the adjusted models substituting TPE with

psilocybin dosage to assess robustness to an alternative dose specification. In a second

analysis to assess whether findings were driven by participants with additional dosing, we

restricted the sample to participants who completed a single session during the study period

(n=75), excluding those with multiple sessions (n=13). In a third analysis, we stratified the

sample by concurrent psychiatric medication use and repeated the adjusted models separately

among participants with (n=41) and without (n=47) pharmacotherapy to descriptively assess

whether patterns of outcomes were consistent across medication status.

To explore correlates of dosing decisions, we used univariate linear regression models

to examine predictors of session dose (TPE). Predictors included pre-session PHQ-8, GAD-7,

and WHO-5 scores; age; birth sex; most commonly reported diagnoses: depression, anxiety,

post-traumatic stress disorder (PTSD); concurrent psychiatric medication use; prior psychedelic

experience (Irrmischer et al., 2025). In a separate multivariate regression model, iPPS

subscales were entered simultaneously to assess their independent contributions to TPE. Given

these analyses' exploratory nature, p-values were not adjusted for multiple comparisons.

Analyses were conducted using R/R Studio version 4.4.2 (R Foundation), and included

only participants who completed all pre- and post-session surveys; we set statistical significance

at $p<0.05$.

Results

Participant characteristics

During the study period, n=311 screening applications were received. Of these, n=126

individuals met eligibility criteria for psilocybin services and were invited to participate, n=91

enrolled, and n=88 completed all study components and comprised the analytic sample.

Participants (n=88) ranged in age from 22-79 years (median: 43 years), were approximately

balanced by birth sex and gender identity (n=46, 52.0% male; n=44, 50.0% male-identifying),

and were predominantly White (n=77, 87.5%), educated (n=74, 84.1% with higher education),

and employed (n=59, 67.1%), with only 22.7% (n=20) reporting annual household income below

$50,000 **(Table 1)**. Most participants were Oregon residents (n=47, 53.4%) and 15.9% (n=14)

were military veterans. Depression (n=45, 51.1%), anxiety (n=37, 42.0%), and PTSD (n=17,

19.3%) were the most reported mental health diagnoses. Almost half (n=41, 46.6%) reported

concurrent use of psychiatric medications (medication type and dosage were not systematically

collected) and participation in mental health therapy (n=50, 56.8%). The majority (n=57, 64.8%)

had prior psychedelic experience, and cost was the most frequently reported barrier to seeking

psilocybin services (n=41, 46.6%) (Table 1). Most participants completed individual sessions

(n=50, 56.8%) rather than group sessions (n=38, 43.2%), and 80.0% (n=70) completed at least

one integration session during follow-up.

Mental health outcomes after a psilocybin session

We examined whether participation in a psilocybin session was associated with changes

in mental health outcomes. Pre-session, 69.3% (n=61 of participants reported mild to severe

depression symptoms, 67.0% (n=59) reported mild to severe anxiety symptoms, and 93.2%

n=82) reported low or very-low sense of well-being, reflecting clinically significant symptom

levels across all measures **(Table 2)**. At approximately 30-days post-session, improvements

were observed across measures, with 61.4% (n=54) of participants reporting minimal

depression, 70.5% (n=62) reporting minimal anxiety, and 64.8% (n=57) reporting normal levels

of well-being.

Modeling results similarly reflected significant improvements in mental health outcomes

over time (**Figure 2**, **Table 3)**. PHQ-8 and GAD-7 mean scores significantly decreased pre-

session to 30-days post-session, suggesting robust declines in depression and anxiety

symptoms: PHQ-8 score change of -4.63 (0.57 SE) points; $p<0.001$; $d=0.90$ (95% CI: 0.68,

1.12) large; GAD-7 score change of -4.85 (0.57) points; $p<0.001$; $d=1.04$ (0.80, 1.17) large.

WHO-5 well-being scores also improved at follow-up with a score change of +10.67 (0.59)

points; $p<0.001$; $d=-2.14$ (-2.38, -1.90) very large), indicating significant gains in well-being. Of

the covariates examined, only concurrent psychiatric medication use was positively associated

with post-session PHQ-8 scores ($p=0.040$).

Sensitivity analyses, one substituting psilocybin for TPE as the dose covariate in all

outcomes models and another limited to participants who completed a single administration

session (n=75), yielded results consistent with the primary analysis, with no change in the

direction or magnitude of timepoint effects (**Supplemental Tables 2 and 3)**. In a third sensitivity

analysis, results from stratified models by concurrent psychiatric medication use were largely

consistent with those from the full cohort, with significant post-session improvement across all

outcomes observed in both groups (p<0.001) **(Supplemental Table 4)**. Overall, these findings

support the robustness of the primary results.

Post-session adverse events

Participants (n=35, 39.7%) reported symptoms of HPPD at 1-day post-session, and two

(2.3%) indicated these symptoms caused significant distress and were attributed to the

psilocybin session, potentially meeting diagnostic HPPD criteria. Fewer participants (n=16,

18.2%) reported HPPD symptoms at 30-days post-session, with none reporting the symptoms

caused significant distress. Three (3.4%) participants reported lingering negative effects of the

psilocybin experience at 30-days post-session, noting increased feelings of anxiety or fear,

existential struggle, social disconnections, and difficulty coping with family members.

Session dose and exploratory analyses predicting psilocybin session dose

The average dose (TPE) consumed by participants was 27.8 mg (8.2 SD), and was 24.3

mg (8.3 SD) when restricted to psilocybin **(Table 4).** In exploratory analyses examining

predictors of dose (TPE), the only significant predictor was concurrent use of psychiatric

medications, which was associated with greater administered dose (b=3.44, p<0.05,

Supplemental Table 5); these participants received an average dose of 29.6 mg TPE (26.2 mg

TPE for those not taking psychiatric medications). No iPPS subscales were significantly

associated with dose (**Supplemental Table 6**).

Discussion

This prospective, naturalistic study presents the first results examining the impact of

psilocybin on mental health outcomes within Oregon's state-regulated model. We found that at

30-days post-psilocybin session, participants reported significant reductions in depression and

anxiety symptoms and significant improvements in well-being. These benefits were observed in

a community-based population that included individuals with multiple psychiatric diagnoses and

those actively using psychiatric medications, populations often excluded from research of

psilocybin's impact on mental health. Our findings suggest that psilocybin services delivered

within a structured and supportive framework can produce therapeutic benefits. This

translational evidence is needed to inform policy and clinical implementation as jurisdictions

worldwide consider regulated psilocybin programs.

As a therapeutic intervention, psilocybin is particularly effective when combined with

therapeutic support (Carhart-Harris et al., 2016, 2018; Dorczok et al., 2025; Griffiths et al.,

2016), though therapeutic support is not uniformly included in studies of psilocybin's impact on

mental health (Dorczok et al., 2025). The results of this study were achieved under an

enhanced, supportive model, relative to the minimum requirements set forth under Oregon's

Measure 109, through incorporating comprehensive screening, multiple preparation sessions,

and structured integration support with licensed facilitators of which many held additional clinical

health licenses. Such enhancements potentially influenced the overall significant improvements

in mental health outcomes among study participants, including limited adverse events and few

reports of psychedelic harm, which is consistent with prior studies (Evans et al., 2025).

Research also indicates that integrative psychological support and facilitated processing after

psychedelic-assisted sessions increase the likelihood of sustained benefits (Gorman et al.,

2021; Greń et al., 2024).

Study participants demonstrated clinically significant improvements in self-reported

symptoms of depression, anxiety, and well-being. At 30-days post-session, 65% of participants

reported normal well-being despite approximately one-third continuing to report mild to severe

symptoms of depression or anxiety. Because the WHO-5 captures mental and emotional

domains related to psychological flexibility and adaptive functioning (Topp et al., 2015), these

findings may suggest a growing capacity for change and emerging emotional resilience, which

are domains of functioning that individuals may particularly value (Connell et al., 2014). Other

studies of the impacts of psilocybin on well-being likewise reported increased well-being or life

satisfaction post-psilocybin session (Mans et al., 2021; Wiepking et al., 2023).

Consistent with our results, a recent naturalistic study of U.S. veterans attending

psychedelic retreats outside the U.S. documented clinically significant reductions in depression

with the PHQ-9 one-month post-psychedelic session, though not all participants consumed

psilocybin as the psychedelic substance (some received ayahuasca) (Calnan et al., 2025). A

naturalistic study from Germany similarly reported significant improvements in depression

(PHQ-9) and anxiety (GAD-7) symptoms following a psilocybin session, with an approximate 25

mg psilocybin dose from fresh mushrooms, though follow-up timepoints were at 1 week and 3

months (Irrmischer et al., 2025). This study also included participants with multiple psychiatric

diagnosis, which is less commonly studied (Aaronson et al., 2024; Rosenblat et al., 2024). In

our study, nearly two-thirds of participants had mental health diagnoses, many concomitant, and

few experienced post-session adverse events, adding to the extant evidence that psilocybin

services can be safely delivered to populations experiencing a range of mental health conditions

in real-world settings.

In clinical trials, dose is standardized and often fixed at 25 mg of synthetic psilocybin

(Back et al., 2024; Carhart-Harris et al., 2021; Goodwin, Croal, et al., 2023; Goodwin et al.,

2022; Rosenblat et al., 2024), including the recent COMPASS Pathways Phase 3 COMP005

trial (Compass, 2025). The average doses administered in our study (27.8 mg TPE, 24.3 mg

psilocybin) were comparable to trials' standard dose, and outcomes were similar whether

assessed by TPE or psilocybin content alone. The novel dosing protocol described in our study,

which accounts for both psilocybin and psilocin content, provides a useful framework for other

states developing regulated programs utilizing natural-derived mushroom products, and may

inform policy decisions regarding maximum dose limits. While Oregon's legal maximum is 50

mg, this study demonstrated meaningful improvements in outcomes with limited adverse events

at an average dose below 30 mg TPE, suggesting that lower doses may be both effective and

well-tolerated in real-world settings. Additional research is needed to confirm both safety and

efficacy of this dosing practice, and among a greater diversity of populations than was included

in this study.

Individuals taking psychiatric medications are often excluded from clinical trials to reduce

the risk of a psilocybin blunting effect, allow for uniform dosing, and minimize the risk of

serotonin toxicity (MacCallum et al., 2022; Malcolm and Thomas, 2022). Two psilocybin clinical

trials have examined the impacts on participants taking psychiatric medications, and found

reduced depressive symptoms (Goodwin, Croal, et al., 2023), and reduced adverse effects of

psilocybin (Becker et al., 2022). We found that participants using psychiatric medications

consumed modestly higher average doses. Sensitivity analyses restricted to this subgroup

yielded results consistent with the main findings, suggesting concurrent psychiatric medication

use did not substantially alter treatment response. While this may suggest that psilocybin

services using whole-fruit mushroom products may be effective at improving mental health

outcomes without requiring psychiatric medication discontinuation, additional controlled

research examining psychiatric medications and doses, and explicitly testing moderation effects,

is needed (Erritzoe et al., 2024).

Strengths and limitations

This study has several notable strengths. It represents the first systematic evaluation of

mental health outcomes following psilocybin administration within the first regulated framework

in the US, and the first to quantify both psilocybin and psilocin content in administered doses.

This real-world study design provides evidence of effectiveness in naturalistic settings with

individuals with mental health complexities thereby offering a realistic assessment of treatment

benefits that complement traditional efficacy data (McInnes et al., 2024). Rigorous data

collection and participant follow-up yielded near-complete retention, with data from only three

participants excluded. Within-person pre–post assessments reduced between-person

confounding, and the use of widely implemented validated instruments for depression, anxiety,

and well-being supports comparison with prior studies.

Limitations of the study include its small sample size and the predominance of

participants residing in Oregon, impacting its generalizability. However, the study was

adequately powered to detect medium to large effect sizes in change in mental health

symptoms, supporting the robustness of the observed results. As an observational study without

a control group, causality cannot be inferred, and placebo effects, self-selection bias,

expectancy bias, and unmeasured confounding factors (such as influential participant lifestyle or

situational changes during the study period) cannot be ruled out (Lewis et al., 2026) Also, the

follow-up period was limited to 30-days post-session; although this duration aligns with many

clinical trials and observational studies (Calnan et al., 2025; Davis et al., 2021; Griffiths et al.,

2016; Rosenblat et al., 2024), longer-term effects remain to be determined and should be

investigated in future studies conducted within Oregon's regulated model (Korthuis et al., 2024).

Conclusion

In this prospective, naturalistic study, we found that carefully screened participants who

underwent a supervised psilocybin session within Oregon's regulated framework and consumed

naturally-derived mushroom products experienced significant improvements in depression,

anxiety, and overall well-being with few adverse events, even among those with concurrent use

of psychiatric medications and complex mental health histories. These findings demonstrate that

psilocybin can be delivered safely and effectively in real-world settings, broadening the potential

accessibility of psilocybin services as a tool to bridge clinical and community care. For Oregon,

these data strengthen confidence in the state's pioneering regulatory framework, offering data-

driven guidance for establishing standardized practices in supportive psilocybin service

frameworks, while advancing policy goals to safely expand this emerging mental health field.

Acknowledgements: We thank the participants for their time, trust, and willingness to share their experiences. We also acknowledge the valuable contributions of the Bendable Therapy team members who supported data collection: Brooke Harralson, River Jenkins, Sam Georgi, and Matt Hailey. These individuals did not receive additional compensation for these contributions.

Author Contributions: A.G., R.R. J.J.Q, L.A.A., R.C.H, and J.N.D. designed the study. A.G., R.R., C.M., and J.N.D. collected the data. E.S. conducted data analysis. A.G., E.S., R.R. J.J.Q., and J.N.D. interpreted the data. A.G., E.S., R.R., and J.N.D. wrote the initial draft of the manuscript. All authors critically reviewed the manuscript and approved the final draft. J.N.D. is accountable for all aspects of the study and manuscript.

Statements and declarations

Ethical Considerations: The WCG Institutional Review Board approved our study (approval confirmation: 45583966) on March 15, 2024.

Consent to Participate: Participants gave written consent for review and signature before completing any study components.

Declaration of Conflicting Interest: A.G reported receipt of honoraria for presentations at University of Louisiana, Monroe. R.R. reported ownership of Aboveground Services and receipt of honoraria for consulting with Healing Advocacy Fund and for presentations with Soundmind. J.J.Q. reported owning shares in Osmind. C.M reported employment with Aboveground Services. R.C.H. reported advisor roles for Tryp Therapeutics, Osmind, Red Light Holland, and Otsuka, and receipt of honoraria for lectures, presentations, or educational events for HMP Global, Aspen Institute, Psychedelic Science IRAS, Heartmind, Clearmind, and Psychedelic Coaching Institute. No other disclosures were reported.

Funding Statement: The authors disclosed receipt of the following financial support for the research, authorship, and/or publication for this article: This work was financially supported in part by the Jurvetson Foundation. The funder had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Data Availability: The datasets generated during and/or analyzed during the current study are available from the corresponding author upon reasonable request.

References

Aaronson ST, van der Vaart A, Miller T, et al. (2024) Single-Dose Synthetic Psilocybin With Psychotherapy for Treatment-Resistant Bipolar Type II Major Depressive Episodes: A Nonrandomized Open-Label Trial. *JAMA Psychiatry* 81(6): 555–562.

Angyus M, Osborn S, Haijen E, et al. (2024) Validation of the imperial psychedelic predictor scale. *Psychological Medicine* 54(12): 3539–3547.

Back AL, Freeman-Young TK, Morgan L, et al. (2024) Psilocybin Therapy for Clinicians With Symptoms of Depression From Frontline Care During the COVID-19 Pandemic: A Randomized Clinical Trial. *JAMA Network Open* 7(12): e2449026.

Becker AM, Holze F, Grandinetti T, et al. (2022) Acute Effects of Psilocybin After Escitalopram or Placebo Pretreatment in a Randomized, Double-Blind, Placebo-Controlled, Crossover Study in Healthy Subjects. *Clinical Pharmacology and Therapeutics* 111(4): 886–895.

Calnan M, Blest-Hopley G, Busch C, et al. (2025) Exploring the Therapeutic Effects of Psychedelics Administered to Military Veterans in Naturalistic Retreat Settings. *Brain and Behavior* 15(7): e70660.

Carhart-Harris R, Giribaldi B, Watts R, et al. (2021) Trial of Psilocybin versus Escitalopram for Depression. *New England Journal of Medicine* 384(15). Massachusetts Medical Society: 1402–1411.

Carhart-Harris RL, Bolstridge M, Rucker J, et al. (2016) Psilocybin with psychological support for treatment-resistant depression: an open-label feasibility study. *The Lancet Psychiatry* 3(7): 619–627.

Carhart-Harris RL, Bolstridge M, Day CMJ, et al. (2018) Psilocybin with psychological support for treatment-resistant depression: six-month follow-up. *Psychopharmacology* 235(2): 399–408.

Cohen J (1992) Quantitative Methods in Psychology: A power primer. *Psychological Bulletin* 112(1): 155–159.

Compass (2025) Compass Pathways Successfully Achieves Primary Endpoint in First Phase 3 Trial Evaluating COMP360 Psilocybin for Treatment-Resistant Depression. Available at: https://ir.compasspathways.com/News--Events-/news/news-details/2025/Compass-Pathways-Successfully-Achieves-Primary-Endpoint-in-First-Phase-3-Trial-Evaluating-COMP360-Psilocybin-for-Treatment-Resistant-Depression/default.aspx (accessed 12 October 2025).

Connell J, O'Cathain A and Brazier J (2014) Measuring quality of life in mental health: Are we asking the right questions? *Social Science & Medicine* 120: 12–20.

Davis AK, Barrett FS, May DG, et al. (2021) Effects of Psilocybin-Assisted Therapy on Major Depressive Disorder: A Randomized Clinical Trial. *JAMA Psychiatry* 78(5): 481–489.

de la Salle S, Kettner H, Thibault Lévesque J, et al. (2024) Longitudinal experiences of Canadians receiving compassionate access to psilocybin-assisted psychotherapy. *Scientific Reports* 14(1). Nature Publishing Group: 16524.

Dorczok MC, Mittmann G, Ettl T, et al. (2025) Psilocybin-assisted psychotherapy in adults with depression – A literature review. *Progress in Neuro-Psychopharmacology and Biological Psychiatry* 142: 111508.

Erritzoe D, Barba T, Spriggs MJ, et al. (2024) Effects of discontinuation of serotonergic antidepressants prior to psilocybin therapy versus escitalopram for major depression. *Journal of Psychopharmacology (Oxford, England)* 38(5): 458–470.

Evans J, Aixalà M, Anderson BT, et al. (2025) On Minimizing Risk and Harm in the Use of Psychedelics. *Psychiatric Research and Clinical Practice* 7(1): 4–8.

Goodwin GM, Aaronson ST, Alvarez O, et al. (2022) Single-Dose Psilocybin for a Treatment-Resistant Episode of Major Depression. *New England Journal of Medicine* 387(18). Massachusetts Medical Society: 1637–1648.

Goodwin GM, Croal M, Feifel D, et al. (2023) Psilocybin for treatment resistant depression in patients taking a concomitant SSRI medication. *Neuropsychopharmacology* 48(10): 1492–1499.

Goodwin GM, Aaronson ST, Alvarez O, et al. (2023) Single-dose psilocybin for a treatment-resistant episode of major depression: Impact on patient-reported depression severity, anxiety, function, and quality of life. *Journal of Affective Disorders* 327: 120–127.

Gorman I, Nielson EM, Molinar A, et al. (2021) Psychedelic Harm Reduction and Integration: A Transtheoretical Model for Clinical Practice. *Frontiers in Psychology* 12. Frontiers.

Greń J, Gorman I, Ruban A, et al. (2024) Call for evidence-based psychedelic integration. *Experimental and Clinical Psychopharmacology* 32(2): 129–135.

Griffiths RR, Johnson MW, Carducci MA, et al. (2016) Psilocybin produces substantial and sustained decreases in depression and anxiety in patients with life-threatening cancer: A randomized double-blind trial. *Journal of Psychopharmacology* 30(12). SAGE Publications Ltd STM: 1181–1197.

Irrmischer M, Puxty D, Yıldırım BO, et al. (2025) Moderating factors in psilocybin-assisted treatment affecting mood and personality: A naturalistic, open-label investigation. *Psychopharmacology* 242(4): 725–740.

Jensen ME, Stenbæk DS, Messell CD, et al. (2025) Single-dose psilocybin therapy for alcohol use disorder: Pharmacokinetics, feasibility, safety and efficacy in an open-label study. *Journal of Psychopharmacology* 39(5). SAGE Publications Ltd STM: 463–473.

Korthuis PT, Hoffman K, Wilson-Poe AR, et al. (2024) Developing the Open Psychedelic Evaluation Nexus consensus measures for assessment of supervised psilocybin services: An e-Delphi study. *Journal of Psychopharmacology* 38(8). SAGE Publications Ltd STM: 761–768.

Kroenke K, Spitzer RL and Williams JBW (2001) The PHQ-9. *Journal of General Internal Medicine* 16(9): 606–613.

Lewis BR, Reid MJ, Novick AM, et al. (2026) Challenges with clinical trial participants in studies with classical psychedelics: A position statement from the National Network of Depression Centers' task group on psychedelics and related compounds. *Journal of Psychopharmacology*. SAGE Publications Ltd STM: 02698811251413490.

MacCallum CA, Lo LA, Pistawka CA, et al. (2022) Therapeutic use of psilocybin: Practical considerations for dosing and administration. *Frontiers in Psychiatry* 13: 1040217.

Malcolm B and Thomas K (2022) Serotonin toxicity of serotonergic psychedelics. *Psychopharmacology* 239(6): 1881–1891.

Mans K, Kettner H, Erritzoe D, et al. (2021) Sustained, Multifaceted Improvements in Mental Well-Being Following Psychedelic Experiences in a Prospective Opportunity Sample. *Frontiers in Psychiatry* 12. Frontiers.

McInnes LA, Marton TF and Qian JJ (2024) Embracing pragmatism for ketamine insurance coverage: Leveraging real-world evidence. *Journal of Affective Disorders* 352: 199–200.

Morton E, Sakai K, Ashtari A, et al. (2023) Risks and benefits of psilocybin use in people with bipolar disorder: An international web-based survey on experiences of 'magic mushroom' consumption. *Journal of Psychopharmacology* 37(1). SAGE Publications Ltd STM: 49–60.

Nayak SM, Gukasyan N, Barrett FS, et al. (2021) Classic Psychedelic Coadministration with Lithium, but Not Lamotrigine, is Associated with Seizures: An Analysis of Online Psychedelic Experience Reports. *Pharmacopsychiatry* 54. Georg Thieme Verlag KG: 240–245.

Nygart VA, Pommerencke LM, Haijen E, et al. (2022) Antidepressant effects of a psychedelic experience in a large prospective naturalistic sample. *Journal of Psychopharmacology (Oxford, England)* 36(8): 932–942.

Oregon Health Authority (2024) Public Health Division-Chapter 333, Division 333 Psilocybin, Client Information Form. Available at: https://secure.sos.state.or.us/oard/viewSingleRule.action?ruleVrsnRsn=319978 (accessed 11 October 2025).

Oregon Health Authority (n.d.) Oregon Psilocybin Services. Available at: https://www.oregon.gov/oha/PH/PREVENTIONWELLNESS/Pages/Oregon-Psilocybin-Services.aspx (accessed 11 October 2025).

Oregon Psilocybin Services (2025) Oregon Psilocybin Services: Health and Safety Fact Sheet. Oregon Health Authority. Available at: https://www.oregon.gov/oha/PH/PREVENTIONWELLNESS/Documents/OPS-Health-and-Safety.pdf.

Oregon Psilocybin Services (n.d.) Oregon Psilocybin Services (OPS) Psilocybin Product Potency Information. Oregon Health Authority. Available at:

https://www.oregon.gov/oha/PH/PREVENTIONWELLNESS/Documents/OPS-Psilocybin-Product-Potency-Information.pdf.

Passie T, Seifert J, Schneider U, et al. (2002) The pharmacology of psilocybin. *Addiction Biology* 7(4): 357–364.

Raison CL, Sanacora G, Woolley J, et al. (2023) Single-Dose Psilocybin Treatment for Major Depressive Disorder: A Randomized Clinical Trial. *JAMA* 330(9): 843–853.

Rosenblat JD, Meshkat S, Doyle Z, et al. (2024) Psilocybin-assisted psychotherapy for treatment resistant depression: A randomized clinical trial evaluating repeated doses of psilocybin. *Med* 5(3): 190-200.e5.

Ross S, Bossis A, Guss J, et al. (2016) Rapid and sustained symptom reduction following psilocybin treatment for anxiety and depression in patients with life-threatening cancer: a randomized controlled trial. *Journal of Psychopharmacology* 30(12). SAGE Publications Ltd STM: 1165–1180.

Schlag AK, Aday J, Salam I, et al. (2022) Adverse effects of psychedelics: From anecdotes and misinformation to systematic science. *Journal of Psychopharmacology* 36(3). SAGE Publications Ltd STM: 258–272.

Shin C, Lee S-H, Han K-M, et al. (2019) Comparison of the Usefulness of the PHQ-8 and PHQ-9 for Screening for Major Depressive Disorder: Analysis of Psychiatric Outpatient Data. *Psychiatry Investigation* 16(4): 300–305.

Spitzer RL, Kroenke K, Williams JBW, et al. (2006) A brief measure for assessing generalized anxiety disorder: the GAD-7. *Archives of Internal Medicine* 166(10): 1092–1097.

Tap SC, Thomas K, Páleníček T, et al. (2025) Concomitant use of antidepressants and classic psychedelics: A scoping review. *Journal of Psychopharmacology*. SAGE Publications Ltd STM: 02698811251368360.

Topp CW, Østergaard SD, Søndergaard S, et al. (2015) The WHO-5 Well-Being Index: a systematic review of the literature. *Psychotherapy and Psychosomatics* 84(3): 167–176.

von Rotz R, Schindowski EM, Jungwirth J, et al. (2023) Single-dose psilocybin-assisted therapy in major depressive disorder: a placebo-controlled, double-blind, randomised clinical trial. *eClinicalMedicine* 56: 101809.

Wiepking L, Bruin E de and Ghiță A (2023) The potential of psilocybin use to enhance well-being in healthy individuals – A scoping review. *Journal of Psychedelic Studies* 7(3): 184–199.

Zhou K, de Wied D, Carhart-Harris RL, et al. (2025) Prediction of hallucinogen persisting perception disorder and thought disturbance symptoms following psychedelic use. *PNAS Nexus* 4(4): pgae560.

Tables

Table 1: Participant demographic, mental health, and psychedelic experience characteristics[a]

Demographic Characteristics	Session Participants (n=88)
Age, years (median, range)	43 (22, 79)
Birth sex, No. (%)	
Male	46 (52.0)
Female	41 (46.6)
Gender, No. (%)	
Male	44 (50.0)
Female	40 (45.5)
LGBTQIA+	3 (3.4)
Race[b] , No. (%)	
White	77 (87.5)
American Indian or Alaska Native	1 (1.1)
Mixed/multiple races	3 (3.4)
Ethnicity, No. (%)	
Non-Hispanic/Non-Latinx	77 (87.5)
Hispanic-Latinx	5 (5.7)
Level of education completed, No. (%)	
Graduate or professional degree	31 (35.2)
College or baccalaureate degree	32 (36.4)
Associate's or technical degree	11 (12.5)
High school or some college	13 (14.8)
Employment, No. (%)	
Employed (full-time, part-time, self-employed, or sporadically)	59 (67.1)
Retired	17 (19.3)
Unemployed/on disability	9 (10.2)
Student	1 (1.1)

Annual household income, No. (%)	
≥ $100,000	30 (34.1)
$50,000-$99,999	25 (28.4)
$10,000-$49,000	18 (20.4)
< $10,000	2 (2.3)
Decline to Answer	13 (14.8)
Residence, No. (%)	
Living in Oregon	47 (53.4)
Living in other U.S. state	40 (45.5)
Living internationally	1 (1.1)
Military veteran, No. (%)	14 (15.9)
Mental Health Characteristics	
Reported mental health diagnosis, No. (%)	
Depression	45 (51.1)
Anxiety	37 (42.0)
PTSD	17 (19.3)
Acute Stress	6 (6.8)
OCD	4 (4.5)
Eating disorder	2 (2.3)
Bipolar disorder[c]	1 (1.1)
Other	6 (6.8)
No mental health diagnosis	32 (36.4)
Currently taking psychiatric medications, No. (%)	41 (46.6)
Currently in therapy, No. (%)	50 (56.8)
Psychedelic Characteristics	
Prior psychedelic experience, No. (%)	57 (64.8)
Barriers to seeking psilocybin services, No. (%)	
Cost	41 (46.6)

Availability	16 (18.2)
Time and schedule	21 (23.9)
Limited knowledge and experience of psychedelics	16 (18.2)
Current support systems in place (or not in place)	6 (6.8)
Legal concerns	4 (4.5)
Judgement from family and peers	4 (4.5)
Current mental health concerns	4 (4.5)
No concerns	28 (31.8)
Other	1 (1.1)

[a] Some categories of characteristics may not add to 100% as some participants declined to answer demographic questions and data are missing. All data is from the screening application, which gathered data at baseline (prior to a psilocybin administration session). Abbreviations: LGBTQIA+, Lesbian, Gay, Bisexual, Transgender, Queer, Intersex, Asexual, all other identities not explicitly mentioned; OCD, obsessive compulsive disorder; PTSD, post-traumatic stress disorder.

[b] Race variable was a "select all that apply" field. Participants who selected more than one race were grouped into a "Mixed / Multiple race" option and custom responses that didn't match the standard categories were grouped into "Other".

[c] Participants who listed bipolar disorder as a mental health diagnosis in their screening application were subsequently screened by Bendable. Those who never had a formal diagnosis of bipolar disorder and were otherwise eligible were allowed to continue to a psilocybin session.

Table 2: Participant classification by PHQ-8, GAD-7, and WHO-5 categories pre- and post-psilocybin session[a]

Participant Classification Category	Pre-Session (n=88)	30-days Post-Session (n=88)
PHQ-8 (depression), No. (%)		
Minimal	27 (30.7)	54 (61.4)
Mild-Moderate	45 (51.1)	31 (35.2)
Moderately Severe-Severe	16 (18.1)	3 (3.4)
GAD-7 (anxiety), No. (%)		
Minimal	29 (33.0)	62 (70.5)
Mild-Moderate	44 (50)	25 (28.4)
Severe	15 (17.0)	1 (1.1)
WHO-5 (well-being), No. (%)		
Normal	6 (6.8)	57 (64.8)
Borderline	0 (0)	9 (10.2)
Low-very low	82 (93.2)	22 (25.0)

[a] Participants completed online surveys that contained the 3 mental health questionnaires (PHQ-8, GAD-7, WHO-5) at two timepoints: pre-session and approximately 30-days post-psilocybin administration session. Pre-session surveys were completed, on average, 75 days prior to the psilocybin administration session; 30-day post-session surveys were completed, on average, 34 days after the psilocybin administration session. Abbreviations: GAD-7, Generalized Anxiety Disorder-7; PHQ-8, Patient Health Questionnaire-8; WHO-5, World Health Organization-5 Well-Being index.

Table 3: Changes in PHQ-8, GAD-7, and WHO-5 scores from pre-session to 30-days post-psilocybin session (n=88 participants)[a]

Measure	Pre-session M (SE)	30-days post-session M (SE)	Change pre-session to post-session M (SE) [95% CI]	p-value	t(df)	Cohen's d [95% CI]
Unadjusted						
PHQ-8	9.15 (0.56)	4.52 (0.56)	4.63 (0.57) [3.48, 5.77]	<0.001	8.06 (87)	0.89 [0.67, 1.11]
GAD-7	8.22 (0.51)	3.36 (0.51)	4.85 (0.57) [3.71, 5.99]	<0.001	8.48 (87)	1.02 [0.78, 1.26]
WHO-5	4.81 (0.53)	15.5 (0.53)	-10.67 (0.59) [-11.85, -9.49]	<0.001	-17.98 (87)	-2.15 [-2.39, -1.92]
Adjusted						
PHQ-8	11.07 (1.54)	6.44 (1.54)	4.63 (0.57) [3.48, 5.77]	<0.001	8.06 (87)	0.90 [0.68, 1.12]
GAD-7	9.88 (1.35)	5.03 (1.35)	4.85 (0.57) [3.71, 5.99]	<0.001	8.48 (87)	1.04 [0.80, 1.17]
WHO-5	4.3 (1.46)	15.0 (1.46)	-10.67 (0.59) [-11.85, -9.49]	<0.001	-17.98 (87)	-2.14 [-2.38, -1.90]

[a] Results reported as mean (SE). Covariates included birth sex (male/female), age (years), concurrent psychiatric medication use (yes/no), and total psilocybin equivalents (TPE, mg) as a continuous dosage. Only concurrent psychiatric medication use (pre-session) was positively associated with PHQ-8 scores at 30-days post-session (p=0.040). No covariates were significant in the models with GAD-7 and WHO-5. Cohen's d interpretations: small d=0.20-0.49; moderate d=0.50-0.79; large d≥0.80; very large d≥1.2. Abbreviations: GAD-7, Generalized Anxiety Disorder-7 questionnaire; PHQ-8, Patient Health Questionnaire-8; WHO-5, World Health Organization Well-Being Index.

Table 4: TPE and psilocybin doses consumed by participants during a legal psilocybin session in Oregon[a]

Dose Type	Dose, mg
TPE dose, mean (SD)	27.8 (8.2)
TPE dose, median (min, max)	29.9 (13.9, 50.2)
Psilocybin dose, mean (SD)	24.3 (8.3)
Psilocybin dose, median (min, max)	22.0 (9.1, 44.6)

[a] TPE, Total Psilocybin Equivalents, is defined as psilocybin (mg) + 1.39 × psilocin (mg). This metric assumes complete conversion of psilocybin to psilocin based on molecular weight.

Figure 1 Title. Study components within the framework of a legal psilocybin services program in Oregon

Figure 1 Footnote: Study components were integrated into an existing legal psilocybin services program in Oregon, Bendable Therapy. Oregon regulatory minimums require one facilitator-client preparation session 24-hours in advance of a psilocybin administration session and to offer an optional integration session post-administration. The study site psilocybin program included additional comprehensive screening, multiple preparation sessions, and structured integration support sessions with licensed facilitators of which many held additional clinical health licenses. Abbreviations: GAD-7, Generalized Anxiety Disorder-7 questionnaire; HPPD, Hallucinogen Persisting Perception Disorder; PHQ-8, Patient Health Questionnaire-8; iPPS, Imperial Psychedelic Predictor Scale; WHO-5, World Health Organization Well-Being-5 Index.

Figure 2 Title: PHQ-8, GAD-7, and WHO-5 estimated marginal mean scores before and 30-days following a legal psilocybin session in Oregon

Figure 2 Footnote: Scores from the PHQ-8 (depression), GAD-7 (anxiety), and WHO-5 (well-being) questionnaires are represented as estimated marginal means with error bars indicating standard errors. Adjusted linear mixed effects models evaluated change in outcomes, with all models significant at p<0.001. Score data were collected from surveys emailed to participants prior to a psilocybin session (completed on average 75 days before the session) and approximately 30-days after the psilocybin session (completed on average 34 days post-session). Abbreviations: GAD-7, Generalized Anxiety Disorder-7; PHQ-8, Patient Health Questionnaire-8; WHO-5, World Health Organization Well-Being Index-5.