Time, talk, and teamwork: Perceptions of personalised dementia care planning conversations in primary care

AUTHORS:

Sarah Griffiths[1], Danielle Wyman[2], Mike Clark[3], Greta Rait[1], Nathan Davies[2]

AFFILIATIONS:

1. Research Department of Primary Care and Population Health, University College London,

   Upper third floor, Royal Free Campus, London, NW3 2PF, UK

2. Centre for Psychiatry and Mental health, Wolfson Institute of Population Health, Faculty of Medicine and Dentistry, Queen Mary University of London, Yvonne Carter Building, London E1 2AB

3. Care Policy & Evaluation Centre, London School of Economics and Political Science

   Houghton Street, London WC2A 2AE

CORRESPONDING AUTHOR:

Sarah Griffiths, Email s.a.griffiths@ucl.ac.uk

ABSTRACT

Background

Dementia affects over 57 million people worldwide. UK and international policy position personalised, conversation-based care planning as central to post-diagnostic support. However, delivery in primary care is inconsistent, and many practitioners lack dementia-specific communication training. Existing evidence focuses on single roles or settings, leaving a gap in understanding how communication operates across the primary care workforce.

Aims

To identify what helps and hinders effective communication for integrated dementia care planning and determine the support and training needs of the wider primary care workforce.

Methods

- Semi-structured interviews - 11 people with dementia, 13 family carers, and 19 primary care practitioners from diverse roles, exploring experiences of care planning conversations
- Reflexive thematic analysis

Results

Three themes were developed, progressing from micro-level communication practices (Theme 1: Beyond the tick-box), through triadic dynamics (Theme 2: Balancing voices in the conversation), to organisational influences (Theme 3: From silos to meaningful shared care planning). Time and Conversation as intervention cut across all themes, shaping trust and disclosure. Participants reported reliance on tick box approaches, inconsistent preparation, and uncertainty about care plan purpose and ownership. Non-clinical roles were commonly viewed as well placed to support meaningful conversations, but were often described as constrained by unclear remit and weak integration.

Conclusions

A persistent gap remains between policy ambitions and everyday practice. Time-pressured, checklist-driven encounters and fragmented systems undermine shared decision-making. The expanded primary care workforce offers untapped potential to address these gaps, but this requires clearer roles, formal integration, and targeted investment in communicative skills.

BACKGROUND

Dementia affects around 57 million people worldwide, with nearly 10 million new cases each year [1]. As needs become increasingly complex, holistic care planning is essential for maintaining quality of life, autonomy, and dignity. The World Alzheimer's Report 2022 [2] and NHS England [3] position personalised care and support planning as a core element of post-diagnostic dementia care, based on conversations about what matters most. People with dementia should have a personalised care and support plan, outlining their needs and how they will be addressed, reviewed annually and shared across services [3].

Key principles of personalised care planning include prioritising conversation over form-filling, ensuring adequate preparation, and supporting partnership and shared decision-making [4]. Recent UK policy reinforces this emphasis, calling for co-produced, holistic care plans alongside a shift to integrated, community-based services and shared digital access to care plans [5].

Despite these ambitions, routine dementia care planning in primary care is often inconsistent. The Quality and Outcomes Framework (QOF) annual dementia review [6] is a potential general practice mechanism for personalised care planning, yet evidence indicates wide variation in the quality and delivery [7, 8]. Policy therefore encourages wider team

involvement, including roles such as social prescribers, care coordinators, and dementia

advisors [3]. While these roles provide personalised, strengths-based support, they often

operate separately from QOF reviews, limiting continuity and shared ownership. In England,

the Additional Roles Reimbursement Scheme (ARRS), funded by NHS England, has

expanded the primary care workforce by more than 26,000 staff [9], many of which could

support dementia care planning. However, implementation is uneven, with reported gaps in

preparation, retention, and team integration [9].

Communication in dementia care is complicated by cognitive, linguistic, and motor speech

changes [10], and many practitioners report low confidence when communicating with

people with dementia [11]. While general personalised care training exists [12,13], there is

little dementia-specific guidance, creating a gap between policy expectations and available

support.

Existing research offers useful insights into the communication experiences and training

needs of practitioners delivering personalised dementia care, but the evidence base remains

limited. A recent review [14] found little evidence on communication strategies for

personalised dementia care in primary care, with most studies focusing on advanced

dementia in care homes. Studies of social prescribers show reliance on carers for

communication and a tendency to consult carers before engaging directly with people with

dementia [15], while research on dementia support workers highlights challenges in

delivering remote personalised care, including difficulties using visual communication tools

[16]. Although valuable, these studies focus on single practitioner groups or specific delivery

modes, and do not capture the range of communication experiences across the expanding

primary care dementia workforce or routine face-to-face care planning. This gap is significant

given that personalised care planning depends on practitioners' ability to engage people with

dementia and carers in meaningful, values-based conversations. As dementia care is

increasingly distributed across teams, a broader, multi-perspective understanding of what

supports or hinders effective communication is needed. This study therefore examines

personalised dementia care planning conversations from the perspectives of people with

dementia, carers, and diverse primary care practitioners to inform workforce development.

AIMS

109   This study aimed to: (1) identify what helps and hinders effective communication for

110   integrated dementia care planning, and (2) determine the support and training needs of the

111   wider primary care workforce to facilitate high-quality personalised dementia care planning

112   conversations.

METHODS

114   We conducted a qualitative interview study using reflexive thematic analysis [17]. Qualitative

115   methods are reported in line with COREQ [18].

Ethics

The study received Health Research Authority approval and ethical approval from Wales

REC 4

(24/WA/0167) on 17 June 2024. Capacity to consent among participants with dementia was

determined by an experienced researcher (SG) in line with the Mental Capacity Act [19].

Written or verbal informed consent was obtained from all participants. In line with the

approved procedure, where verbal consent was obtained, this was audio recorded,

witnessed and documented in writing by a researcher. The document and audio file were

stored on a secure university data management system.

Public Involvement

This study was co-developed with a lived experience group [20]. Two people with dementia, one carer, and one former carer collaborated with the team throughout, advising on ethics, recruitment, methods, and materials, and contributing to data interpretation, guided by research on involving people with dementia in analysis [21].

Recruitment

Recruitment took place between 22nd August 2024 and 3rd April 2025. We aimed to recruit up to 12 people with dementia, 12 carers, and 26 primary care practitioners across England. Practitioners were eligible if they worked in or with primary care and were involved in personalised dementia care planning. The sample size was chosen to ensure sufficient 'information power' [22] across groups to address the research questions. Eligibility criteria are shown in Table 1.

Table 1: Eligibility Criteria

Participant Group	Inclusion Criteria	Exclusion Criteria
People with Dementia	• Clinical diagnosis of dementia, confirmed by practitioner at study site • Aged ≥18 • Mild–moderate dementia with capacity to consent • Able to communicate in English • Has engaged with a primary care practitioner for a care and support planning conversation* within the past month (timeframe selected to counteract recall difficulties)	• Under 18 • Lacking capacity to consent • Unable to communicate in English
Carers	• Adult family member or friend providing unpaid care to a participating person with dementia • Able to communicate in English • Willing and able to provide written or verbal informed consent • Has engaged with a primary care practitioner for care and support planning with the person with dementia within the past month	• Under 18 • Carers with a cognitive impairment • Paid carers not related to the person with dementia
Primary Care Practitioners	• Practitioners involved in personalised dementia care planning in primary care* (e.g., GP, practice nurse, dementia advisor, social prescriber, specialist dementia nurse, care coordinator) • Able to communicate fluently in English • Willing and able to provide written or verbal informed consent	• Under 18

*This was explained to potential participants as any conversations designed to gain an understanding of the support needs and strengths of a person with dementia +/- their carers.

141   Recruitment sources included Primary Care Networks (groups of general practices sharing

142   resources to deliver integrated care), charities supporting people with dementia, professional

143   networks and social enterprises employing practitioners such as social prescribers and

144   dementia advisors.

Data collection

Semi-structured interviews were conducted in person or via video call, and individually or as

dyads, depending on participant preference and practical considerations. Topic guides

(Supplementary File) were developed by the research team and informed by literature [14,

23], policy [3] and input from the lived experience group. Interviews with people with

dementia and carers explored their experiences of personalised care planning

conversations, including what mattered to them, helpful or unhelpful communication

approaches, preferences, and perceived gaps. Practitioner interviews focused on

experiences of facilitating these conversations, including challenges, strategies, cross-

service integration, and training needs.

Data analysis

Interviews were audio-recorded, transcribed verbatim, and analysed using reflexive thematic

analysis, led by SG. Following Braun and Clarke's guidance [17], analysis moved iteratively

through familiarisation, inductive coding, and the development, review, and refinement of

themes, with repeated transcript readings to ensure deep engagement. Initial codes were

generated systematically across the dataset.

The project adopted an experiential, constructionist stance, treating participants' accounts as

meaningful while acknowledging their contextual shaping. Coding attended to both semantic

and latent meanings. To enhance rigour, DW independently reviewed a subset of transcripts

and coding to support interpretation and reflexive discussion [24]. Coding and theme

development were refined through regular team discussions with all authors and the lived

experience group, whose reflections informed analytic decisions and strengthened the

coherence and credibility of the final themes.

RESULTS

Participants

Forty-three semi-structured interviews were conducted with 11 people with dementia and 13

family carers with varied diagnoses and living arrangements, and 19 practitioners from a

range of roles. Participants took part individually, as dyads, or in one triad (Tables 2–3). The

practitioner sample mostly had under five years' experience, reflecting the relative newness

of many primary care roles (Table 3).

Table 2: Participant characteristics – people with dementia and carers

Participant	ID	Region	Sex	Age	Ethnicity	Type of dementia (Interviewee or cared for person)	IMD quintile	Individual interview/dyad
Person with dementia	DE-01	SE England	F	76-80	White Other	FTD language variant - Primary progressive aphasia	5	Dyad
Carer	CE-01	SE England	M	71-75	White British		Same as PWD	
Person with dementia	DE-02	North England	M	61-65	White British	FTD language variant - Primary progressive aphasia	5	Dyad
Carer	CE-02	North England	F	61-65	White British		Same as PWD	
Person with dementia	DE-03	North England	M	71-75	White Irish	Vascular	1	Individual interview
Person with dementia	DE-04	North England	M	71-75	White British	Alzheimer's	1	Individual interview
Person with dementia	DL-03	SE England	M	81-85	White British	Alzheimer's	4	Dyad
Carer	CL-03	SE England	M	56-60	White British		5	
Person with dementia	DL-06	SE England	M	76-80	White British	Unspecified	4	Dyad
Carer	CL-	SE	F	76-	White British		4	

	06	England		80				
Person with dementia	DL-01	SE England	F	76-80	White Irish	Unspecified	5	Individual interview
Person with dementia	DL-04	SE England	M	86-90	White Other	Mixed	3	Triad
Carer	CL-04	SE England	F	86-90	White Other		Same as PWD	
Carer	CL-05	SE England	F	56-60	White Other		3	
Person with dementia	DE-06	South England	M	61-65	White British	Lewy Body	4	Individual interview
Person with dementia	DD-06	SW England	M	91-95	White British	Vascular	4	Dyad
Carer	CD-06	SW England	F	86-90	White British		Same as PWD	
Person with dementia	DE-05	North England	F	46-50	British Asian – Pakistani	Alzheimer's	1	Individual interview
Carer	CL-01	SE England	M	36-40	British Asian – Bangladeshi	Unspecified	3	Individual interview
Carer	CD-01	SW England	M	71-75	White British	Unspecified	2	Individual interview
Carer	CD-02	SW England	F	76-80	White British	Fronto-temporal behavioural variant	3	Individual interview
Carer	CD-03	SW England	F	81-85	White British	Mixed	2	Individual interview
Carer	CL-02	SE England	F	56-60	White Irish	Mixed	3	Individual interview
Carer	CL-07	SE England	F	51-55	White British	Alzheimer's	1	Individual interview

180   *Table 3: Participant characteristics – Practitioners*

Role	ID	Region	Sex	Ethnicity	Work setting	Years experience	Ethnicity
Social Prescriber	PE-01	Southeast England	F	White Irish	GP surgeries	<5	White Irish
Dementia Advisor	PL-01	Southeast England	M	Black British	Charity	11-20	Black British
Social Prescriber	PE-02	Southwest England	F	White British	GP surgeries	<5	White British
Social Prescriber	PE-03	North of England	F	White British	GP surgeries	<5	White British
Social Prescriber	PD-01	Southwest England	F	White British	Social Enterprise working with GP surgeries	<5	White British
Social Prescriber	PD-02	Southwest England	F	White British	Social Enterprise working with GP surgeries	5-10	White British
GP	PL-02	Southeast England	F	White British	GP surgery	5-10	White British
GP	PL-03	Southeast England	M	Mixed Caribbean	GP surgery	<5	Mixed Caribbean
GP	PL-04	Southeast England	F	White other	GP surgery	<5	White other
Dementia Advisor	PL-05	Southeast England	F	White British	Charity	<5	White British
Admiral Nurse	PE-04	North of England	F	White British	GP surgeries	<5	White British
Dementia Advisor	PL-06	Southeast England	F	White British	Charity	<5	White British
Care co-ordinator	PL-07	Southeast England	F	White British	GP surgery	5-10	White British
Admiral Nurse	PE-05	Southeast England	F	White Other	GP surgeries	<5	White Other
GP	PL-09	Southeast England	F	White British	GP surgery	>20	White British
Dementia Advisor	PL-08	Southeast England	F	White British	Charity	<5	White British

GP	PL-10	Southeast England	F	White British	GP surgery		White British
Trainee GP and Clinical Academic Fellow	PD-05	Southwest England	F	White British	GP surgery	<5	Whtie British
Dementia Advisor	PE-07	North of England	F	British Pakistani	Charity	<5	British Pakistani

181
182

Findings

Three core themes and eight sub-themes were developed (Fig. 1), broadening in scope from

micro-level communication practices (Theme 1: Beyond the tick-box), through the dynamics

of balancing voices (Theme 2: Balancing voices in the conversation), to wider contextual

influences on care planning (Theme 3: From silos to meaningful shared care planning).

'Time' and 'Conversation as intervention' were identified as cross-cutting themes and are

integrated across the findings.

Figure 1: Themes

Theme 1: Beyond the tick box

This theme examines how checklist-driven approaches often limit connection and

personalisation. Participants felt meaningful conversations happened when both sides had

time and preparation, communication adaptations were adopted to support communication

(e.g., visual resources), deepening understanding and supporting genuine partnerships.

More than a checklist: why conversations matter

Most practitioners relied on standard checklists, which practitioners, people with dementia

and carers recognised as impersonal and limited in capturing the complexity and fluctuation

of dementia.

'It was an entire tick box exercise whereby it was merely yes or no, and even if it

was a no, it wouldn't have made any difference, there was no follow-on from the

no.' (PWD: DE-06)

'If you go in with too much structure some people just won't engage…you won't

get the information you need. So, you've got to do it in a…communicative way.'

(Social prescriber: PE-02)

Participants across all groups felt that meaningful conversations required time, a

relaxed pace, and genuine curiosity, with unhurried encounters helping them feel

known, creating space for honest reflection, and enabling deeper discussion through

open-ended questions.

'Questions like, "And you're coping OK?". They're dead ends. Better to ask,

"What's your day like?" or "When did you last do something for yourself?"'

(Carer:CL-02)

They also noted that setting shaped interaction, with home visits offering fuller context,

clinics fewer distractions, and remote consultations often feeling rushed. Flexibility was

valued over any one format.

'When I have done home visits the person is much more relaxed…they're able to

talk about themselves more. On the telephone sometimes people with dementia

just clam up.' (Dementia Advisor:PL-06)

Time enabled people to share their stories, helping them process change, build trust,

and enabling practitioners to better identify strengths and support needs. When asked

what practitioners should ask to understand his needs and strengths, one person with

dementia said:

'My life history -where I was born, who I married, and what happened to them.'

(PWD:DL-03)

Reassurance was a valued outcome of conversation, with affirmation and

encouragement reducing anxiety and guilt for both people with dementia and carers.

'I worry and feel guilty about doing the right thing…She said buy him what he'll

eat at this stage. It was totally reassuring and so lovely.' (Carer:CL-01)

People with dementia and carers felt checklists often missed important areas like their

everyday lives, how well they understand their medications, or planning for contingencies,

reinforcing a sense of being unseen.

'I get asked, have you had any falls? But they could ask - Am I still able to go

shopping? …Am I still able to use the computer?' (PWD:DE-04)

Moving away from checklists towards a conversational approach enabled collaborative

problem-solving, with gentle, scenario-based examples helping to normalise support

and make change feel manageable. Speaking about a social prescriber, one person

with dementia reflected:

'He didn't just say, "Get a carer." He said, "How would you feel about someone

coming in with a cup of tea, to check you're alright?" It made it easier to accept.'

(PWD:DE-05)

Adaptive and non-verbal connection

Participants across all groups emphasised that non-verbal cues shaped whether

people felt safe and understood, with small gestures and a calm presence helping to

build trust.

'She's very friendly… she'll just sort of like…touches you… she really looks at

you.' (Carer:CE-02)

For some people with dementia, emotional tone mattered more than the specific content of

what was said, serving as an immediate indicator of whether an interaction felt safe or

threatening.

'When I'm speaking to somebody, straight away I'm trying to analyse: are they a

threat?' (PWD:DE-03)

Some practitioners described adapting their communication by simplifying language

and using visual or creative strategies to make abstract information more concrete,

support comprehension, and provide reassurance:

'If I'm talking about something new or abstract… I print off details… if there's

photographs of people [in a support group], that might give reassurance.' (Social

Prescriber:PD-01)

Others described wanting more training in these approaches, alongside clearer

permission and remit to use them in practice.

'Training on… pictorial stuff…would [be] really helpful…knowing what we can

use.' (GP:PL-02)

'A creative [resource] with…a map of their life to show also they're still

significant… I'd love to do it but from a commissioners standpoint, they [wouldn't]

think that's your remit. (Social prescriber:PD-01)

Cultural and linguistic bridges

Culture and language shaped whether conversations felt safe and inclusive, with

participants valuing practitioners who showed cultural humility and respect.

'If you were to give them respect by saying 'Salam Alaikum' because you knew

they were Pakistani, you would see that smile.' (Dementia Advisor: PE-07)

Some practitioners avoided potentially sensitive subjects, such as sexuality or gender, for

fear of causing offence.

'…we exclude questions…about being gay, because these kinds of things are

forbidden in Islam.' (Dementia Advisor:PE-07)

While well-intentioned, this could lead to important parts of identity being overlooked,

suggesting the need for respectful and more conversational ways of exploring identity.

Language barriers were a further challenge. Telephone interpreters were often

inaccessible, face-to-face bilingual support inconsistent, and practitioners relied on

creative strategies such as involving relatives to sustain conversation or using drawing

and pictures.

'One lovely guy …was telling me about when he was young…I caught the word

and looked it up, showed him pictures. It was beautiful…like a big castle.' (Social

Prescriber: PE-01)

Practitioners who shared cultural lived experiences with people with dementia were also

seen as powerful bridges, reducing dementia stigma within communities.

Being prepared for meaningful conversations

Preparation was seen as essential for conversation that feels respectful, supportive, and

genuinely personalised. People with dementia and carers valued knowing what to expect

and having time to reflect before reviews, helping them identify priorities and participate

actively rather than reactively.

Practitioners emphasised the importance of preparing for the encounter, by setting

expectations: introducing their role and the purpose of the conversation to avoid confusion

and build trust.

'I explain to them I'm not a nurse or a doctor… I just say, I've come to see how

you are.' (Dementia Advisor:PD-07)

Knowledge of dementia subtypes and trajectories was also viewed as part of being

adequately prepared for personalised care planning, with gaps in understanding

undermining trust.

Carer: *'They all think he's got Alzheimer's cos dementia is just Alzheimer's'*

PWD: *'Yeah, that's true.'* **Carer**: *'It's PPA and they don't understand that. They*

should do but they don't' (Dyad: DE-02; CE02)

Overall, this theme underscores the importance of time and the idea that conversation itself

is an intervention and a foundation for personalised care. Participants valued care-planning

not as a procedural task, but as a flexible, relational process through which reassurance,

shared understanding, and personalised support are built.

Theme 2: Balancing voices in the conversation

Central to this theme is the challenge of ensuring both people with dementia and carers feel

heard during care-planning conversations.

Reliance on carers

Time pressures and inaccessible formats (such as phone calls) often led practitioners to

defer to carers, sidelining people with dementia. Carers felt pressured to speak on their

behalf and worried about misrepresenting their views.

'…on the phone…you're having to be really careful cos it's my interpretation of

what's going on…it's not necessarily correct, is it?' (Carer:CE-02)

Some practitioners described mistrusting carer input in sensitive areas, adding complexity to

triadic care planning, as concerns (e.g., about driving or behaviour) were sometimes

minimised to avoid upsetting the person with dementia or triggering consequences such as

sedatives or care transfers.

Theme one explored how language barriers shape conversations. When family members

support communication, practitioners questioned whose voice was being heard when

relatives declined professional interpretation without consulting the patient:

'Is everything you're saying via the family member being directly translated or is it

their take?... Sometimes we say, "Would you like me to get a telephone

translator?" but instead of that being translated to the patient, the answer is just

no.' (GP:PL-01)

Language barriers not only complicate communication but can also contribute to the

exclusion of person with dementia from care planning conversations, amplifying issues of

voice and representation in care.

Balancing perspectives

Carers often felt rushed or conflicted when their views differed from the person with

dementia's or when appointments left little space for their own needs. Both parties

could struggle with honesty in each other's presence, highlighting the need for sensitive

management, including awareness of family dynamics, separate conversations, and

skills to balance perspectives.

'I say, "Oh, is that definitely what's happening at the moment…it might have been

like in the past, but what about now, and…invite more of a group discussion…

there are times where…it's easier to speak to them separately.' (GP: PL-02)

Carers also wanted recognition that their cared for person's account may not fully

capture the situation, without feeling they must work hard to justify their perspective.

'it's being able to have two conversations… some [practitioners] are skilful at

that. You don't come away thinking "Oh, God, my mother said X, Y, and Z, and

that's totally not the case." You don't know if they've… taken that…as gospel.'

(Carer:CL-02)

Some practitioners encouraged families to discuss issues privately first, supporting

preparation for care planning (Theme one) and enabling later separate conversations.

They emphasised the importance of careful question framing in triadic interactions, to

protect dignity and balance the needs of all parties.

'When a [practitioner] is directing questions at the carer in front of the person

with dementia, they should be sensitive… especially when…delicate, regarding

incontinence. They could frame [as] "What are your worries for your loved one

regarding this area?" (Carer:CL-02)

Practitioner accounts of complex situations highlight the skilled work required for listening to

and acknowledging both perspectives:

'Maybe 10 years ago they said don't ever let me go into a nursing home…they're

now realising they can't cope…listening to them both and advocating for what's

safe and best for the patient can be difficult…Listen to what people are saying

and try to guide them in whichever direction you feel they want to go, and you

think is best for them.' (GP:PD-05)

However, some practitioners described facing challenges when carers decline support

despite signs of strain or risk, and when carers struggled to accept changes in the person

with dementia or interpreted symptoms as deliberate behaviour.

'Sometimes carers… think [PWD is] doing something on purpose… They might

not recognise there's been a change… and this is the reason for it.' (Dementia

Advisor:PL-08)

Including carers as equal partners in care planning therefore involves recognising their level

of understanding, acceptance and emotional strain.

Effective care planning depends on taking time to negotiate how both the person with

dementia and the carer are heard. Conversation itself can be experienced as an intervention

that validates and supports each perspective.

Theme 3: From silos to meaningful shared care planning

This theme shows how aspirations for personalised dementia care are undermined by

persistent fragmentation. *'Lost in the system'* concerns how this fragmentation is felt, through

communication gaps that leave people overwhelmed and uncertain about care planning and

who should be involved. '*Speaking across silos'* examines the communicative mechanisms

through which fragmentation is sustained and shows how communication that bridges

organisational boundaries is fundamental to meaningful, person-centred care planning.

Lost in the system

Siloed working and disjointed systems were described as the norm across the sample. GPs

felt overwhelmed by workload and time pressures, limiting collaboration despite recognising

its value, while non-clinical practitioners reported feeling disconnected and unclear about

roles and how to work together.

'Understanding how all the different services work together would have been

useful … it needs to be much more cohesive and joined up for it to be really

helpful.' (Dementia Advisor:PL-06)

The absence of a named point of contact for patients/carers compounded this sense of

disconnection. Non-clinical team members often assumed coordination roles by default,

trying to bridge systemic gaps and provide reassurance, but without formal recognition or

support.

'I'm finding…the carers in particular struggling so much they're overwhelmed

with contacts. So, I'm the one person who'll be the link to all the other contacts,

they will call me first.' (Social Prescriber:PE-01)

For some, this carried an emotional toll, requiring repeated advocacy within an unresponsive

system.

'It's a frustration. You just have to keep poking the bear until somebody actually

listens …sometimes it may be too late…the patient may have had a fall…it's

really a stopping of…an ultimate horrific end.' (Care Co-ordinator: PL-07)

For people with dementia and carers, the consequences of fragmentation were evident in

everyday encounters. They described interacting with multiple practitioners without knowing

who they were or how they were connected, creating a sense of being processed rather than

held within a coherent care network.

*'**Carer:** We had this guy phone from…What did they call him? He was from the*

clinic…but he wasn't a doc-…Not a therapist, even… He phoned twice and had

*a chat with you and then discharged you. **Person with dementia:** It didn't do*

anything. It was just filling up the NHS "Oh, now this is somebody else that

you'll be speaking to."' (Dyad:DL-06 and CL-06)

There was a general conceptual confusion over care plans across all participants. Many

people with dementia and carers did not recall ever having seen a care plan or participated

in a dementia review. Care plans were rarely referred to or explained in accessible terms.

'I wonder if it was that lady that came here…she might have been doing the care

plan…I think they need to be clearer; written down.' (Carer: CL-04)

Some practitioners were also unclear about the concept of a care plan and whether what

they were doing constituted care planning.

'*I've never heard of anyone talk about that before to be honest.*' (PD-02: Social

Prescriber)

People with dementia called for more collaborative, two-way models that recognise their

ongoing agency; models that, in principle, should already be in place.

'Monitoring should be a two-way engagement…knowing that if we've had issues

we would go back and ask for a meeting together and then it would grow from

there.' (PWD:DE-06)

These accounts show that people with dementia, carers and practitioners are not just

navigating a fragmented system but often feel lost within it, with communication gaps and

unclear roles and processes leaving them overwhelmed, frustrated, and unsupported.

Speaking across silos

Across accounts, Collaboration within primary care and with other services was constrained

by incompatible systems and limited time and space for multidisciplinary dialogue. People

with dementia noticed when practitioners could access their records, avoiding repetitive or

impersonal assessments and improving interaction quality.

'It went well…[they] knew about me so didn't have to go through…repetitive

statements which means…they haven't really read the medical record

beforehand.' (PWD:DE-06)

However, even when practitioners shared access to electronic care records, they lamented

the lack of conversations with each other, describing fragmented, task-oriented exchanges

that limited shared sense-making.

'It doesn't feel joined up in terms of a multidisciplinary team…like an urgent email

referral into older people's mental health team…they look at my notes, and I see

their notes…but there would be no direct communication…we don't have time to

review [electronic] feedback from every service we refer to… that…loop of

communication just physically can't happen.' (GP:PD-05)

As a result, care plans were rarely dynamic or genuinely shared, becoming annual,

template-driven reconstructions disconnected from evolving needs.

Many practitioners described teams working in parallel, with heavy caseloads and little

visibility of each other's work, making it hard to sustain a shared sense of direction.

'… other teams are just as busy…I haven't always got time to check what

they're doing... if we had clear care plans that would be so much better because

we'd all know…the direction of travel.' (Admiral Nurse:PE-05)

Referrals between services were inconsistent and often lacked essential details, placing the

burden of discovery on those picking up new cases.

'We often get limited information…It just says can you speak to this person with

dementia…You pick up the telephone…that's when you start to realise what

you're dealing with.' (Dementia Advisor:PL-06)

Formal structures for shared planning were limited, and information-sharing often depended

on ad hoc communication.

'It's more corridor conversations… "Oh, I've seen so-and-so, they might be

struggling a bit with that."' (Admiral Nurse:PE-05)

Structural silos also shaped how the capacities of non-clinical roles were used or

overlooked. Many described being invisible within the system, with GPs unaware of their

existence even when efforts had been made to introduce their roles.

'Half [the GPs]… don't even know we exist…even though I left leaflets in the

surgery.' (Dementia Advisor:PE-07)

Local variation meant some people with dementia and carers formed strong

relationships with non-clinical practitioners, while others never encountered them. Even

within multi-disciplinary teams (MDTs), these roles were often treated as peripheral,

despite their capacity to build trust and support values-based conversations.

Practitioners expressed a desire for greater connection, clearer expectations, and more

meaningful involvement.

'Social prescribers have time… I'd like us to be more connected to

clinicians…Our job is to make their job better… We sit and talk…most doctors

would love that luxury. If they guide us on what they want… what do you want

us to do? What are you trying to achieve here …getting it nailed what this care

plan is supposed to achieve.' (Social Prescriber: PE-01)

Despite this potential, non-clinical roles often lacked access to records or formal channels to

feed their insights into care planning.

Where co-location, shared technology, or regular MDT meetings existed, practitioners

reported more fluid collaboration, clearer communication, and greater confidence in

shared decisions.

'The GP might ask us to cover a particular topic with somebody…we can go to

the house…sit with them…help them think through scenarios.' (Social

Prescriber:PE-03)

Together, these accounts show that meaningful shared care planning depends on both

structural coordination and time and space for practitioners to talk, reflect, and plan together.

What was often missing was sustained conversation, between teams and with people with

dementia and carers, capable of turning fragmented activity into coherent, person-centred

care. Where enabled, such dialogue can bring clearer direction, greater confidence, and

reduced emotional strain, positioning conversation itself as a therapeutic and integrative

intervention.

500 DISCUSSION

Findings illuminate what helps and hinders effective communication for integrated dementia

care planning and identify the support and training needs of the wider primary care

workforce. There is a persistent gap between policy expectations for personalised care and

everyday practice. Although guidelines prioritise conversation over form-filling [4], care

planning was often experienced as checklist-driven, constraining meaningful dialogue. Given

people with dementia may conceal or downplay difficulties to avoid stigma and preserve

dignity [25], such formats risk obscuring what matters most.

Time operated as a critical enabler of effective communication, allowing unhurried

discussions in which participants could share stories, reflect on change, and build trust.

Conversation itself was experienced as a form of intervention, offering reassurance,

partnership, and emotional support rather than merely gathering information. These time-rich

interactions were often linked to non-clinical roles perceived as having greater relational

capacity, yet their informal and marginalised status limited their impact on integrated care

planning.

Despite being central to personalised care [4], preparation for care planning conversations

was often inadequate, leaving their purpose, scope, and roles unclear and undermining trust.

Participants stressed that meaningful involvement of people with dementia requires adapted

communication to support memory and comprehension, yet many lacked the time, training,

or tools to provide this. Wider workforce roles were often seen as better positioned to offer

such support, but without formal endorsement or training, their potential remains

underdeveloped.

Participants also noted cultural and linguistic dimensions of care planning. Small, culturally

attuned gestures were seen to support rapport, while language barriers and avoidance of

sensitive identity-related topics risked excluding what mattered most.

Practitioners valued a conversational, collaborative approach to care planning but described

structural fragmentation hindering its delivery, including incompatible IT systems, unclear

roles and weak communication pathways. These gaps left non-clinical staff underused, GPs

overstretched, and people with dementia and carers unsupported by a joined-up approach.

Our findings align with previous research positioning personalised dementia care as

relational and shaped by communicative competence and organisational context [14]. The

cross-cutting themes of time and conversation as intervention highlight that communication

is not an adjunct but the foundation of care planning. Yet the organisational conditions

required to sustain this work remain inconsistent and fragile.

Evidence on ARRS role integration highlights uncertainty around remit, communication

pathways, and access to patient information [26,27]. We extend this by showing the

communicative consequences of such ambiguity, including reduced continuity, fewer

opportunities for shared understanding, and limited co-creation of meaningful care plans.

While previous work notes reliance on carers to facilitate communication [11], our findings

show that carers often experience this as a burden, particularly when they feel responsible

for interpreting needs or when inaccessible formats such as phone calls are used. This

aligns with earlier research showing that language barriers, including reliance on telephone

interpreters, can hinder rapport and the communication of empathy [28].

Across all groups, participants were also uncertain about what a 'care plan' is, whether one

existed, and who was responsible for it, reinforcing evidence that care plans are often

invisible in practice [8]. Greater conceptual clarity is therefore foundational to meaningful

personalised planning and shared ownership.

Implications for practice

Policy frameworks [29,30] prioritise personalised conversations, shared decision-making,

and adaptable care plans to prevent unnecessary admissions. However, our findings

suggest the communicative foundations required to realise these ambitions remain weak.

NHS *Fit for the Future* [5] emphasises digital upskilling. While this may reduce administrative

burden, it risks sidelining workforce training in communicative competencies. Despite

guidance from the Personalised Care Institute [12] and the Dementia Training Standards

Framework [31], practitioners reported uncertainty about adapting communication and

managing triadic interactions. Digital transformation should be accompanied by investment

in conversational skills.

Cultural competence must extend beyond awareness to include communicative strategies

that preserve dignity and sustain participation. Clearer shared understanding of care plan

purpose and ownership is also needed. Co-designed, accessible preparatory tools could

support this.

Crucially, effective use of the expanded workforce requires capitalising on untapped

potential. Non-clinical roles were widely perceived as having the time and relational

orientation required for personalised dementia care yet remained structurally underused.

Their integration into multidisciplinary care planning is therefore essential.

Policy implications

566   Policy ambitions for personalised dementia care depend on communicative practices that

567   remain under-resourced. While workforce expansion has been prioritised, integration has

568   lagged behind. Policies should move beyond role creation to formal inclusion, giving non-

569   clinical staff the authority, access, and training to lead care planning. Without this, the

570   relational capacity of the workforce will continue to be underused.

Strengths and limitations

A principal strength of this study is its multi-perspective design, capturing how care planning

conversations are experienced by people with dementia, informal carers, and a diverse

range of practitioners. By foregrounding communicative processes in primary care and

earlier-stage dementia, the study addresses key evidence gaps and advances

understanding of the relational foundations of personalised care. A limitation is the

underrepresentation of marginalised groups, particularly people from minority ethnic

backgrounds and diverse sexual orientations, thereby constraining transferability. Future

research should prioritise underserved populations, use observational methods to examine

real-time care planning, and evaluate system-level innovations such as shared digital care

plans to support communicative continuity.

Conclusion

This study highlights a persistent gap between policy ambitions for personalised dementia care and everyday practice. Time-poor, checklist-driven encounters and fragmented systems undermine the trust-building conversations that enable disclosure and shared decision-making. Closing this gap requires investment in communicative skills alongside structural reform, particularly through integrating the wider primary care workforce and its underused relational capacity.

REFERENCES

1. World Health Organization. *Dementia*. Geneva: World Health Organization; 2023. Available from*: https://www.who.int/news-room/fact-sheets/detail/dementia*

2. Gauthier S et al. World Alzheimer Report 2022: Life after diagnosis: Navigating treatment, care and support. 2022; *Alzheimer's Disease International*: London, England.

3. NHS England. Dementia: good care planning – information for primary care and commissioners; *https://www.england.nhs.uk/publication/dementia-good-care-planninginformation-for-primary-care-and-commissioners*. 2017.

4. Roberts S et al. The Year of Care approach: developing a model and delivery programme for care and support planning in long term conditions within general practice. *BMC Fam Pract* 2019; 20(1): 153.

5. DHSC. Fit for the future: 10 year health plan for England. *NHS England;* 2005.

6. NHS England. Quality and Outcomes Framework guidance for 2025/26 2025.

7. Wheatley, A et al., *The primary care annual dementia review: a qualitative study of the views and experiences of service users and providers.* medRxiv, 2022: p. 2022.04.26.22274255.

8.   Spencer E et al. Access to personalised dementia care planning in primary care: a

mixed methods evaluation of the PriDem intervention. *BMJ open* 2024; 14(7):

e082083.

9.   Bramwell D et al. Implementing the Additional Roles Reimbursement Scheme in

seven English Primary Care Networks: a qualitative study. *Br J Gen Pract* 2024;

74(742):. e323-e329.

10.   Volkmer A. Assessment and therapy for language and cognitive communication

difficulties in dementia and other progressive diseases, ed. J.R. Pres. 2013.

11.   Di Lorito C et al. What are the challenges that social prescribers face when

supporting people within dementia and how can these be addressed? A qualitative

study. PLoS One 2025; 20(1): p. e0317749.

12.   Personalised Care Institute. Available from:

https://www.personalisedcareinstitute.org.uk/.

13.   E-Learning for Healthcare. Social Prescribing: Learning for Link Workers. Available

from: https://www.e-lfh.org.uk/programmes/social-prescribing/.

14.   Sunjaya AM et al. Communication strategies for delivering personalised dementia

care and support: a mixed-methods systematic review and narrative synthesis. *Age*

Ageing 2025; 54(5).

15.   Pick A et al. Social prescribing in primary care for people living with dementia: a

qualitative exploration of different roles and services in England. *BMC Prim Care*

2025; 26(1): 346.

16.   Wheat H et al. Practitioners' ability to remotely develop understanding for

personalised care and support planning: a thematic analysis of multiple data sources

from the feasibility phase of the Dementia Personalised Care Team (D-PACT)

intervention. *Dementia* 2023; 22(7):1461-1486.

17.   Braun V, Clarke V. Reflecting on reflexive thematic analysis. Qualitative Research in

Sport, Exercise and Health 2019; 11(4): 589-597.

18.   Tong A et al. Consolidated criteria for reporting qualitative research (COREQ): a 32-

634   item checklist for interviews and focus groups. *Int J Qual Health Car* 2007; 19: 349-

635   357.

636   19.   Mental Capacity Act. 2005.

637   20.   Griffiths S et al. Reflections on co-production: Developing a dementia research

638   funding application with a diverse lived experience group. *Dementia* 2024; 23(5):

639   709-723.

640   21.   Stevenson M, Taylor BJ. Involving individuals with dementia as co-researchers in

641   analysis of findings from a qualitative study. *Dementia* 2019; 18(2): 701-712.

642   22.   Malterud K et al. Sample Size in Qualitative Interview Studies: Guided by Information

643   Power. *Qual Health Res* 2016; 26(13): 1753-1760.

644   23.   Sharkiya S.H. Quality communication can improve patient-centred health outcomes

645   among older patients: a rapid review. *BMC Health Services Research* 2023; Aug

646   22;23(1):886.

647   24.   Levitt HM et al. Journal article reporting standards for qualitative primary, qualitative

648   meta-analytic, and mixed methods research in psychology: The APA Publications and

649   Communications Board task force report. *American Psychological Association: US*

650   2018; p. 26-46.

651   25.   Kohl G et al. People With Dementia Disclosing Their Diagnosis to Social Networks: A

652   Systematic Review and Meta-Synthesis. *Gerontologist* 2025; 65(3).

653   26.   Jones B et al. Challenges and enablers to implementation of the Additional Roles

654   Reimbursement Scheme in primary care: a qualitative study. *British Journal of*

655   *General Practice* 2024; 74(742):e315-e322.

656   27.   Pescheny J V et al. Facilitators and barriers of implementing and delivering social

657   prescribing services: a systematic review. *BMC Health Serv Res* 2018; 18(1): 86.

658   28.   Burnand A et al. Providing inclusive care and empowering people with dementia as a

659   clinical pharmacist: a qualitative study of clinical pharmacist's experiences. *Aging &*

660   *Mental Health* 2025; 29(8): 1407-1413.

661   29.   NHS England. Comprehensive model of personalised care. Available from:

662   *https://www.england.nhs.uk/personalisedcare/comprehensive-model-of-personalised-*

663   *care/.*

664   30.   NHS England. Neighbourhood Health Guidelines. Available from:

665   *https://www.england.nhs.uk/long-read/neighbourhood-health-guidelines-2025-26/.*

666   31.   NHS England. Dementia training standards framework. Available from:

667   *https://www.hee.nhs.uk/our-work/dementia-awareness/core-skills.*