Explainable advanced electrocardiography predicts coronary artery disease on coronary computed tomography angiography

Manoj Rajamohan MBBS[1], Daniel E Loewenstein MD[1,2], Maren Maanja MD PhD[2], Zaidon Al-Falahi MBBS[1], Abhinayeni Kuhasri[1,3], Kevin Yang MBBS[1,3], Chosita Cheepvasarach MBBS[4], Thomas Lindow MD PhD[2,5], Todd T Schlegel MD[2,6],

Yalu Wen PhD[7], Patrick A Gladding MBBS PhD[7,8],

Martin Ugander MD PhD[1,2,4], Rebecca Kozor MBBS PhD[1,4]

1. Kolling Institute, Faculty of Medicine and Health, University of Sydney, Sydney, Australia

2. Department of Clinical Physiology, Karolinska University Hospital, and Karolinska Institutet, Stockholm, Sweden

3. University of New South Wales, Sydney, Australia

4. Royal North Shore Hospital, Sydney, Australia

5. Department of Respiratory Medicine, Allergology, and Palliative Medicine, Clinical Sciences, Lund University, Lund, Sweden

6. Nicollier-Schlegel SARL, Trélex, Switzerland

7. University of Auckland, Auckland, New Zealand

8. North Shore Hospital, Waitemata District Health Board, Auckland, New Zealand

Corresponding author:
Dr Rebecca Kozor

Royal North Shore Hospital

Reserve Road

St Leonards, NSW, 2065

Australia

Email: rebecca.kozor@sydney.edu.au

ABSTRACT

BACKGROUND: Conventional electrocardiography (ECG) has limited diagnostic accuracy for detecting coronary artery disease (CAD) in patients with stable chest pain. Advanced electrocardiography (A-ECG) may improve diagnostic performance. The study aimed to derive, externally validate, and prognostically validate an explainable A-ECG score for detecting CAD on coronary computed tomography angiography (CCTA).

METHODS: Participants attending an outpatient rapid access chest pain clinic (RACC) underwent a standard 12-lead ECG and CCTA. Any CAD was defined as any calcified or non-calcified plaque. Elastic net with nested resampling was used to derive an A-ECG score using measures from the conventional ECG, derived vectorcardiography, and measures of waveform complexity. **RESULTS:** In the derivation cohort (n=171, age 59±13 years, 60% male), n=99 (58%) had any CAD on CCTA. A seven parameter A-ECG score to detect any CAD was derived. In an external validation cohort (n=762, age 57±12 years, 49% male, n=428 (56%) with any CAD), the score had an area under the receiver operating characteristic curve [95% confidence interval] of 0.72 [0.68–0.76], sensitivity 90 [88–92]%, specificity 31 [29-33]%, positive predictive value 63 [61–64]%, negative predictive value 70 [68–71]%, positive likelihood ratio 1.3 and inverse negative likelihood ratio 3.1. In the UK Biobank (n=27,239, 966 events, follow-up 1.9 [0.7–4.4] years, age 66±8 years, 50% female), higher A-ECG scores were associated with cardiovascular events after adjusting for age, sex and cardiovascular risk factors (p<0.001). **CONCLUSIONS:** Explainable A-ECG predicts any CAD on CCTA with good overall diagnostic and prognostic performance. This scalable, low-cost approach may aid triage and risk stratification in chest pain pathways.

Keywords: advanced electrocardiography, coronary artery disease, coronary computed

tomography angiography, rapid access chest pain clinic, screening tool

Introduction

Chest pain is one of the most common presentations to emergency departments with substantial cost associated with admission to hospital for further diagnostic assessment and treatment (1). By obviating the need for hospitalisation, the rapid access chest pain clinic (RACC) model has been shown to be a valuable outpatient clinical pathway for the safe and efficient assessment, diagnosis and treatment of patients presenting with symptoms suspicious of cardiac chest pain and at low to intermediate risk (2, 3). A RACC seeks to reduce the financial burden associated with unnecessary hospital admission for management of patients with low to intermediate risk chest pain, through the use of risk stratification algorithms such as the HEART score (4) and highly accurate diagnostic tests including coronary computed tomography angiography (CCTA)(5).

Standard 12-lead electrocardiography (ECG) is the initial diagnostic test used in the risk stratification of chest pain. However, ECG has a relatively low sensitivity and specificity for the diagnosis of clinically significant coronary artery disease (CAD) outside of the diagnosis of myocardial infarction (6, 7). The use and interpretation of the ECG has been largely unchanged in the past few decades. However, recent improvements in digital technology, specifically in signal processing and biomedical computing, have enabled newer innovative techniques of ECG analysis which have demonstrated significant promise, collectively termed advanced electrocardiography (A-ECG).

A-ECG refers to the integration of conventional ECG parameters with other ECG analysis techniques including, singular value decomposition measures of waveform complexity and derived vectorcardiographic (VCG) measures. All these measures are quantified from the standard resting 12-lead ECG. In clinical practice, A-ECG analysis can be performed on digitally acquired ECG files, both rapidly and inexpensively (8).

We hypothesised that A-ECG can improve the diagnostic assessment of intermediate risk chest pain in the RACC. The aim of our study was to create an A-ECG score for any CAD as detected by CCTA in a RACC patient population.

Methods

Study Design

This was a prospective study to derive and validate an A-ECG score for the detection of any CAD, as defined by the presence of any calcified or non-calcified plaque or luminal stenosis in any epicardial coronary artery on CCTA. The derivation cohort consisted of patients who were prospectively referred for assessment at an academic hospital RACC between February 2017 and January 2022. External validation was subsequently performed in an independent external validation cohort drawn retrospectively from a separate healthcare institution, as well as a prognostic cohort. The study was approved by local ethics committees and all participants either provided written informed consent or a retrospective waiver of consent was obtained.

Derivation cohort

The derivation cohort inclusion criteria were age >18 years, referred to the outpatient RACC at Royal North Shore Hospital (Sydney, Australia) following presentation to the emergency department and assessed between February 2017 and January 2022, and underwent digitally recorded ECG and CCTA as part of the chest pain pathway work-up. Exclusion criteria were patients with known prior CAD, prior acute myocardial infarction, and previous revascularisation, including percutaneous coronary intervention and coronary artery bypass grafting.

Validation cohort

The external validation cohort consisted of 762 patients (370 [49%] men, age 57±12 years) from North Shore and Waitakere Hospitals, Te Whatu Ora Waitematā Health, New Zealand, having undergone CCTA and ECG for assessment of CAD in setting of chest pain.

Prognosis cohort

The prognostic performance of the A-ECG score for detecting any CAD was evaluated in a cohort from the United Kingdom (UK) Biobank – a large-scale database of volunteer participants from the UK(9). Prognosis was assessed as a composite of time to first cardiovascular event or all-cause mortality following the date of the index ECG. Included subjects had undergone a resting 12-lead ECG as well as an extensive questionnaire detailing demographic, medical, family and psychosocial history. Included subjects were divided into those without pre-existing cardiovascular disease (CVD) and those with either self-reported or pre-existing diagnosis of CVD, including stroke, myocardial infarction or angina. Subjects were excluded if they had ECG abnormalities which would limit analysis, such as significant atrial or ventricular arrhythmia or QRS duration >120 ms.

ECG acquisition and analysis

Resting 12-lead ECG data for each subject were collected from the local ECG storage systems and exported into xml files. A-ECG semi-automatic software developed in-house was used to analyse the xml files. A-ECG analysis encompassed: conventional ECG measures derived automatically, including all major waveform intervals, axes and voltages; A-ECG measures, obtained after ECG signal averaging; vectorcardiographic measures obtained from the Frank X, Y and Z leads derived from the 12-lead ECG using Kors'

regression transformation, and QRS and T-wave complexity measures quantified from eigenvector lead waveforms calculated using singular value decomposition of the 12-lead ECG. Further exclusion criteria included non-sinus rhythm, bundle branch block (QRS>120ms), pre-excitation, heart rate > 100bpm, and corrupted or uninterpretable ECG files.

CCTA analysis

Each clinically acquired CCTA was analysed by a study investigator (MR) blinded to the A-ECG results for the presence, distribution, and severity of CAD. Any coronary artery disease was defined as the presence of any coronary plaque (calcified or non-calcified) or luminal stenosis. Significant CAD was defined as luminal stenosis >50% in any coronary artery.

Statistical Analysis

Statistical analysis was performed in R (R Foundation for Statistical Computing, version 4.1.2, Vienna, Austria). Continuous variables are reported as mean±SD and count data as number and frequency. Baseline characteristics were compared using the t test, or Mann-Whitney U-test, as appropriate depending on normal distribution. A multivariable A-ECG score for detecting any CAD was derived using elastic net logistic regression. Nested resampling was applied to determine cross-validated diagnostic accuracy and obtain 95% confidence intervals (CI). Nested resampling was used as a method of cross validation and model evaluation to minimise the risk of overfitting by optimisation bias in a predictive model. The derived A-ECG score was then used to estimate the probability of a given subject in the validation cohort having any CAD. Prognosis was evaluated by Kaplan-Meier. Multivariable cubic spline analysis was generated to demonstrate changes in hazard ratio with

increasing AECG score as a continuous variable, with adjustment of age, sex, hypertension, diabetes mellitus, dyslipidaemia, family history, smoking, and elevated body mass index.

Results

Study population

The final analysis of the derivation population included 171 patients (60% male, age 59±13 years). Table 1 summarises the baseline patient characteristics. Of these, 72 (42%) had normal coronary arteries and 99 (58%) had any CAD. The total cohort was deemed low-to-intermediate risk for major cardiac events as indicated by a HEART score of 3±1 (range 1-6). Any CAD group had a higher risk score compared to the normal coronaries group (HEART score 4±1 vs 3±1; p<0.001). The any CAD group was older (64±12 vs 51±12 years, p<0.001) and had a higher proportion of males (67% vs 51%, p=0.04). They were also more likely to be hypertensive (46% vs 22%, p=0.001, hyperlipidaemic (59 vs 26%, p=<0.001) and smokers (42 vs 25%, p=0.02). Rates of diabetes did not differ between the two groups, and both groups were overweight as indicated by body mass index (BMI) (28±6 vs 27±5 kg/m2, p=0.46).

Table 1: Baseline patient characteristics of the derivation cohort.

Parameters	All	Normal coronaries	Any coronary disease	p-value
Number of patients, n (%)	171 (100)	72 (42)	99 (58)	
Age, years	59 ± 13	51 ± 12	64 ± 12	<0.001
Male sex, n (%)	103 (60)	37 (51)	66 (67)	0.04
HEART score	3 ± 1	3 ± 1	4 ± 1	<0.001
Body mass index, kg/m^2	27 ± 5	27 ± 5	28 ± 6	0.46
Hypertension, n (%)	62 (36)	16 (22)	46 (46)	0.001

Diabetes, n (%)	18 (11)	4 (6)	14 (14)	0.1
Smoker, n (%)	60 (35)	18 (25)	42 (42)	0.02
Hyperlipidaemia, n (%)	77 (45)	19 (26)	58 (59)	<0.001

Advanced ECG analysis

Derivation analysis: A-ECG analysis of the derivation cohort identified seven measures that had the strongest association with any CAD and included: (1) Increasing age, (2) the percent of the total QRS loop area in the frontal plane that subtends the left lower quadrant of that frontal plane, (3) the elevation angle (degrees) of the 3-dimensional QRS loop when three eighths of the way into the loop, (4) the elevation angle (degrees) of the 3-dimensional QRS loop when six eighths of the way into the loop, as the ST segment loop extends from the end of the QRS wave to the end of the T wave (5) T-wave complexity, defined as the sum of T-wave eigenvectors 3 to 8 divided by (eigenvector 1 minus eigenvector 2), (6) voltage sum of T-wave eigenvectors 4 to 8, and (7) sex.

These features were combined to derive a seven parameter A-ECG score to detect any CAD with an area under the receiver operating characteristic (ROC) curve of [95% confidence interval] of 0.78 [0.77–0.78], sensitivity 82 [81–83]%, specificity 54 [53-55]%, positive predictive value 71 [71–72]%, negative predictive value 70 [68–71]%, positive likelihood ratio 1.8 and inverse negative likelihood ratio 3.0. The measures included in the final seven-parameter A-ECG score, and the intercept and coefficients for the regression equations are presented in Tables 2 and 3. Diagnostic performance measures are shown in Table 4.

Table 2: A-ECG measures in the score.

	Total (n=171)	Normal coronaries (n=72)	Any stenosis (n=99)	p-value
Age	62.1(52.8-73.1)	54.7(44.3-62.5)	69.2(60.3-76.3)	<0.001

Total QRS loop area in the frontal plane subtending the left lower quadrant, (%)	0.90(0.05-5.4)	0.94(0.05-7.1)	0.89(0.06-4.7)	0.01
Elevation angle of the 3D QRS loop when 3/8 of the way into the loop, degrees	28.8(15.4-38.9)	33.1(20.3-41.4)	26.2(13.4-35.5)	<0.001
Elevation angle of the 3D QRS loop when 6/8 of the way into the loop, as the ST segment loop extends from the end of the QRS to end of the T-wave, degrees	31.9(25.1-41.2)	35.8(28.8-44.6)	29.8(19.7-38.9)	0.004
T-wave complexity, defined as the sum of T-wave eigenvectors 3 to 8 divided by (eigenvector 1 minus eigenvector 2), unitless ratio	0.09(0.06-0.16)	0.09(0.06-0.14)	0.10(0.07-0.21)	0.02
voltage sum of T-wave eigenvectors 4 to 8, mV	17.8(15.1-20.9)	16.8(14.3-19.7)	18.5(15.6-21.1)	0.01
Male sex, n (%)	103 (60)	37 (51)	66 (67)	0.04

Table 3: A-ECG estimates

Term	estimate	SD
(Intercept)	3.39465903	0
Age	-0.05681443	-0.7536
Total QRS loop area in the frontal plane subtending the left lower quadrant	0.004451245	0.03346
Elevation angle of the 3D QRS loop when 3/8 of the way into the loop	0.000747439	0.014837
Elevation angle of the 3D QRS loop when 6/8 of the way into the loop, as the ST segment loop extends from the end of the QRS to end of the T-wave	0.001766589	0.030392
T-wave complexity, defined as the sum of T-wave eigenvectors 3 to 8 divided by (eigenvector 1 minus eigenvector 2)	-0.1309184	-0.02902
voltage sum of T-wave eigenvectors 4 to 8	-0.00208173	-0.01101

Male sex	-0.1892255	-0.09288

Table 4: Performance markers in the A-ECG score to detect any CAD.

Performance Marker	Derivation cohort	Validation cohort
AUC (95% CI)	0.78 [0.77–0.78]	0.72 [0.68–0.76]
Sensitivity, %	82 [81–83]%	90 [88–92]%
Specificity, %	54 [53-55]%	31 [29-33]%
Positive predictive value, %	71 [71–72]%	63 [61–64]%
Negative predictive value, %	70 [68–71]%,	70 [68–71]%
Positive likelihood ratio	1.8	1.3
Inverse Negative likelihood ratio	3.0	3.1

Abbreviations: A-ECG, advanced electrocardiography; CAD, coronary artery disease; AUC, area under the curve; CI, confidence interval

Validation analysis: Sensitivity analyses were undertaken in the validation cohort. Table 5 summarises the patient characteristics of the validation cohort. Diagnostic performance measures for the derivation and validation cohorts are shown in Table 4. The A-ECG score's diagnostic utility was retained for the detection of any CAD. The sensitivity showed a substantial numerical increase, whilst specificity, positive predictive value and positive likelihood ratio were reduced. Negative predictive value and inverse negative likelihood ratio remained similar.

Table 5: Patient characteristics of the validation cohort.

Parameters	All	Normal coronaries	Any coronary disease	p-value
Number of patients, n (%)	762 (100)	334 (44)	428 (56)	
Age (mean (SD))	57 ± 12	53 ± 14	61 ± 10	<0.001
Male sex, n (%)	370 (49)	136 (41)	234 (55)	<0.001
Body mass index, kg/m^2	29 ± 6	29 ± 7	29 ± 6	0.5
Hypertension, n (%)	275 (36)	79 (24)	196 (46)	<0.001
Diabetes, n (%)	111 (15)	39 (12)	72 (17)	0.06
Hyperlipidaemia, n (%)	109 (14)	38 (11)	71 (17)	0.05

Prognosis analysis: The UK Biobank population for prognostic validation consisted of 27,239 subjects, mean age 66±8 years, 50% male. The mean follow-up duration was 1.9 [0.7-4.4] years with 966 cardiovascular events occurring over this period. Of these subjects, 23,051 (85%) had no previous cardiac history at the time of their inclusion. The baseline characteristics of the UK Biobank cohort are presented in Table 6. The A-ECG score is associated with prognosis: cumulative rate of cardiovascular hospitalisation or all-cause death (figure 1) and hazard ratio for cardiovascular events (Figure 2).

Table 6: UK Biobank cohort patient characteristics

Characteristics	UK Biobank population, all subjects	Group with no prior history of CVD	Group with prior history of CVD
n (%)	27,239 (100)	23,051 (85)	4,188 (15)
Age, years	66 ± 8	66 ± 8	69 ± 7
Height, cm	170 ± 9	170 ± 9	171 ± 9
Weight, kg	76 ± 15	75 ± 15	78 ± 15
BSA, m2	1.89 ± 0.22	1.88 ± 0.22	1.92 ± 0.22
Male sex, n (%)	13,528 (50)	11,015 (48)	2,513 (60)
Family history of CVD, n (%)	11,760 (43)	9,755 (42)	2,005 (48)
BMI, kg/m2	26.5 ± 4.5	26.4 ± 4.4	27.2 ± 4.6
Physically inactive, n (%)	2,994 (11)	2,458 (11)	536 (13)
Diabetes, n (%)	1,667 (6)	1,188 (5)	479 (11)
Hypertension, n (%)	8,030 (30)	5,833 (25)	2,197 (53)
Hyperlipidaemia, n (%)	5,548 (20)	3,650 (16)	1,898 (45)
Medications	-	-	-
Lipid lowering, n (%)	4,590 (17)	3,109 (14)	1,481 (35)
Anti-hypertensives, n (%)	4,120 (15)	2,871 (13)	1,249 (30)
Insulin, n (%)	134 (1)	94 (0)	40 (1)

Abbreviations: UK, United Kingdom; CVD, cardiovascular disease; BSA, body surface area;

BMI, body mass index;

Figure 1: Cumulative rate of cardiovascular hospitalisation or all-cause mortality per quartile of A-ECG score

Figure 2: Restricted cubic spline of adjusted hazard ratio for cardiovascular events and all-cause mortality

Discussion

The main finding of the study is that explainable A-ECG analysis of the resting standard 12-lead ECG can identify the presence of CAD on CCTA with good diagnostic accuracy and an independent prognostic association. A seven-parameter model integrating age, sex and five A-ECG measures achieved an AUC of 0.78 for detecting any coronary artery disease (CAD) in a low-to-intermediate risk chest pain derivation cohort and AUC 0.72 in an external validation cohort . Importantly, the score also stratified long-term cardiovascular prognosis adjusted for age, sex and cardiovascular risk factor in a large external population cohort, supporting both its diagnostic and prognostic utility.

This is the first study to derive an explainable A-ECG score using CCTA-defined CAD, and it was done using a contemporary low-to-intermediate risk chest pain population. While previous investigations have shown that A-ECG improves the detection of catheterization-confirmed CAD and structural heart disease compared with conventional ECG interpretation, most have focused on invasive angiographic endpoints or functional ischemia. For example, prior studies have shown the ability of A-ECG to detect the presence of catheterization-proven or stress perfusion imaging-proven CAD with higher sensitivity and specificity than conventional ECG interpretation (8, 10). Additionally, other studies have demonstrated the utility of A-ECG in identifying left ventricular systolic dysfunction, left ventricular hypertrophy and other cardiac conditions (11-13). In contrast, the present study leverages CCTA as the reference standard, reflecting the current guideline-endorsed first-line imaging modality for stable chest pain evaluation (14-16). This is clinically important, as CCTA identifies both obstructive and non-obstructive plaque, which are increasingly recognised as prognostically meaningful.

Clinical implications in the RACC setting

Patients presenting with low-to-intermediate risk chest pain represent a major diagnostic and resource burden. Although RACC pathways incorporating CCTA are cost-effective and safe, a substantial proportion of patients ultimately demonstrate no CAD, raising the need for upstream screening strategies to refine further testing referral. The present A-ECG score offers a rapid, low-cost, and scalable approach that could be applied at the point of first assessment.

The external validation cohort showed a modest inverse negative likelihood ratio of 3.1 for any CAD, suggesting that a score greater than 50% was approximately three times more likely to exclude CAD as a result of a negative test result, and this can potentially be useful to justify no further testing. This supports its potential role as an adjunct to clinical risk scores and a screening tool, enabling more targeted use of CCTA or other imaging.

Mechanistic insights from A-ECG features

The discriminative capacity and explainability of the A-ECG model is underpinned by physiologically meaningful features. Two of the seven measures reflect T-wave waveform complexity, a marker of ventricular repolarization heterogeneity. Increased T-wave complexity has consistently been associated with adverse cardiovascular outcomes and long-term mortality(17, 18), arrhythmic risk (increased risk for sudden cardiac death, appropriate therapies in patients with new ICD implants and identification of long QT syndrome compared to healthy controls(19, 20)), and has been shown to detect subclinical ischemia in asymptomatic populations undergoing exercise stress testing(21). Therefore, the inclusion of T-wave measures supports the concept that early ischemic or structural myocardial alterations manifest as subtle repolarization abnormalities, even in the absence of overt ECG changes.

Three further ECG measures relate to QRS loop orientation and spatial geometry, reflecting ventricular depolarization patterns. Alterations in QRS vector direction and loop morphology have long been associated with regional myocardial injury, ischemia, and adverse prognosis (23–32). Our findings align with prior work demonstrating that vectorcardiographic metrics provide incremental diagnostic information beyond conventional axis and voltage measurements, particularly in patients with functionally significant CAD or regional wall motion abnormalities.

Finally, the inclusion of age and male sex reflects these well-established demographic risks for CAD(22, 23). Their integration with electrocardiographic biomarkers enables a hybrid physiological-clinical risk model, which may enhance generalisability and interpretability. Importantly, the score retains a prognostic association despite adjusting for age and sex, confirming the incremental prognostic contribute of the A-ECG measures beyond age and sex alone.

External validation and prognostic relevance

While specificity, positive predictive value and positive likelihood ratio declined in the external validation cohort, sensitivity improved. Importantly, AUC, negative predictive value and inverse negative likelihood ratio remained similar. This suggests that the model may preferentially exclude subclinical atherosclerotic disease in heterogeneous populations. Importantly, in the UK Biobank cohort, the A-ECG score was associated with cardiovascular events and mortality even after adjustment for age, sex and cardiovascular risk factors,

indicating that the electrocardiographic signatures captured by the model reflect biologically meaningful myocardial substrate rather than purely anatomical plaque burden.

Limitations

Several limitations merit consideration. The derivation cohort was modest in size and consisted mostly of non-obstructive CAD, reflecting real-world RACC populations, but limiting precision for detecting high-grade stenoses. However, nested resampling was employed to mitigate overfitting, and the performance was evaluated in a robustly sized validation cohort (n=762). CCTA was used as the reference standard, and therefore microvascular dysfunction or vasospastic disease may have been misclassified as "normal." Finally, the study was observational and clinical utility will require larger-scale prospective studies.

Conclusions

A-ECG can predict the presence of any CAD on CCTA with an overall good diagnostic and prognostic performance, using a seven-parameter score including age, sex and five ECG measures derived from A-ECG analysis of the standard 12-lead ECG. This approach provides an explainable and mechanistically interpretable, scalable, and low-cost screening tool that could complement existing clinical pathways by refining downstream testing selection. Prospective trials are now warranted to evaluate its impact on diagnostic efficiency, cost-effectiveness, and patient outcomes.

Funding

RK was supported by a grant from New South Wales Health - EMC Round 2 Cardiovascular disease.

Declaration of competing interests

TTS is a principal of Nicollier-Schlegel SARL, a company that performs ECG research consultancy using the software used in the present study. MU and TTS are owners of Advanced ECG Systems, which is a company developing commercial clinical applications of Advanced ECG analyses of the kind that has been evaluated in this study. RK has a financial interest in Advanced ECG Systems through being married to MU.

References

1.   Skinner JS, Smeeth L, Kendall JM, Adams PC, Timmis A. NICE guidance. Chest pain of recent onset: assessment and diagnosis of recent onset chest pain or discomfort of suspected cardiac origin. Heart (British Cardiac Society). 2010;96(12):974-8.

2.   Yu C, Sheriff J, Ng A, Brazete S, Gullick J, Brieger D, et al. A Rapid Access Chest Pain Clinic (RACPC): Initial Australian Experience. Heart, lung & circulation. 2018;27(11):1376-80.

3.   Black JA, Cheng K, Flood JA, Hamilton G, Parker S, Enayati A, et al. Evaluating the benefits of a rapid access chest pain clinic in Australia. Medical journal of Australia. 2019;210(7):321-5.

4.   Backus BE, Six AJ, Kelder JC, Bosschaert MAR, Mast EG, Mosterd A, et al. A prospective validation of the HEART score for chest pain patients at the emergency department. International journal of cardiology. 2013;168(3):2153-8.

5.   Kozor R, Mooney J, Lowe H, Kritharides L, Altman M, Klimis H, et al. Rapid Access Chest Pain Clinics: An Australian Cost-Benefit Study. Heart, lung & circulation. 2022;31(2):177-82.

6.   Sox HCJ, Garber AM, Littenberg B. The resting electrocardiogram as a screening test: a clinical analysis. Annals of internal medicine. 1989;111(6):489-502.

7.   Ashley EA, Raxwal V, Froelicher V. An evidence-based review of the resting electrocardiogram as a screening technique for heart disease. Progress in cardiovascular diseases. 2001;44(1):55-67.

8.   Schlegel TT, Kulecz WB, Feiveson AH, Greco EC, DePalma JL, Starc V, et al. Accuracy of advanced versus strictly conventional 12-lead ECG for detection and screening of coronary artery disease, left ventricular hypertrophy and left ventricular systolic dysfunction. BMC cardiovascular disorders. 2010;10(1):28-.

9.   Sudlow C, Gallacher J, Allen N, Beral V, Burton P, Danesh J, et al. UK biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. PLoS medicine. 2015;12(3):e1001779-e.

10.   Zong Y, Maanja M, Chaireti R, Schlegel TT, Ugander M, Antovic JP. Substantial prevalence of subclinical cardiovascular diseases in patients with hemophilia A evaluated by advanced electrocardiography. Journal of electrocardiology. 2020;58:171-5.

11.   Sapra R, Hallqvist L, Schlegel TT, Ugander M, Bell M, Maanja M. Predicting peri-operative troponin elevation by advanced electrocardiography. Journal of electrocardiology. 2021;68:1-5.

12.   Maanja M, Wieslander B, Schlegel TT, Bacharova L, Abu Daya H, Fridman Y, et al. Diffuse Myocardial Fibrosis Reduces Electrocardiographic Voltage Measures of Left Ventricular Hypertrophy Independent of Left Ventricular Mass. Journal of the American Heart Association. 2017;6(1):n/a.

13.   Maanja M, Schlegel TT, Kozor R, Lundin M, Wieslander B, Wong TC, et al. The electrical determinants of increased wall thickness and mass in left ventricular hypertrophy. Journal of electrocardiology. 2020;58:80-6.

14.   Excellence NIfHaC. Clinical guideline CG95 "Chest pain of recent onset: assessment and diagnosis.". NICE. London, England2010.

15.   Rochitte CE, Gottlieb I, de Roos A, Cox C, Bush DE, Hoe J, et al. Diagnostic performance of coronary angiography by 64-row CT angiography. The New England journal of medicine. 2008;359(22):2324.

16.   Mowatt G, Cummins E, Waugh N, Walker S, Cook J, Jia X, et al. Systematic review of the clinical effectiveness and cost-effectiveness of 64-slice or higher computed tomography angiography as an alternative to invasive coronary angiography in the investigation of coronary artery disease. Health technology assessment (Winchester, England). 2008;12(17):iii-iv.

17.   Okin PM, Devereux RB, Fabsitz RR, Lee ET, Galloway JM, Howard BV. Principal Component Analysis of the T Wave and Prediction of Cardiovascular Mortality in American Indians: The Strong Heart Study. Circulation (New York, NY). 2002;105(6):714-9.

18.   Zabel M, Malik M, Hnatkova K, Papademetriou V, Pittaras A, Fletcher RD, et al. Analysis of T-Wave Morphology From the 12-Lead Electrocardiogram for Prediction of Long-Term Prognosis in Male US Veterans. Circulation (New York, NY). 2002;105(9):1066-70.

19.   Shi B, Harding S, Larsen P. Analysis of ECG Measures of Cardiac Repolarization in Relation to Arrhythmic Events in an Implantable Cardioverter Defibrillator Population. Annals of noninvasive electrocardiology. 2017;22(2):e12390-n/a.

20.   Priori SG, Mortara DW, Napolitano C, Diehl L, Paganini V, Cantù F, et al. Evaluation of the Spatial Aspects of T-Wave Complexity in the Long-QT Syndrome. Circulation (New York, NY). 1997;96(9):3006-12.

21.   Al-Zaiti SS, Runco KN, Carey MG. Increased T wave complexity can indicate subclinical myocardial ischemia in asymptomatic adults. Journal of electrocardiology. 2011;44(6):684-8.

22.   Dawber TR, Moore FE, Mann GV. Coronary heart disease in the Framingham study. American Journal of Public Health and the Nation's Health. 1957;47(4 Pt 2):4.

23.   Lerner DJ, Kannel WB. Patterns of coronary heart disease morbidity and mortality in the sexes: A 26-year follow-up of the Framingham population. The American heart journal. 1986;111(2):383-90.