Implementation of the genome-informed risk assessment (GIRA) may lead to large disruptions to the health system

Sandra Lapinska[1], Xinzhe Li[1], Ravi Mandla[1], Zhuozheng Shi[1], Veronica Tozzo[2], Alexander Flynn-Carroll[1], Marylyn D. Ritchie[3], Daniel J. Rader[3,5], Penn Medicine Biobank, Bogdan Pasaniuc[3,4]

[1]Graduate Group in Genomics and Computational Biology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA
[2]Department of Computational Medicine, David Geffen School of Medicine at University of California, Los Angeles, Los Angeles, CA, USA
[3]Department of Genetics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA
[4]Department of Computer and Information Sciences, School of Engineering and Applied Sciences, University of Pennsylvania, Philadelphia, PA, USA
[5]Division of Translational Medicine and Human Genetics, Department of Medicine, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA, USA

To whom correspondence should be addressed sandra.lapinska@pennmedicine.upenn.edu, bogdan.pasaniuc@pennmedicine.upenn.edu

Abstract

The Genome Informed Risk Assessment (GIRA) report from eMERGE has become a standard approach to implement genomic precision medicine at scale. Here, we assess GIRA's utility and impact in a health care system independent of eMERGE, focusing on 9 adult conditions using the Penn Medicine Biobank (PMBB, n=48,279). We find a large number of patients – 50.1% (n=24,185) – were deemed by GIRA as high-risk for at least one of the 9 conditions with 30.4% (n=14,676) due to polygenic and/or monogenic risk. Stratifying by ancestry revealed significant differences in high-risk proportions, with higher rates in African/African American (AFR) (56.6% vs. 50.1%, $p=7.43 \times 10^{-36}$) and lower rates in East (42.0%) and South Asian (40.0%). Increased high-risk rates were observed in the lowest quartile of social deprivation index, highlighting the influence of environmental factors and access to care on GIRA's utility. GIRA was a good predictor of prevalent cases (in-line with the eMERGE GIRA reported results); incident case prediction was substantially attenuated for 5 of the 9 conditions (e.g., OR of 2.36 vs. HR of 1.31 for atrial fibrillation (AFIB)). We find demographic compositions of high-risk patients differed from the incident cases for some of the conditions; for example, high-risk for AFIB individuals where enriched for European ancestries in contrast with incident AFIB cases that were enriched for AFR ancestries. Overall, our results show the accuracy of GIRA as a biomarker to stratify high-risk patients for precision medicine and highlight implementation challenges in its impact on the health system if implemented at scale.

Introduction

Current screening practices for identifying individuals at high-risk of disease rely on clinical risk factors and family history (FH), with assessment of monogenic variants being incorporated for select cancer and cardiometabolic conditions as it can confer several-fold increased disease risk for carriers[1–5]. Family history may reflect rare pathogenic variants of heredity, shared exposures to environmental and lifestyle factors, as well as common genetic susceptibility, thus providing an indirect measure of overall genetic predisposition to disease[6–10]. More recently, polygenic risk scores (PGS) that aggregate the effects of many common variants into a single measure of risk have demonstrated great potential in risk stratification as individuals in the upper tail of distribution can attain risk equivalent to clinical risk factors and monogenic variants[4,11–13]. Thus, combining PGS with family history and monogenic risk provides a more comprehensive genetic risk assessment for common, complex diseases, enabling earlier identification of high-risk individuals who may benefit from preventative interventions[1]. Additionally, individuals carrying the same monogenic variant demonstrate wide variation in disease expression and penetrance[14]. Considering polygenic background improves risk estimation, as it can modify the penetrance and expressivity of monogenic variants[13]. Moreover, for most conditions, a positive family history with a high PGS increases disease risk, whereas a low PGS can offset the risk associated with a positive family history[6,15]. Recognizing that each of these genetic risk factors provides independent and complementary information into inherited disease susceptibility, the Electronic Medical Records and Genomics (eMERGE) Network[16,17] has developed the genome-informed risk assessment (GIRA) report as part of their multi-site clinical study. eMERGE deployed cross-ancestry and ancestry-calibrated PGS, together with monogenic risk and family history, across 11 conditions (9 adult) to evaluate health outcomes among high-risk individuals[1,18,19]. High-risk status for each condition was determined according to criteria established by eMERGE with high-risk PGS thresholds chosen to be equivalent to an odds ratio ≥ 2.

Evaluating GIRA in healthcare systems independent of eMERGE – where the patient populations may differ from those used to calibrate GIRA is a critical step in broadly adoption of genomic precision medicine[20]. Specifically, GIRA's predictive ability may be diminished due to differences in ancestry proportions, environmental exposures, disease prevalence, or case ascertainment[1,18,21]. For instance, in the University of Pennsylvania Health System (UPHS), 55.0% of the patient population self-reported as 'White', 18.2% as 'African American', and 4.1% as 'Asian', with 4.7% self-reporting ethnicity as 'Hispanic'[22]. The Penn Medicine Biobank (PMBB), an electronic health record (EHR)-linked biobank embedded within the UPHS, includes 23.2% individuals of African/African American (AFR) ancestry, the largest representation within any single-institutional medical biobank in the US, providing a unique opportunity to validate GIRA's population-specific performance. More specifically, even though PMBB has a higher proportion of individuals who self-reported as 'White' (66.8%) and 'African American' (24.6%), it is largely representative of the UPHS and a valuable patient population for validation. The clinical utility of the GIRA report also depends on its ability to identify individuals at high-risk of developing the condition in the future as it enables more targeted interventions[23–25]. Evaluation of GIRA for prevalent conditions may overstate GIRA's predictive ability, since a strong association with prevalent conditions does not necessarily reflect how well the tool predicts

future condition events[26]. For instance, PGS strongly associate with prevalent conditions but
have lower predictive accuracy in incident conditions[27]. Cox proportional hazard models for
incidence disease (defined as individuals that develop condition after being evaluated for GIRA),
on the other hand, directly quantify prospective risk by estimating how risk components within
genetic risk assessments influence the likelihood of developing the disease, providing a more
clinically meaningful assessment of prediction. To date, however, no studies have evaluated
GIRA longitudinally for incident disease prediction.

We leveraged longitudinal data in PMBB to investigate the predictive accuracy of GIRA to
assess its ability to accurately capture high-risk patients[22]. Prevalent cases were defined as any
diagnosis recorded before, at, or after entry in biobank (baseline), whereas incident cases were
diagnosed after PMBB entry only. We implemented validated phenotype algorithms for nine
adult-onset conditions - atrial fibrillation (AFIB), breast cancer (BC), chronic kidney disease
(CKD), coronary heart disease (CHD), hypercholesterolemia (HC), obesity, prostate cancer
(PC), type 2 diabetes (T2D), and colorectal cancer (CC) – and evaluated the ability of GIRA to
identify disease cases. In total, 50.1.% (17,747) of unique PMBB patients were labeled by GIRA
as high-risk due to polygenic, monogenic, and/or family history risk for at least one condition
(28.7%, 2.3%, and 28.8%, respectively). Stratifying PMBB patients by ancestry found significant
differences in high-risk proportions when compared to all patients, most notably AFR individuals
were labeled by GIRA-risk at elevated rates (56.6% vs 50.1%, $p=7.43\times10^{-36}$) than in other
ancestries suggesting a need for ancestry-specific calibration. Lower GIRA-risk rates in East
and South Asian ancestry were driven solely by family history suggesting a larger impact from
factors like reporting pattern variability than differences in estimated genetic risk. Of the 17,747
GIRA high-risk individuals, 65.0% were diagnosed with at least one condition before PMBB
recruitment (defined as prevalent cases) and 28.3% developed at the condition after PMBB
entry (defined as incident cases). Effect sizes of GIRA to identify prevalent cases were largely
consistent with those reported by eMERGE ranging from 2.28 to 3.97 for BC and CKD. In
contrast, hazard ratios (HRs) for developing condition after entry in biobank (incident cases)
were significantly lower for 5 out of the 9 conditions, suggesting odds ratios may overstate the
clinical utility of GIRA (e.g. CKD OR: 3.63 (95% CI: 3.16-4.17) vs HR: 1.82 (95% CI: 1.61-2.05)).
GIRA high-risk individuals were of similar ancestry composition to incident cases across all
conditions except AFIB, CKD, and obesity suggesting GIRA may have miscalibration across
ancestries. Finally, elevated GIRA-risk rates were observed in the lowest quartile of social
deprivation index highlighting the potential miscalibration of GIRA due to environmental factors
or healthcare access. In summary, GIRA showed considerable promise for identifying high-risk
individuals in an independent real-world healthcare system. However, its underperformance for
certain conditions highlights the need for condition- and context-specific validation and
refinement to ensure equitable clinical implementation.

Results

Phenotyping nine adult GIRA conditions in the Penn Medicine Biobank (PMBB)

We analyzed 48,279 PMBB patients aged 18 to 75 years at sample collection after excluding
patients enrolled in the Penn Medicine Cancer Risk Evaluation Program (CREP). Majority of
these individuals were genetically similar to European American (EUR) (71.7%) and
African/African American (AFR) (23.2%) populations (**Figure 1; Table S1; Methods**). PMBB
has the largest sample size of AFR individuals among single-institutional medical biobanks,
making it valuable for evaluating the generalizability of GIRA across diverse populations[28–30]. To
closely align our analysis with eMERGE, we established case and control status for nine adult
GIRA conditions using validated phenotyping algorithms from the Phenotype Knowledgebase
(PheKB) when applicable; otherwise, the Chronic Condition Warehouse or the Polygenic Risk
Methods Development (PRIMED) Consortium were used (**Methods**). We observed higher
prevalence estimates than previously reported population-based estimates for all traits except
BC, CC, PC, and obesity (**Figure 1; Table S1, S3**). For example, population-based prevalence
estimates for CHD, T2D, and AFIB are approximately 6.4%[31], 6.28%[32], and 13.7%[33],
respectively, significantly lower than compared to 13.7%, 14.5%, and 17.2% in PMBB. In
contrast, CKD prevalence in PMBB (11.5%) was lower than reported population estimates of
14%[34]. Comparison of disease prevalence between PMBB and the original GIRA validation
cohort revealed statistically significant higher prevalence in PMBB for 3 conditions and lower
prevalence for 5 conditions (**Table S3**). For example, prevalence was significantly lower in
PMBB than in eMERGE for AFIB (17.2% vs. 23.0%; P = $1.13x10^{-104}$) and CKD (11.5% vs.
17.5%; P = $6.95x10^{-220}$) whereas CHD (13.7% vs. 9.2%; P = $1.41x10^{-146}$) and HC (12.1% vs.
8.1%; P = $4.05x10^{-48}$) were more prevalent in PMBB. Significant differences were also
observed for all cancer conditions, with the lower prevalence in PMBB being explained by
exclusion of PMBB participants enrolled in CREP. Prevalence estimates in PMBB generally fell
between population-based estimates and those observed in eMERGE. Higher prevalence for
most conditions is likely due to volunteer bias in biobank participants which has been shown to
distort prevalence estimates relative to the target population[35,36]. As expected, incidence
estimates, defined as receiving diagnosis after sample collection in PMBB, were significantly
lower than corresponding prevalence estimates. Observation time was defined as the earliest of
the following times: initial disease diagnosis, last clinical encounter, or death. Incidence
estimates ranged from 0.5% for CC to 7.2% for obesity. PMBB incidence estimates differed to
population-level incidence; however, AFIB has reported incidence rates of 23.7 per 1000
person-years (PY) over 1-year follow-up[35] (compared with 5.69 per 1000PY over 1-year follow-
up in PMBB) while BC has 130.8 per 100,000 women[37] (compared with 227.0 per 100,000
women per year in PMBB), reflecting cohort-specific differences in ascertainment (**Figure 1;
Table S2, S3**).

Demographic characteristics of prevalent and incident disease in PMBB

Next, to disentangle biological effects from healthcare disparities and ascertainment biases, we
compared ancestry, age and sex enrichments in prevalent versus incident cases for each
condition. We found AFR individuals were enriched in both prevalent and incident obesity, T2D,
CKD, CHD, and PC whereas EUR individuals were enriched in BC and CC, largely in-line with
existing epidemiological studies[38–42]. Differences in ancestry enrichment patterns were observed
between prevalent and incident cases for AFIB and HC. We found an enrichment of AFR

164 individuals in incident AFIB (OR: 1.14 (95% CI: 1.03-1.26)) whereas prevalent AFIB was
165 enriched in EUR individuals (OR: 1.39 (95% CI: 1.31-1.48)). In contrast, incident HC was
166 enriched for EUR individuals (OR: 1.10 (95% CI: 1.00-1.21)) while prevalent HC was enriched
167 for AFR individuals (OR: 1.16 (95% CI: 1.08-1.24)). These enrichment patterns suggest possible
168 participation and recruitment biases in PMBB that could be impacted by factors like survival
169 bias, disease severity, healthcare access and utilization, or issues regarding inclusivity (**Figure**
170 **2; Table S7**)[43]. Biobanks are known to suffer from inclusion bias with several studies comparing
171 participants to the background population using publicly available census data[44,45]. Next, we
172 found incident T2D and BC were enriched for older individuals, specifically those aged 45-65
173 and aged 65-100 (e.g. incident T2D OR: 1.27 (95% CI: 1.15-1.41) vs prevalent T2D OR: 0.68
174 (95% CI: 0.63-0.72)), compared to prevalent cases. Incident CC also showed enrichment for
175 individuals aged 65-100. Prevalent AFIB, CKD, CC, CHD, and PC were enriched for individuals
176 aged 45-65 whereas incident cases were not (**Supplementary Figure 1; Table S8**). These
177 findings highlight that incident disease tends to skew toward older individuals, likely reflecting
178 longer survival trends among adults aged 60 or older with common, not immediately fatal
179 conditions like T2D, AFIB, or HC[47,48]. Conversely, enrichment among individuals aged 18-45
180 was observed only for prevalent CC (OR: 1.25 (95% CI: 0.99-1.58)), mirroring trends of
181 increasing diagnoses among individuals under age 55[49], likely driven by earlier screening
182 practices and resulting in greater biobank enrollment among younger individuals with elevated
183 CC risk factors. Lastly, all conditions were enriched for males except prevalent and incident
184 obesity and HC, consistent with previous epidemiological studies[50–54] , aside from CKD, which is
185 more prevalent globally among women[34,55]. This may be due to potential under-screening and
186 underdiagnosis of CKD among women within healthcare systems[56]. BC and PC were not
187 evaluated for sex enrichment as GIRA returns results to self-reported sex at birth of female and
188 male, respectively (**Supplementary Figure 2; Table S9**). Together, these findings demonstrate
189 how ancestry, age, and sex enrichments reflect known epidemiologic patterns while also
190 revealing potential ascertainment biases between prevalent and incident cases that need to be
191 considered in downstream analyses.

192 **GIRA accurately identifies prevalent disease cases in PMBB**

193 We next assessed the clinical utility of the PGS risk component of the GIRA report by evaluating
194 its ability to identify prevalent disease in PMBB (**Methods**). To validate the PGS performance
195 for eight adult-onset conditions, we associated each condition-specific PGS with its
196 corresponding prevalent condition to find comparable ORs per standard deviation (SD) to those
197 from eMERGE (**Table S6, Methods**). EUR individuals in PMBB had largely similar direction of
198 effects associated with having a score above the specific threshold ('High Risk-PGS') compared
199 to the eMERGE reported ORs[55] across prevalent conditions (**Figure 3**; **Table S13**). The 95%
200 confidence interval for all prevalent conditions either included or exceeded an OR of 2. CHD
201 individuals in the top 5% of the PGS had higher ORs in PMBB compared to eMERGE (EUR:
202 3.07 (95% CI: 2.67-3.53) vs 2.36 (95% CI: 2.12-2.62), respectively). In contrast, prevalent CKD
203 and HC had lower ORs; however, AFR individuals for HC had higher ORs (AFR PMBB: (4.36
204 (95% CI: 3.09- 6.22)) vs AFR eMERGE (2.98 (95% CI: 1.88-4.60)); **Table S13**). This shows that
205 GIRA's cross-ancestry PGS along with its 'High Risk-PGS' thresholds can identify individuals

with increased risk of condition and yield broadly reproducible risk estimates in an independent
biobank. However, the variability in ORs for EUR and AFR individuals in PMBB relative to
eMERGE reported ORs suggests that the effect of high-risk GIRA may vary across different
patient populations to which results will be returned to. More specifically, differences in sample
characteristics and case-control definitions between the eMERGE GIRA study and PMBB may
influence predictive performance when applied to cohorts from healthcare systems independent
of eMERGE.
213
Next, all GIRA high genetic risk components were aggregated into a binary combined risk score
metric, referred to as 'High Risk - All', to assess the clinical utility of the 'High Risk - All' metric
by comparing individuals at high-risk for at least one risk component to those classified as not
high-risk. We found ORs ranging from 3.42 (95% CI: 2.81-4.15) for prevalent T2D to 3.63 (95%
CI: 3.16-4.17) for prevalent CKD to 4.76 (95% CI: 2.86-7.70) for prevalent CC across
ancestries. Prior studies reported that individuals in the top 2% of the genome-wide polygenic
score were associated with a 2.66–4.93-fold increased risk for CKD across ancestries[57] and the
top 2% of PGS distribution had a 2.5-4.5-fold increased risk for T2D across ancestries[56,57].
Except for CHD, all conditions had ORs larger than 2 thus confirming the utility of GIRA as
biomarker to identify prevalent cases in a health system biobank (**Figure 3**).

GIRA underperforms in identifying incident disease cases in PMBB

Next, we evaluated GIRA's ability to identify individuals at risk of developing the condition in the
future to find that GIRA-'High Risk' were predictive of incident disease, yielding hazard ratios
(HRs) directionally consistent with the corresponding ORs observed in prevalent cases, albeit
with smaller magnitude than ORs (**Figure 3, Table S11**). This attenuation is consistent with the
expectation that incidence risk prediction reflects more conservative effect estimates than
prevalent associations as it addresses survival or ascertainment biases. The most significant
reductions were observed for CKD (OR: 3.63 (95% CI: 3.16-4.17) vs HR: 1.82 (95% CI: 1.61-
2.05)) and AFIB (OR: 2.36 (95% CI: 2.10-2.66) vs HR: 1.31 (95% CI: 1.05-1.63)). Thus, GIRA
but is much weaker at predicting who will newly develop these conditions over time and thus
less informative for future disease prediction. Indeed, across all conditions except AFIB, the 15-
year cumulative incidence was substantially higher in individuals at high-risk, highlighting that
accounting for time through HRs and cumulative incidence offers a more accurate
representation of risk compared to ORs (**Figure 4)**.

Genetic risk estimation accuracy varies by ancestry and condition

The differential performance of PGS across populations is one of the main practical challenges
for their use in clinical settings[58,59]. We observed largely similar direction of effects in both
incident and prevalent cases across ancestries, though the magnitudes varied greatly across
ancestries (**Table S10, S11, Supplementary Figure 3**). For example, ORs among AFR
individuals for HC (OR: 4.98 (95% CI: 3.60-6.96)) were higher than in EUR individuals (OR: 3.91
(95% CI: 3.25-4.72)). Furthermore, AFIB, CHD, T2D, and obesity showed 95% confidence
intervals or ORs that did not reach a threshold of OR > 2 in AFR individuals, highlighting

reduced performance in non-EUR populations and underscoring the need for additional
ancestry-specific validation for certain conditions (**Figure 3, Table S11**). Results were
consistent with eMERGE for 5 conditions in EUR and AFR ancestry but variable in other
populations, likely due to the relatively low sample size and large estimation uncertainties
(**Table S13**). Finally, we found variability in the predictiveness of the three risk components of
GIRA (polygenic, monogenic, family history) across EUR and AFR individuals. For example,
EUR individuals with 'High Risk-Monogenic' or 'High Risk-Family History' for incident BC had
higher HRs ('High Risk-Monogenic' HR: 3.86 (95% CI: 2.29-6.48); 'High Risk-Family History'
HR: 2.37 (95% CI: 1.94-2.88)) compared to those with 'High Risk-PGS' (HR: 1.79 (95% CI:
1.25-2.55)). The same pattern was observed for HC and AFIB between 'High Risk-PGS' and
'High Risk-Monogenic'. Furthermore, AFR individuals had larger HRs for incident BC for 'High
Risk-PGS' and 'High Risk-Monogenic', but not for 'High Risk-Family History' (2.75 (95% CI:
1.60-4.71); 8.16 (95% CI: 3.31-20.1); 1.73 (95% CI: 1.24-2.41), respectively) compared to EUR
individuals. However, HRs for incident CHD were lower for 'High Risk-PGS' and 'High Risk-
Monogenic' and higher for 'High Risk-Family History' in AFR individuals. Incident PC showed
elevated HRs for 'High Risk-Family History' in AFR individuals (**Figure 5, Table S12**).

GIRA-high-risk population demographics differ from general biobank population

Having established the predictive performance of GIRA across the nine adult-onset conditions,
we next turned into contextualizing the individuals labeled as high-risk by GIRA. 24,148 (50.1%)
of all individuals in PMBB were labeled as high-risk for at least one condition, of which 13,870
(28.7%) due to high PGS risk, 1,111 (2.3%) had monogenic risk, and 13,873 (28.8%) reported a
family history of disease. 14,676 (30.4%) were at high-risk due to the polygenic and/or
monogenic risk components larger than the eMERGE expected results (25%[58]). This shows
GIRA's potential to identify a larger proportion of the population that could benefit from early
intervention while also highlighting the disruptive impact and burden on the health system if
GIRA is implemented in clinical practice. Among high-risk individuals, the proportion of those
who "already have disease" ranged from 2.5% for PC to 11.0% for obesity. More specifically,
2.5% of the 8,102 individuals at high PC risk, 12.9% of 2,311 at high CKD risk, and 11.0% of
1,449 at high obesity risk developed the corresponding condition during follow-up. These
relatively low percentages suggest that most high-risk individuals have not yet manifested
clinically recognizable disease within the observation time, which aligns with risk assessments
ability to identify risk before disease onset[60]. A large proportion of high-risk individuals fell into
the "status not ascertained" category as they could not be definitively classified as a case or
control with our data. Of the individuals labeled as high-risk by GIRA's criteria across all
conditions, 5,023 (20.8%) individuals developed at least one condition during follow-up. Lastly,
there were 6,965 PMBB individuals who developed at least one condition during follow-up but
were not at high-risk. Thus, these findings indicate that case proportions among high-risk
individuals likely underestimate the true burden of disease due to these individuals having not
yet developed the disease or remain unclassified due to phenotyping uncertainty.

Next, to understand variability in high-risk rates in PMBB subpopulations, we compared the
high-risk proportion among individuals in different ancestries and social deprivation index
quartiles to those for all PMBB individuals. We found significantly higher high-risk rates among

individuals of AFR ancestry (56.6% vs 50.1%, Z=-12.5, p=7.43x10^{-36}, respectively) and lower-
risk rates for EUR, EAS, and SAS individuals (e.g. SAS: 40.0% vs 50.1%, Z = 5.25, p = 1.49x10^{-7}
). We found that the lower rates in East and South Asian ancestries were driven solely by
family history indicating a large impact from factors like environment, recall bias, or reporting
behavior than underlying genetic risk. Interestingly, higher rates in AFR individuals were not
driven by monogenic risk, which saw lower high-risk rates compared to all high monogenic risk
individuals (1.6% vs 2.3%, Z=4.77, p=1.85x10^{-6}, respectively) (**Table S15-16**). Second, to
understand how socioeconomic factors may impact GIRA's utility, we determined high-risk
proportions among social deprivation index (SDI) quartiles. Compared to all high-risk PMBB
individuals, there were significantly higher high-risk rates among the most deprived (SDI quartile
4) (53.7% vs 50.1%, Z=-6.77, p=1.30x10^{-11}, respectively), which were driven by high PGS risk
and high family history risk. Additionally, lower high PGS risk rates were observed in the least
deprived quartile (SDI quartile 1). Monogenic risk, however, had lower high-risk rates at SDI
quartile 4 (1.7% vs. 2.3%, Z = 3.98, p=6.79x10^{-5}) (**Table S17**), suggesting the GIRA's polygenic
risk and family history performance is more impacted by lower socioeconomic status than
monogenic risk as illustrated in previous studies[61,62]. Lastly, we evaluated the demographic
composition of the GIRA high-risk group versus the incident case population across the nine
conditions. Except obesity, CKD, and AFIB, ancestry enrichments among high-risk individuals
largely mirrored those observed in incidence rates. Specifically, obesity showed no enrichment
for EUR or AFR individuals among high-risk individuals, whereas incident obesity was
overrepresented in AFR individuals and underrepresented in EUR individuals. High-risk
individuals for CKD were enriched for EUR and depleted for AFR individuals, while the opposite
pattern was seen for incident CKD (**Figure 7**). This under-enrichment of AFR individuals within
the GIRA-classified high-risk individuals may be due to the underperformance of genetic risk in
AFR individuals relative to EUR individuals[63]. As a secondary analysis, we restricted to
individuals not lost during follow-up (**Methods**) to find our results remained largely consistent
(**Supplementary Figures 8-10)**.

Discussion

The Penn Medicine Biobank as institutional biobank embedded within the University of
Pennsylvania's Healthcare System (UPHS) provided an ideal setting for validation of the
condition-specific high-risk GIRA criteria in a real-world health system. We showed that GIRA
can identify individuals with a 2.81 to 3.41-fold increased risk depending on the prevalent
condition, largely in line with the eMERGE reported results. The clinical translation of GIRA
relies on its ability to predict future disease cases rather than the strength of the association with
prevalent conditions. We presented the first study that evaluated GIRA's high-risk criteria
longitudinally showing that hazards ratios show reduced performance of GIRA for various
conditions. We demonstrated differences in high-risk classification across ancestry and social
deprivation index quartiles. Finally, we report that GIRA high-risk individuals differed in their
demographic composition compared to incident cases suggesting that additional factors, such
as social determinants of health or environmental exposures, may contribute to disease
development beyond genetic risk. This was further stressed by lower high-risk rates in East and
South Asian ancestries being driven by family history and higher high-risk rates in the most

deprived social deprivation index quartile. Our analysis demonstrates that though GIRA is
largely generalizable for prevalent conditions, real-world implementation will require careful
choice of metrics to convey magnitude of risk, potential condition-specific calibration, and
ancestry-stratified performance metrics to accurately communicate risk across diverse
populations.

Half of PMBB participants were identified as high-risk for at least one condition, highlighting the
substantial burden that returning GIRA results could place on the healthcare system. 28.7% of
PMBB participants had at least one high-risk PGS result, which is greater than the 23.2%[62,63] in
the first 2,500 processed samples in the GIRA study. However, the Genomic Medicine at
Veterans Affairs (GenoVA) study, which is providing insights for evaluating the clinical utility of
monogenic and PGS implementation for six common diseases in primary care settings,
estimated an expected 33% at high-risk PGS[66,67]. More specifically, in their first 840 processed
samples, 2% were at high-risk monogenic while 37.1% had at least one high-risk PGS result.
We also observed 30.4% PMBB individuals with a high-risk PGS or monogenic result while
eMERGE expected 25%. The proportion of high-PGS risk individuals in PMBB fell in between
those reported by the GIRA and GenoVa studies, while the 2.3% for high-risk monogenic
closely mirrored the reported 2% by the GenoVa study suggesting that the risk assessments are
largely generalizable and that differences in high-risk proportions among different studies likely
reflect differences in cohort composition, PGS thresholds, and the specific PGS implemented,
rather than differences in underlying genetic risk.

We conclude with several limitations of our work. First, even with current efforts to enroll a
diverse group of individuals into PMBB, the relatively small sample size for Admixed American,
East Asian, and South Asian individuals caused large estimation uncertainties. Even with
differences in the proportion of diverse individuals, we observed OR > 2 for EUR and AFR
individuals in PMBB; however, the increase in risk varied by population as compared to
eMERGE, likely reflecting differential performance of these score to predict disease risk across
health care systems with different sample characteristics[11,68]. Validating GIRA in a cohort with
different ancestry proportions remains valuable, as it evaluates its performance and potential
biases in a setting more representative of real-world healthcare systems. PMBB provided such
a setting as the biobank is embedded within the UPHS. More specifically, PMBB participants
have comparable age and gender distributions, but a higher proportion of individuals who self-
identify as African American (24.6% vs 18.2%) or White (66.8% vs 55.0%) compared with the
University of Pennsylvania Healthcare System[65,66]. Second, PMBB is composed of individuals
recruited from the UPHS, a population potentially exhibiting a higher disease burden than the
general population. This inflates prevalence estimates and monogenic variant penetrance,
which carries bias and may limit the generalizability of these findings. Third, lack of precision in
family history information in the EHR due to recall bias and sensitivity to wording in queries may
lead to inaccurate risk assessment[69,70].

Figures

Figure 1. Distribution of nine adult GIRA conditions across ancestry, age, and sex in PMBB. Left panel shows the prevalence (defined as participants diagnosed at dates prior to, at, and post enrollment) compared to the incidence (defined as participants receiving diagnosis after sample collection) of each condition within PMBB. Right shows the distribution of cases by genetically defined ancestry groupings with distribution in the overall biobank at the top. Bars are annotated with ancestry proportions for participants of EUR or AFR ancestry only, as these groups contain the majority of the participants. The ancestry proportions are shown relative to the total PMBB participants included in the analysis. We included all PMBB participants aged 18 to 75 at sample collection who were not enrolled in the Penn Medicine Cancer Risk Evaluation Program.

Abbreviations: GIRA: Genome Informed Risk Assessment, PMBB: Penn Medicine Biobank, T2D: Type 2 Diabetes, AFIB: Atrial Fibrillation, BC: Breast Cancer, CKD: Chronic Kidney Disease, CC: Colorectal Cancer, CHD: Coronary Heart Disease, HC: Hypercholesterolemia, PC: Prostate Cancer, EUR: European American ancestry, AFR: African/African American ancestry, AMR: Admixed American ancestry, EAS: East Asian ancestry, SAS: South Asian ancestry

Figure 2. Ancestry enrichments within nine conditions as compared to PMBB participants. Enrichment within prevalent (lighter shade) versus incident (darker shade) cases across the nine adult GIRA conditions is estimated using odds ratios and their standard errors. N_p (N_i) denotes the number of prevalent (incident) cases for each condition among biobank participants aged 18 to 75 at the time of sample collection who were not enrolled in the Penn Medicine Cancer Risk Evaluation Program (N_{total}=48,279). Odds ratios are plotted on the $\log_{10}$ scale. Cutoff bars extend beyond the displayed range and were truncated at x-axis limits of 0.25 and 6 for visualization.

Abbreviations: GIRA: Genome Informed Risk Assessment, PMBB: Penn Medicine Biobank, T2D: Type 2 Diabetes, AFIB: Atrial Fibrillation, BC: Breast Cancer, CKD: Chronic Kidney Disease, CC: Colorectal Cancer, CHD: Coronary Heart Disease, HC: Hypercholesterolemia, PC: Prostate Cancer, EUR: European American ancestry, AFR: African/African American ancestry, AMR: Admixed American ancestry, EAS: East Asian ancestry, SAS: South Asian ancestry

406
Figure 3. GIRA evaluation in PMBB across nine conditions considered by the eMERGE
study. PMBB participants were labeled as high-risk for each condition using the eMERGE
Genome Informed Risk Assessment (GIRA) together with their condition-specific criteria for
each risk indicator. All risk indicators are aggregated into a combined risk score to distinguish
individuals at high genetic risk—based on the GIRA-recommended criteria for PGS, monogenic
risk, and family history—from those without high genetic risk status. Odds/ Hazard ratios were
estimated in a model adjusting for sex, age, age^2, and first ten genotype or exome PCs. For
each hazard ratio model, observation time was defined as the earliest of the following times:
date of first diagnosis, last clinical encounter, or death. Odds ratios were stratified by ancestry
for prevalent cases (left). Hazard ratios for incident cases were compared to prevalent cases for
all PMBB participants, irrespective of ancestry (right).
Abbreviations: PMBB: Penn Medicine Biobank, T2D: Type 2 Diabetes, AFIB: Atrial Fibrillation,
BC: Breast Cancer, CKD: Chronic Kidney Disease, CC: Colorectal Cancer, CHD: Coronary
Heart Disease, HC: Hypercholesterolemia, PC: Prostate Cancer, EUR: European American
ancestry, AFR: African/African American ancestry, AMR: Admixed American ancestry, EAS:
East Asian ancestry, SAS: South Asian ancestry
423
424

425
Figure 4. Cumulative incidence curves with stratification by GIRA high-risk versus the
rest ('Not High Risk'). Observation time was defined as the earliest of the following times: date
of first diagnosis, last follow-up date within a 15 year follow-up window from sample collection.
Participants whose high-risk status for each condition was triggered by one or more of the three
GIRA genetic risk factors were aggregated into a 'High Risk - All' group.
Abbreviations: GIRA: Genome Informed Risk Assessment, T2D: Type 2 Diabetes, AFIB: Atrial
Fibrillation, BC: Breast Cancer, CKD: Chronic Kidney Disease, CC: Colorectal Cancer, CHD:
Coronary Heart Disease, HC: Hypercholesterolemia, PC: Prostate Cancer

Figure 5. Hazard Ratios for GIRA high risk stratified by the three genetic risk components of the GIRA report: Polygenic Risk Score (PGS), Monogenic, and Family History. High Risk - PGS models were adjusted for sex, age, age^2, and first ten genotype PCs. High Risk - Monogenic models were adjusted for sex, age, age^2, and first exome PCs. High Risk - Family History models were adjusted for sex, age, and age^2. For each model, observation time was defined as the earliest of the following times: date of first diagnosis, last clinical encounter, or death. The disease status was 1 if the individual developed the condition and 0 otherwise.

Abbreviations: T2D: Type 2 Diabetes, AFIB: Atrial Fibrillation, BC: Breast Cancer, CKD: Chronic Kidney Disease, CC: Colorectal Cancer, CHD: Coronary Heart Disease, HC: Hypercholesterolemia, PC: Prostate Cancer, EUR: European American ancestry, AFR: African/African American ancestry

Figure 6. Characterization of the PMBB participants labeled as High-Risk using GIRA guidelines for nine conditions. The left panel shows the percentage of unique PMBB participants classified as high-risk based on one or more of the three genetic risk components in the GIRA report. Grey bars indicate the overall proportion of high-risk individuals across all and within each condition, while colored bars show the contribution of each specific component–PGS, monogenic risk, or family history–to high-risk status. The right panel shows the disease status of each high-risk individual across conditions, categorized as case ("Already Have Disease") or control ("Currently Don't Have Disease"). Prevalent cases (lighter bar) are compared to incident cases (darker bar). Individuals whose status could not be definitively determined based on the phenotype algorithm are labeled as "Status Not Ascertained."

Abbreviations: PMBB: Penn Medicine Biobank, GIRA: Genome Informed Risk Assessment, T2D: Type 2 Diabetes, AFIB: Atrial Fibrillation, BC: Breast Cancer, CKD: Chronic Kidney Disease, CC: Colorectal Cancer, CHD: Coronary Heart Disease, HC: Hypercholesterolemia, PC: Prostate Cancer

Figure 7. Ancestry enrichments among high-risk PMBB participants across nine conditions compared with overall PMBB enrichments.

Enrichment within incident (lighter shade) cases versus PMBB participants labeled as high-risk (darker shade) across the nine adult GIRA conditions is estimated using odds ratios and their standard errors. High Risk – All error bar for CC in EAS ancestry individuals is not plotted due to small sample size. Odds ratios are plotted on the $\log_{10}$ scale. Cutoff bars extend beyond the displayed range and were truncated at x-axis limits of 0.25 and 6 for visualization.

Abbreviations: PMBB: Penn Medicine Biobank, GIRA: Genome Informed Risk Assessment, T2D: Type 2 Diabetes, AFIB: Atrial Fibrillation, BC: Breast Cancer, CKD: Chronic Kidney Disease, CC: Colorectal Cancer, CHD: Coronary Heart Disease, HC: Hypercholesterolemia, PC: Prostate Cancer, EUR: European American ancestry, AFR: African/African American ancestry, AMR: Admixed American ancestry, EAS: East Asian ancestry, SAS: South Asian ancestry

Methods

Study population

The Penn Medicine Biobank (PMBB) is an electronic health record linked biobank at the
University of Pennsylvania that integrates an academic health system that serves south-central
Pennsylvania and New Jersey as well as northern Delaware with genomic and exome data[22].
PMBB agnostically recruits individuals from outpatient clinical settings from whom it obtains
consent to their EHR data as well as to collection and storage of biospecimens and genetic
sequencing data. These participants underwent genome and exome sequencing at Regeneron
Genetics Center. Clinical information from 57,170 sequenced individuals, of which
approximately 26.3% of participants are of non-European ancestry, makes PMBB one of the
more diverse medical biobanks. Clinical diagnosis, procedural, visit, laboratory, prescription,
and demographic information was used to define each condition.

Genotype quality control and imputation

Detailed information on genotyping and imputation for Freeze 2.0 can be found in a previous
publication[22]. Briefly, in Freeze 2.0, 43,913 PMBB participants were genotyped using a custom
genotyping array constructed from the Global Screening Array. An additional 13,257 individuals
were genotyped via genotype by sequencing capture as part of Freeze 3.0. To ensure data
quality, SNPs with more than 5% missingness were excluded. Samples with over 10%
missingness or sex discrepancies were also removed using PLINK 2.0. Following genotype
quality control, imputation was conducted using the TOPMed version R3 reference panel via the
Michigan TOPMed Imputation Server 2 with minimac v4.1.6. Phasing was carried out using
Eagle v2.4 and the GRCh38/hg38 genome build. The data were imputed in separate batches,
so the final quality control step involved merging all batches into a single dataset. An OR logic
was then applied to filter variants based on the following criteria: an average R^2 greater than
0.2, sequenced in Freeze 2.0, or sequenced in Freeze 3.0.

Exome sequencing and quality control

Whole exome sequencing (WES) on 57,170 individuals in the Penn Medicine Biobank was
performed at the Regeneron Genetics Center (Tarrytown, NY) as described previously[67]. This
total consists of 43,913 samples from Freeze 2.0 and an additional 13,257 sequenced in 2024
as part of Freeze 3.0. DNA libraries were prepared using a custom NEDNext Ultra II FS DNA
library prep kit to generate fragments at an average size of 200 base pairs. During library
amplification with KAPA HiFi polymerase, a Y-shaped adapter was ligated and asymmetric 10
base pair barcodes were added to enable exome capture and sequencing. DNA samples in
Freeze 2.0 were processed for exome capture on a custom IDT xGen probe library. Freeze 3.0
used the RGC-developed Twist Comprehensive Exome panel for exome capture. Samples from
both freezes were then sequenced on an Illumina NovaSeq 6000 system using S4 flow cells.

Sequencing reads were aligned to the human reference genome build GRCh38 using the
Burrows-Wheeler Aligner - MEM algorithm. To ensure consistency given the differences in
exome capture products, the datasets were reprocessed and jointly recalled across the
combined set of their respective calling regions. Variants were jointly called using GLnexus
v1.4.3 with the DeepVariantWES configuration. Samples with contamination more than 5%, less
than 80% of target bases at 20x coverage, gender discordance, plating errors, duplicates, and
no genotyping array data were removed.

Population stratification

To infer genetically defined ancestry groups, principal component analysis (PCA) was
performed using the *smartpca* module from the EIGENSOFT package. PCA was conducted on
a linkage disequilibrium (LD)-pruned set of 182,701 common SNPs that had undergone quality
control, including removal of SNPs that failed Hardy-Weinberg equilibrium test (p < 1e-6), had a
minor allele frequency below 0.1, and related individuals. The study population was assigned to
genetically inferred ancestry (GIA) groups - European American (EUR), Admixed American
(AMR), African/African American (AFR), East Asian (EAS), and South Asian (SAS) – using
quantitative discriminant analysis (QDA) based on the top 20 principal components. The
continental ancestry populations from 1000 Genomes Project were used as a reference. QDA
assigned each PMBB individual to the closest ancestry group based on genetic similarity in
principal component space.

For ancestry-specific analysis, participants with complete information on the defined condition
and covariates were stratified by European American and African American ancestry groups,
which had sufficient sample sizes for analysis.

Phenotype definition

We adapted validated electronic phenotype algorithms from the Phenotype KnowledgeBase
(PheKB) to define case and control status for four conditions – Breast Cancer,
Hypercholesterolemia, Colorectal Cancer, and Chronic Kidney Disease. PheKb is a
collaborative online environment created by the eMERGE network to serve as a central
repository for the development, shareability, and validity of algorithms using health care
generated data across implementation sites.[71] Details on the phenotyping algorithms are
provided in the **Supplementary Methods**. We adapted the chronic kidney disease algorithm to
focus on cases that had an estimated glomerular filtration rate (eGFR) below 60 ml/min/1.73m2
or received a dialysis or kidney transplant without the incorporation of staging CKD based on
albuminuria or eGFR.[63,72]

Atrial fibrillation (AFIB) algorithm was adapted from ICD-9 and 10 diagnosis code-based case
definition outlined in the Chronic Conditions Data Warehouse (CCW). The CCW algorithms
were developed from claims data using research literature, expert panels, and criteria used by
other federal sources and have been previously applied as phenotyping methods to EHR data
to ascertain chronic disease status.[73] For an individual to be classified as having AFIB based on
the CCW algorithm, they must have at least 1 inpatient claim or 2 hospital outpatient claims with
diagnosis codes that are at least 30 days apart but within 1 year. A previous study found 1
inpatient or 2 outpatient codes separated by >30 days but within 1 year had a PPV of 78.7%
and sensitivity of 92.5% in their prevalent approach while their incident approach had a PVV of
69.7% and sensitivity of 93.6%.[74]

Coronary Heart Disease, Prostate Cancer, and Obesity algorithms were adapted from the
Polygenic Risk Methods in Diverse Populations (PRIMED) Consortium. Lastly, we implemented
the most up to date phenotyping algorithm for type 2 diabetes from the Szczerbinski et al.
study[75]. This study developed and validated a phenotyping algorithm for type 2 diabetes,
modified from the Northwestern University algorithm in PheKB[76], in the All of Us cohort using

electronic health records as well as a universal algorithm to identify individuals without
diabetes.[75]
565
Case or control status for breast and prostate cancer were assigned to only females and males,
respectively. For each of the conditions, we considered the earliest date at which the diagnosis
was ascertained as the diagnosis date. All participants diagnosed at dates prior to, at, or post
sample collection (baseline) in PMBB were considered prevalent, while participants diagnosed
after sample collection were considered incident. Additional details on the ascertainment of
each condition are provided in the **Supplementary Methods**.
572
Monogenic risk score classification
Genetic variants were annotated using Ensembl's Variant Effect Predictor (VEP) version 113.0[77]
to access population allele frequencies from the gnomAD Genome Aggregation Database,
clinical significance as reported by ClinVar, and predicted gene impact. The analysis focused on
rare, high-quality variants located within protein-coding regions of 16 genes associated with the
six monogenic conditions included in the GIRA study: atrial fibrillation (*LMNA*); breast cancer
(*BRCA1, BRCA2, PALB2, PTEN, TP53, STK11*); colorectal cancer (*EPCAM, MLH1, MSH2,*
MSH6, PMS2, STK11, PTEN, TP53); coronary heart disease (*APOB, LDLR, LDLRAP1,*
PCSK9); hypercholesterolemia (*APOB, LDLR, LDLRAP1, PCSK9*); and prostate cancer
(*BRCA1, BRCA2, EPCAM, MLH1, MSH2, MSH6, PMS2*). Variants with a minor allele frequency
(MAF) greater than 0.005 in any of the following gnomAD racial subpopulations—Non-Finnish
European, African, or Latino—were excluded from the analysis as well as synonymous variants.
We did not consider superpopulation maximum allele frequencies. We further selected variants
classified in ClinVar with designations indicating pathogenicity, including *Pathogenic*,
Pathogenic/Likely pathogenic, *Likely pathogenic*, *Pathogenic/Likely pathogenic/Pathogenic, low*
penetrance, *Pathogenic/Likely pathogenic | risk factor*, *Pathogenic | association | protective*,
Likely pathogenic | risk factor, *Pathogenic/Likely pathogenic | other*, and *Pathogenic | other*.
Only variants submitted by multiple contributors or reviewed by an expert panel were retained.
591
Polygenic risk score derivation and ancestry calibration
To ensure our validation of GIRA's high-risk criteria in PMBB closely mirrored the original
implementation decisions made in the eMERGE study, we used the same polygenic risk scores
that were employed in their study. Each condition's PGS was calculated by multiplying individual
risk allele dosages by their corresponding weights that were downloaded from the eMERGE
GitHub repository (https://github.com/broadinstitute/eMERGE-implemented-PRS-models-
Lennon-et-al) using pgsc_calc. These scores are also available on the UCSC Genome Browser
(https://genome.ucsc.edu/s/Max/emerge), and PGS Catalog IDs for most scores are detailed in
the Lennon et al. publication[29]. As the SNP weights used for obesity in the GIRA study were not
publicly available, we evaluated the performance of all published obesity PGS from the PGS
Catalog within the All of Us Research Program (AoU) to identify the best-performing score[74].
Within all participants of AoU v7, we applied each published obesity PGS, then standardized
and residualized out ancestry PCs from each PGS as described previously[78]. We then
estimated the area under the receiver operator curve (AUC) for logistic regression models when
predicting obesity status based on the PGS, adjusting for age, sex, 10 PCs, and sequencing

site. The obesity PGS with the highest AUC was selected for downstream analyses within PMBB.

Supplementary Table 14 reports the number of single nucleotide polymorphisms (SNPs) in the base file for each published PGS and the proportion available in imputed PMBB data, showing ≥75% overlap depending on the PGS. Performance of each PGS in PMBB was consistent with the odds ratios per standard deviation reported by eMERGE.

To achieve comparable score distributions across ancestry groups, the raw PGSs were calibrated using genetic principal components, implemented through the –run_ancestry method in pgsc_calc. In brief, this adjustment accounts for differences in the mean and variance of PGS distributions between ancestries, following the calibration of ancestry for PGS approach used in the eMERGE Network's GIRA report. As a result, similar proportions of individuals in each ancestry group fall above a given percentile threshold. For each condition, the GIRA report defined ancestry-independent thresholds above which individuals were classified as "high risk." These thresholds differed by condition. After calibrating the PGS for ancestry, scores were converted into a categorical variable to distinguish individuals at high polygenic risk, based on the GIRA-recommended percentile cutoff, from those below the threshold.

Family history classification
Family history (FamHx) information from the EHR was used to identify individuals with a family history for five GIRA conditions where family history is considered a risk factor. To identify FamHx of specific conditions, we extracted the following FamHx information:
- Atrial fibrillation: Individuals whose parents had a history of 'Atrial fibrillation'
- Breast cancer: Individuals whose first and second-degree relatives had a history of 'breast cancer', 'Bilateral breast cancer', 'Postmenopausal breast cancer', 'Premenopausal breast cancer', 'Left Breast Cancer', and 'Right Breast Cancer
- Chronic kidney disease: Individuals whose first-degree relatives had a history of 'Kidney Disease', 'Kidney Disorder', 'Chronic Kidney Disease', and 'Polycystic Kidney Disease'
- Coronary heart disease: Individuals whose first-degree relatives had a history of 'Heart Disease'
- Prostate cancer: Individuals whose first-degree male relatives had a history of 'Prostate Cancer'

First-degree relatives included an individual's parents, siblings, and children. Second-degree relatives included an individual's grandparents, grandchildren, uncles, aunts, nephews, nieces, and half-siblings.
Monogenic risk and family history components of the GIRA report could not be validated, as eMERGE did not release these corresponding results.

Statistical Analysis
All analysis used Python 3 or R 4.2.1.

To align our population with the eMERGE GIRA study, we restricted our analysis to adults aged 18 to 75 at enrollment. We also excluded individuals enrolled in the Penn Medicine Cancer Risk Evaluation Program, as their increased genetic cancer risk could bias results for cancer-related conditions. Logistic regression models, both across and stratified by ancestry, were used to

estimate associations with high-risk status for each condition, adjusting for the first ten genotype
or exome principal components, age, age², and sex. Models were run separately for each GIRA
risk component, PGS, monogenic risk, and family history, as well as for a combined risk score,
labeled as 'High Risk - All', aggregating all high-risk indicators to distinguish individuals at high
genetic risk from those without. We focused on the odds ratios for each high-risk component in
EUR and AFR ancestry groups, given the limited sample sizes of other ancestry groups across
conditions.

Cox proportional hazard models were performed for each condition to estimate hazard ratios
comparing those at high-risk based on each GIRA risk component or the combined risk score
with those not at high risk. We report hazard ratios for EUR and AFR groups for each high-risk
component. For each hazard ratio model, observation time was defined as the earliest of the
following times: date of first diagnosis, last clinical encounter, or death. The disease status was
1 if the individual developed the condition and 0 otherwise. We estimate time-dependent
disease incidences for each condition stratified by high-risk status versus those not at high-risk
using 1 minus the Kaplan-Meier estimator. Observation time was defined as the earliest of the
following times: date of first diagnosis, last follow-up date within a 15-year follow-up window
from the time of sample collection, or death. High-risk status was assigned to participants with
at least one of the three GIRA genetic risk indicators.

Odds ratios and 95% confidence intervals for ancestry, age, or sex enrichment were estimated
using logistic regression models comparing individuals in one context group to those in all the
other context groups for both prevalent and incidence cases. Ancestry enrichment models were
adjusted for age, age², sex, and batch. Age enrichment models were adjusted for sex and
batch. Sex enrichment models were adjusted for age, age², and batch. The same models were
repeated to assess enrichment among high-risk PMBB participants across nine conditions.
Since we only consider females or males for breast cancer and prostate cancer, respectively,
we don't estimate the odds ratios for sex enrichment.
675
To compare high-risk proportions across ancestry strata and social deprivation index (SDI)
quartiles with those observed in the overall PMBB population, we performed two-proportion z-
tests to evaluate whether high-risk percentages in specific subpopulations differed significantly
from the overall high-risk percentage. Ancestry-specific high-risk proportions were calculated by
determining the proportion classified as 'High Risk – All' and the high-risk proportion for each
genetic risk component within each ancestry stratum. P values were adjusted for multiple testing
across 24 comparisons (four high-risk categories across six ancestry strata). Ancestry
enrichment for each high-risk categories was also estimated using logistic regression models
comparing individuals in one ancestry strata to all other ancestries (**Table S16**). The models
were adjusted for age, age², sex, and batch. SDI scores were categorized into quartiles. High-
risk proportions within each SDI quartile were calculated using the same approach as for
ancestry strata. P values were adjusted for multiple testing across 16 comparisons (four high-
risk categories across four SDI quartiles).
689

690 In secondary analysis, we excluded individuals who were lost during follow-up. Individuals not
691 lost during follow-up were defined as those who had at least one encounter in the two years
692 prior to the last encounter date.
693
694 **Data Availability**
695 Researchers can access the Penn Medicine Biobank individual-level genotype, exome, and
696 phenotype data following its data application procedures.
697
Acknowledgements

We acknowledge the Penn Medicine BioBank (PMBB) for providing data and thank the patient-
participants of Penn Medicine who consented to participate in this research program. We would
also like to thank the Penn Medicine BioBank team and Regeneron Genetics Center for
providing genetic variant data for analysis. The PMBB is approved under IRB protocol# 813913
and supported by Perelman School of Medicine at University of Pennsylvania, a gift from the
Smilow family, and the National Center for Advancing Translational Sciences of the National
Institutes of Health under CTSA award number UL1TR001878.

References

1. Returning integrated genomic risk and clinical recommendations: The eMERGE study.

Genetics in Medicine **25**, 100006 (2023).

2. Daly, M. B. *et al.* NCCN Guidelines Insights: Genetic/Familial High-Risk Assessment:

Breast, ovarian, and pancreatic, version 1.2020. *J. Natl. Compr. Canc. Netw.* **18**, 380–391

(2020).

3. Gupta, S. *et al.* NCCN Guidelines Insights: Genetic/Familial High-Risk Assessment:

Colorectal, version 3.2017. *J. Natl. Compr. Canc. Netw.* **15**, 1465–1475 (2017).

4. Khera, A. V. *et al.* Genome-wide polygenic scores for common diseases identify individuals

with risk equivalent to monogenic mutations. *Nature Genetics* **50**, 1219–1224 (2018).

5. Abul-Husn, N. S. *et al.* Genetic identification of familial hypercholesterolemia within a single

U.S. health care system. *Science* **354**, aaf7000 (2016).

6. Mars, N. *et al.* Systematic comparison of family history and polygenic risk across 24

common diseases. *American journal of human genetics* **109**, (2022).

7. Lu, T., Forgetta, V., Richards, J. B. & Greenwood, C. M. T. Capturing additional genetic risk

from family history for improved polygenic risk prediction. *Commun. Biol.* **5**, 595 (2022).

8. Hujoel, M. L. A., Gazal, S., Loh, P.-R., Patterson, N. & Price, A. L. Liability threshold

modeling of case-control status and family history of disease increases association power.

Nat. Genet. **52**, 541–547 (2020).

9. Guttmacher, A. E., Collins, F. S. & Carmona, R. H. The family history--more important than

ever. *N. Engl. J. Med.* **351**, 2333–2336 (2004).

10. Valdez, R., Yoon, P. W., Qureshi, N., Green, R. F. & Khoury, M. J. Family history in public

health practice: a genomic tool for disease prevention and health promotion. *Annu. Rev.*

Public Health **31**, 69-87 1 p following 87 (2010).

11. Conti, D. V. *et al.* Trans-ancestry genome-wide association meta-analysis of prostate

cancer identifies new susceptibility loci and informs genetic risk prediction. *Nat. Genet.* **53**,

65–75 (2021).

12. Ge, T. *et al.* Development and validation of a trans-ancestry polygenic risk score for type 2

diabetes in diverse populations. *Genome Med.* **14**, 70 (2022).

13. Fahed, A. C. *et al.* Polygenic background modifies penetrance of monogenic variants for

tier 1 genomic conditions. *Nat Commun* **11**, 3635 (2020).

14. Murray Leech, J. *et al.* Common genetic variants modify disease risk and clinical

presentation in monogenic diabetes. *Nat. Metab.* **7**, 1819–1829 (2025).

15. Saadatagah, S. *et al.* Polygenic Risk, Rare Variants, and Family History: Independent and

Additive Effects on Coronary Heart Disease. *JACC Adv* **2**, 100567 (2023).

16. McCarty, C. A. *et al.* The eMERGE Network: A consortium of biorepositories linked to

electronic medical records data for conducting genomic studies. *BMC Medical Genomics* **4**,

1–11 (2011).

17. The Electronic Medical Records and Genomics (eMERGE) Network: past, present, and

future. *Genetics in Medicine* **15**, 761–771 (2013).

18. Lennon, N. J. *et al.* Selection, optimization and validation of ten chronic disease polygenic

risk scores for clinical implementation in diverse US populations. *Nature Medicine* **30**, 480–

487 (2024).

19. Kullo, I. J. Clinical use of polygenic risk scores: current status, barriers and future

750 directions. *Nature Reviews Genetics* 1–18 (2025).

20. Collins, G. S., Reitsma, J. B., Altman, D. G. & Moons, K. G. M. Transparent Reporting of a

752 multivariable prediction model for Individual Prognosis or Diagnosis (TRIPOD): the TRIPOD

753 statement. *Ann. Intern. Med.* **162**, 55–63 (2015).

21. Xiang, R. *et al.* Recent advances in polygenic scores: translation, equitability, methods and

755 FAIR tools. *Genome Med.* **16**, 33 (2024).

22. Verma, A. *et al.* The Penn Medicine BioBank: Towards a Genomics-Enabled Learning

757 Healthcare System to Accelerate Precision Medicine in a Diverse Population. *Journal of*

758 *Personalized Medicine* **12**, 1974 (2022).

23. Pfeiffer, R. M. & Gail, M. H. Two criteria for evaluating risk prediction models. *Biometrics*

760 **67**, 1057–1065 (2011).

24. McHugh, J. K. *et al.* Assessment of a polygenic risk score in screening for prostate cancer.

762 *N. Engl. J. Med.* **392**, 1406–1417 (2025).

25. Chatterjee, N., Shi, J. & García-Closas, M. Developing and evaluating polygenic risk

764 prediction models for stratified disease prevention. *Nat. Rev. Genet.* **17**, 392–406 (2016).

26. Kraft, P. *et al.* Beyond odds ratios--communicating disease risk based on genetic profiles.

766 *Nat. Rev. Genet.* **10**, 264–269 (2009).

27. Elliott, J. *et al.* Predictive accuracy of a polygenic risk score-enhanced prediction model vs

768 a clinical risk score for coronary artery disease. *JAMA* **323**, 636–645 (2020).

28. Haas, R. *et al.* Diverse genomes, shared health: Insights from a health system biobank.

770 *medRxiv* (2025) doi:10.1101/2025.06.11.25329386.

29. All of Us Research Program Genomics Investigators. Genomic data in the All of Us

Research Program. *Nature* **627**, 340–346 (2024).

30. Piekos, J. A. *et al.* Evaluating the relationships between genetic ancestry and the clinical

phenome. *Pac. Symp. Biocomput.* **29**, 389–403 (2024).

31. Heart disease prevalence. https://www.cdc.gov/nchs/hus/topics/heart-disease-

prevalence.htm (2024).

32. Khan, M. A. B. *et al.* Epidemiology of type 2 diabetes - Global Burden of Disease and

forecasted trends. *J. Epidemiol. Glob. Health* **10**, 107–111 (2020).

33. https://jamanetwork.com/journals/jamanetworkopen/fullarticle/2801355.

34. CDC. Chronic kidney disease in the United States, 2023. *Chronic Kidney Disease*

https://www.cdc.gov/kidney-disease/php/data-research/index.html (2025).

35. van Alten, S., Domingue, B. W., Faul, J., Galama, T. & Marees, A. T. Reweighting UK

Biobank corrects for pervasive selection bias due to volunteering. *Int. J. Epidemiol.* **53**,

(2024).

36. Pimplaskar, A. *et al.* Inclusion bias affects common variant discovery and replication in a

health-system linked biobank. *medRxiv* (2025) doi:10.1101/2025.04.04.25325131.

37. Cancer of the breast (female) - cancer stat facts. *SEER*

https://seer.cancer.gov/statfacts/html/breast.html.

38. Murphy, C. C., Wallace, K., Sandler, R. S. & Baron, J. A. Racial disparities in incidence of

young-onset colorectal cancer and patient survival. *Gastroenterology* **156**, 958–965 (2019).

39. Ng, M. C. Y. Genetics of type 2 diabetes in African Americans. *Curr. Diab. Rep.* **15**, 74

(2015).

40. Rostand, S. G., Kirk, K. A., Rutsky, E. A. & Pate, B. A. Racial differences in the incidence of

treatment for end-stage renal disease. *N. Engl. J. Med.* **306**, 1276–1279 (1982).

41. https://jamanetwork.com/journals/jamanetworkopen/fullarticle/2788170.

42. https://jamanetwork.com/journals/jamanetworkopen/fullarticle/2835519.

43. Rodosthenous, R. S. *et al.* Recontacting biobank participants to collect lifestyle, behavioural

and cognitive information via online questionnaires: lessons from a pilot study within

FinnGen. *BMJ Open* **12**, e064695 (2022).

44. Gajda, M., Kowalska, M. & Zejda, J. E. Impact of two different recruitment procedures

(random vs. Volunteer selection) on the results of seroepidemiological study (SARS-CoV-

2). *Int. J. Environ. Res. Public Health* **18**, 9928 (2021).

45. Fry, A. *et al.* Comparison of sociodemographic and health-related characteristics of UK

Biobank participants with those of the general population. *Am. J. Epidemiol.* **186**, 1026–

1034 (2017).

46. Kathiresan, N. *et al.* Representation of race and ethnicity in the contemporary US health

cohort all of Us Research Program. *JAMA Cardiol.* **8**, 859–864 (2023).

47. Divo, M. J., Martinez, C. H. & Mannino, D. M. Ageing and the epidemiology of

multimorbidity. *Eur. Respir. J.* **44**, 1055–1068 (2014).

48. Le Couteur, D. G. & Thillainadesan, J. What is an aging-related disease? An

epidemiological perspective. *J. Gerontol. A Biol. Sci. Med. Sci.* **77**, 2168–2174 (2022).

49. Mt, 63% N. H. V. T. Colorectal cancer screening* (%), adults 45 years and older by state,

2020. https://www.cancer.org/content/dam/cancer-org/research/cancer-facts-and-

statistics/colorectal-cancer-facts-and-figures/colorectal-cancer-facts-and-figures-2023.pdf.

50. Dai, H. *et al.* Global, regional, and national prevalence, incidence, mortality, and risk factors

for atrial fibrillation, 1990-2017: results from the Global Burden of Disease Study 2017. *Eur.*

Heart J. Qual. Care Clin. Outcomes **7**, 574–582 (2021).

51. Xi, Y. & Xu, P. Global colorectal cancer burden in 2020 and projections to 2040. *Transl.*

Oncol. **14**, 101174 (2021).

52. Gwira, J. A., Fryar, C. D. & Gu, Q. Prevalence of total, diagnosed, and undiagnosed

diabetes in adults: United States, August 2021-August 2023. *NCHS Data Brief* (2024)

doi:10.15620/cdc/165794.

53. Rodgers, J. L. *et al.* Cardiovascular risks associated with gender and aging. *J. Cardiovasc.*

Dev. Dis. **6**, 19 (2019).

54. Shapira, N. Women's higher health risks in the obesogenic environment: a gender nutrition

approach to metabolic dimorphism with predictive, preventive, and personalised medicine.

EPMA J. **4**, 1 (2013).

55. Xie, K. *et al.* Global, regional, and national burden of chronic kidney disease, 1990-2021: a

systematic analysis for the global burden of disease study 2021. *Front. Endocrinol.*

(Lausanne) **16**, 1526482 (2025).

56. Chesnaye, N. C., Carrero, J. J., Hecking, M. & Jager, K. J. Differences in the epidemiology,

management and outcomes of kidney disease in men and women. *Nat. Rev. Nephrol.* **20**,

7–20 (2024).

57. Khan, A. *et al.* Genome-wide polygenic score to predict chronic kidney disease across

ancestries. *Nat. Med.* **28**, 1412–1420 (2022).

58. Martin, A. R. *et al.* Clinical use of current polygenic risk scores may exacerbate health

disparities. *Nat. Genet.* **51**, 584–591 (2019).

59. Managing differential performance of polygenic risk scores across groups: Real-world

experience of the eMERGE Network. *The American Journal of Human Genetics* **111**, 999–

1005 (2024).

60. Baptista, P. V. Principles in genetic risk assessment. *Ther. Clin. Risk Manag.* **1**, 15–20

(2005).

61. Wang, J. Y. *et al.* Three open questions in polygenic score portability. *Nat. Commun.* **17**,

942 (2026).

62. Hughes Halbert, C. *et al.* Social determinants of family health history collection. *J.*

Community Genet. **7**, 57–64 (2016).

63. Khan, A. *et al.* Polygenic risk alters the penetrance of monogenic kidney disease. *Nature*

Communications **14**, 1–10 (2023).

64. Rashmi, R. & Mohanty, S. K. Examining chronic disease onset across varying age groups

of Indian adults using competing risk analysis. *Sci. Rep.* **13**, 5848 (2023).

65. Breast cancer risk in American women. https://www.cancer.gov/types/breast/risk-fact-sheet

(2020).

66. Vassy, J. L. *et al.* The GenoVA study: Equitable implementation of a pragmatic randomized

trial of polygenic-risk scoring in primary care. *Am. J. Hum. Genet.* **110**, 1841–1852 (2023).

67. Hao, L. *et al.* Development of a clinical polygenic risk score assay and reporting workflow.

Nature Medicine **28**, 1006–1013 (2022).

68. Ge, T. *et al.* Validation of a trans-ancestry polygenic risk score for type 2 diabetes in

diverse populations. *medRxiv* (2021) doi:10.1101/2021.09.11.21263413.

69. Conway-Pearson, L. S. *et al.* Family health history reporting is sensitive to small changes in

wording. *Genet. Med.* **18**, 1308–1311 (2016).

70. Wilson, B. J. Systematic review: Family history in risk assessment for common diseases.

Ann. Intern. Med. **151**, 878 (2009).

71. Kirby, J. C. *et al.* PheKB: a catalog and workflow for creating electronic phenotype

algorithms for transportability. *Journal of the American Medical Informatics Association :*

JAMIA **23**, 1046 (2016).

72. Shang, N. *et al.* Medical records-based chronic kidney disease phenotype for clinical care

and "big data" observational and genetic studies. *npj Digital Medicine* **4**, 1–13 (2021).

73. Voss, R. W. *et al.* Comparing ascertainment of chronic condition status with problem lists

versus encounter diagnoses from electronic health records. *Journal of the American*

Medical Informatics Association : JAMIA **29**, 770 (2022).

74. Chamberlain, A. M. *et al.* Identification of Incident Atrial Fibrillation From Electronic Medical

Records. *Journal of the American Heart Association* (2022) doi:10.1161/JAHA.121.023237.

75. Szczerbinski, L. *et al.* Algorithms for the identification of prevalent diabetes in the All of Us

Research Program validated using polygenic scores. *Scientific Reports* **14**, 1–11 (2024).

76. Type 2 Diabetes Mellitus. https://phekb.org/phenotype/type-2-diabetes-mellitus.

77. McLaren, W. *et al.* The Ensembl Variant Effect Predictor. *Genome Biology* **17**, 1–14 (2016).

78. Khera, A. V. *et al.* Whole-genome sequencing to characterize monogenic and polygenic

contributions in patients hospitalized with early-onset myocardial infarction. *Circulation* **139**,

1593–1602 (2019).

880