Remote Patient Monitoring in Heart Failure: A Systematic Review, Meta-Analysis, and Trial Sequential Analysis

Title: Remote Patient Monitoring in Heart Failure: A Systematic Review, Meta-Analysis, and Trial Sequential Analysis

Authors: Vicky Muller Ferreira, MD[1]; Victor Ayres Muller, MD[1]

[1]Independent Researcher, Rio de Janeiro, Brazil

Corresponding Author: Vicky Muller Ferreira, MD. Email:

vickymuller@heartlandprotocol.org

Word Count: ~6,000

Keywords: remote patient monitoring; heart failure; telemonitoring; meta-analysis; trial sequential analysis

Registration: PROSPERO CRD420261323628

Abstract

Whether the cumulative evidence for remote patient monitoring (RPM) in heart failure (HF) is robust to sequential monitoring, and whether trials report geographic access modifiers, remains uncertain. We conducted a systematic review, meta-analysis, and trial sequential analysis (TSA) of 65 RCTs (59 poolable; ~23,000 participants) across four databases through February 2026, encompassing structured telephone support (15 trials), non-invasive telemonitoring (33), and invasive hemodynamic monitoring (11). Random-effects meta-analysis used REML with Hartung-Knapp-Sidik-Jonkman adjustment. RPM significantly reduced all-cause mortality (RR 0.911, 95% CI 0.842–0.985; P=0.021; I^2=0%; k=41; NNT 104/year; prediction interval 0.840–0.988). TSA suggested that accrued evidence exceeded the required information size under the 15% relative risk reduction assumption, supporting a stable mortality signal. HF hospitalization was reduced (RR 0.781, 95% CI 0.710–0.859; P<0.001; k=39; NNT 18/year), though the prediction interval crossed 1.0 (0.586–1.040), indicating that in some settings the effect may be attenuated. No statistically significant interaction by RPM type was detected (all-cause mortality $P_{interaction}$=0.80; HF hospitalization $P_{interaction}$=0.14). GRADE certainty was moderate for mortality and low for HF hospitalization. A descriptive geographic access analysis revealed that only 2 of 59 poolable trials reported formal rural/urban subgroups, precluding conclusions about whether RPM differentially benefits underserved populations.

Highlights

- Trial sequential analysis supports a stable RPM mortality signal under prespecified assumptions

- All-cause mortality reduced 9% (NNT 104/yr, prediction interval narrowly excludes null)

- HF hospitalization reduced 22% (NNT 18/yr), though prediction interval crosses 1.0

- Large-trial sensitivity remained significant but attenuated for HF hospitalization (RR 0.837)

- Only 2 of 59 poolable trials reported formal rural/urban subgroups

Introduction

Heart failure (HF) affects over 64 million people worldwide and remains a leading cause of hospitalization, disability, and death [1]. Despite advances in pharmacological therapy, HF hospitalization rates remain high, with 30-day readmission rates exceeding 20% in many health systems [2]. The burden is disproportionately borne by patients in rural and underserved areas, where access to specialized HF care is limited [3].

Remote patient monitoring (RPM) encompasses a spectrum of technologies designed to extend clinical oversight beyond the hospital or clinic setting. Three broad categories have been studied in HF: structured telephone support (STS), in which nurses or automated systems contact patients at regular intervals to assess symptoms and guide self-management; non-invasive telemonitoring (TM), which involves the electronic transmission of physiological data (weight, blood pressure, heart rate, oxygen saturation) from the patient's home to a clinical team; and invasive hemodynamic monitoring, using implanted sensors (such as the CardioMEMS pulmonary artery pressure sensor) or cardiac implantable electronic device (CIED) algorithms to detect congestion before clinical decompensation.

The evidence base for RPM in HF has grown substantially over the past two decades. The 2015 Cochrane review by Inglis et al. established that both STS and non-invasive TM reduced all-cause mortality compared with usual care, with mortality risk ratios of 0.87 for STS and 0.80 for non-invasive TM [4]. Since then, several landmark trials have been published. The TIM-HF2 trial demonstrated that structured remote management reduced the percentage of days lost to unplanned cardiovascular hospitalization or death [5], while the CHAMPION and MONITOR-HF trials showed that hemodynamic-guided management with the CardioMEMS sensor reduced

HF hospitalizations [6,7]. The GUIDE-HF trial, however, yielded neutral overall results, with benefit confined to a pre-COVID sensitivity analysis [8]. The recently published PROACTIVE-HF trial expanded invasive monitoring to HFpEF [9], and the MIGHTY-HEART trial tested a smartphone-based coaching intervention in a diverse urban population, though its primary composite endpoint was neutral [10].

Despite this growing evidence, important questions remain unanswered. First, no prior meta-analysis has applied trial sequential analysis (TSA) to evaluate whether the cumulative evidence for RPM is robust to repeated testing as trials accrue. Second, the most recent comprehensive meta-analysis predates several important trials (PROACTIVE-HF, MIGHTY-HEART, PRADOC, MESSAGE-HF, MONITOR-HF). Third, whether the benefits of RPM extend to geographically underserved populations — where RPM could theoretically provide the greatest incremental value — has not been systematically examined, despite evidence from TIM-HF2 suggesting that patients farther from cardiologists derive greater benefit [5]. Fourth, HF phenotype-specific data (HFrEF, HFpEF, HFmrEF) remain limited, as most trials enrolled mixed populations.

We therefore conducted a systematic review, meta-analysis, and TSA to evaluate the effects of RPM on mortality, hospitalization, and patient-reported outcomes in adults with HF, with the primary objectives of (1) assessing whether the cumulative evidence is robust to sequential monitoring and (2) describing geographic access reporting and exploratory effect modification.

Materials and Methods

Protocol and Registration

This systematic review was registered with PROSPERO (CRD420261323628) and conducted in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2020 guidelines [11]. Searches were completed before registration; this timing and subsequent protocol amendments are documented in the online data repository (DOI: 10.5281/zenodo.19865261). The completed PRISMA 2020 checklist is available in the same repository.

Eligibility Criteria

We included randomized controlled trials that enrolled adults (age $\geq$18 years) with an established diagnosis of HF (any LVEF phenotype: HFrEF $\leq$40%, HFmrEF 41–49%, or HFpEF $\geq$50%), randomized to any form of RPM (structured telephone support, non-invasive telemonitoring, or invasive hemodynamic monitoring) versus usual care or standard care, with a minimum follow-up of 3 months, reporting at least one outcome of interest (all-cause mortality, HF hospitalization, all-cause hospitalization, cardiovascular mortality, composite outcomes, emergency department visits, or quality of life). Exclusion criteria included observational studies, single-arm trials, studies with fewer than 20 participants, conference abstracts without a full publication, and studies where the RPM intervention could not be isolated from other components (e.g., bundled cardiac rehabilitation programs).

Information Sources and Search Strategy

We searched PubMed/MEDLINE, Cochrane CENTRAL, ClinicalTrials.gov, and the WHO International Clinical Trials Registry Platform (ICTRP) from inception through February 15, 2026, without language restrictions. The search strategy combined Medical Subject Headings (MeSH) and free-text terms for remote monitoring technologies (including telemonitoring,

telehealth, structured telephone support, CardioMEMS, home monitoring, mHealth, wearable devices) and heart failure, with an RCT filter. The full electronic search strategies for all four databases are provided in eTable 1 of the Supplementary Material. The protocol was registered with English-language restriction; however, the search was subsequently conducted without language restriction to maximize sensitivity (protocol amendment #6). Embase and CINAHL were not searched because institutional access was unavailable. To partially mitigate this limitation, we relied on Cochrane CENTRAL, which aggregates records from multiple sources including Embase: 42.9% of the 2,138 CENTRAL records retrieved in our search carried an Embase identifier. PubMed combined with CENTRAL captures approximately 97% of relevant RCTs in therapeutic systematic reviews [12,13]. A post-hoc cross-referencing analysis against three contemporary meta-analyses that searched additional databases [14–16] was conducted to assess the potential impact of this omission (eText 1).

Study Selection

Records were imported, combined, and deduplicated using a multi-pass approach (exact DOI/PMID matching, normalized title matching, and fuzzy matching with manual adjudication). Title and abstract screening used a semi-automated approach with predefined exclusion rules validated against a reference set of known eligible trials. Records flagged by the algorithm (n=219, 8.3%) underwent manual adjudication with final consensus decisions retained in the screening dataset. A post-hoc sensitivity audit of the screening algorithm identified two false negatives due to missing search terms (telemanagement, CHF), which were rescued and included. Records passing initial screening underwent a structured audit to exclude registry-only entries, protocols, conference abstracts, and non-RCT designs. All remaining records were assessed at full-text level using structured eligibility tiers. The archived screening datasets

preserve final consensus decisions rather than complete independent reviewer-level decisions; therefore, the process should be interpreted as semi-automated screening with manual adjudication and audit rather than fully independent dual screening. The PRISMA 2020 flow diagram is presented in Figure 1.

Data Extraction

Data were extracted using a two-reviewer process. One reviewer (VMF) performed the initial structured extraction with AI-assisted tabulation support, and the second reviewer (VAM) independently reviewed the extracted data against source reports; discrepancies were resolved by consensus before analysis. The archived extraction dataset reports the final consensus values used for analysis; reviewer-level intermediate disagreement fields were not retained for every study. Quality checks identified and corrected three errors during the consensus process: MONITOR-HF HF hospitalization HR (0.82 corrected to 0.56, verified against original publication Table 2), CHAMPION PMID, and the CO-0056 mortality denominator (corrected from the pooled intervention denominator to the telephone arm denominator). Extracted effect estimates were checked against original publication tables and supplementary materials where available; extraction notes and a consensus log documenting reviewer roles, correction flags, and final verification status for all 65 studies are available in the Zenodo repository. Data were extracted using a standardized form encompassing 185 variables, including study characteristics, participant demographics, intervention details, geographic context (rural/urban classification, distance to care), outcomes (event counts for binary outcomes; means and standard deviations for continuous outcomes; hazard ratios with 95% confidence intervals when reported), and risk of bias domains. RPM interventions were classified into three pre-specified categories: structured telephone support (STS), non-invasive telemonitoring, and invasive hemodynamic monitoring.

Risk of Bias Assessment

Risk of bias was assessed using the Cochrane Risk of Bias 2 (RoB 2) tool [17] across five domains: randomization process (D1), deviations from intended interventions (D2), missing outcome data (D3), measurement of the outcome (D4), and selection of the reported result (D5). All RPM trials are inherently open-label for participants and clinical staff; therefore, D2 ratings reflect whether knowledge of group assignment could have influenced co-interventions or adherence beyond the intended effect.

Effect Measures and Statistical Analysis

The primary outcomes were all-cause mortality, HF hospitalization, and all-cause hospitalization. Secondary outcomes included cardiovascular mortality, composite of all-cause mortality and hospitalization, emergency department visits, and quality of life (Minnesota Living with Heart Failure Questionnaire [MLHFQ] and Kansas City Cardiomyopathy Questionnaire [KCCQ]).

For binary outcomes with event counts, risk ratios (RR) were calculated using the Mantel-Haenszel method. Rows in which event counts represented recurrent events or episodes rather than participants with events (events exceeding denominators) were excluded from binary RR pooling and from control event rate/NNT calculations. Studies reporting only hazard ratios (HR) with confidence intervals — without extractable event counts — were not included in the primary RR analysis but were pooled separately using the generic inverse-variance method with log-transformed HRs and corresponding standard errors. This yielded two parallel analyses for each primary outcome: a primary RR analysis maximizing the number of contributing studies with valid participant-level event counts (e.g., k=41 for all-cause mortality) and a sensitivity HR-only

analysis restricted to time-to-event data (e.g., k=16 for all-cause mortality). Continuous outcomes were pooled as mean differences (MD). RR and HR estimates are reported separately throughout and are not combined in a single model, as mixing effect measures can introduce bias [18]. For trials reporting outcomes at multiple time points, events at the protocol-specified primary endpoint were extracted (e.g., CHAMPION at 6 months rather than extended follow-up at 15–18 months). The cluster-randomized trial (CHAT) was adjusted using the design effect method (DE = 1.35, based on ICC = 0.02 and mean cluster size 18.4), with both event counts and denominators reduced per Cochrane Handbook Section 23.1.3 [19].

Random-effects meta-analysis was performed using restricted maximum likelihood (REML) estimation for the between-study variance (τ^2) with the Hartung-Knapp-Sidik-Jonkman (HKSJ) adjustment for confidence intervals [20,21]. Statistical heterogeneity was assessed using I^2, τ^2, Cochran's Q test, and 95% prediction intervals [22,23].

Subgroup and Sensitivity Analyses

Pre-specified subgroup analyses examined the influence of RPM type (STS, non-invasive TM, invasive), HF phenotype (HFrEF, HFpEF, HFmrEF, mixed), geographic context (rural/remote, urban, mixed, not reported), country income level (high-income versus other), follow-up duration ($6 months versus > 6 months$), publication era ($2015 versus >2015), and risk of bias (high versus low/some concerns). Interaction P values were calculated using the Q-test for between-subgroup differences.

Ten pre-specified sensitivity analyses were performed for each primary outcome: (1) leave-one-out analysis, (2) fixed-effect (common-effect) model, (3) exclusion of active comparator arms, (4) exclusion of invasive RPM studies, (5) exclusion of cluster-randomized trials, (6) exclusion

of high risk of bias studies, (7) exclusion of flagged outliers, (8) restriction to HR-reporting studies only, (9) restriction to follow-up $3 months, and (10) restriction to large trials (N200$). Meta-regression examined the association of treatment effect with total sample size, publication year, follow-up duration, mean age, and mean LVEF.

Publication Bias

Publication bias was assessed visually using funnel plots and statistically using Egger's regression test [24], Begg's rank correlation test, and Peters' regression test. The trim-and-fill method was applied to estimate the impact of potential missing studies [25]. Contour-enhanced funnel plots were generated to distinguish asymmetry due to publication bias from asymmetry due to other causes. A large-trial sensitivity analysis (N$$200) was performed to assess whether small-study effects drove the overall signal.

Trial Sequential Analysis

Trial sequential analysis (TSA) was performed using α=0.05, β=0.20 (80% power), and relative risk reductions (RRR) of 15% and 20%, with O'Brien-Fleming spending boundaries [26]. TSA was used as a sensitivity framework to assess whether cumulative evidence was robust to repeated testing under the specified assumptions.

Certainty of Evidence

The certainty of evidence was assessed using the Grading of Recommendations, Assessment, Development, and Evaluations (GRADE) framework [27] across five domains: risk of bias, inconsistency, indirectness, imprecision, and publication bias.

All analyses were performed in R (version 4.5.2) using the meta [28] and metafor [29] packages. Statistical significance was defined as P < 0.05 (two-sided).

Results

Study Selection

The systematic search identified 4,037 records across four databases: PubMed (n=1,387), Cochrane CENTRAL (n=2,138), ClinicalTrials.gov (n=348), and WHO ICTRP (n=164). After identifying 1,405 duplicate records and rescuing one that had been incorrectly flagged, 1,404 records were removed, yielding 2,633 unique records for title and abstract screening. Of these, 1,616 were excluded, and 1,017 records proceeded to full-text assessment. A structured audit excluded 701 records (registry entries, protocols, conference abstracts, non-RCTs), while 26 records were added or reclassified into the full-text pool (Tier B candidates, manual review survivors, and pipeline rescues), yielding 342 records for full-text review across four tiers (A–D). After detailed eligibility evaluation, 65 studies met all inclusion criteria (including two records rescued from title/abstract screening via post-hoc algorithm sensitivity analysis), of which 59 were poolable for quantitative synthesis (Figure 1).

Study Characteristics

Table 1. Characteristics of Included Studies

Study	Trial Name	RPM Type	Design	N (Int)	N (Ctrl)	Follow-up (mo)	Age	Female (%)	HF Type	LV EF (%)	NYHA III-IV (%)	Setting	Country	RoB Overall
CHAMPION (Abraham 2011)	CHAMPION	Invasive	Parallel RCT	270	280	6.0	61.6 ± 12.8	27	Mixed	23	100	NR	USA	Some concerns
REDUCEhf (Adamson 2011)	REDUCEhf	Invasive	Parallel RCT	202	198	12.0	55.0 ±	30	HFrEF	23 ± 8	51	Outpatient	USA	Some concerns

Study	Trial Name	RPM Type	Design	N (Int)	N (Ctrl)	Follow-up (mo)	Age	Female (%)	HF Type	LVEF (%)	NYHA III-IV (%)	Setting	Country	RoB Overall
							15.0							
EVOLVO (Landolina 2012)	EVOLVO	Invasive	Parallel RCT	99	101	16.0	NR	18	HFrEF	NR	17	Outpatient	Italy	Some concerns
IN-TIME (Hindricks 2014)	IN-TIME	Invasive	Parallel RCT	333	331	12.0	65.3 ± 9.3	18	HFrEF	26 ± 6	57	Tertiary hospital (36 centres, 7 countries: Aus...	Germany	Some concerns
CONNECT-OptiVol (Lüthje 2015)	CONNECT-OptiVol	Invasive	Parallel RCT	87	89	15.0	66.0 ± 12.0	20	HFrEF	33 ± 11	43	Inpatient implant / outpatient follow-up (singl...	Germany	High
OptiLink HF (Böhm 2016)	OptiLink HF	Invasive	Parallel RCT	505	497	22.8	66.1 ± 10.1	23	HFrEF	27 ± 6	81	Outpatient	Germany	Some concerns
TELECART (Sardu 2016)	TELECART	Invasive	Parallel RCT	96	95	12.0	71.8 ± 8.5	28	HFrEF	NR	55	Multicentre (3 Italian centres); outpatient+dev...	Italy	Some concerns
MORE-CARE (Boriani 2017)	MORE-CARE	Invasive	Parallel RCT	437	428	24.0	66.0 ± 11.0	21	HFrEF	27 ± 7	63	Outpatient	Italy (multinational)	Some concerns
RESULT (Tajstra 2020)	RESULT	Invasive	Parallel RCT	299	301	12.0	NR	18	HFrEF	NR	22	Outpatient	Poland	Some concerns
GUIDE-HF (Lindenfeld 2021)	GUIDE-HF	Invasive	Parallel RCT	497	503	12.0	71.0	38	Mixed	38	71	Outpatient	USA/Canada	Some concerns
MONITOR-HF (Brugts 2023)	MONITOR-HF	Invasive	Parallel RCT	176	172	48.0	66.9 ± 11.4	22	Mixed	30	100	Outpatient (25 hospitals, Netherlands)	Netherlands	Some concerns
Jerant 2001 (3-arm) (Jerant 2001)	Jerant 2001 (3-arm)	Non-invasive TM	Parallel RCT	13	12	6.0	NR	NR	NR	NR	NR	Post-discharge	USA	High
Benatar (2003)	—	Non-invasive TM	Parallel RCT	108	108	3.0	62.9 ± 13.2	64	Mixed	38	NR	Outpatient/Post-discharge	USA	Some concerns
Jerant 2003 (3-arm) (Jerant 2003)	Jerant 2003 (3-arm)	Non-invasive TM	3-arm RCT	13	12	6.0	NR	NR	NR	NR	NR	Post-discharge	USA	High

Study	Trial Name	RPM Type	Design	N (Int)	N (Ctrl)	Follow-up (mo)	Age	Female (%)	HF Type	LVEF (%)	NYHA III-IV (%)	Setting	Country	RoB Overall
TEN-HMS (Cleland 2005)	TEN-HMS	Non-invasive TM	3-arm RCT	173	85	8.0	67.0 ± 13.0	20	HFrEF	25 ± 8	31	Outpatient/Post-discharge (patients recently di...	Germany/Netherlands/UK	Some concerns
Antonicelli (2008)	—	Non-invasive TM	Parallel RCT	28	29	12.0	77.0 ± 8.0	43	Mixed	35 ± 6	42	Post-discharge	Italy	Some concerns
Dansky 2008 (Dansky 2008)	Dansky 2008	Non-invasive TM	Parallel RCT	172	112	4.0	77.1	NR	NR	NR	NR	Post-discharge	USA	High
Kashem 2008 (Kashem 2008)	Kashem 2008	Non-invasive TM	Parallel RCT	24	24	12.0	53.0 ± 10.0	28	HFrEF	25 ± 3	NR	Outpatient	United States	Some concerns
Dansky 2009 (Health Buddy) (Dansky 2009)	Dansky 2009 (Health Buddy)	Non-invasive TM	Parallel RCT	64	46	6.0	76.9	NR	NR	NR	NR	Post-discharge	USA	High
Home-HF (Dar 2009)	Home-HF	Non-invasive TM	—	91	91	6.0	70.0 ± 12.8	32	Mixed	NR	NR	Post-discharge	UK	Some concerns
Giordano (2009)	—	Non-invasive TM	Parallel RCT	230	230	12.0	57.0 ± 10.0	NR	NR	NR	NR	Outpatient	Italy	High
HHH (Mortara 2009)	HHH	Non-invasive TM	—	301	160	12.0	60.0 ± 12.0	14	HFrEF	28 ± 7	NR	Home-based	UK (multinational)	High
MOBITEL (Scherr 2009)	MOBITEL	Non-invasive TM	Parallel RCT	66	54	6.0	65.0	26	Mixed	25	87	Post-discharge	Austria	High
Tompkins 2010 (Tompkins 2010)	Tompkins 2010	Non-invasive TM	Parallel RCT	193	197	6.0	76.7 ± 8.2	42	NR	NR	NR	Outpatient/Home	USA	Some concerns
TIM-HF (Koehler 2011)	TIM-HF	Non-invasive TM	Parallel RCT	354	356	26.0	66.9 ± 10.8	20	HFrEF	27 ± 6	50	Outpatient ambulatory	Germany	Some concerns
THCM (Wade 2011)	THCM (Wade 2011)	Non-invasi	Parallel RCT	164	152	6.0	75.8	58	NR	NR	NR	Post-discharge	USA	High

Study	Trial Name	RPM Type	Design	N (Int)	N (Ctrl)	Follow-up (mo)	Age	Female (%)	HF Type	LVEF (%)	NYHA III-IV (%)	Setting	Country	RoB Overall
(Wade 2011)		ve TM												
TEHAF (Boyne 2012)	TEHAF	Non-invasive TM	Parallel RCT	197	185	12.0	71.0 ± 11.9	42	Mixed	36	43	Outpatient HF clinic (3 centres: Maastricht, He...	Netherlands	Some concerns
TEMA-HF 1 (Dendale 2012)	TEMA-HF 1	Non-invasive TM	Parallel RCT	80	80	6.0	75.9 ± 9.6	38	Mixed	35 ± 15	NR	Post-discharge	Belgium	Some concerns
Gámez-López 2012 (Gámez-López 2012)	Gámez-López 2012	Non-invasive TM	4-arm RCT	52	52	11.0	71.0	NR	Mixed	NR	40	Post-discharge	Spain	Some concerns
Seto 2012 (Seto 2012)	Seto 2012	Non-invasive TM	Parallel RCT	50	50	6.0	55.1 ± 13.7	18	HFrEF	27 ± 8	46	Outpatient heart function clinic	Canada	Some concerns
Delaney 2013 (Delaney 2013)	Delaney 2013	Non-invasive TM	Parallel RCT	46	47	3.0	77.6	67	NR	NR	100	Post-discharge / Home care	USA	Some concerns
Pedone 2015 (Pedone 2015)	Pedone 2015	Non-invasive TM	Parallel RCT	50	46	6.0	79.9 ± 6.8	53	Mixed	44 ± 13	68	Mixed	Italy	High
TIM-HF2 (Koehler 2018)	TIM-HF2	Non-invasive TM	Parallel RCT	765	773	12.0	70.0 ± 11.0	30	Mixed	41 ± 13	47	Outpatient (post-hospitalization)	Germany	Some concerns
Renewing Health (Olivari 2018) (Olivari 2018)	Renewing Health (Olivari 2018)	Non-invasive TM	Parallel RCT	229	110	12.0	NR	NR	Mixed	NR	NR	Post-discharge	Italy (multinational)	Some concerns
Pekmezaris 2019 (Pekmezaris 2019)	Pekmezaris 2019	Non-invasive TM	Parallel RCT	46	58	3.0	58.4 ± 15.2	41	Mixed	NR	70	Post-discharge	USA	Some concerns
OSICAT (Galinier 2020)	OSICAT	Non-invasive TM	Parallel RCT	482	455	18.0	70.0 ± 12.4	27	Mixed	39 ± 14	49	Outpatient/post-discharge (hospitalised for acu...	France	Some concerns
Yanicelli 2021 (Yanicelli 2021)	Yanicelli 2021	Non-invasive TM	Parallel RCT	20	20	3.0	NR	34	HFrEF	36 ± 12	30	Outpatient	Argentina	High

Study	Trial Name	RPM Type	Design	N (Int)	N (Ctrl)	Follow-up (mo)	Age	Female (%)	HF Type	LVEF (%)	NYHA III-IV (%)	Setting	Country	RoB Overall
Johnson 2022 (Johnson 2022)	Johnson 2022	Non-invasive TM	Parallel RCT	16	15	3.0	60.4 ± 12.9	38	Mixed	NR	NR	Inpatient → post-discharge (academic hospital)	USA	High
AMULET (Krzesiński 2022)	AMULET	Non-invasive TM	Parallel RCT	300	305	12.0	67.0 ± 16.0	21	Mixed	32 ± 15	25	Outpatient ambulatory clinics (9 sites)	Poland	Some concerns
ControlVit (Achury-Saldaña 2024)	ControlVit	Non-invasive TM	Parallel RCT	70	70	6.0	64.1 ± 12.0	30	HFrEF	30 ± 10	11	Outpatient	Colombia	Some concerns
Arnar 2025 (Arnar 2025)	Arnar 2025	Non-invasive TM	Parallel RCT	86	89	12.0	64.7 ± 13.2	34	Mixed	44 ± 13	13	Outpatient HF clinic (Landspitali —The National ...	Iceland	Some concerns
M-Cardio (ERICA-HF) (Rustambekova 2025)	M-Cardio (ERICA-HF)	Non-invasive TM	—	127	117	12.0	60.8 ± 7.2	37	HFmrEF	42 ± 12	78	Post-discharge	Kyrgyzstan	Some concerns
Vietnam HFrEF RCT (Tran 2025)	Vietnam HFrEF RCT	Non-invasive TM	Parallel RCT	87	83	6.0	60.9 ± 15.3	26	HFrEF	33 ± 6	37	Outpatient	Vietnam	Some concerns
TeleMedBot (Zheleznykh 2025)	TeleMedBot	Non-invasive TM	Parallel RCT	57	43	6.0	61.5 ± 13.1	26	Mixed	46 ± 12	NR	Post-discharge	Russia	High
Riegel (2002)	—	STS	Parallel RCT	130	228	6.0	72.5 ± 13.1	46	Mixed	42 ± 17	98	Post-discharge	United States	Some concerns
Laramee (2003)	—	STS	Parallel RCT	141	146	3.0	70.6 ± 11.4	42	Mixed	NR	38	Post-discharge	United States	Some concerns
DIAL (GESICA investigators 2005)	DIAL	STS	Parallel RCT	760	758	24.0	64.8 ± 13.9	27	Mixed	30	50	Outpatient (ambulatory); 51 centers public and ...	Argentina	Some concerns
HFHC (Heart Failure Home Care Trial) (Soran 2008)	HFHC (Heart Failure Home Care Trial)	STS	—	160	155	6.0	76.9 ± 7.1	69	HFrEF	24 ± 9	42	Post-discharge	USA	Some concerns

Study	Trial Name	RPM Type	Design	N (Int)	N (Ctrl)	Follow-up (mo)	Age	Female (%)	HF Type	LVEF (%)	NYHA III-IV (%)	Setting	Country	RoB Overall
VA TeleHF (Wakefield) (Wakefield 2008)	VA TeleHF (Wakefield)	STS	3-arm RCT	47	49	12.0	71.8 ± 10.2	0	Mixed	44	72	Post-discharge	USA	Some concerns
Wakefield 2009 (3-arm) (Wakefield 2009)	Wakefield 2009 (3-arm)	STS	3-arm RCT	47	49	6.0	69.0 ± 10.0	1	Mixed	NR	NR	Post-discharge	USA	Some concerns
Tele-HF (Chaudhry 2010)	Tele-HF	STS	Parallel RCT	826	827	6.0	61.0	44	Mixed	35	57	Outpatient cardiology practices (33 sites, USA)	USA	Some concerns
SPAN-CHF II (Weintraub 2010)	SPAN-CHF II	STS	Parallel RCT	95	93	3.0	69.5 ± 14.2	37	Mixed	32 ± 17	53	Multicenter (4 sites: NEMC, RIH, Lahey, BIDMC)	USA	Some concerns
CHAT (Krum 2013)	CHAT	STS	Cluster RCT	139	161	12.0	73.0 ± 10.0	38	Mixed	37 ± 14	42	Primary care (general practice)	Australia	Some concerns
BEAT-HF (Ong 2016)	BEAT-HF	STS	Parallel RCT	715	722	6.0	73.0	47	Mixed	43	75	Hospital (post-discharge)	USA	Low
Negarandeh 2019 (Negarandeh 2019)	Negarandeh 2019	STS	—	40	40	2.0	NR	NR	NR	NR	NR	NR	Iran	Some concerns
DAA/Feijó Brazil (Feijó 2021)	DAA/Feijó Brazil	STS	Parallel RCT	99	107	3.0	64.0 ± 12.0	42	Mixed	33 ± 12	44	Outpatient (HF clinic, Hospital de Clínicas de ...	Brazil	Some concerns
MESSAGE-HF (Rohde 2024)	MESSAGE-HF	STS	Parallel RCT	352	347	6.0	61.3 ± 14.6	32	HFrEF	27	39	Outpatient HF clinics (post-HF hospitalization,...	Brazil	Some concerns
PRADOC (Roubille 2024)	PRADOC	STS	—	202	202	12.0	75.0 ± 12.0	30	Mixed	42 ± 16	90	Post-discharge	France	Some concerns
Ribeiro (2025)	—	STS	Parallel RCT	70	57	6.0	63.6 ± 10.7	50	Mixed	NR	NR	Post-discharge	Brazil	Some concerns

The 65 included RCTs encompassed approximately 23,000 participants across 20 countries.

RPM interventions were classified as structured telephone support (15 trials), non-invasive

telemonitoring (33 trials), or invasive hemodynamic monitoring (11 trials), with 6 trials excluded from quantitative synthesis because they used active comparators, had single-arm or before/after designs, or reported only outcomes incompatible with pooling (eTable 6). The median sample size was 244 (range 25–1,653) and the median follow-up was 6 months (range 2–48 months; one study [Negarandeh 2019] had 2-month follow-up, retained by investigator decision as a borderline case). HF phenotype was predominantly HFrEF or mixed; no trials enrolled HFpEF patients exclusively among the 59 poolable studies.

Risk of Bias

The majority of studies (44 of 59) were judged as having "some concerns" for overall risk of bias, primarily driven by the open-label design inherent to RPM interventions (D2: deviations from intended interventions). Thirteen studies were rated as high risk of bias, and two as low risk. Detailed RoB 2 assessments are presented in eFigures 1–2 and eTable 2.

Primary Outcomes

All-Cause Mortality

Table 2. Summary of Meta-Analysis Results

Category	Outcome	Measure	k	Effect [95% CI]	p-value	I2 (%)	Prediction Interval
Primary outcomes (RR, Mantel-Haenszel)	All-cause mortality (primary)	RR	41	0.911 [0.842, 0.985]	0.0208	0.0	[0.840, 0.988]
Primary outcomes (RR, Mantel-Haenszel)	HF hospitalization (primary)	RR	39	0.781 [0.710, 0.859]	0.0000	47.2	[0.586, 1.040]
Primary outcomes (RR, Mantel-Haenszel)	All-cause hospitalization (primary)	RR	28	0.959 [0.892, 1.031]	0.2462	54.3	[0.809, 1.137]
Primary sensitivity (HR subset, metagen)	All-cause mortality	HR	16	0.880 [0.784, 0.989]	0.0335	0.0	[0.783, 0.989]
Primary sensitivity (HR subset, metagen)	HF hospitalization	HR	13	0.766 [0.698, 0.841]	0.0000	1.1	[0.698, 0.841]

Category	Outcome	Measure	k	Effect [95% CI]	p-value	I2 (%)	Prediction Interval
Primary sensitivity (HR subset, metagen)	All-cause hospitalization	HR	10	0.870 [0.711, 1.064]	0.1514	71.0	[0.505, 1.497]
Secondary outcomes (ratio)	CV Mortality	RR	9	0.814 [0.668, 0.992]	0.0433	0.0	—
Secondary outcomes (ratio)	ED Visits	RR	6	0.795 [0.550, 1.149]	0.1699	41.4	—
Secondary outcomes (ratio)	Composite ACM+Hospitalization	HR	25	0.827 [0.765, 0.894]	0.0000	37.5	—
Secondary outcomes (continuous, MD)	MLHFQ	MD	11	-6.53 [-10.28, -2.79]	0.0030	93.6	—
Secondary outcomes (continuous, MD)	KCCQ	MD	3	8.14 [-24.03, 40.30]	0.3902	83.7	—

RPM significantly reduced all-cause mortality across 41 RCTs (RR 0.911, 95% CI 0.842–0.985; P=0.021; I^2=0%; τ^2=0) (Figure 2, Table 2). The 95% prediction interval was 0.840–0.988, excluding the null under the random-effects model. The number needed to treat was 104 per year. In sensitivity analysis restricted to studies reporting hazard ratios (k=16), the result was consistent (HR 0.880, 95% CI 0.784–0.989; P=0.034).

HF Hospitalization

RPM significantly reduced HF hospitalization across 39 RCTs (RR 0.781, 95% CI 0.710–0.859; P<0.001; I^2=47.2%; τ^2=0.018) (Figure 3). The 95% prediction interval was 0.586–1.040, crossing 1.0, consistent with moderate heterogeneity. The NNT was 18 per year. In HR sensitivity (k=13), the effect was more pronounced (HR 0.766, 95% CI 0.698–0.841; P<0.001; I^2=1.1%).

All-Cause Hospitalization

RPM did not significantly reduce all-cause hospitalization across 28 RCTs (RR 0.959, 95% CI 0.892–1.031; P=0.246; I^2=54.3%) (Figure 4). The prediction interval was 0.809–1.137.

Secondary Outcomes

Cardiovascular mortality was significantly reduced (RR 0.814, 95% CI 0.668–0.992; P=0.043; I^2=0%; k=9; low certainty after downgrading for imprecision). The composite of all-cause mortality and hospitalization was significantly reduced (HR 0.827, 95% CI 0.765–0.894; P<0.001; I^2=37.5%; k=25; low certainty after downgrading for indirectness due to mixed effect measures and inconsistency). Emergency department visits showed a non-significant reduction (RR 0.795, 95% CI 0.550–1.149; P=0.17; I^2=41.4%; k=6). Quality of life measured by MLHFQ improved (MD –6.53 points, 95% CI –10.28 to –2.79; P=0.003; I^2=93.6%; k=11), favoring RPM (lower scores indicate better quality of life), though the very high heterogeneity makes the pooled mean difference difficult to interpret. KCCQ showed a non-significant improvement (MD 8.14, 95% CI –24.03 to 40.30; P=0.39; I^2=83.7%; k=3).

Subgroup Analyses

No statistically significant interaction by RPM type was detected for any primary outcome, though the analysis was likely underpowered to detect moderate subgroup differences given the imbalanced subgroup sizes (STS k=10, TM k=20, Invasive k=11 for ACM). For all-cause mortality, the test for subgroup differences yielded $P_{interaction}$=0.80, with similar point estimates across subgroups (non-invasive TM RR 0.871; invasive monitoring RR 0.925; STS RR 0.928). For HF hospitalization, $P_{interaction}$=0.14, and for all-cause hospitalization, $P_{interaction}$=0.22 (Table 2, Figure 7).

HF phenotype showed a significant interaction for all-cause hospitalization ($P_{interaction}$=0.002), with mixed HF phenotype trials showing a directionally favorable estimate (RR 0.940) compared with HFrEF-only trials (RR 1.037). This was not significant for all-cause mortality

($P_{interaction}$=0.46) and was borderline for HF hospitalization ($P_{interaction}$=0.056). The single trial classified as HFmrEF in the subgroup analysis (M-Cardio [ERICA-HF], mean LVEF 42%) contributed a directionally favorable but imprecise estimate, though this subgroup comprises only one study and should be interpreted cautiously.

Geographic context was examined as a descriptive subgroup. Among 41 trials reporting all-cause mortality, 13 were classified as rural/remote and 28 as not reported. No significant interaction was observed ($P_{interaction}$=0.90). The sparse reporting of geographic context — only two trials (TIM-HF2 and CHAT) provided formal rural/urban subgroup analyses — limited the interpretability of this analysis.

No significant interaction was observed for follow-up duration ($6 vs > 6 months$), $publication era$ (≤ 2015 vs >2015), country income level (eFigure 8), or risk of bias for any primary outcome (all $P_{interaction}$>0.05).

In multivariable meta-regression for HF hospitalization, total sample size was borderline rather than statistically significant after excluding recurrent-count rows from the binary RR analysis (β=0.00026, P=0.063). Mean LVEF was associated with treatment effect (β=–0.032, P=0.0049), but this ecological finding should be interpreted cautiously because study-level covariates can be confounded by intervention type, era, and patient selection. No covariates were significant predictors for all-cause mortality.

Sensitivity Analyses

Table 3. Sensitivity Analyses for Primary Outcomes

Outcome	Analysis	k	RR	LCI	UCI	P	I2 (%)	Note
ACM	10_large_trials_n200	24	0.918	0.849	0.993	0.0331	0.0	N>=200 only (n excl.=17)

Outcome	Analysis	k	RR	LCI	UCI	P	I2 (%)	Note
ACM	11_incl_active_comparators	42	0.918	0.848	0.995	0.0372	0.0	Post-hoc: including MIGHTY-HEART active comparator (CO-0570)
ACM	2_fixed_effect	41	0.900	0.832	0.974	0.0087	0.0	Fixed-effect (common); no HKSJ correction
ACM	3_usual_care_only	37	0.910	0.839	0.986	0.023	0.0	Comparator type = usual care only (n excl.=4)
ACM	4_excl_invasive	30	0.906	0.826	0.994	0.0382	0.0	Excl. Invasive RPM (n excl.=11)
ACM	5_excl_cluster_rct	40	0.907	0.838	0.981	0.0165	0.0	Excl. CHAT (CO-0944) cluster RCT
ACM	6_low_rob_only	35	0.914	0.843	0.992	0.0314	0.0	Excl. High RoB (n excl.=6)
ACM	7_excl_outliers	36	0.930	0.868	0.998	0.043	0.0	Excl. flagged outliers (n excl.=5)
ACM	8_hr_only_subset	16	0.880	0.784	0.989	0.0335	0.0	HR-reporting studies only (metagen); k=16
ACM	9_fu_ge3mo	41	0.911	0.842	0.985	0.0208	0.0	FU >=3 months only (n excl.=0)
HF_hosp	10_large_trials_n200	23	0.837	0.779	0.899	<0.001	21.8	N>=200 only (n excl.=16); excluded recurrent-count rows incompatible with binary RR (n=2)
HF_hosp	11_incl_active_comparators	40	0.790	0.721	0.865	<0.001	46.7	Post-hoc: including MIGHTY-HEART active comparator (CO-0570); excluded recurrent-count rows incompatible with binary RR (n=2)
HF_hosp	2_fixed_effect	39	0.805	0.763	0.850	<0.001	47.2	Fixed-effect (common); no HKSJ correction
HF_hosp	3_usual_care_only	33	0.793	0.723	0.870	<0.001	43.7	Comparator type = usual care only (n excl.=7); excluded recurrent-count rows incompatible with binary RR (n=1)
HF_hosp	4_excl_invasive	30	0.725	0.629	0.835	<0.001	53.0	Excl. Invasive RPM (n excl.=11)
HF_hosp	5_excl_cluster_rct	38	0.779	0.706	0.859	<0.001	48.6	Excl. CHAT (CO-0944) cluster RCT; excluded recurrent-count rows incompatible with binary RR (n=2)
HF_hosp	6_low_rob_only	31	0.797	0.724	0.877	<0.001	46.3	Excl. High RoB (n excl.=8); excluded recurrent-count rows incompatible with binary RR (n=2)
HF_hosp	7_excl_outliers	33	0.812	0.750	0.878	<0.001	33.3	Excl. flagged outliers (n excl.=6); excluded recurrent-count rows incompatible with binary RR (n=2)
HF_hosp	8_hr_only_subset	13	0.766	0.698	0.841	<0.001	1.1	HR-reporting studies only (metagen); k=13
HF_hosp	9_fu_ge3mo	39	0.781	0.710	0.859	<0.001	47.2	FU >=3 months only (n excl.=0); excluded recurrent-count rows incompatible with binary RR (n=2)

All 10 sensitivity analyses were directionally consistent with the primary results for both all-cause mortality and HF hospitalization (Table 3). For all-cause mortality, leave-one-out analysis showed pooled RR ranging from 0.901 to 0.928, with no single study disproportionately influencing the result. The fixed-effect model was consistent (RR 0.900, P=0.009). Excluding invasive RPM studies yielded RR 0.906 (P=0.038), excluding high RoB studies yielded RR 0.914 (P=0.031), and restricting to large trials (N$$200) yielded RR 0.918 (P=0.033; k=24). For

HF hospitalization, all sensitivities remained statistically significant, including the large-trial sensitivity (RR 0.837, 95% CI 0.779–0.899; k=23).

Publication Bias

For all-cause mortality, Egger's test was significant (P=0.040) and Begg's test was significant (P=0.014), suggesting funnel plot asymmetry. Peters' regression test was non-significant (P=0.246), and the large-trial sensitivity analysis (N$$200; k=24) yielded RR 0.918 (P=0.033), indicating that the mortality reduction persisted when small studies were excluded. Trim-and-fill analysis estimated 6 missing studies on the right side and yielded an adjusted RR of 0.925 (95% CI 0.851–1.005; P=0.064), attenuating the mortality estimate to borderline/non-significance (Figure 6). The contour-enhanced funnel plot suggested that asymmetry was driven primarily by small studies in regions of statistical significance, consistent with small-study effects; selective non-publication cannot be excluded.

For HF hospitalization, all three tests were significant (Egger P<0.001, Begg P<0.001, Peters P=0.0002), with trim-and-fill estimating 12 missing studies and an adjusted RR of 0.830 (95% CI 0.737–0.934; P=0.003). The substantial asymmetry was consistent with small-study effects and potentially greater treatment effects in smaller, more intensive interventions.

Trial Sequential Analysis

For all-cause mortality with RRR=15%, the accrued information size (AIS=18,738 patients) exceeded the sequentially adjusted required information size (9,880 patients), and the cumulative Z-curve crossed the O'Brien-Fleming monitoring boundary, supporting a stable mortality benefit under this assumption (Figure 8). For HF hospitalization, the AIS did not reach the

heterogeneity-adjusted benchmark at RRR=15%, but crossed the monitoring boundary at RRR=20%, indicating that the TSA result is assumption-dependent.

NNT and GRADE

For all-cause mortality, the NNT was 93 overall and 104 per year (based on a control event rate of 12.0% over a median follow-up of 13.5 months among studies contributing valid participant-level event counts). For HF hospitalization, the NNT was 17 overall and 18 per year (control event rate 26.9%, median follow-up 13.2 months), after excluding two recurrent-count rows from CER/NNT calculation. NNT annualization assumes a constant hazard rate, which may overestimate annual benefit for outcomes with non-proportional hazards; these estimates should be interpreted with this caveat. Furthermore, the prediction interval for HF hospitalization crosses 1.0, meaning that in some clinical settings the NNT could be substantially larger or even infinite. GRADE certainty was moderate for all-cause mortality (downgraded for suspected publication bias), low for HF hospitalization (downgraded for inconsistency and suspected publication bias), very low for all-cause hospitalization, low for cardiovascular mortality (downgraded for very serious imprecision: borderline significance, small k=9, optimal information size not met), and low for the composite outcome (downgraded for indirectness due to mixed effect measures and inconsistency) (eTable 3).

Discussion

This systematic review and meta-analysis — among the most comprehensive to date on RPM in HF, encompassing 59 poolable RCTs — demonstrates that RPM significantly reduces all-cause mortality by 9% (RR 0.911; NNT 104 per year) and HF hospitalization by 22% (RR 0.781; NNT 18 per year), with no significant effect on all-cause hospitalization. These findings were

directionally robust across 10 sensitivity analyses, while publication-bias and small-study signals require cautious interpretation of the headline effects.

The mortality reduction is statistically consistent ($I^2=0\%$), with a narrow prediction interval (0.840–0.988). This statistical homogeneity should not be overinterpreted as clinical interchangeability across RPM modalities, eras, and healthcare systems. The HKSJ method provides more reliable confidence interval coverage than the DerSimonian-Laird approach, particularly with a moderate number of studies [20,21].

Although no statistically significant interaction by RPM technology type was detected, the analysis had limited power to detect moderate differences given the subgroup sample sizes. Ezimoha et al. reported a significant RPM-type interaction (P=0.007) in a smaller analysis (k=15) [16]. The non-significant interaction in our larger analysis should not be interpreted as proof of equivalence among technologies. Nonetheless, the directional consistency across all three RPM categories suggests that the mechanism of benefit may relate less to any single device and more to the clinical feedback loop that RPM enables — the systematic detection of deterioration, timely clinical response, and guided self-management. This interpretation aligns with the TIM-HF2 rationale, where the structured management protocol was considered as important as the monitoring technology [5].

Our results update and extend the Cochrane reviews by Inglis et al., which reported STS mortality RR 0.87 (95% CI 0.77–0.98) and TM mortality RR 0.80 (95% CI 0.68–0.94) in 2015 [4]. Our overall RR of 0.911 is broadly consistent with but less favorable than these earlier estimates, reflecting the inclusion of several large neutral trials published since 2015 (BEAT-HF, OSICAT, PRADOC, MESSAGE-HF) and the dilution of earlier positive signals by more

pragmatic, larger-scale designs. With an NNT of 104 per year for mortality and 18 per year for HF hospitalization, the absolute benefit remains potentially meaningful, but the HF hospitalization estimate should be interpreted alongside the large-trial sensitivity and publication-bias analyses.

Three contemporary meta-analyses published in 2025 provide relevant context. De Lathauwer et al. pooled 41 RCTs (16,312 patients) with access to Embase and reported an OR of 0.81 (95% CI 0.69–0.95) for mortality, directionally consistent with our RR of 0.911 [14]. Their component analysis identified video communication as an effectiveness enhancer — a finding our data could not evaluate but that warrants future investigation. Dobre et al. reported 105 studies (45,072 patients) across multiple databases [15]; however, their substantially larger yield reflects inclusion of non-randomized designs and post-hoc analyses, whereas our review was restricted to RCTs with extractable outcome data. Ezimoha et al. (15 RCTs) reported a significant RPM-type interaction for mortality (P=0.007) [16], contrasting with our non-significant interaction; however, their smaller sample (k=15 vs our k=41) and different subgroup definitions limit direct comparison. The concordance of the mortality benefit direction across independent reviews strengthens confidence in the overall signal, although the exact effect size remains sensitive to analytic choices and small-study effects.

The invasive monitoring evidence deserves specific commentary. The CHAMPION trial showed a striking 28% reduction in HF hospitalization with CardioMEMS-guided management [6], subsequently replicated in MONITOR-HF (HR 0.56) [7]. However, the GUIDE-HF trial yielded neutral overall results (HR 0.88, P=0.16), with benefit confined to a pre-COVID sensitivity analysis [8] — the pandemic reduced event rates in both arms, narrowing the between-group

difference. The MORE-CARE trial, which evaluated a different mechanism (remote monitoring of CRT-D device diagnostics rather than direct hemodynamic measurement), was similarly neutral. When pooled, invasive monitoring did not significantly reduce all-cause mortality (RR 0.925, P=0.35), though HF hospitalization was significantly reduced (RR 0.846, P=0.006). This pattern is consistent with the mechanism of action: hemodynamic monitoring primarily targets congestion (the driver of hospitalization) rather than arrhythmia or pump failure (drivers of mortality).

IN-TIME warrants specific discussion as a potential outlier. This trial of Biotronik Home Monitoring using ICD/CRT-D devices reported an extreme mortality hazard ratio of 0.36 (10 vs 27 deaths; n=664) [30]. Leave-one-out analysis demonstrated that removing any single trial shifted the pooled all-cause mortality RR within a narrow range (0.901–0.928), confirming that no single trial, including IN-TIME, disproportionately drives the overall mortality conclusion.

HF phenotype showed limited interaction effects. The predominance of HFrEF or mixed populations in the included trials reflects the historical focus of RPM research. None of the 59 poolable trials enrolled HFpEF patients exclusively; PROACTIVE-HF, which targeted HFpEF, was excluded from quantitative synthesis owing to its feasibility design without a usual-care control arm for the primary endpoint. The significant interaction for all-cause hospitalization by HF phenotype (P=0.002) should be interpreted cautiously given the small number of subgroup-specific trials and the potential for confounding by intervention type and era.

Geographic access was examined as a secondary descriptive analysis, motivated by the expectation that RPM could provide the greatest incremental benefit in populations with limited access to in-person HF care. The findings were informative primarily as a reporting gap. Only

two of 59 poolable trials reported formal rural/urban subgroup analyses: TIM-HF2 found no significant interaction (P=0.66), and CHAT was specifically designed for rural and remote patients. Laramee et al. (2003) reported substantial differences between rural (2% HF rehospitalization) and urban (14%) subgroups, but this was a single-center study. The overwhelming majority of trials did not report geographic context at all, making patient-level geographic effect modification infeasible. If RPM is to be deployed as a strategy to address healthcare disparities, trials must be designed and powered to detect geographic access interactions.

The publication bias findings warrant cautious interpretation. Both Egger's (P=0.040) and Begg's (P=0.014) tests were significant for all-cause mortality, while Peters' regression test was non-significant (P=0.246). Trim-and-fill analysis imputed 6 studies, yielding an adjusted RR of 0.925 (95% CI 0.851–1.005; P=0.064), attenuating the mortality estimate to borderline/non-significance. The large-trial sensitivity analysis (N$$200; k=24) yielded RR 0.918 (P=0.033), demonstrating that the mortality signal persists when small studies are excluded, though with a smaller effect size. The contour-enhanced funnel plots suggest that asymmetry is driven primarily by small studies in significance regions, consistent with small-study effects, but selective non-publication cannot be excluded. For HF hospitalization, the publication bias signal was more pronounced: all three tests were significant (Egger P<0.001, Begg P<0.001, Peters P=0.0002), and trim-and-fill imputed 12 studies (31% of k=39), yielding an adjusted RR of 0.830. This degree of asymmetry warrants caution in interpreting the HF hospitalization estimate.

The large-trial sensitivity complements the publication bias analysis and suggests that the HF hospitalization benefit may be closer to the attenuated large-trial estimate (RR 0.837, sensitivity

#10) than the overall pooled estimate (RR 0.781) in pragmatic settings. In multivariable meta-regression, total sample size was borderline (P=0.063) rather than statistically significant after excluding recurrent-count rows from binary pooling. No covariates were significant predictors for all-cause mortality, consistent with the low statistical heterogeneity (I^2=0%) observed for this outcome.

This review has several strengths. It is among the most comprehensive RPM meta-analyses of RCTs, including 59 poolable trials across all three RPM categories. The search encompassed four databases without language restrictions. The analytic approach used REML+HKSJ, included 10 sensitivity analyses per outcome, applied TSA as a sequential-monitoring sensitivity framework, and used GRADE to rate certainty. The geographic access analysis highlights an important reporting gap.

Several limitations should be acknowledged. First, the open-label design inherent to RPM trials introduces potential performance and detection bias, though hard endpoints (mortality, hospitalization) are less susceptible to this bias than patient-reported outcomes. Second, heterogeneity was moderate for HF hospitalization (I^2=47.2%) and all-cause hospitalization (I^2=54.3%), likely reflecting variation in intervention intensity, healthcare system context, and patient populations across decades of research. Third, geographic access data were sparse, limiting interpretation of this secondary descriptive analysis. Fourth, some older trials had small sample sizes, contributing to the small-study effects detected in publication bias analysis. Fifth, the mix of effect measures (RR from event counts, HR from time-to-event analyses) required parallel sensitivity analyses rather than a single unified model; additionally, recurrent-event counts could not be pooled as binary participant-level risks and were excluded from RR/NNT calculations. Sixth, the median follow-up of 6 months may not capture longer-term effects of

sustained RPM programs. Seventh, the search was limited to four databases (PubMed/MEDLINE, Cochrane CENTRAL, ClinicalTrials.gov, and WHO ICTRP); Embase and CINAHL were not searched due to absence of institutional access, a recognized deviation from Cochrane Handbook recommendations [19]. However, several factors mitigate the potential impact of this omission. Cochrane CENTRAL partially aggregates Embase-indexed content: 42.9% of the 2,138 CENTRAL records retrieved in our search carried an Embase identifier, and Sampson et al. found that CENTRAL captures a growing proportion of Embase RCTs [13]. Slobogean et al. demonstrated that PubMed combined with CENTRAL identifies approximately 97% of relevant RCTs in therapeutic systematic reviews [12]. Of our 65 included studies, 63 (96.9%) had PubMed identifiers, suggesting minimal unique Embase-only yield for this topic. Furthermore, cross-referencing our study list against three contemporary meta-analyses that searched Embase (De Lathauwer et al., 41 RCTs; Dobre et al., 105 studies; Ezimoha et al., 15 RCTs) [14–16] revealed concordant mortality direction across reviews (eText 1). Nevertheless, the possibility that Embase-only European or Asian RCTs were missed cannot be excluded, and this limitation is reflected in the GRADE publication bias domain assessment. Eighth, screening used semi-automated rules with manual adjudication and audit, and archived files do not preserve complete independent reviewer-level decisions. Ninth, although extraction was completed using a two-reviewer consensus process, the archived dataset retains final consensus values rather than a complete row-level history of all initial reviewer disagreements.

Clinical Implications

The NNT of 18 per year for HF hospitalization, while not directly comparable due to differences in patient populations, outcome definitions, and follow-up durations, is in a broadly similar range to NNTs reported for SGLT2 inhibitors in DAPA-HF and EMPEROR-Reduced (~19–21/year for

the composite of CV death and HF hospitalization) and mineralocorticoid receptor antagonists in EMPHASIS-HF (~22/year). Combined with the mortality benefit (NNT 104/year), these results support considering RPM programs as an adjunct to guideline-directed medical therapy in HF, particularly where clinical response pathways are well defined. The absence of statistically significant differential effect by RPM type — while acknowledging the limited power to detect moderate differences — suggests that healthcare systems can consider selecting the technology most appropriate to their resources and infrastructure — an important consideration for low- and middle-income countries and for rural health systems where invasive monitoring may not be feasible.

Future research should prioritize geographic access-enriched trials to determine whether RPM differentially benefits patients in rural and underserved settings, HFpEF-specific RPM trials (currently underrepresented), cost-effectiveness analyses comparing RPM modalities, and longer-term follow-up studies to assess sustainability of benefit. Given the cumulative mortality signal, future work may be more informative as comparative effectiveness and implementation research than as simple RPM-versus-no-monitoring trials.

Conclusions

In this systematic review, meta-analysis, and trial sequential analysis of 59 poolable RCTs encompassing approximately 23,000 patients, remote patient monitoring significantly reduced all-cause mortality (RR 0.911, 95% CI 0.842–0.985; I^2=0%; GRADE moderate certainty), with a prediction interval (0.840–0.988) that excludes the null. Trial sequential analysis supported a stable mortality signal under the prespecified 15% relative risk reduction assumption, although publication-bias sensitivity attenuated the estimate. HF hospitalization was significantly reduced

(RR 0.781, 95% CI 0.710–0.859; NNT 18/year; GRADE low certainty), though the prediction interval (0.586–1.040) indicates that in some contexts the effect may be attenuated or absent, consistent with the moderate heterogeneity observed. No statistically significant differential effect by RPM type was detected. Geographic access context remains a reporting gap: only 2 of 59 poolable trials reported formal rural/urban subgroups, and dedicated geographic access-enriched trials are needed to determine whether RPM can address disparities in HF care delivery.

Data Availability

All data, analysis code, and materials necessary to reproduce the findings are publicly available in the Zenodo repository (DOI: 10.5281/zenodo.19865261).

References

1.   Savarese G, Becher PM, Lund LH, et al. Global burden of heart failure: A comprehensive and updated review of epidemiology. Cardiovascular Research. 2023; 118(17) :3272–3287.

2.   Heidenreich PA, Bozkurt B, Aguilar D, et al. 2022 AHA/ACC/HFSA guideline for the management of heart failure. Journal of the American College of Cardiology. 2022; 79(17) :e263–e421.

3.   Knorring S von, Salomonsson S, et al. Geographic access to heart failure care: A systematic review. ESC Heart Failure. 2021; 8(3) :1722–1738.

4.   Inglis SC, Clark RA, Dierckx R, Prieto-Merino D, Cleland JGF. Structured telephone support or non-invasive telemonitoring for patients with heart failure. Cochrane Database of Systematic Reviews. 2015; (10) :CD007228.

5.   Koehler F, Koehler K, Deckwart O, et al. Efficacy of telemedical interventional management in patients with heart failure (TIM-HF2): A randomised, controlled, parallel-group, unmasked trial. Lancet. 2018; 392(10152) :1047–1057.

6.   Abraham WT, Adamson PB, Bourge RC, et al. Wireless pulmonary artery haemodynamic monitoring in chronic heart failure: A randomised controlled trial. Lancet. 2011; 377(9766) :658–666.

7.   Brugts JJ, Raber L, et al. Remote haemodynamic monitoring of pulmonary artery pressures in patients with chronic heart failure (MONITOR-HF): A randomised clinical trial. Lancet. 2023; 401(10394) :2113–2123.

8.   Lindenfeld J, Zile MR, Desai AS, et al. Haemodynamic-guided management of heart failure (GUIDE-HF): A randomised controlled trial. Lancet. 2021; 398(10304) :991–1001.

9.   Guichard JL, Leclercq C, Abraham WT, et al. Hemodynamic-guided management of heart failure with preserved ejection fraction: The PROACTIVE-HF trial. JACC: Heart Failure. 2024; 12(10) :1728–1740.

10.   Masterson Creber R et al. Remotely monitoring and coaching heart failure patients after hospital discharge: The MIGHTY-HEART randomized clinical trial. JAMA Internal Medicine. 2025;

11.   Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. BMJ. 2021; 372 :n71.

12.   Slobogean GP, Verma A, Giustini D, Slobogean BL, Mulpuri K. MEDLINE, EMBASE, and Cochrane index most primary studies but not abstracts included in orthopedic meta-analyses. Journal of Clinical Epidemiology. 2009; 62(12) :1261–1267.

13.   Sampson M, Barrowman NJ, Moher D, Klassen TP, Pham B, Lang D, et al. Should meta-analysts search Embase in addition to Medline? Journal of Clinical Epidemiology. 2003; 56(10) :943–955.

14.   De Lathauwer C, Betz J, Shah SJ, et al. Remote monitoring programs in heart failure: A systematic review and meta-analysis of randomized controlled trials. European Journal of Heart Failure. 2025;

15.   Dobre C, Milič P, et al. Impact of remote patient monitoring on outcomes in heart failure: A systematic review and meta-analysis. American Journal of Therapeutics. 2025;

16.   Ezimoha RN et al. Efficacy of remote patient monitoring in heart failure management: A systematic review and meta-analysis. Cureus. 2025; 17(3) :e92812.

17.   Sterne JAC, Savović J, Page MJ, et al. RoB 2: A revised tool for assessing risk of bias in randomised trials. BMJ. 2019; 366 :l4898.

18.   Tierney JF, Stewart LA, Ghersi D, Burdett S, Sydes MR. Practical methods for incorporating summary time-to-event data into meta-analysis. Trials. 2007; 8 :16.

19.   Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, et al., editors. Cochrane handbook for systematic reviews of interventions. 2023. 6.4. Cochrane; 2023.Available from: https://training.cochrane.org/handbook

20.   Hartung J, Knapp G. On tests of the overall treatment effect in meta-analysis with normally distributed responses. Statistics in Medicine. 2001; 20(12) :1771–1782.

21.   Veroniki AA, Jackson D, Viechtbauer W, et al. Methods to estimate the between-study variance and its uncertainty in meta-analysis. Research Synthesis Methods. 2016; 7(1) :55–79.

22.   Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. BMJ. 2003; 327(7414) :557–560.

23.   Riley RD, Higgins JPT, Deeks JJ. Interpretation of random effects meta-analyses. BMJ. 2011; 342 :d549.

24.   Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. BMJ. 1997; 315(7109) :629–634.

25.   Duval S, Tweedie R. Trim and fill: A simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. Biometrics. 2000; 56(2) :455–463.

26.   Wetterslev J, Thorlund K, Brok J, Gluud C. Trial sequential analysis may establish when firm evidence is reached in cumulative meta-analysis. Journal of Clinical Epidemiology. 2008; 61(1) :64–75.

27.   Guyatt GH, Oxman AD, Vist GE, et al. GRADE: An emerging consensus on rating quality of evidence and strength of recommendations. BMJ. 2008; 336(7650) :924–926.

28.   Balduzzi S, Rücker G, Schwarzer G. How to perform a meta-analysis with R: A practical tutorial. Evidence-Based Mental Health. 2019; 22(4) :153–160.

29.   Viechtbauer W. Conducting meta-analyses in R with the metafor package. Journal of Statistical Software. 2010; 36(3) :1–48.

30.   Hindricks G, Taborsky M, Glikson M, et al. Implant-based multiparameter telemonitoring of patients with heart failure (IN-TIME): A randomised controlled trial. Lancet. 2014; 384(9943) :583–590.

Figures

IDENTIFICATION

Additional records from
other sources (n = 0)

Records identified from databases
(n = 4,037):
PubMed/MEDLINE (n = 1,387)
Cochrane CENTRAL (n = 2,138)
ClinicalTrials.gov (n = 348)
WHO ICTRP (n = 164)

SCREENING

Duplicates removed (n = 1,404 net)
(1,405 removed; 1 Tele-HF record restored)

Records screened
(n = 2,633)

Records excluded at title/abstract screening
(n = 1,616 net):
Automated exclusion (n = 1,514)
Review queue excluded (n = 77)
Manual review excluded (n = 26)
MONITOR-HF rescue (+ 1)

Records sought for retrieval
(n = 1,017)
(incl. 1 MONITOR-HF rescue)

Records not retrieved
(n = 0)

Full-text audit (Phase 4b):
(n = 1,017 assessed)

Records excluded at full-text audit
(n = 699 net):
Pass 1 — registry/tier filters (n = 595)
Pass 2 — cross-tier rules (n = 82)
REVIEW_MANUAL excluded (n = 24)
REVIEW_MANUAL added (+ 2)

ELIGIBILITY

Full-text articles assessed
for eligibility (n = 342)
(Tier A: 60 | Tier B: 21 | Tier C: 187 | Tier D: 1)

Full-text articles excluded (n = 279):
Tier A (n = 48):
Not RCT / no results (n = 26)
Sub-analysis (n = 13)
Review/editorial (n = 4)
Not RPM primary (n = 3)
Other (n = 2)
Tier B (n = 21):
Sub-analysis (n = 8)
Review/editorial (n = 4)
Not HF population (n = 3)
Other (n = 6)
Tier C (n = 136):
Sub-analysis (n = 22)
Mixed population (n = 19)
Not RPM primary (n = 17)
Review/editorial (n = 15)
No RCT design (n = 14)
Feasibility/pilot (n = 11)
Other (n = 38)
Tier D (n = 1)
Audit survivors excl. at FT (n = 73)

INCLUDED

Studies included in review
(n = 65)*:
Tier A — dedicated trials (n = 12)
Tier C — additional RCTs (n = 53)
* Includes 2 records rescued from T/A screening
via post-hoc algorithm sensitivity analysis

Poolable RCTs in meta-analysis
(n = 59):
Structured Telephone Support (STS)
Non-Invasive Telemonitoring (TM)
Invasive (CardioMEMS / ICD-based)
Note: 6 excluded from pool (single-arm,
before/after, no outcome, active comparator)

Figure 1. PRISMA 2020 flow diagram of study selection.

Figure 1. PRISMA 2020 flow diagram of study selection. A total of 4,037 records were identified across four databases. After deduplication, screening, and full-text assessment, 65 studies were included and 59 were poolable for quantitative synthesis.

Figure 2. Forest plot — all-cause mortality (RR).

Figure 2. Forest plot of all-cause mortality. Risk ratios with 95% confidence intervals from random-effects meta-analysis (REML + HKSJ). The diamond represents the pooled estimate (RR 0.911, 95% CI 0.842–0.985; k=41).

Figure 3. Forest plot — HF hospitalization (RR).

Figure 3. Forest plot of heart failure hospitalization. Pooled RR 0.781 (95% CI 0.710–0.859; I^2=47.2%; k=39).

Figure 4. Forest plot — all-cause hospitalization (RR).

Figure 4. Forest plot of all-cause hospitalization. Pooled RR 0.959 (95% CI 0.892–1.031; I^2=54.3%; k=28).

Figure 5. Forest plot — composite ACM + hospitalization (HR).

Figure 5. Forest plot of composite all-cause mortality and hospitalization. Pooled HR 0.827 (95% CI 0.765–0.894; I^2=37.5%; k=25).

Figure 6. Funnel plot — ACM with trim-and-fill.

Figure 6. Funnel plot with trim-and-fill analysis for all-cause mortality. Open circles represent observed studies; filled circles represent imputed studies (k_0=6). Adjusted RR 0.925 (95% CI 0.851–1.005).

Figure 7. Subgroup forest — RPM type x ACM.

Figure 7. Subgroup analysis of all-cause mortality by RPM type. No significant interaction was observed ($P_{interaction}$=0.80). STS: structured telephone support; TM: non-invasive telemonitoring.

Figure 8. Trial sequential analysis — ACM (RRR=15%).

Figure 8. Trial sequential analysis for all-cause mortality (RRR=15%, α=0.05, β=0.20, O'Brien-Fleming boundaries). The cumulative Z-curve (blue) crosses the monitoring boundary, and the accrued information size (18,738) exceeds the sequentially adjusted required information size (9,880), supporting a stable mortality benefit under this assumption.