Personalizing neuromodulation for chronic pain: A connectivity-guided randomized trial

Enrico De Martino[1], Margit Midtgaard Bach[1], Bruno Nascimento Couto[1], Anne Jakobsen[1], Pedro Nascimento Martins[2], Stian Ingemann-Molden[1], Adenauer G. Casali[3], Thomas Graven-Nielsen[1], Daniel Ciampi de Andrade[1]

[1] Center for Neuroplasticity and Pain (CNAP), Department of Health Science and Technology, Faculty of Medicine, Aalborg University, Aalborg, Denmark
[2] Federal University of Juiz de Fora, Juiz de Fora, Brazil
[3] Institute of Science and Technology, Federal University of São Paulo, São Paulo, Brazil

Corresponding author

Daniel Ciampi de Andrade

Center for Neuroplasticity and Pain (CNAP)

Department of Health Science and Technology

Faculty of Medicine, Aalborg University

Selma Lagerløfs Vej 249, 9260 Gistrup, Denmark

Email: dca@hst.aau.dk

Short title: Cortical connectivity-guided trial

Keywords: Non-invasive cortical stimulation; neuromodulation; cortical connectivity; biomarkers of treatment response, precision medicine, neuropathic pain, nociplastic pain, nociceptive pain.

Abstract

In this randomized, double-blind, controlled trial of 8 weeks of repetitive transcranial magnetic stimulation (rTMS) for chronic pain, we compared the classic primary motor cortex (M1) rTMS with a novel target-selection strategy based on pre-therapy cortical connectivity. Guided by principles of homeostatic plasticity, we tested whether stimulating the cortical site with the lowest pre-therapy global connectivity would be more effective than two active comparators: stimulating the site with the highest pre-therapy global connectivity or stimulating M1 independent of connectivity. Before starting rTMS treatment, TMS-evoked EEG potentials were recorded from four cortical targets: M1, the dorsolateral prefrontal cortex, the anterior cingulate cortex, and the posterosuperior insular cortex. For each target, global connectivity was quantified using a distance-weighted, phase-based index (debiased weighted phase lag index, wPLI) derived from pre- and post-TMS-evoked EEG activity, capturing both the magnitude and spatial extent of TMS-induced oscillatory phase locking across cortical regions. Target allocation in the Low- and High-Connectivity groups was based on this global connectivity measure. Ninety patients with chronic pain were randomized to Low-Connectivity, High-Connectivity, or Classic-M1 groups. Treatment consisted of 12 rTMS sessions delivered over 8 weeks to the assigned target. The primary outcome was the proportion of patients achieving $\geq$ 30% reduction in pain intensity. Secondary outcomes included continuous change in pain intensity, pain interference, sleep, fatigue, mood, quality of life, and patient global impression of change. No between-group differences were observed for primary or secondary outcomes ($p > 0.05$). In exploratory analyses, we examined whether pre-therapy local connectivity (within-target wPLI) predicted treatment response. In the Classic-M1 group, lower pre-therapy local M1 connectivity was associated with a greater reduction in pain intensity ($r = 0.50$, $p = 0.005$). This association was not observed in the Low- or High-Connectivity groups. A regression model including group-by-connectivity interaction indicated that the relationship between local connectivity and pain reduction differed between the Classic-M1 and High-Connectivity groups ($p = 0.038$). The results of this clinical trial showed that connectivity-based target allocation using global connectivity did not improve clinical outcomes. However, lower local M1 connectivity was associated with greater pain reduction following Classic-M1 stimulation, suggesting that local M1 connectivity may serve as a potential biomarker of response.

Introduction

Chronic pain is one of the most prevalent and disabling medical conditions[1], and its therapeutic management is challenging, as most treatments show modest to moderate effects in clinical trials[2]. In the absence of methods to individualize treatments and offer interventions to those more likely to respond, a 'trial-and-error' approach remains the predominant strategy in clinical practice, often leading to delays in effective pain control[3]. These limitations indicate the need for a more individualized and precision-medicine approach to pain management.[4]

Repetitive transcranial magnetic stimulation (rTMS) has emerged as a promising intervention for alleviating chronic pain.[5] TMS uses brief, high-intensity magnetic pulses to induce electric currents, thereby stimulating neurons by depolarizing myelinated axons in the cortex,[6] with low side effects. rTMS targeting the primary motor cortex (M1) is the most extensively studied approach in clinical trials,[7] due to its effects on pain-modulatory networks.[8] Several studies have shown analgesic effects of rTMS M1 in peripheral neuropathic pain and fibromyalgia, with a response rate between 35% and 50% of patients.[9–12] A smaller number of studies have reported analgesic effects of rTMS targeting three non-motor cortical regions: the left dorsolateral prefrontal cortex (DLPFC), anterior cingulate cortex (ACC), and posterosuperior insula (PSI). The DLPFC is a key region for mood and cognitive-appraisal aspects of pain.[13] rTMS of the DLPFC has been used with some success for the treatment of chronic pain associated with co-morbid depressive symptoms,[14] migraine,[15] burning mouth syndrome,[16] and peripheral neuropathic pain.[7] ACC is a key region of the limbic system and is involved in cognitive and affective aspects of pain processing.[17,18] rTMS has shown that targeting the ACC can alleviate chronic pain in fibromyalgia,[19] while also leading to a reduction in negative mood and anxiety scores.[20] PSI is a major cortical recipient of spinothalamic projections and plays a key role in processing incoming sensory-discriminative nociceptive signals and modulating nociceptive gain.[21] rTMS of the PSI produced analgesia in patients with chronic peripheral neuropathic pain.[22] Despite these findings, response rates to rTMS across all four stimulation targets generally remain below 50%, with responders experiencing an average pain reduction of approximately 30%.[9–12] This modest efficacy may reflect substantial inter-individual variability in pain processing, as cortical areas can be differentially engaged and modulated by pain experience across individuals.[23]

In this context, we propose that cortical connectivity measures derived from TMS can

provide mechanistic insight into the inter-individual variability in rTMS response. By

combining TMS with electroencephalography (TMS-EEG), we found that phase-based

connectivity metrics derived from TMS-evoked potentials (TEPs) may be relevant for

treatment optimization. Specifically, we showed that reduced phase-based connectivity

measures, including inter-trial phase coherence (ITC)[24] and weighted Phase Lag Index

(wPLI),[25] were associated with individual pain sensitivity,[26] and that rTMS targeting M1

increased both measures, with effects persisting beyond the stimulation session.[27]

These findings led us to investigate cortical phase-based connectivity as a potential

neurophysiological predictor of rTMS response. In the feasibility pilot study of this clinical

trial, we found that pre-therapy cortical connectivity, quantified using a distance-weighted

debiased wPLI index, showed the potential to predict the pain-reducing response to M1

rTMS[28]. People with chronic pain with lower pre-therapy global connectivity were more

likely to respond to ten sessions of M1 rTMS with reduced pain, suggesting that reduced

connectivity before treatment may indicate a greater capacity for rTMS to modulate cortical

dynamics.[28] Given the established pain-modulatory effects of M1 stimulation, enhanced M1

responsiveness to rTMS may translate into more effective pain reduction, supporting the

potential use of pre-therapy TMS-EEG to guide rTMS target selection.

Building on this rationale, the present study aimed to determine whether rTMS efficacy could

be improved by guiding target selection based on pre-therapy cortical connectivity. We

conducted a trial using TMS-EEG to quantify connectivity among four candidate targets: M1,

left DLPFC, ACC, and PSI. Patients were randomized to one of three groups. In the Low-

Connectivity and High-Connectivity groups, the pre-therapy connectivity index was

calculated for each target and used to identify, respectively, the cortical site with the lowest

or highest global connectivity as the stimulation target. In the Classic-M1 group, rTMS was

delivered to M1 independent of the pre-therapy connectivity measures. We hypothesized that

the use of the cortical areas with the lowest pre-therapy global connectivity as a target for

rTMS would lead to a higher number of responders compared to two active-control groups:

one in which rTMS was delivered to the target with the highest connectivity, and a second in

which therapy was delivered to the Classic-M1 target, independently of the results of pre-

therapy global connectivity measures.[7,10,12]

Materials and methods

Participants

The study was conducted at the Center for Neuroplasticity and Pain, Aalborg University, Aalborg (Denmark) from April 2024 to June 2025. The study was approved by the local ethics committee "North Denmark Region Committee" (N-20230076), registered on ClinicalTrials.gov (NCT06395649), and the protocol was registered on the Open Science Framework website (https://osf.io/b2e8h/overview).[29] Ninety patients with chronic pain participated in a double-blind, randomized, controlled, three-arm, parallel-group trial. Patients were eligible to participate if they were between 18 and 80 years old and had chronic pain, defined as pain present on most days for more than 3 months. Patients were required to report a mean pain intensity of 3-9 on a numeric rating scale (NRS; 0=no pain; 10=worst pain imaginable). Participants were excluded if they were pregnant or breastfeeding, had uncontrolled major depression as the primary diagnosis, or had current or recent history of substance abuse. Furthermore, exclusion criteria included an inability to cooperate with study procedures or complete study assessments due to cognitive difficulties, as well as any contraindication to TMS, such as a history of epilepsy, the presence of cranial ferromagnetic implants (e.g., intracranial neurostimulators), and tattoos or permanent makeup containing metallic pigments in the facial region. Patients were also excluded if they were unable to fulfill the criteria in the Transcranial Magnetic Stimulation Adult Safety Screen questionnaire.[30] All potential patients underwent a standardized screening process prior to enrolment, during which pain mechanisms (neuropathic, nociceptive, or nociplastic)[31] and ICD-11 pain categories (primary or secondary pain)[32] were determined.

Sample size estimation

The sample size calculation was informed by previous studies demonstrating analgesic effects of rTMS applied to the four cortical targets examined in this protocol. In a trial on neuropathic pain, Attal et al. reported that M1-rTMS had a 44% response rate and decreased pain intensity by 1.5 ± 0.8 numerical rating scale (NRS) points after 8 sessions, compared with 0.8 ± 1.5 NRS points under sham stimulation.[7] Short et al. found that 10 sessions of left DLPFC rTMS in fibromyalgia patients yielded an average daily pain intensity reduction of around 29% relative to baseline, compared with around 4% in the sham group.[14]

In an rTMS study targeting the ACC in fibromyalgia, Tzabazis et al. observed 31–56% reductions in average pain intensity after 20 sessions of 10-Hz rTMS, compared with 15%

reductions with sham stimulation.[19] Dongyang et al. demonstrated that five sessions of 10-Hz
rTMS to PSI in peripheral neuropathic pain resulted in a 58% response rate, with reduced
visual analogue scale (VAS) scores from 68 to 34 mm (≈50% reduction) compared with 66 to
52 mm (≈24% reduction) under sham.[22] Based on these findings, the analgesic effects of
rTMS across the four cortical targets in chronic pain can be characterized as moderate
(Cohen's d = 0.4–0.5). The present study assumed a standardized mean difference (Cohen's
d) of 0.4 for the contrast between cortical regions with low and high connectivity, with a
negligible therapeutic effect in the latter condition (i.e., sham stimulation). For the three-arm
parallel design, a two-sided significance level of $\alpha = 0.05$ was used, with a step-down
Dunnett correction to control the family-wise error rate across the two experimental vs
control comparisons. The disjunctive power was set at $1 - \beta = 0.80$ under the global
alternative hypothesis, with clinically relevant effect sizes defined as $\delta = 0.65$ (clinically
meaningful improvement, corresponding to 30% reduction in pain intensity) and $\delta = 0.12$
(negligible improvement). Assuming equal allocation across all three arms, these assumptions
yield a required sample size of n=79 patients, with a family-wise error rate of 0.05 and a
disjunctive power of 0.80. Anticipating a dropout rate of approximately 15%, we therefore
planned to recruit 90 patients, divided into three groups of 30 patients each.

Procedure

The patients were allocated in a 1:1:1 ratio to one of three treatment arms: Low-Connectivity,
High-Connectivity, or Classic-M1. Each participant was randomized to one of these three
groups prior to inclusion using a computerized random sequence generator
(https://www.randomizer.org), ensuring balanced allocation with a planned sample of 30
patients per arm. For participants assigned to the Low-Connectivity group, rTMS was applied
to the individually identified cortical region with the lowest global connectivity. For those
assigned to the High-Connectivity group, rTMS was applied to the cortical region with the
highest global connectivity. Participants assigned to the Classic-M1 group received
stimulation over M1 according to established protocols,[7] without consideration of
connectivity, although still assessed. Patients and assessors were blinded to group assignment
throughout the trial. All patients underwent an assessment 1 week before starting rTMS
treatment, including questionnaires and a connectivity assessment using TMS-EEG.

For the rTMS intervention, the induction phase consisted of five consecutive daily rTMS
sessions delivered over one week. This was followed by a 7-week maintenance phase with

one rTMS session per week (12 sessions in total). Each session lasted 20 minutes, including

15 minutes of active stimulation and 5 minutes of questionnaires. Target locations remained

the same throughout the treatment. This protocol was similar to previous rTMS studies

demonstrating analgesic effects in patients with chronic pain.[7,33] Patients were allowed to

keep their concomitant analgesic drug treatment throughout the study.

Transcranial Magnetic Stimulation evoked electroencephalographic

responses

Single-pulse TMS was applied in a randomized order to the four cortical regions, M1,

DLPFC, ACC, and PSI (Figure 1), following previously established procedures.[23,27,28]

Stimulation was delivered using a biphasic stimulator (MagPro R30, MagVenture A/S,

Denmark) equipped with a figure-of-eight coil (Cool-B65) and a double-cone coil (Cool-D-

B80). TMS-evoked potentials were acquired with a TMS-compatible passive 63-channel

EEG cap (g.tec medical engineering GmbH, Austria), with the ground electrode placed on the

right zygoma, Cz at the vertex, and the reference on the right mastoid[27,34]. The electrode

impedances were kept below 5 kΩ, and the data were sampled at 4.800 Hz (g.HIamp

amplifier, g.tec medical engineering GmbH).[35] To minimize auditory evoked potentials from

the TMS coil click, patients wore noise-cancelling in-ear headphones (ER3C, Etymotic

Research, Schaumburg, USA) in combination with the TMS Auditory Artifact Control

(TAAC) masking toolbox.[36] To minimize somatosensory contamination from coil vibration,

two net caps (GVB-geliMED GmbH, Germany) and a thin plastic stretch wrap were placed

over the EEG cap.[23,27] Coil positioning was guided using neuronavigation (InVesalius

Navigator, version 3.1), with the participant's head and the TMS coil calibrated through an

optical infrared tracking system (Polaris Vicra, NDI, Ontario, Canada).[37] The navigated TMS

system continuously monitored coil location and maintained targeting accuracy below 3

mm.[38] The rt-TEP toolbox supported real-time visualization and quality control of TEPs.[39]

For M1, PSI, and ACC stimulation was delivered over the hemisphere contralateral to the

side of greatest pain intensity. If no side predominance was reported, the left hemisphere was

selected. The M1 hotspot for the hand was determined by identifying the coil position that

elicited the largest visible hand twitch, typically corresponding to C3 (left) or C4 (right). The

rest motor threshold (rMT) was defined as the lowest intensity producing a visible hand

twitch.[40] The left DLPFC target was located in the middle frontal gyrus using the method

described previously.[41] This location corresponds to the area below the F3 electrode, and an

intensity of 110% of the hand rMT was used.[42] For the ACC, the target was localized 4 cm

anterior to the tibialis anterior (TA) muscle motor hotspot,[43] corresponding to Fc1 (left) or

Fc2 (right). The TA muscle hotspot was identified as the coil position eliciting a visible leg

muscle twitch. The TA rMT was defined as the lowest intensity required to elicit this

response. ACC TMS-EEG was performed at 90% of the TA rMT. The PSI target was

determined using a validated fast-PSI localization formula.[44] The electrodes C5 or C6 were

nearest to the stimulation site. PSI stimulation was delivered at 90% of the TA rMT. To

ensure clear TEPs, defined as a peak-to-peak amplitude between 6 µV and 25 µV within the

20–100 ms window, the coil orientation and stimulation intensity were adjusted for each

cortical target, and the averaged TEPs were calculated across the first 30 stimuli. This process

was used to evoke a clear local cortical response while minimizing excessive cortical spread.

EEG preprocessing was performed using a custom-designed Python pipeline (Python

Software Foundation). Continuous EEG data were segmented into epochs from −800 to +800

ms relative to the TMS trigger.[28] To correct for the TMS-induced artifact, the signal segment

from 0 to 15 ms (for M1 and DLPFC stimulation) and from 0 to 25 ms (for ACC and PSI

stimulation) was replaced with a mirrored copy of the pre-stimulus data.[27] Epochs and

channels contaminated by decay artifacts, 50 Hz line noise, ocular activity, or muscle activity

were visually inspected and manually removed.[28] Independent component analysis (fast-ICA)

was subsequently applied to eliminate any remaining artifacts.[45,46] The data were then band-

pass filtered between 1 and 45 Hz using a third-order Butterworth filter, downsampled to

1200 Hz, and re-referenced to the average reference[28], and cleaned by interpolating removed

channels using spherical spline interpolation.[47]

233 Classification of the targets using the connectivity index

The global connectivity index used to classify targets was extracted from the pre- and post-

TMS stimulus periods for all electrode pairs using the debiased wPLI.[48] EEG data were

downsampled to 300 Hz, and time–frequency decomposition was performed using Morlet

wavelets with 3.5 cycles, spanning 15 logarithmically spaced frequencies from 8 to 35 Hz.

Pre-TMS stimulus connectivity was estimated within a window from −500 to −50 ms relative

to the TMS pulse, whereas post-TMS connectivity was assessed between 30 and 130 ms.

Connectivity changes were first computed as the normalized mean difference between

connectivity at pre- and post-TMS periods. To quantify the spatial propagation of TMS-

induced changes in connectivity, a distance-weighted wPLI propagation index was computed.

For each electrode pair, the peak TMS-induced change in connectivity was identified as the

95[th] percentile of the post–pre difference. These values were then weighted by the Euclidean

distance between electrode pairs, and the median of the weighted peaks across all pairs was

calculated. This global connectivity index was developed in a pilot study[28] as a standardized

measure integrating both the magnitude and spatial extent of TMS-induced connectivity,

thereby reducing the likelihood that the metric would be disproportionately driven by local

response amplitude and allowing direct comparisons across distinct stimulation targets.

To assess whether local connectivity values across targets were primarily driven by

differences in local cortical excitability, local mean field power (LMFP) was also computed

within stimulation-specific EEG clusters (Supplementary Table 1), and a Spearman

correlation analysis was performed to assess whether the connectivity index correlated with

LMFP. LMFP was defined as the root-mean-square of TEPs across predefined electrode

clusters surrounding the stimulation site,[49] within the same time window used to compute the

connectivity index.

Treatment by repetitive transcranial magnetic stimulation

The therapeutic intervention consists of 10-Hz rTMS applied to one of the four targets. Target

locations were determined during the TMS–EEG session and recorded under neuronavigation

using three anatomical distances from external landmarks (distance from the vertex, distance

from the tragus, and diagonal distance).[50] The rTMS treatment consisted of thirty trains, each

lasting 10 seconds, with 20-second inter-train intervals, and delivered in accordance with

standard procedures. Each train contained 100 pulses, for a total of 3000 administered over

15 minutes. Stimulation intensity varied according to the cortical target: 90% of the rMT of

the hand muscle for M1,[7] 110% of the rMT of the hand muscle for DLPFC,[51] and 90% of the

leg muscle PSI[22] and ACC.[19] These protocols have been used in previous studies involving

patients with chronic pain and have demonstrated efficacy and safety.[7,14,19,22]

Blinding assessment

At the end of the last treatment session, blinding efficacy was evaluated using a structured set

of questions administered to patients and rTMS operators. The assessment aimed to

determine whether blinding was maintained throughout the trial. Participants and operators

were asked whether they believed they could identify the treatment sequence to which the

participant had been allocated, and which treatment condition they believed had been

administered. Previous studies have demonstrated that this type of structured inquiry provides

a robust evaluation of blinding integrity.[52]

Primary outcome

The primary outcome was the proportion of participants who reported a ≥30% reduction in VAS pain intensity at the last visit (week 8) compared with the average VAS pain intensity during the 7 days prior to study initiation. Pain intensity was measured using a VAS, where 0 represents "no pain" and 100 represents "the worst pain imaginable."[53]

Secondary outcome

Secondary outcomes included the proportion of participants who reported a ≥50% reduction in VAS pain intensity at the last visit compared with the average VAS pain intensity during the 7 days prior to study initiation. Furthermore, the following outcomes were assessed before and at the end of the study: pain intensity over the last 24 hours and the last 7 days (assessed by VAS), Brief Pain Inventory (BPI), sleep quality, fatigue, mood and anxiety symptoms (Hospital Anxiety and Depression Scale, HADS), health-related quality of life (EuroQol-5D, EQ-5D), and global impression of change assessed by both patients (PGIC) and operators (OGIC). The BPI was administered to quantify both pain severity and pain interference. Pain severity was calculated as the mean of four intensity items (worst, least, average, and current pain), and pain interference was calculated as the mean of seven items assessing the impact of pain on general activity, mood, walking ability, normal work, relationships with others, sleep, and enjoyment of life.[54] The spatial distribution of pain was quantified using body maps,[55] which were divided into 29 predefined anatomical regions. The total number of regions marked as painful was calculated to assess the extent of the pain.[56] Mood and anxiety symptoms were assessed using the Hospital Anxiety and Depression Scale (HADS), a 14-item self-report measure covering anxiety (7 items) and depression (7 items), with higher scores indicating greater symptom severity.[57] Health-related quality of life was evaluated using the EuroQol-5D (EQ-5D), which assesses five dimensions: mobility, self-care, usual activities, pain/discomfort, and anxiety/depression.[58] Finally, sleep quality and fatigue were measured using two VAS. For sleep, participants rated "How did you sleep last night?" on a scale from 0 (bad sleep) to 100 (good sleep). For fatigue, they rated "How tired have you been in the last 24 hours?" from 0 (no tiredness) to 100 (worst imaginable tiredness).[59] Patient Global Impression of Change (PGIC) and Operator Global Impression of Change (OGIC) were assessed at the end of the maintenance phase using standard 7-point Likert scales ranging from "very much worse" to "very much improved," and were compared across treatment groups.

Explorative analyses

Using the wPLI values to compute the connectivity index for patient classification, we additionally derived a measure of local connectivity at the target site. This was calculated within stimulation-specific EEG electrode clusters used for the LMFP as the average debiased wPLI across all electrode pairs within each cluster, thereby reflecting within-cluster phase-based connectivity.

Statistical analyses

Descriptive statistics were used to characterize the distribution of all variables. Continuous variables were summarized using means and standard deviations (SD) or medians and interquartile ranges, as appropriate, based on their distribution (Shapiro–Wilk test), while categorical variables were summarized using absolute and relative frequencies. A two-sided significance level of $\alpha = 0.05$ was used for all analyses.

A primary intention-to-treat analysis was conducted, with all randomized participants analyzed in their assigned treatment group. The primary outcome, responder status, was analyzed using a chi-squared test to compare proportions between treatment groups. In addition, an explorative intention-to-treat sensitivity analysis was conducted, including only participants who attended at least four induction sessions. Furthermore, to assess whether contrasts between the randomized stimulation target and M1 predicted treatment response, the connectivity index and local-to-M1 difference (LTM1D) variable were computed as the difference between target-area and M1 connectivity. Treatment response was analyzed using an ordinal logistic regression model with pain VAS reduction at the end of the trial as the outcome.

Secondary clinical outcomes were analyzed using baseline-adjusted regression models. Ordinal outcomes were analyzed using ordinal logistic regression, while continuous outcomes were analyzed using linear regression. All models included fixed effects for treatment group, time, and a treatment-by-time interaction, as well as a nonlinear term for baseline pain intensity to account for potential nonlinearity in baseline–outcome relationships. PGIC and OFIC were analyzed using the Kruskal–Wallis' test.

Exploratory analyses examined associations between local connectivity values and changes in VAS pain intensity from baseline to the end of treatment, assessed over the previous 24 hours and the previous week. Linear regression models were fitted to assess the relationship

339   between connectivity measures and pain reduction (percentage reduction from baseline),

340   including interaction terms to evaluate whether these associations differed by stimulation

341   target. This analysis was conducted in an exploratory, hypothesis-generating manner.

Results

Patient characteristics

A total of 132 patients were screened for eligibility, of whom 90 were included (Figure 2). Two patients had the baseline assessment but did not receive any rTMS treatment. Baseline demographic and clinical characteristics are summarized in Table 1. The three arms were broadly comparable in age, sex distribution, pain mechanism, ICD-11 pain classification, and years with pain. In the High-Connectivity arm, 57% of participants were assigned to PSI, followed by ACC (23%) and M1 (17%), with DLPFC selected in only one case (1%). In contrast, in the Low Connectivity arm, 53% of participants were assigned to DLPFC, followed by M1 (27%), with PSI and ACC each accounting for 10%. Most participants across groups received rTMS on the left hemisphere. Patient´s primary pain diagnoses are shown in Supplementary Table 2. Across all three groups, patients exhibited a broad range of chronic pain conditions, typical of clinical chronic pain populations. Diagnoses were described as either primary or secondary pain syndromes according to the ICD-11 classification and also included the mechanistic classification of pain: neuropathic conditions (e.g., radiculopathies, polyneuropathies, nerve lesions), nociceptive and degenerative musculoskeletal disorders (e.g., osteoarthritis, spondyloarthropathies, arthropathies), and nociplastic pain syndromes (e.g., fibromyalgia, chronic primary cervical pain, and chronic widespread pain). Medication used before and after rTMS treatment is summarized in Supplementary Table 3. The medication was similar across the three groups before and after the rTMS treatment, and no substantial between-group differences were observed. Across all groups, the most frequently used medications were acetaminophen, NSAIDs, pregabalin/gabapentin, and weak opioids.

Transcranial magnetic stimulation intensities, compliance, and side effects

TMS intensities used for TMS-EEG and rTMS are reported in Supplementary Table 4. Out of the 12 total rTMS sessions, the average number of rTMS sessions attended was 11.2±2.0 (96%) in the Low-Connectivity group, 11.4 ± 1.2 (96%) in the High-Connectivity group, and 11.8±0.3 (98%) in the Classic-M1 group. One patient in the High-Connectivity group missed six consecutive sessions (from weeks 3 to 8), and one in the Low-Connectivity group missed five consecutive sessions (from weeks 4 to 8). All other missed sessions were isolated single sessions or two consecutive missed sessions (7 in the High-Connectivity group, 9 in the Low-

Connectivity group, and 6 in the Classic-M1 group). The most common reasons for missed

sessions were seasonal diseases or personal reasons. Side effects reported during the rTMS

treatment period are summarized in Supplementary Table 5. Overall, rTMS was well-

tolerated across all three treatment arms, with no serious adverse events reported. The most

common side effect was headache, occurring in 30% of participants in the Classic-M1 group,

40% in the High-Connectivity group, and 37% in the Low-Connectivity group. Dizziness was

also relatively frequent, reported by 13% of participants in the Classic-M1 group, 27% in the

High-Connectivity group, and 37% in the Low-Connectivity group. Other side effects, such

as twitching, numbness, or electrical sensations, unusual sensory experiences, or feelings of

confusion or drowsiness, were reported infrequently ($\leq$13% in all groups), with no significant

differences between treatment arms (all p>0.17). Reports of unusual smells, tastes, or

emotions were uncommon, occurring in 0–10% of participants across groups. No significant

side effects were observed between the treatment groups, and all adverse events were mild

and transient. Most side effects (59%) were reported during the induction week, and 41%

during the maintenance phase.

Blinding assessment

389   Forty-two participants (38%) reported they could identify the treatment group to which they

390   had been allocated. However, only 9 of these correctly identified their assigned group.

391   Regarding therapist blinding, in 12 treatment cases (13%), the rTMS therapist reported being

392   able to determine the patient's group. However, only 4 of these guesses were correct.

Primary outcome

Fourteen patients in the Low-Connectivity group (48%), 15 in the High-Connectivity group

(50%), and 11 in the Classic-M1 group (37%) showed a $\geq$30% reduction in pain VAS scores

at the end of treatment. The percentage reduction in pain was 33$\pm$55% in the Low-

Connectivity group, 32$\pm$35% in the High-Connectivity group, and 27$\pm$33% in the Classic-M1

group (Figure 3). In the intention-to-treat analysis, there was no significant difference in

responder rates between treatment groups for either the categorical or continuous analysis

(Table 2). Results were unchanged in a modified intention-to-treat sensitivity analysis that

included participants who attended at least 4 induction sessions (p = 0.475). No significant

associations were observed between the global connectivity index and changes in VAS pain

ratings, and no significant interactions with Time or treatment group were detected

(Supplementary Table 6). Similarly, in the analysis examining the LTM1D difference, no

significant main effects or interactions were found (Supplementary Table 6). Global
connectivity values are reported in Supplementary Table 7.

Secondary outcomes

Using a ≥50% reduction in pain VAS scores at the end of the treatment, twelve patients in the Low-Connectivity group (41%), eleven in the High-Connectivity group (37%), and ten in the Classic-M1 group (33%) were responders. Across all secondary outcomes, no between-group differences reached significance (Table 2 and Supplementary Fig. 1). For VAS pain intensity over the last 24 hours, mean pain reduction was 29±49% in the Low-Connectivity group, 32 ± 34% in the High-Connectivity group, and 27±5% in the Classic-M1 group, with no significant group differences (p=0.188). A similar pattern was observed for VAS pain intensity over the last 7 days, with reductions of 18±53%, 13±30%, and 13±28%, respectively (p=0.346). No significant between-group differences were observed for pain severity (p=0.581), pain interference (p=0.954), sleep quality (p=0.180), fatigue (p=0.724), or mood symptoms (p=0.286). Health-related quality of life did not differ significantly across the three treatment arms (p=0.259). PGIC ratings were comparable across groups (p=0.427), as were OGIC ratings (p=0.936).

Exploratory analyses

Pain VAS reduction after therapy over the last 7 days was associated with local connectivity in the Classic-M1 group (r=0.527, p=0.003; Figure 4A), but not in the Low-Connectivity (r=0.037, p=0.847) or High-Connectivity groups (r=−0.065, p=0.743). Similarly, pain VAS reduction after therapy over the last 24 hours was associated with local connectivity in the Classic-M1 group (r=0.495, p=0.005; Figure 4B), but not in the Low-Connectivity (r=0.279, p=0.136) or High-Connectivity groups (r=−0.205, p=0.296). Interaction analyses of the correlation slope revealed a difference between the Classic-M1 and High-Connectivity groups in regression slopes between local connectivity and pain VAS reduction over the last 24 hours after therapy (p=0.037; Supplementary Table 8).

Correlation between connectivity index and cortical excitability

The values of the connectivity index and LMFP are reported in Supplementary Table 9. No significant correlation between the connectivity index and LMFP was found (ρ=0.12, p=0.25; Supplementary Fig. 2).

Discussion

The primary aim of this randomized, double-blind, parallel trial was to determine whether rTMS target allocation based on pre-therapy lower cortical connectivity could identify one of four potential cortical targets associated with a higher probability of response in chronic pain. The results did not show a higher proportion of responders with this strategy compared to classic uninformed M1-rTMS therapy, as no significant differences in pain improvement were observed among the Low-Connectivity, High-Connectivity, and Classic-M1 groups. These findings indicate that a global connectivity index measured after stimulation of four discrete cortical sites was insufficient to capture target-specific neurophysiological markers of rTMS responses in chronic pain. Secondary clinical outcomes also did not differ significantly across the three groups, providing no evidence of a superior analgesic effect for any target.

While global responses elicited by local cortical probing did not inform target selection, exploratory analyses showed that local oscillatory responses following M1 probing did carry predictive value. Specifically, lower pre-therapy local connectivity, measured in the EEG cluster around the stimulation site, was associated with greater pain relief only in the Classic-M1 group. This suggests that local connectivity features of M1 may be more relevant for predicting responsiveness to rTMS than global connectivity measures. These findings open two key research questions in the development of neuromodulation-based brain biomarkers for pain relief: (i) are cortical connectivity biomarkers of rTMS response inherently site-specific? (ii) do local connectivity measures offer greater predictive value than global connectivity indices?

In the present trial, the proportion of patients showing a clinically meaningful reduction in pain intensity was 37% in the Classic-M1 group (11/30 patients), 50% in the High Connectivity group (15/30 patients), and 48% in the Low Connectivity group (14/29 patients). These response rates are consistent with those reported in previous rTMS clinical trials, suggesting that the overall therapeutic efficacy observed in the current study aligns with the existing literature and that connectivity-based target allocation did not enhance analgesic efficacy. Hodaj et al. reported a response rate of 47% (27/57 patients),[12] Quesada et al. reported a response rate of 47% (17/36 patients)[10] after M1 rTMS using 30% pain reduction as the definition of responder. Using a 50% of pain reduction threshold, Attal et al.

reported a response rate of 44% (22/49 patients).[7] Investigations of other cortical targets have

reported response rates within a comparable range. For PSI stimulation, Dongyang et al.

reported clinically meaningful pain reduction in 58% of patients with peripheral neuropathic

pain,[22], whereas Galhardoni et al. observed no clear analgesic effects in patients with central

neuropathic pain.[20] Regarding ACC stimulation, Tzabazis et al. reported a mean pain

reduction of 43% in fibromyalgia patients following rTMS,[19] whereas a larger study in

central neuropathic pain failed to demonstrate an analgesic effect.[20] Results from DLPFC

stimulation have also been heterogeneous, with some studies reporting analgesic

benefits[16,51,60–62] and others showing no therapeutic effect.[7] Therefore, the responder rates of

35–50% and an average pain reduction of approximately 30±40% observed in the current

study underscore the persistent challenge of identifying reliable predictors of treatment

response, consistent with findings from all previous studies.

A relevant clinical finding of this study concerns the response rate across different chronic

pain etiologies. In the present trial, we included patients with different chronic pain

conditions, represented by the three mechanistic descriptors of pain: nociceptive, nociplastic,

and neuropathic. Yet the observed response rates were comparable to those reported in

previous rTMS trials focusing on specific pain conditions such as neuropathic pain and

fibromyalgia.[9,10] This finding suggests that the analgesic effects of rTMS may generalize

across chronic pain conditions, rather than being limited to defined mechanistic categories. It

also suggests that interindividual differences in cortical connectivity may be more important

in predicting response to rTMS than the etiology of pain or its mechanistic descriptors. This

is, in itself, not different from other non-pharmacological treatments for chronic pain, which

have shown trans-etiological positive effects.[63]

Using TME-EEG, experimental pain models[26] and chronic pain conditions[64] have been

associated with reductions in cortical excitability and connectivity compared with non-painful

conditions or healthy individuals, reflecting altered oscillatory synchronization within pain-

related networks, as observed in many resting-state EEG studies in chronic pain.[65] By

probing different cortical regions with TMS, such as M1 and DLPFC, reductions in cortical

excitability and connectivity differ across individuals, [23] and these differences correlate with

individual pain sensitivity.[26,66] In contrast, rTMS to M1 and PSI have been shown to increase

local excitability and connectivity,[27,67] suggesting that rTMS-induced modulation of cortical

local connectivity may be therapeutically relevant, particularly in cortical regions

characterized by reduced connectivity.[67] Based on these results, we hypothesized that cortical

targets with the lowest global connectivity would be more susceptible to rTMS-induced
analgesia, assuming connectivity–clinical response relationships would be similar across all
cortical regions. This hypothesis was not confirmed. Nevertheless, important mechanistic
insights emerged from the exploratory analyses. In the Classic-M1 group, pain relief was
associated with low local connectivity. This finding suggests that local network organization
within M1 may play an important role in determining susceptibility to rTMS-induced
modulation. Specifically, lower local phase-based synchronization within M1 may reflect a
cortical state with greater capacity for plastic change, thereby facilitating analgesic responses
to rTMS. Here, local connectivity was quantified using the debiased wPLI, a phase-based
measure that captures the degree to which neuronal populations oscillate with consistent non-
zero phase relationships, thereby indexing functional synchronization while minimizing the
influence of volume conduction[48]. Lower local connectivity values indicate reduced local
phase synchronization, reflecting a more desynchronized cortical network state. In a
secondary analysis of the present dataset, comparing responders and non-responders across
M1 stimulation, irrespective of arm allocation, and using complementary neurophysiological
measures such as ITC and event-related spectral perturbation in the alpha frequency band, we
observed that responders to M1 stimulation were characterized by lower pre-therapy local
oscillatory synchronization.[68] Taken together, these findings may indicate that lower pre-
therapy local connectivity and oscillatory synchronization within the stimulated M1 region
are associated with greater analgesic responses to rTMS, supporting the notion that local M1
network properties may be important to determine M1 rTMS responsivity. Future clinical
studies are needed to further investigate and validate this hypothesis.

This study has several notable strengths and limitations. This is the first large, randomized,
mechanism-based clinical trial aiming to improve the efficacy of rTMS for chronic pain by
using neurophysiological assessment to guide individualized target selection. Whereas most
previous studies have focused on comparing active versus sham stimulation or contrasting
cortical targets, the present approach aimed to move beyond response-rate estimation toward
personalization of treatment based on individual cortical network properties. Second, the
exploratory analyses generate testable hypotheses for future trials, suggesting that TMS-
EEG-derived local connectivity measures within M1 may help identify patients more likely to
respond to rTMS. This study may represent a concrete step toward biomarker-informed
patient stratification in non-invasive neuromodulation trials.

Important limitations should also be acknowledged. The study did not include a sham
stimulation control group because this was not the trial's primary objective. The analgesic
efficacy of rTMS in chronic pain has been demonstrated in multiple prior sham-controlled
studies, and the responder rates observed here are consistent with those reported for active
treatment (35-50%). Furthermore, M1 rTMS is currently recommended by international
guidelines,[69] and the use of sham can be ethically questioned. A further limitation concerns
whether differences in connectivity values across stimulation targets were driven by
variations in local cortical excitability. However, LMFP was not significantly associated with
the connectivity index, suggesting that local excitability did not substantially confound this
measure. In addition, global connectivity may inherently differ across stimulation targets,
potentially influencing the distribution of DLPFC and PSI targets between connectivity
groups. This highlights the potential value of more localized connectivity measures to
mitigate such bias.

In conclusion, this study showed that rTMS target allocation based on a global cortical
connectivity index derived from TMS–evoked EEG potentials did not significantly identify a
cortical target associated with a higher probability of treatment response. Exploratory
analyses revealed that lower pre-therapy local connectivity within the stimulated M1 region
was associated with greater pain relief. Future studies should prospectively test M1-local
connectivity–based stratification to better optimize individualized neuromodulation strategies
for chronic pain.

Competing interests

The authors declare that they have no known competing financial interests or personal relationships that could have influenced the work reported in this paper.

Acknowledgements

The current study was supported by an ERC Horizon Europe Consolidator grant (PersoNINpain 101087925) and a Novo Nordisk grant (Grant NNF21OC0072828). The Center for Neuroplasticity and Pain was supported by the Danish National Research Foundation (DNRF121). TGN receives funding from the Lundbeck Foundation (R441-2023-232).

Author contribution

Enrico De Martino: Conceptualization, Methodology, Formal analysis, Investigation, Data curation, Visualization, Writing- Original draft preparation; Project administration; **Margit Midtgaard Bach**: Investigation, Data curation, Writing - Review & Editing; **Anne Jakobsen**: Investigation, Writing - Review & Editing; **Bruno Andry Nascimento Couto**: Software, Formal analysis, Data Curation, Visualization, Writing - Review & Editing; **Pedro Nascimento Martins**: Visualization, Statistical analysis, Writing - Review & Editing; **Adenauer Girardi Casali**: Software; Writing - Review & Editing; **Stian Ingemann-Molden**: Investigation, Writing - Review & Editing; **Thomas Graven-Nielsen**: Conceptualization, Methodology, Writing - Review & Editing; **Daniel Ciampi de Andrade**: Conceptualization, Methodology, Visualization, Writing - Review & Editing; Project administration, Funding acquisition.

Data availability

The data of this study are available from the corresponding author upon reasonable request.

References

1. Cohen SP, Vase L, Hooten WM. Chronic pain: an update on burden, best practices, and
new advances. *The Lancet*. 2021;397(10289):2082-2097. doi:10.1016/S0140-
6736(21)00393-7

2. Smith SM, Fava M, Jensen MP, et al. John D. Loeser Award Lecture: Size does matter,
but it isn't everything: The challenge of modest treatment effects in chronic pain
clinical trials. *Pain*. 2020;161:S3-S13. doi:10.1097/j.pain.0000000000001849

3. Ciampi de Andrade D, García-Larrea L. Beyond trial-and-error: Individualizing
therapeutic transcranial neuromodulation for chronic pain. *European Journal of Pain*.
2023;27(9):1065-1083. doi:10.1002/ejp.2164

4. Wang J, Doan L V. Clinical pain management: Current practice and recent innovations
in research. *Cell Rep Med. Cell Press*. 2024;5(10). doi:10.1016/j.xcrm.2024.101786

5. Lefaucheur JP, Antal A, Ahdab R, et al. The use of repetitive transcranial magnetic
stimulation (rTMS) and transcranial direct current stimulation (tDCS) to relieve pain.
Brain Stimul. 2008;1(4):337-344. doi:10.1016/j.brs.2008.07.003

6. Siebner HR, Funke K, Aberra AS, et al. Transcranial magnetic stimulation of the brain:
What is stimulated? – A consensus and critical position paper. *Clinical*
Neurophysiology. 2022;140:59-97. doi:10.1016/j.clinph.2022.04.022

7. Attal N, Poindessous-Jazat F, De Chauvigny E, et al. Repetitive transcranial magnetic
stimulation for neuropathic pain: a randomized multicentre sham-controlled trial.
Brain. 2021;144(11):3328-3339. doi:10.1093/brain/awab208

8. Moisset X, De Andrade DC, Bouhassira D. From pulses to pain relief: An update on the
mechanisms of rTMS-induced analgesic effects. *European Journal of Pain (United*
Kingdom). Blackwell Publishing Ltd. 2016;20(5):689-700. doi:10.1002/ejp.811

9. Mhalla A, Baudic S, Ciampi de Andrade D, et al. Long-term maintenance of the
analgesic effects of transcranial magnetic stimulation in fibromyalgia. *Pain*.
2011;152(7):1478-1485. doi:10.1016/j.pain.2011.01.034

10. Quesada C, Pommier B, Fauchon C, et al. New procedure of high-frequency repetitive
transcranial magnetic stimulation for central neuropathic pain: A placebo-controlled
randomized crossover study. *Pain*. 2020;161(4):718-728.
doi:10.1097/j.pain.0000000000001760

11. Hosomi K, Sugiyama K, Nakamura Y, et al. A randomized controlled trial of 5 daily
sessions and continuous trial of 4 weekly sessions of repetitive transcranial magnetic
stimulation for neuropathic pain. *Pain*. 2020;161(2):351-360.
doi:10.1097/j.pain.0000000000001712

12.   Hodaj H, Payen JF, Hodaj E, et al. Long-term treatment of chronic orofacial, pudendal,
and central neuropathic limb pain with repetitive transcranial magnetic stimulation of
the motor cortex. *Clinical Neurophysiology*. 2020;131(7):1423-1432.
doi:10.1016/j.clinph.2020.03.022

13.   Seminowicz, Moayedi M. The Dorsolateral Prefrontal Cortex in Acute and Chronic
Pain. *Journal of Pain*. 2017;18(9):1027-1035. doi:10.1016/j.jpain.2017.03.008

14.   Short EB, Borckardt JJ, Anderson BS, et al. Ten sessions of adjunctive left prefrontal
rTMS significantly reduces fibromyalgia pain: A randomized, controlled pilot study.
Pain. 2011;152(11):2477-2484. doi:10.1016/j.pain.2011.05.033

15.   Brighina F, Piazza A, Vitello G, et al. rTMS of the prefrontal cortex in the treatment of
chronic migraine: A pilot study. *J Neurol Sci*. 2004;227(1):67-71.
doi:10.1016/j.jns.2004.08.008

16.   Umezaki Y, Badran BW, DeVries WH, Moss J, Gonzales T, George MS. The Efficacy of
Daily Prefrontal Repetitive Transcranial Magnetic Stimulation (rTMS) for Burning
Mouth Syndrome (BMS): A Randomized Controlled Single-blind Study. *Brain Stimul*.
2016;9(2):234-242. doi:10.1016/j.brs.2015.10.005

17.   Rolls ET. The cingulate cortex and limbic systems for emotion, action, and memory.
Brain Struct Funct. *Springer*. 2019;224(9):3001-3018. doi:10.1007/s00429-019-01945-
2

18.   Fuchs PN, Peng YB, Boyette-Davis JA, Uhelski ML. The anterior cingulate cortex and
pain processing. *Front Integr Neurosci*. *Frontiers Research Foundation*. 2014;8(MAY).
doi:10.3389/fnint.2014.00035

19.   Tzabazis A, Aparici CM, Rowbotham MC, Schneider MB, Etkin A, Yeomans DC. Shaped
magnetic field pulses by multi-coil repetitive transcranial magnetic stimulation (rTMS)
differentially modulate anterior cingulate cortex responses and pain in volunteers and
fibromyalgia patients. *Mol Pain*. 2013;9(1). doi:10.1186/1744-8069-9-33

20.   Galhardoni R, Da Silva VA, García-Larrea L, et al. Insular and anterior cingulate cortex
deep stimulation for central neuropathic pain disassembling the percept of pain.
Neurology. 2019;92(18):E2165-E2175. doi:10.1212/WNL.0000000000007396

21.   Nagai M, Kishi K, Kato S. Insular cortex and neuropsychiatric disorders : A review of
recent literature. *European Psychiatry*. 2007;22:387-394.
doi:10.1016/j.eurpsy.2007.02.006

22.   Dongyang L, Fernandes AM, da Cunha PHM, et al. Posterior-superior insular deep
transcranial magnetic stimulation alleviates peripheral neuropathic pain — A pilot
double-blind, randomized cross-over study. *Neurophysiologie Clinique*.
2021;51(4):291-302. doi:10.1016/j.neucli.2021.06.003

23.   De Martino E, Casali A, Casarotto S, et al. Acute pain drives different effects on local
and global cortical excitability in motor and prefrontal areas: insights into
interregional and interpersonal differences in pain processing. *Cerebral Cortex*.
2023;33(18):9986-9996. doi:10.1093/cercor/bhad259

24.   Makeig S, Debener S, Onton J, Delorme A. Mining event-related brain dynamics.
Trends Cogn Sci. 2004;8(5):204-210. doi:10.1016/j.tics.2004.03.008

25.   Vinck M, Oostenveld R, Van Wingerden M, Battaglia F, Pennartz CMA. An improved
index of phase-synchronization for electrophysiological data in the presence of
volume-conduction, noise and sample-size bias. *Neuroimage*. 2011;55(4):1548-1565.
doi:10.1016/j.neuroimage.2011.01.055

26.   De Martino E, Casali A, Casarotto S, et al. Evoked oscillatory cortical activity during
acute pain: Probing brain in pain by transcranial magnetic stimulation combined with
electroencephalogram. *Hum Brain Mapp*. 2024;45(6). doi:10.1002/hbm.26679

27.   De Martino E, Casali AG, Nascimento Couto BA, Graven-Nielsen T, Ciampi de Andrade
D. Increase in beta frequency phase synchronization and power after a session of high
frequency repetitive transcranial magnetic stimulation to the primary motor cortex.
Neurotherapeutics. 2025;22(1):e00497. doi:10.1016/j.neurot.2024.e00497

28.   De Martino E, Nascimento Couto BA, Jakobsen A, et al. Pretreatment TMS-EEG
connectivity assessment as a potential predictor of rTMS effectiveness in chronic pain:
A feasibility pilot study. *Clinical Neurophysiology*. 2025;176:2010681.
doi:10.1016/j.clinph.2025.02.268

29.   Bach MM, De Martino E, Ciampi De Andrade D, Nascimento Couto AB, Graven-Nielsen
T. Personalized pain treatment by transcranial magnetic stimulation in patients with
chronic pain: a study on mechanisms. *OSF*. Published online October 2025.
doi:https://osf.io/b2e8h/overview

30.   Rossi S, Antal A, Bestmann S, et al. Safety and recommendations for TMS use in
healthy subjects and patient populations, with updates on training, ethical and
regulatory issues: Expert Guidelines. *Clinical Neurophysiology*. 2021;132(1):269-306.
doi:10.1016/j.clinph.2020.10.003

31.   Fitzcharles MA, Cohen SP, Clauw DJ, Littlejohn G, Usui C, Häuser W. Nociplastic pain:
towards an understanding of prevalent pain conditions. *The Lancet. Elsevier B.V.*
2021;397(10289):2098-2110. doi:10.1016/S0140-6736(21)00392-5

32.   Treede RD, Rief W, Barke A, et al. A classification of chronic pain for ICD-11. *Pain.*
Lippincott Williams and Wilkins. 2015;156(6):1003-1007.
doi:10.1097/j.pain.0000000000000160

33.   Mhalla A, de Andrade DC, Baudic S, Perrot S, Bouhassira D. Alteration of cortical
excitability in patients with fibromyalgia. *Pain*. 2010;149(3):495-500.
doi:10.1016/j.pain.2010.03.009

34.   Seeck M, Koessler L, Bast T, et al. The standardized EEG electrode array of the IFCN.
Clinical Neurophysiology. Elsevier Ireland Ltd. 2017;128(10):2070-2077.
doi:10.1016/j.clinph.2017.06.254

35.   De Martino E, Casali A, Casarotto S, Hassan G, Rosanova M, Ciampi de Andrade D.
Acute pain drives different effects on local and global cortical excitability in motor and
prefrontal areas: Insights into interregional and interpersonal differences in pain
processing. *bioRxiv* . Published online 2023. doi:10.1101/2023.05.26.542414

36.   Russo S, Sarasso S, Puglisi GE, et al. TAAC - TMS Adaptable Auditory Control: A
universal tool to mask TMS clicks. *J Neurosci Methods*. 2022;370:109491.
doi:10.1016/j.jneumeth.2022.109491

37.   Souza VH, Matsuda RH, Peres ASC, et al. Development and characterization of the
InVesalius Navigator software for navigated transcranial magnetic stimulation. *J
Neurosci Methods*. 2018;309:109-120. doi:10.1016/j.jneumeth.2018.08.023

38.   D'Ambrosio S, Jiménez-Jiménez D, Silvennoinen K, et al. Physiological symmetry of
transcranial magnetic stimulation-evoked EEG spectral features. *Hum Brain Mapp*.
2022;43(18):5465-5477. doi:10.1002/hbm.26022

39.   Casarotto S, Fecchio M, Rosanova M, et al. The rt-TEP tool: real-time visualization of
TMS-Evoked Potentials to maximize cortical activation and minimize artifacts. *J
Neurosci Methods*. 2022;370:109486. doi:10.1016/j.jneumeth.2022.109486

40.   Rossini PM, Burke D, Chen R, et al. Non-invasive electrical and magnetic stimulation of
the brain, spinal cord, roots and peripheral nerves: Basic principles and procedures for
routine clinical and research application. An updated report from an I.F.C.N.
Committee. *Clinical Neurophysiology*. 2015;126(6):1071-1107.
doi:10.1016/j.clinph.2015.02.001

41.   Mylius V, Ayache SS, Ahdab R, et al. Definition of DLPFC and M1 according to
anatomical landmarks for navigated brain stimulation: Inter-rater reliability, accuracy,
and influence of gender and age. *Neuroimage*. 2013;78:224-232.
doi:10.1016/j.neuroimage.2013.03.061

42.   Beam W, Borckardt JJ, Scott T. Reeves MD, Mark S. George MD. An efficient and
accurate new method for locating the F3 position for prefrontal TMS applications
William. *Brain Stimul*. 2009;1(2):50-54.
doi:10.1097/MCA.0000000000000178.Endothelial

43. Kreuzer PM, Lehner A, Schlee W, et al. Combined rTMS treatment targeting the
Anterior Cingulate and the Temporal Cortex for the Treatment of Chronic Tinnitus. *Sci*
Rep. 2015;5. doi:10.1038/srep18028

44. da Cunha PHM, Tanaka H, Lapa JD da S, et al. The fast-posterior superior insula (Fast-
PSI): A neuronavigation-free targeting method for non-invasive neuromodulation.
Brain Stimul. Elsevier Inc. 2022;15(5):1178-1180. doi:10.1016/j.brs.2022.08.009

45. Rogasch NC, Thomson RH, Farzan F, et al. Removing artefacts from TMS-EEG
recordings using independent component analysis: Importance for assessing
prefrontal and motor cortex network properties. *Neuroimage*. 2014;101:425-439.
doi:10.1016/j.neuroimage.2014.07.037

46. Korhonen RJ, Hernandez-Pavon JC, Metsomaa J, Mäki H, Ilmoniemi RJ, Sarvas J.
Removal of large muscle artifacts from transcranial magnetic stimulation-evoked EEG
by independent component analysis. *Med Biol Eng Comput*. 2011;49(4):397-407.
doi:10.1007/s11517-011-0748-9

47. Delorme A, Makeig S. EEGLAB⬚: an open source toolbox for analysis of single-trial EEG
dynamics including independent component analysis. *J Neurosci Methods*.
2004;134:9-21. doi:10.1016/j.jneumeth.2003.10.009

48. Vinck M, Oostenveld R, Van Wingerden M, Battaglia F, Pennartz CMA. An improved
index of phase-synchronization for electrophysiological data in the presence of
volume-conduction, noise and sample-size bias. *Neuroimage*. 2011;55(4):1548-1565.
doi:10.1016/j.neuroimage.2011.01.055

49. Kähkönen S, Komssi S, Wilenius J, Ilmoniemi RJ. Prefrontal TMS produces smaller EEG
responses than motor-cortex TMS: Implications for rTMS treatment in depression.
Psychopharmacology (Berl). 2005;181(1):16-20. doi:10.1007/s00213-005-2197-3

50. Sparing R, Buelte D, Meister IG, Pauš T, Fink GR. Transcranial magnetic stimulation and
the challenge of coil placement: A comparison of conventional and stereotaxic
neuronavigational strategies. *Hum Brain Mapp*. 2008;29(1):82-96.
doi:10.1002/hbm.20360

51. Short EB, Borckardt JJ, Anderson BS, et al. Ten sessions of adjunctive left prefrontal
rTMS significantly reduces fibromyalgia pain: A randomized, controlled pilot study.
Pain. 2011;152(11):2477-2484. doi:10.1016/j.pain.2011.05.033

52. Berlim MT, Broadbent HJ, Van Den Eynde F. Blinding integrity in randomized sham-
controlled trials of repetitive transcranial magnetic stimulation for major depression:
A systematic review and meta-analysis. *International Journal of*
Neuropsychopharmacology. 2013;16(5):1173-1181. doi:10.1017/S1461145712001691

53. Patel K V., Allen R, Burke L, et al. Evaluation of composite responder outcomes of pain
intensity and physical function in neuropathic pain clinical trials: An ACTTION
individual patient data analysis. *Pain*. 2018;159(11):2245-2254.
doi:10.1097/j.pain.0000000000001324

54. Tan G, Jensen MP, Thornby JI, Shanti BF. Validation of the brief pain inventory for
chronic nonmalignant pain. *Journal of Pain*. 2004;5(2):133-137.
doi:10.1016/j.jpain.2003.12.005

55. Southerst D, Côté P, Stupar M, Stern P, Mior S. The reliability of body pain diagrams in
the quantitative measurement of pain distribution and location in patients with
musculoskeletal pain: A systematic review. *J Manipulative Physiol Ther*.
2013;36(7):450-459. doi:10.1016/j.jmpt.2013.05.021

56. Murphy DA, Ali SM, Boudreau SA, Dixon W, Wong D, van der Veer SN. Summary and
Analysis of Digital Pain Manikin Data in Adults With Pain Experience: Scoping Review. *J
Med Internet Res*. 2025;27:e69360. doi:10.2196/69360

57. Zigmond AS, Snaith RP. The Hospital Anxiety and Depression Scale. *Acta Psychiatr
Scand*. 1983;67(6):361-370. http://www.hqlo.com/content/1/1/29

58. EuroQol Group. EuroQol--a new facility for the measurement of health-related quality
of life. *Health Policy (New York)*. 1990;16(3):199-206. doi:10.1016/0168-
8510(90)90421-9

59. Zisapel N, Tali N. Determination of the minimal clinically significant difference on a
patient visual analog sleep quality scale. *J Sleep Res*. 2003;12:291-298.

60. Brighina F, Piazza A, Vitello G, et al. rTMS of the prefrontal cortex in the treatment of
chronic migraine: A pilot study. *J Neurol Sci*. 2004;227(1):67-71.
doi:10.1016/j.jns.2004.08.008

61. Nardone R, Höller Y, Langthaler PB, et al. RTMS of the prefrontal cortex has analgesic
effects on neuropathic pain in subjects with spinal cord injury. *Spinal Cord*.
2017;55(1):20-25. doi:10.1038/sc.2016.87

62. Freigang S, Lehner C, Fresnoza SM, et al. Comparing the Impact of Multi-Session Left
Dorsolateral Prefrontal and Primary Motor Cortex Neuronavigated Repetitive
Transcranial Magnetic Stimulation (nrTMS) on Chronic Pain Patients. *Brain Science*.
2021;11:961. doi:10.3390/brainsci

63. Wang Y, Aaron R, Attal N, Colloca L. An update on non-pharmacological interventions
for pain relief. *Cell Rep Med. Cell Press*. 2025;6(2). doi:10.1016/j.xcrm.2025.101940

64. Jakobsen A, De Martino E, Couto BAN, et al. Reduced alpha-band phase coherence
and cortical complexity in fibromyalgia: a TMS-EEG exploratory study. *bioRxiv*.
Published online September 30, 2025. doi:10.1101/2025.09.29.678190

65.   Ploner M, Sorg C, Gross J. Brain Rhythms of Pain. *Trends Cogn Sci*. 2017;21(2):100-
787   110. doi:10.1016/j.tics.2016.12.001

788   66.   Chowdhury NS, Chiang AKI, Millard SK, et al. Combined transcranial magnetic
789   stimulation and electroencephalography reveals alterations in cortical excitability
790   during pain. *Elife*. 2023;12. doi:10.7554/eLife.88567

791   67.   Chowdhury NS, Millard SK, De Martino E, et al. Posterior-superior insula repetitive
792   transcranial magnetic stimulation reduces experimental tonic pain and pain-related
793   cortical inhibition in humans. *Pain*. 2025;166(6):1314-1327.
794   doi:10.1097/j.pain.0000000000003488

795   68.   De Martino E, Bach MM, Jakobsen A, et al. Lower pre-treatment TMS-evoked cortical
796   reactivity and alpha-band oscillatory dynamics predict efficacy of primary motor
797   cortex neuromodulation for chronic pain. *MedRxiv*. Published online October 14,
798   2025. doi:10.1101/2025.10.11.25337793

799   69.   Moisset X, Bouhassira D, Attal N. French guidelines for neuropathic pain: An update
800   and commentary. *Rev Neurol (Paris)*. *Elsevier Masson s.r.l.* 2021;177(7):834-837.
801   doi:10.1016/j.neurol.2021.07.004

Figure legend

Figure 1: Grand-average TMS-evoked EEG potentials (TEPs).

(**A**) primary motor cortex (M1), (**B**) dorsolateral prefrontal cortex (DLPFC), (**C**) anterior cingulate cortex (ACC), and (**D**) posterior superior insula (PSI) for left- and right-hemisphere. For each cortical target, the lower panels display butterfly plots of TEPs from all EEG channels averaged across patients. The red vertical line indicates the TMS pulse at 0 ms. Above each butterfly plot, topographical maps illustrate the spatial distribution of TEP amplitudes at the characteristic post-stimulation time.

Figure 2: CONSORT flow diagram.

132 patients were assessed for eligibility; 43 were excluded, and 90 were randomized in a 1:1:1 ratio to the Low-Connectivity, High-Connectivity, or Classic-M1 arm. All participants allocated to the High-Connectivity and Classic-M1 groups were included in the final analysis, whereas one participant in the Low-Connectivity group was excluded.

Figure 3: Pain intensity scores on a visual analogue scale.

Data points represent individual participants, and square markers indicate group means with error bars representing standard deviations. Pain intensity VAS scores (0-100) were assessed at baseline time points (7 days before TMS–EEG, 24 hours before TMS–EEG, and 24 hours before rTMS), during the induction phase (D1–D5; yellow shaded area), and throughout the maintenance phase, with pain assessed at the time of each visit (W2–W8; green shaded area).

Figure 4: Correlation between local connectivity and percentage pain reduction.

(**A**) Correlation between local (within-cluster) connectivity and percentage change in VAS pain intensity over the last 7 days (negative values indicate pain reduction). (**B**) Correlation between local (within-cluster) connectivity and VAS pain reduction in the last 24 hours (negative values indicate pain reduction).

Table 1

Baseline characteristics of the study groups.

Characteristic		Classic-M1 N = 30	High-Connectivity N = 30	Low-Connectivity N = 30
Age (years)[1]		51 42-62)	55 (46, 60)	57 (48, 61)
Sex	Male	12 (40%)	11 (37%)	9 (30%)
	Female	17 (57%)	19 (63%)	21 (70%)
	Intersex	1 (3.3%)	0 (0%)	0 (0%)
Mechanism-based classification	Neuropathic	10 (33%)	9 (30%)	10 (33%)
	Nociceptive	10 (33%)	11 (37%)	9 (30%)
	Nociplastic	10 (33%)	10 (33%)	11 (37%)
IDC-11 classification	Primary pain	11 (37%)	10 (33%)	10 (33%)
	Secondary pain	19 (63%)	20 (67%)	20 (67%)
Years with pain		9 (5, 15)	7 (3, 15)	9 (4, 16)
rTMS target	M1	30 (100%)	5 (17%)	8 (27%)
	DLPFC	0 (0%)	1 (3%)	16 (53%)
	ACC	0 (0%)	7 (23%)	3 (10%)
	PSI	0 (0%)	17 (57%)	3 (10%)
Hemisphere contralateral to most pain	Left	24 (80%)	22 (73%)	20 (67%)
	Right	6 (20%)	8 (27%)	10 (33%)

[1]Age is presented as median (interquartile range).

All other variables are presented as n (%), with percentages calculated within each group.

Table 2

Primary and secondary outcomes

Outcome	Before rTMS treatment			After rTMS treatment			p-value
	Classic-M1	High-Connectivity	Low-Connectivity	Classic M1	High connectivity	Low connectivity	
Primary outcome (number of responders)	-	-	-	11 (37%)	15 (50%)	14 (48%)	0.530[1]
Secondary outcomes							
Last 7 days (VAS)	64 (59, 75)	68 (65, 73)	64 (57, 73)	58 (40, 72)	62 (40, 76)	47 (35, 64)	0.346
Last 24 h (VAS)	67 (57, 76)	69 (61, 75)	64 (52, 75)	63 (34, 72)	62 (35, 78)	54 (37, 65)	0.188
BPI							
Pain severity	22 (18, 25)	24 (21, 28)	22 (18, 26)	20 (14, 25)	20 (17, 26)	18 (14, 22)	0.581
Pain interference	45 (34, 50)	37 (27, 46)	43 (32, 47)	38 (28, 45)	36 (18, 40)	34 (27, 46)	0.954
Sleep VAS	37 (26, 65)	41 (24, 75)	34 (21, 65)	56 (37, 80)	67 (33, 77)	72 (40, 85)	0.180
Fatigue VAS	70 (58, 76)	66 (29, 74)	66 (52, 77)	60 (40, 66)	58 (30, 70)	60 (50, 69)	0.724
HADS	14 (11, 18)	11 (3, 15)	11 (7, 15)	12 (6, 15)	9 (5, 15)	9 (4, 15)	0.286
EQ-5D	12 (11, 15)	12 (10, 13)	12 (10, 14)	12 (10, 13)	11 (10, 13)	11 (9, 12)	0.259
PGIC	-	-	-	3 (3, 4)	3 (3, 4)	3 (3, 4)	0.427
OGIC	-	-	-	4 (2, 4)	3 (3, 4)	3 (2, 4)	0.936

Median (Q1, Q3)

[1]Chi-squared test used for the number of participants showing ≥30% reduction in VAS at the end of the treatment from baseline compared with the average VAS pain intensity during the 7 days prior to study initiation.
Kruskal-Wallis test used for PGIC and OGIC. Global p-values obtained via baseline-adjusted ordinal logistic models for all the other variables.