1 **Copy Number Analysis in Congenital Nevi: Concordance and Diagnostic**
2 **Limitations of aCGH, sWGS, and Methylation Profiling**

3 Anton Karelin[1], Ines B. Brecht[2], Michaela Pogoda[1,3], German Demidov[1], Michael Abele[2], Dominik T.
4 Schneider[4], Daniel Aldea[5], Heather C Etchevers[5], Susana Puig[6,7], Matthias Hahn[8], Christopher
5 Schroeder[1], Stephan Forchhammer[8] on behalf of the MELCAYA consortium

7
1 Institute of Medical Genetics and Applied Genomics, Eberhard Karls University of Tübingen,
Tübingen, Germany

2 Pediatric Hematology and Oncology, Children's Hospital, Eberhard Karls University of Tübingen,
Tübingen, Germany

3 NGS Competence Center Tübingen, Tübingen, Germany

4 Clinic of Pediatrics, Klinikum Dortmund, University Witten/Herdecke, Dortmund, Germany

5 Aix Marseille University, INSERM, Marseille Medical Genetics, Marseille, France.

6 Dermatology Department, Hospital Clínic de Barcelona, Universitat de Barcelona, Institut de
investigacions biomediques August Pi I Sunyer (IDIBAPS), Barcelona, Spain

7 Centro de Investigación Biomédica en Red de Enfermedades Raras (CIBERER), Instituto de Salud
Carlos III, Barcelona, Spain

8 Department of Dermatology, Eberhard Karls University of Tübingen, Tübingen, Germany
20

____

Corresponding author

Dr. Stephan Forchhammer

Universitäts-Hautklinik, Eberhardt Karls Universität, Tübingen

Liebermeisterstrasse 25, 72076 Tübingen, Germany

E-mail: stephan.forchhammer@med.uni-tuebingen.de

Data availability statement

The data presented in this study are available on request from the corresponding author.

Funding statement

This publication is funded by the European Union through the Horizon Research and Innovation
program under Grant Agreement No 101096667.

NGS methods were performed with the support of the DFG-funded NGS Competence Center
Tübingen (INST 37/1049–1), funded by the Deutsche Forschungsgemeinschaft (DFG, German
Research Foundation) – Project-ID 286/2020B01—428994620.

The STEP registry is funded by the Deutsche Kinderkrebsstiftung.

ORCID:

GD: 0000-0001-9075-4276

MA: 0000-0002-9780-603X

DA: 0000-0001-5101-0194

HCE: 0000-0003-0201-3799

MH: 0000-0001-7511-7447

SF: 0000-0002-1274-2613

Conflict of interest disclosure:

AK: None

IBB: None

MP: None

GD: None

MA: None

DTS: None

DA: None

HCE: None

SP: None

MH: None

CS: CS reports institutional grants from Illumina Inc. and Oxford Nanopore Technologies and research
grants from BMS Stiftung Immunonkologie and Westdeutsche Studiengruppe GmbH outside the
submitted work.

SF: SF received personal honoraria from Kyowa Kirin, Stemline and Recordati Rare Diseases
(speaker's honoraria, advisory board), as well as institutional grants from BioNTech SE, Neracare and
SkylineDX. All outside the submitted work.

Ethics approval statement

The study was approved by the Ethics Committee of the Medical Faculty of the University of
Tübingen (Project ID 399/2023BO1) and conducted in accordance with the Declaration of Helsinki.

Patient consent statement

Written informed consent was given by all patients or their legal guardians. Consent was waived for
deceased patients or samples older than 5 years.

Contributors statement

AK: data collection, analysis and interpretation of results, writing-review and editing.

CS: project administration, data collection, analysis and interpretation of results, writing-review and
editing.

IBB: writing-review and editing

GD: writing-review and editing

MP: methodological development, sample preparation, writing methods, review

MA: writing-review and editing

DS: writing-review and editing

DA: writing-review and editing

HCE: writing-review and editing

SP: writing-review and editing

MH: data collection, writing-review and editing.

CS: project administration, data collection, analysis and interpretation of results, writing-review and
editing.

SF: conceptualization, data collection, analysis and interpretation of results, project administration,
writing-original draft, writing-review and editing.

Acknowledgments

The findings and conclusions in this report are those of the authors and do not necessarily represent
the official position of the European Commission.

The study was funded by an EU Horizon Grant (Melanoma in Childhood, Adolescents, and Young
Adults - MELCAYA, HORIZON-MISS-2021-CANCER-02, ref. 101096667).

ABSTRACT

Background: Distinguishing benign proliferative nodules (PNs) from melanoma arising within congenital melanocytic nevi remains a major diagnostic challenge. Copy number alteration (CNA) analysis is widely used to support classification, but current criteria were developed using array comparative genomic hybridization (aCGH). The performance of alternative platforms such as shallow whole-genome sequencing (sWGS) and methylation arrays in this setting is poorly defined.

Objectives: The objective of this study is to compare CNA profiles obtained from aCGH, sWGS, and methylation arrays in atypical nodules arising within congenital nevi, and to correlate these molecular findings with clinical outcomes.

Methods: Sixteen samples from fourteen patients were retrospectively analyzed using all three platforms. CNAs were cataloged, concordance across methods was quantified using the Jaccard index, and molecular classifications were compared. Clinical follow-up was reviewed to provide clinical context.

Results: aCGH detected 39 CNAs, sWGS 60, and methylation profiling 66. Concordance was highest between sWGS and methylation (mean Jaccard 0.67), followed by aCGH versus sWGS (0.64) and aCGH versus methylation (0.49). Cases with high aneuploidy demonstrated strong cross-platform agreement, whereas low-burden lesions exhibited greater variability between methods. Divergent molecular classifications were observed in six cases.

Conclusions: While all methods reliably detect broad chromosomal changes, sWGS and methylation arrays identify many additional focal CNAs that may not align with CGH-based diagnostic criteria. Until platform-specific thresholds are established, aCGH remains the most conservative and clinically validated approach for evaluating proliferative nodules in congenital nevi.

SIGNIFICANCE

Accurate molecular classification of melanocytic proliferations in congenital nevi is essential but challenging, particularly in patients with multiple proliferative nodules. This study provides the first systematic comparison of aCGH, sWGS, and methylation-based CNA profiling in this setting. We show that higher-resolution platforms detect substantially more focal aberrations, which can lead to discordant and potentially overcalled malignancy assessments when applying CGH-derived criteria. Our findings highlight the need for platform-adapted diagnostic frameworks and support continued use of CGH as the most conservative and clinically validated method for risk stratification.

  GRAPHICAL ABSTRACT

162 1. INTRODUCTION

Pediatric melanoma is a rare neoplasm, with an estimated incidence of four cases per million
children and adolescents under 20 years of age annually [1,2]. Incidence rises markedly during
puberty and adolescence, where melanoma accounts for approximately 7% of all malignancies
among individuals aged 15–19 years [3]. Most cases occur in adolescents, whereas melanoma in
children under 10 years remains exceptionally uncommon [2,4].

Pediatric melanomas comprise several biologically distinct subtypes, including Spitz melanomas,
conventional melanomas, melanomas arising within large or giant congenital melanocytic nevi
(CMN), and rare variants. Prognosis varies widely between these subgroups. Spitz melanocytomas
and Spitz melanomas show a high rate of nodal involvement but low melanoma-specific mortality [5–
7]. In contrast, conventional melanomas and melanomas arising in congenital nevi behave more
aggressively. Although rare, melanomas arising in CMN account for a disproportionate number of
fatal pediatric melanoma cases, particularly in children under five years of age [2,6,7].

Diagnosis of melanoma arising in CMN remains challenging. Giant CMN may contain multiple,
sometimes rapidly enlarging nodules, and distinguishing benign proliferative nodules (PN) from
melanoma can be difficult based on clinical and histologic criteria alone. Histopathologic assessment
typically evaluates mitotic activity, ulceration, cytologic atypia, and architectural features to
differentiate PNs from melanoma [2,8–10].

Molecularly, PNs and melanomas differ mainly in their patterns of copy number alterations (CNAs).
Benign PNs typically exhibit whole-chromosome gains or losses, whereas melanomas show multiple
segmental CNAs, including partial chromosomal gains and deletions [11–13]. Comparative genomic
hybridization (CGH) is widely regarded as the reference standard for CNA analysis in this setting
because it reliably discriminates between simple numerical aberrations and the complex segmental
patterns typical of malignancy [11–13]. However, array-CGH is technically demanding and limited to
specialized laboratories.

Alternative CNA detection approaches have become increasingly available. Shallow whole-genome
sequencing (sWGS) provides genome-wide CNA profiles at low sequencing depth and is suitable for
FFPE tissue, with diagnostic sensitivity comparable to CGH in melanocytic tumors [14,15]. CNAs can
also be inferred from DNA methylation arrays using computational tools such as Conumee or
Epicopy, enabling genome-wide CNA profiles from the same data used for methylation-based tumor
classification [16,17].

Despite their growing use, no prior study has systematically compared sWGS- and methylation-
derived CNA profiles with CGH specifically in proliferative nodules and melanomas arising within
congenital nevi. Here, we compare CNA detection across all three platforms and correlate molecular
findings with clinical outcome.

2. METHODS

2.1 Study cohort

Children and young adults (0–30 years) with atypical melanocytic proliferations arising within a
congenital melanocytic nevus between 2005 and 2024 were included. Cases originated either from
referral specimens evaluated for expert dermatopathological consultation or from patients treated at
the University Hospital Tübingen. Archival FFPE samples were re-evaluated, and 14 of 28 eligible
cases were included after excluding samples with insufficient material or DNA yield. Sixteen lesions
from these 14 patients were analyzed, as one patient had three nodules excised at different time
points. Written informed consent was obtained where applicable; archival cases older than five years
were included under an institutional consent waiver.

2.2 Histology and molecular classification

Only lesions in which morphology alone did not allow reliable distinction between PN and melanoma
were included. All cases were jointly assessed by at least two dermatopathologists specialized in
melanocytic tumors. Molecular classification followed the criteria of Bastian et al. [12]. Lesions
without CNAs or with whole-chromosome gains/losses and a single non–melanoma-typical arm-level
aberration were classified as benign. Single melanoma-typical segmental aberrations were

considered borderline. Multiple (≥2) segmental or arm-level aberrations defined malignant molecular
profiles.

2.3 Array comparative genomic hybridization

Comparative genomic hybridization (CGH) analyses were performed in two distinct settings. For
cases 1–7a and 7b, CGH was newly conducted retrospectively. DNA was extracted from formalin-
fixed, paraffin-embedded (FFPE) tissue samples, which were processed following standardized
protocols for microdissection and DNA isolation. Microarray analyses were carried out using the
SurePrint G3 Human CGH Microarray Kit 4x180K (Agilent Technologies, Santa Clara, USA).
Hybridizations were scanned on an Agilent SureScan DX Microarray Scanner, with human reference
DNA (female or male, Agilent Technologies) serving as the comparative reference. Data analysis was
performed using Agilent Cytogenomics software (v5.1.2.1).

For cases 7c–14, existing CGH data from routine diagnostics, generated between 2009 and 2021,
were utilized. Tumor tissue was manually microdissected from four to six 25-µm-thick paraffin
sections, deparaffinized, and digested with proteinase K. DNA extraction and labeling were
performed using genomic DNA universal linkage system labeling kits (Agilent Technologies, Santa
Clara, USA). Hybridization employed the Agilent Human Genome CGH 105K microarray (Agilent
Technologies). Arrays were scanned using an Agilent DNA microarray scanner, and image data were
analyzed with Feature Extraction software (v10.5.1.1), DNA Analytics (v4.0.85), and Genomic
Workbench Lite Edition (v6.5.0.18).

2.4 sWGS

DNA was extracted from formalin-fixed, paraffin-embedded (FFPE) tissue samples, which were
processed following standardized protocols for microdissection and DNA isolation. The DNA was
quantified using the Qubit Fluorometer and dsDNA High sensitivity assay (both Thermo Fisher
Scientific, Waltham, USA). The genomic fragment lengths of the samples were assessed with the
Tapestation 4200 and the Genomic DNA ScreenTape Analysis (Agilent Technologies, Santa Clara,
USA). Since some of the DNA samples showed discolorations, all of them were cleaned up with
paramagnetic beads (SPRISelect, Beckman Coulter, Brea, USA) to reduce potential contaminants. 100
ng of the DNA were subsequently sheared and prepared for Next Generation Sequencing (NGS) using
the NEBNext UltraShear FFPE DNA Library Prep Kit with NEBNext Unique Dual Index UMI Adaptors
(both New England Biolabs, Ipswich, USA) with slight adjustments to the fragmentation incubation
and the PCR protocol to accommodate for differences in the original sample qualities. The
concentrations of the libraries were assessed using the Qubit Fluorometer and dsDNA High
Sensitivity Assay-Kit (both Thermo Fisher Scientific, Waltham, USA), while fragment sizes were
determined with the Fragment Analyzer 5300 and the Fragment Analyzer DNA HS NGS fragment
Fragment kit (Agilent Technologies, Santa Clara, USA).

Sequencing was performed on a NovaSeq 6000 (Illumina, San Diego, CA, USA) in paired-end mode
(2x154 bp) with an average of 177 million reads per sample.

2.5 Methylation

DNA was extracted from formalin-fixed, paraffin-embedded (FFPE) tissue samples, which were
processed following standardized protocols for microdissection and DNA isolation. The DNA was
quantified using the Qubit Fluorometer and dsDNA High sensitivity assay (both Thermo Fisher

Scientific, Waltham, USA). The genomic fragment lengths of the samples were assessed with the
Tapestation 4200 and the Genomic DNA ScreenTape (both Agilent Technologies, Santa Clara, USA).

200 ng of DNA were prepared using the Twist Targeted Methylation Sequencing Workflow (Twist
Bioscience, South San Francisco, USA) including the NEBNext Enzymatic Methyl-seq Library
Preparation Protocol (New England Biolabs, Ipswich, USA) according to the manufacturers' protocol
with 8-plex pre-capture pooling and targeted enrichment using the Twist Human Methylome
enrichment panel. The concentrations of the libraries and pools were assessed using the Qubit
Fluorometer and dsDNA High Sensitivity Assay-Kit (both Thermo Fisher Scientific, Waltham, USA),
while fragment sizes were determined with the TapeStation 4200 and the High Sensitivity D1000
ScreenTape (both Agilent Technologies, Santa Clara, USA).

Sequencing was performed on a NovaSeq 6000 (Illumina, San Diego, CA, USA) in paired-end mode
(2x154 bp) with an average of 227 million reads per sample.

2.6. Sequencing

Shallow whole-genome sequencing (sWGS) and methylation sequencing (EM-seq) data as well as
quality control were analyzed with an in-house pipeline called megSAP (version archived at Zenodo:
https://zenodo.org/records/15063428). Sequencing reads were mapped to human genome
reference GRCh38 using BWA-MEM2 for sWGS and Bismark for EM-seq [18,19].

Mean coverage was 6.78x (range: 5.57-7.62) and 78.01x (range: 19.48-137.31) for sWGS and EM-seq
respectively.

2.7. Copy-Number Analysis

Copy-number alteration (CNA) analysis from sWGS and EM-seq data was performed using ClinCNV, a
segmentation-based algorithm optimized for low-coverage whole-genome and targeted sequencing
data [20]. As no matched normal samples were available, CNA calling was performed using a tumor-
only approach, assuming diploidy across the majority of the genome.

Coverage tracks were generated at 10 kb resolution, and GC correction was applied. CNA detection
was executed with tumor-only mode using the following parameters:

--onlyTumor, --scoreS 1000, --lengthS 10, --minimumNumOfElemsInCluster 15, --purityStep 5, --
clonePenalty 1000, --minimumPurity 5, --filterStep 0, with hg38 as the reference genome.
Pseudoautosomal regions on chromosome X were excluded from the analysis to avoid sex-related
copy number bias.

All detected CNA profiles were assessed by manual inspection. CNAs were categorized as whole
chromosome or chromosome arm gains and deletions.

2.8. Statistics

Statistical analyses were performed using Microsoft Excel (version 16.0). Descriptive statistics were
calculated as appropriate. Concordance between CNA profiles obtained by aCGH, sWGS, and
methylation arrays was quantified using the Jaccard similarity coefficient, defined as the ratio of
shared to total aberrations for each method pair. Heatmap visualizations were generated using
Morpheus (Broad Institute; https://software.broadinstitute.org/morpheus/).

2.9. Ethics statement

The study was approved by the Ethics Committee of the University of Tübingen (Project ID
399/2023BO1) and conducted in accordance with the Declaration of Helsinki.

3. RESULTS

3.1. Epidemiology, clinical and histological presentation

The cohort included 14 patients aged 8 days to 26 years (nine males, five females). Most lesions
arose within giant CMN located on the back or trunk. Two cases involved congenital blue nevi and
two involved medium- or small-sized CMN with atypical features. Histopathologically, most nodules
represented proliferative nodules (PN), including spitzoid variants requiring molecular workup for
classification . Follow-up ranged from 2 to 94 months. Three patients died from their disease, seven
remained disease-free, and four lacked follow-up (Table 1).

Table 1. Clinical and histopathological characteristics of the study cohort. **Abbreviations:** CMN, (giant unless otherwise specified) congenital melanocytic nevus; PN, proliferative nodule; d, days; mo, months; y, years; D, deceased; A, alive without disease; n.a., not available.

Case	Age	Sex	Clinical	Histology	Follow-up
1	12- 17 y	m	Nodule in CMN, head/neck	PN on CMN	82 mo, D
2	0-5 y	m	CMN, scalp	PN on CMN	14 mo, D
3	0-5 y	f	Nodule in CMN, right hip	PN on CMN	17 mo, A
4	0-5 y	f	Dark pigmented nodule in CMN, right shoulder	PN on CMN	48 mo, A
5	12- 17 y	m	Medium-sized CN with nodular areas on the back of the left hand	Atypical spitzoid PN on CMN	87 mo, A
6	0-5 y	f	CMN with multiple nodular areas, buttock	PN on CMN	49 mo, A
7a	0-5 y	f	CMN with multiple nodular areas, back	PN on CMN	2 mo, D (after excision of the lymph node metastasis)
7b	0-5 y		CMN with multiple nodular areas, back	PN on CMN	
7c	0-5 y		Nodular swelling, left axilla	lymph node metastasis	
8	0-5 y	f	CMN with multiple nodular areas, back	PN on CMN	36 mo, A
9	0-5 y	f	CMN with darker areas, right shoulder	PN on CMN	26 mo, A
10	0-5 y	m	CMN with nodular areas, back	PN on CMN	n.a.
11	6-11 y	f	n.a.	PN on congenital blue nevus	n.a.
12	12- 17 y	m	n.a.	PN on CMN	94 mo, A
13	6-11 y	m	n.a.	PN on large congenital blue nevus	n.a.
14	24-29 y	m	Small, papillomatous CMN	Spitzoid PN on CMN	n.a.

3.2 Comparative Analysis of Copy Number Alteration Detection by aCGH, sWGS, and Methylation
Profiling

All lesions were analyzed by aCGH, sWGS, and methylation-derived CNA profiling, and all detected
aberrations are listed in Table 2. Across the cohort, the platforms revealed a broadly consistent
pattern of chromosomal gains and losses, but with marked differences in the number and resolution
of detected events. aCGH identified 39 CNAs, whereas sWGS and methylation profiling detected 60
and 66, respectively. Despite this, substantial overlap existed: 34 CNAs were shared between aCGH
and sWGS, 34 between aCGH and methylation, and 45 between sWGS and methylation. A core set of
32 CNAs was concordantly detected across all three methods.

Table 2. Copy number alterations detected by aCGH, sWGS, and methylation-based CNA profiling across all cases. For each lesion, all chromosomal gains and losses identified by array comparative genomic hybridization (aCGH), shallow whole-genome sequencing (sWGS), and methylation-derived CNA analysis are listed. **Abbreviations:** del, deletion; gain, copy number gain; qter, terminal end of the long arm; pter, terminal end of the short arm; none, no detectable CNAs.

Case	CGH	sWGS	Methylation
1	del(5, 7, 10, 12)	del(5, 7, 10, 12)	del(5p, 7, 10q, 12q)
2	gain(1q, 13q12.2–q32.1); del(3, 5p10–p12, 5q12.1–q14.1, 9, 10, 11, 12, 14, 17, 21)	gain(1, 2, 4, 6p, 7q, 13q12.12–q21.32, 13q22.1–q32.1, 15, 22); del(3, 5p10–p12, 5q12.1–q35.1, 9, 10, 11, 12, 14, 17, 21)	gain(1, 2, 4, 6p, 7, 13q12.12–q21.32, 13q22.1–q32.1, 15, 22); del(3p, 5q12.1–q14.1, 9, 10, 11p, 12, 14, 17, 21)
3	gain(2, 8, 13, 20, 22)	gain(2, 8, 13, 20, 22)	gain(2, 8, 13, 20, 22)
4	gain(6, 8, 9, 16, 18, 19, 20)	gain(6, 8, 9, 16, 18, 19, 20)	gain(6, 8, 9, 16, 18, 19, 20)
5	del(4q22.2–q28.2, 9)	gain(1p22.2–p36.13, 3p12.3–p25.3, 3q11.2–qter, 5p12–p13.3, 5q, 8q12.1–q24.23, 11q, 12q12–q23.3, 15q12–q25.2); del(1p36.13–pter, 2, 4q22.2–q28.2, 6q12–q22.31, 9, 16p)	gain(3p12.3–p25.3, 3q11.2–qter, 5p12–p15.3, 5q, 11p11.2–p15.4, 11q, 12q12–q23.3); del(1p36.13–pter, 4q22.2–q28.2, 9, 16)
6	gain(4, 8, 9, 12, 13, 14, 16, 19, 20, 21, 22); del(5, 10, 18)	gain(4, 8, 9, 12, 13, 14, 16q11.2–q24.1, 21); del(5, 10, 18)	gain(4, 9, 12, 13, 14, 16q11.2–q24.1, 19q13.31–q13.33, 21); del(1p32.2–pter, 5, 6p12.2–p22.2, 10, 18)
7a	gain(7, 8, 9, 12, 13, 16q, 17, 19, 20, 22); del(5, 10)	gain(7, 8, 9, 12, 13, 16q, 17q, 22); del(5, 10)	gain(7, 8, 9, 12, 13, 16p11.2–pter, 16q12.2–qter, 17, 19p13.11–pter, 19q13.11–q13.33, 22); del(5, 10)
7b	del(7)	gain(5q11.1–q33.1); del(7, 8q11.1–q23.3, 16p12.1–p13.3)	del(7)
7c	gain(1q, 7, 8, 20); del(3q13.3–qter, 5, 11, 16q, 17, 18)	gain(1q, 7, 8, 20); del(3q12.2–qter, 5, 11, 16q, 17, 18)	gain(1q, 7, 8, 20); del(3q12.2–qter, 5, 11, 16q, 17, 18)
8	gain(1q, 8, 16, 20); del(3, 5, 7, 11, 12, 14, 19, 21)	gain(1q, 8, 16, 20); del(3, 5, 7, 11, 12, 14, 19, 21)	gain(1q, 8, 16, 20); del(3, 5, 7, 11, 12, 14, 19, 21)
9	none	none	del(20q11.21–qter)
10	gain(7, 8, 12, 13, 16, 18, 20, 21); del(3)	gain(7, 8, 12, 13, 16, 18, 20, 21); del(3)	gain(7, 8, 12, 13, 16, 18, 20, 21); del(3)
11	none	none	del(8q)
12	gain(5, 15, 18, 20, 21, 22)	gain(5, 15, 18, 20, 21, 22)	gain(5, 15, 18, 20, 21, 22)
13	gain(6p21.1–pter)	gain(6p22.2–pter)	gain(6p22.2–pter); del(8q11.1–q24.23, 13q12.13–q33.3)
14	none	none	del(20q11.21–qter)

As expected, aCGH predominantly captured broad whole-chromosome alterations, while sWGS and
methylation profiling revealed numerous additional subchromosomal segmental and focal events.
Methylation profiling, in particular, identified recurrent terminal and arm-level deletions (e.g.,
del(20q11.21–qter), del(8q)) that were not detected by either aCGH or sWGS, suggesting increased
sensitivity to focal or possibly subclonal aberrations. Jaccard similarity coefficients demonstrated the
highest concordance between sWGS and methylation profiling (mean 0.67), followed by aCGH versus
sWGS (0.64) and aCGH versus methylation (0.49). Cases with extensive aneuploidy—such as cases 3,
4, 7c, 8, 10, and 12—showed perfect concordance across all methods, whereas cases with few
aberrations exhibited greater variability, largely driven by additional focal events detected by
methylation arrays.

To visualize CNA burden across the cohort, a heatmap summarizing the number of CNAs per case and
platform was generated, together with representative CNA profiles illustrating platform-specific
resolution differences (Figure 1). Beyond the raw CNA counts, we assessed how platform-specific
variation influenced molecular classification (Table 3). Based on established CGH-derived criteria,
where lesions with no CNAs or exclusively whole-chromosome gains or losses are considered benign,
and profiles with two or more segmental or arm-level aberrations are classified as malignant, aCGH
classified most lesions as benign (13/16), whereas sWGS and methylation profiling resulted in
malignant classifications in 5/16 and 7/16 cases, respectively. Six cases showed complete
concordance across all platforms, four showed minor discrepancies, and six displayed major
discrepancies that shifted the diagnostic category. Minor discrepancies were defined as differences
in the number or type of detected CNAs that did not alter the resulting molecular classification,
whereas major discrepancies were defined as discordances that shifted the diagnostic category
(benign, borderline, or malignant). These divergences were primarily driven by additional segmental
or focal CNAs detected by sWGS or methylation profiling, underscoring the challenge of applying
CGH-based criteria to higher-resolution platforms.

Figure 1: Cross-platform comparison of CNV detection by aCGH, sWGS, and methylation profiling. (**A**) Heatmap showing the number of chromosomal, arm-level, and segmental CNVs detected per case across all three platforms. High-burden cases demonstrate strong cross-platform concordance, whereas low-burden lesions display additional focal alterations, predominantly detected by methylation arrays.

(**B**) Exemplary CNA profiles of chromosome 16 from case 6. While aCGH identifies a whole-chromosome gain, both sWGS and methylation profiling detect only a long segmental gain on 16q and do not call the gain on 16p, illustrating platform-specific differences in resolution and segmentation. CGH data are displayed on a $\log_2$ ratio scale, whereas for sWGS and methylation profiling the y-axis represents estimated absolute copy number on a linear scale, with 2 corresponding to diploidy.

3.3 Correlation of clinical outcome

Clinical follow-up was reviewed to assess the relationship between molecular classifications and
patient outcomes. Interpretation was limited in patients with giant congenital nevi containing
multiple proliferative nodules, as the excised lesion might not represent the biologically relevant
precursor of metastasis.

This issue was most evident in patient 7, from whom three distinct nodules were sampled over time
(cases 7a, 7b, and 7c). Cases 7a and 7b showed differing CNA patterns across platforms but were
classified as benign by aCGH (and by methylation for 7b). In contrast, case 7c represented the lymph
node metastasis and exhibited a highly complex CNA profile characteristic of melanoma. Neither 7a
nor 7b shared the CNA pattern of the metastatic lesion, suggesting that the true primary melanoma
node was not sampled. Thus, the molecular classifications of 7a and 7b cannot be directly correlated
with the patient's adverse clinical course; these findings suggest they represent independent events.

Other discordant cases showed similar diagnostic uncertainty. In case 1, aCGH and sWGS classified

the lesion as benign, whereas methylation profiling detected arm-level deletions consistent with a

malignant profile. The patient ultimately died; however, due to the multifocal nature of the

congenital nevus, it remains unclear whether the analyzed nodule represented the lesion that gave

rise to metastasis. In contrast, cases 5 and 6 showed benign CNA profiles by aCGH but malignant-

appearing profiles by sWGS and methylation analysis; both patients remained disease-free. Case 13

showed borderline results by aCGH and sWGS but a malignant CNA profile derived from methylation

sequencing; no clinical follow-up was available.

Across the cohort, three patients experienced fatal outcomes. Because several of these patients had

multiple nodules and the true primary melanoma focus often could not be unequivocally identified,

formal performance metrics such as sensitivity and specificity could not be reliably established at the

lesion level. Qualitatively, aCGH tended to classify fewer lesions as malignant than sWGS and

methylation profiling, particularly in patients with an overall benign clinical course, whereas the

higher-resolution platforms more often yielded malignant CNA patterns, including in lesions from

patients who remained disease-free. These observations highlight both the increased sensitivity of

sWGS and methylation-based CNA analysis and the difficulty of attributing biological significance to

small focal aberrations in multifocal congenital nevi where the causative melanoma precursor may

remain unsampled.

Table 3. Molecular classifications based on CNA patterns detected by aCGH, sWGS, and methylation

profiling. Shown are the numbers of whole-chromosome (chr), chromosome-arm (arm), and

segmental (seg) alterations per platform, the resulting molecular classification (B = benign, BL =

borderline, M = malignant), and the degree of discrepancy between methods. **Abbreviations:** chr –

chromosome; arm – chromosome arm; seg – segment; aCGH – array comparative genomic

hybridization; sWGS – shallow whole-genome sequencing; B – benign; BL – borderline; M –

malignant.

Case	CGH	sWGS	Methylation	Discrepancy
1	4 chr – B	4 chr – B	1 chr, 3 arm – M	major
2	8 chr, 1 arm, 3 seg – M	13 chr, 2 arm, 4 seg – M	12 chr, 3 arm, 3 seg – M	minor
3	5 chr – B	5 chr – B	5 chr – B	no
4	7 chr – B	7 chr – B	7 chr – B	no
5	1 chr, 1 seg – B	1 chr, 3 arm, 9 seg – M	2 chr, 2 arm, 7 seg – M	major
6	14 chr – B	10 chr, 1 seg – B	9 chr, 4 seg – M	major
7a	11 chr, 1 arm – B	8 chr, 2 arm – M	8 chr, 4 seg – M	major
7b	1 chr – B	1 chr, 3 seg – M	1 chr – B	major
7c	7 chr, 2 arm, 1 seg – M	7 chr, 2 arm, 1 seg – M	7 chr, 2 arm, 1 seg – M	no
8	11 chr, 1 arm – B	11 chr, 1 arm – B	11 chr, 1 arm – B	no
9	none – B	none – B	1 seg – B	minor
10	9 chr – B	9 chr – B	9 chr – B	no
11	none – B	none – B	1 arm – B	minor
12	6 chr – B	6 chr – B	6 chr – B	no

13	1 seg – **BL**	1 seg – **BL**	3 seg – **M**	major
14	none – **B**	none – **B**	1 seg – **B**	minor

4. DISCUSSION

Although aCGH, sWGS, and methylation-based CNA analysis have often been described as
comparable approaches for the detection of larger CNAs, our study demonstrates that relevant
differences emerge when these methods are applied to atypical melanocytic proliferations in
congenital nevi. Previous studies have reported high concordance between aCGH and next-
generation sequencing (NGS)–based approaches for detecting broad CNAs in well-characterized
genomic regions [21–25]. In contrast, we observed substantial variability in the detection of
segmental and focal aberrations, which in several cases resulted in divergent molecular
classifications. These discrepancies likely reflect differences in technical resolution, genomic
coverage, and analytical sensitivity across the platforms [21–25].

Interpreting molecular findings in this clinical setting is inherently challenging, particularly in patients
with giant congenital nevi who frequently harbor multiple proliferative nodules. In such cases, it is
often unclear whether the sampled lesion corresponds to the biologically relevant precursor of
metastasis. This issue has been repeatedly emphasized in the literature [10,12] and was illustrated
most clearly in our cohort by patient 7. This aligns with prior reports noting that melanomas arising in
congenital nevi often develop within complex, heterogeneous nodular fields where histology and
molecular profiles may diverge [2,10–12].

In this context, it is important to emphasize that current diagnostic criteria for molecular
classification in congenital nevi and proliferative nodules were developed using CGH [10–13].
Numerical aberrations involving whole chromosomes are considered typical for benign proliferative
nodules, while melanomas are characterized by multiple segmental aberrations [2,10–12]. Because
these criteria were validated in cohorts assessed by CGH, they may not directly translate to higher-
resolution platforms. sWGS and methylation profiling exhibit increased sensitivity for detecting small
CNAs or subclonal aberrations (<100 kb), particularly in FFPE tissue [16,26–28]. While these
additional findings might capture biologically meaningful alterations, they might also represent low-
level or focal changes without prognostic significance, raising the possibility of diagnostic
overinterpretation [16,26–28]. In the other direction, in tissue sections selected for DNA extraction
from ambiguous neoplasms, anomalous mosaic clones may be present at levels too low to meet
detection thresholds, a limitation that has been described for both array-based and low-pass
sequencing approaches [29,30].

Our findings therefore support the need for platform-adapted diagnostic frameworks. High-
resolution platforms such as sWGS and methylation arrays provide technical advantages—including
improved genomic resolution and the ability to extract CNA data from the same sample used for
methylation classification—but their increased sensitivity must be balanced against the risk of
upstaging otherwise benign lesions. Larger, prospectively collected cohorts including unequivocally
benign proliferative nodules and melanomas arising within congenital nevi will be essential to
delineate which CNA patterns detected by higher-resolution methods carry true biological and

prognostic relevance. Targeted validation with orthogonal techniques like digital PCR, FISH or MLPA,
or deeper coverage with targeted NGS will be necessary to establish thresholds of reliability.

This study has several important limitations. First, the cohort size is inherently small due to the rarity
of proliferative nodules and melanomas arising in congenital nevi, which restricts the statistical
power of the analysis. Second, many patients had multiple nodules, and in some cases—including
patient 7—the exact melanoma precursor could not be identified, preventing definitive lesion-level
outcome correlation. Third, inherent differences between the platforms—including resolution, noise
characteristics, and algorithmic approaches—likely contributed to discordances. Finally, the
retrospective use of archival FFPE samples introduces variability in DNA quality and tumor content,
which may influence CNA detection.

In summary, while all three platforms reliably captured broad chromosomal alterations, sWGS and
methylation arrays identified substantially more focal and segmental CNAs. These differences
significantly influenced molecular classification and must be interpreted cautiously, particularly in the
setting of multinodular congenital nevi where the melanoma precursor may remain unsampled. Until
platform-specific diagnostic criteria are established and supported by larger outcome-linked cohorts,
CGH remains the most conservative and clinically validated approach for risk stratification in
proliferative nodules arising within congenital nevi.

Literature

1. Ferrari A, et al. Defining and listing very rare cancers of paediatric age: Consensus of the Joint
Action on Rare Cancers. *Eur J Cancer*. 2019;110:120–126. doi:10.1016/j.ejca.2018.12.030

2. Merkel EA, Mohan LS, Shi K, et al. Paediatric melanoma: Clinical update, genetic basis, and
advances in diagnosis. *Lancet Child Adolesc Health*. 2019;3(9):646–654. doi:10.1016/S2352-
4642(19)30168-4

3. McMullan P, Grant-Kels JM. Childhood and adolescent melanoma: An update. *Clin Dermatol*.
2025;43(1):16–23.

4. Yousif R, Boull C, Gerami P, et al. The demographics and trends in pediatric melanoma in the
United States: An analysis of the National Cancer Database. *Pediatr Dermatol*. 2021;38(5):1191–
1197. doi:10.1111/pde.14742

5. Pampena R, Piccolo V, Muscianese M, et al. Melanoma in children: A systematic review and
individual patient meta-analysis. *J Eur Acad Dermatol Venereol*. 2023;37(9):1758–1776.
doi:10.1111/jdv.19276

6. Hawryluk EB, Moustafa D, Bartenstein D, et al. A retrospective multicenter study of fatal pediatric
melanoma. *J Am Acad Dermatol*. 2020;83(5):1274–1281. doi:10.1016/j.jaad.2019.10.064

7. Pappo AS, McPherson V, Pan H, et al. A prospective, comprehensive registry that integrates
molecular analysis of pediatric and adolescent melanocytic lesions. *Cancer*. 2021;127(20):3825–3831.
doi:10.1002/cncr.33652

8. Phadke PA, Rakheja D, Le LP, et al. Proliferative nodules arising within congenital melanocytic nevi:
A histologic, immunohistochemical, and molecular analysis of 43 cases. *Am J Surg Pathol*.
2011;35(5):656–669. doi:10.1097/PAS.0b013e31821243bd

9. Vergier B, Laharanne E, Prochazkova-Carlotti M, et al. Proliferative nodules vs melanoma arising in
giant congenital melanocytic nevi during childhood. *JAMA Dermatol*. 2016;152(10):1147–1151.
doi:10.1001/jamadermatol.2016.2195

10. Pavlova O, et al. 5-hydroxymethylcytosine expression distinguishes proliferative nodules arising in
congenital nevi from malignant melanoma. Am J Surg Pathol. 2016;40(9):1226–1234.
doi:10.1097/PAS.0000000000000671

11. Yélamos O, Arva NC, Obregon R, et al. A comparative study of proliferative nodules and lethal
melanomas in congenital nevi from children. *Am J Surg Pathol*. 2015;39(3):405–415.
doi:10.1097/PAS.0000000000000360

12. Bastian BC, Xiong J, Frieden IJ, et al. Genetic changes in neoplasms arising in congenital
melanocytic nevi: Differences between nodular proliferations and melanomas. *Am J Pathol*.
2002;161(4):1163–1169. doi:10.1016/S0002-9440(10)64394-4

13. North JP, Vemula SS, Bastian BC. Chromosomal copy number analysis in melanoma diagnostics.
Methods Mol Biol. 2014;1102:199–226. doi:10.1007/978-1-62703-727-3_17

14. Alomari AK, Miedema JR, Carter MD, et al. DNA copy number changes correlate with clinical
behavior in melanocytic neoplasms: Proposal of an algorithmic approach. *Mod Pathol*.
2020;33(7):1307–1317. doi:10.1038/s41379-020-0505-9

15. Geiersbach KB, Gliem TJ, Jenkins SM, et al. Single-nucleotide polymorphism array for
histologically ambiguous melanocytic tumors. *J Mol Diagn*. 2022;24(11):1160–1170.
doi:10.1016/j.jmoldx.2022.08.008

16. Cho S, Kim HS, Zeiger MA, Umbricht CB, Cope LM. Measuring DNA copy number variation using
high-density methylation microarrays. *J Comput Biol*. 2019;26(4):295–304.
doi:10.1089/cmb.2018.0175

17. Daenekas B, Pérez E, Boniolo F, et al. Conumee 2.0: Enhanced copy-number variation analysis
from DNA methylation arrays for humans and mice. *Bioinformatics*. 2024;40(2):btae029.
doi:10.1093/bioinformatics/btae029

18. Vasimuddin M, Misra S, Li H, Aluru S. Efficient architecture-aware acceleration of BWA-MEM for
multicore systems. *Proc IEEE Int Parallel Distrib Process Symp*. 2019;314–324.
doi:10.1109/IPDPS.2019.00042

19. Krueger F, Andrews SR. Bismark: A flexible aligner and methylation caller for bisulfite-seq
applications. *Bioinformatics*. 2011;27(11):1571–1572. doi:10.1093/bioinformatics/btr167

20. Demidov G, Ossowski S. ClinCNV: Novel method for allele-specific somatic copy-number
alteration detection. *bioRxiv*. 2019; doi:10.1101/2019.12.20.884791

21. Wang J, Lan Y, Qi HY, et al. Comparison of fluorescence in situ hybridization, next-generation
sequencing, and DNA methylation microarray for copy number variation assessment in gliomas. *Lab
Invest*. 2025;105(7):104168. doi:10.1016/j.labinv.2024.104168

22. Hayes JL, Tzika A, Thygesen H, et al. Diagnosis of copy number variation by Illumina next
generation sequencing is comparable in performance to oligonucleotide array comparative genomic
hybridisation. *Genomics*. 2013;102(3):174–181. doi:10.1016/j.ygeno.2013.06.003

23. Wang N, Tao ZY, Wu T, et al. Benchmarking copy number variation detection with low-coverage
whole-genome sequencing. *Brief Bioinform*. 2025;26(5):bbaf514. doi:10.1093/bib/bbaf514

24. Coutelier M, Holtgrewe M, Jäger M, et al. Combining callers improves the detection of copy
number variants from whole-genome sequencing. *Eur J Hum Genet*. 2022;30(2):178–186.
doi:10.1038/s41431-021-00939-9

25. Scheinin I, Sie D, Bengtsson H, et al. DNA copy number analysis of fresh and formalin-fixed
specimens by shallow whole-genome sequencing with identification and exclusion of problematic
regions. *Genome Res*. 2014;24(12):2022–2032. doi:10.1101/gr.175141.114

26. Mesbah Ardakani N, Thomas C, Robinson C, et al. Comparison of copy number variation detection
in FFPE samples using CGH and methylation arrays. *Pathology*. 2017;49(3):285–291. doi:
10.1016/j.pathol.2016.11.008

27. Bauer J, Bastian BC. Genetic and pathologic aspects of melanocytic neoplasms in congenital nevi.
Dermatol Ther. 2006;19(1):40–49. doi:10.1111/j.1529-8019.2005.00056.x

28. Van der Linden M, Raman L, Vander Trappen A, et al. Evaluation of copy number variation
analysis in pediatric melanocytic tumors using methylation arrays. *Arch Pathol Lab Med*.
2019;143(12):1540–1547. doi:10.5858/arpa.2018-0401-OA

29. Robberecht, C., Fryns, J. P., & Vermeesch, J. R. (2010). Piecing together the problems in
diagnosing low-level chromosomal mosaicism. Genome medicine, 2(7), 47. doi:10.1186/gm168

527
528 **30.** Liu Y, Hao S, Guo X, et al. Accuracy and depth evaluation of clinical low pass genome sequencing
529 in the detection of mosaic aneuploidies and CNVs. *BMC Med Genomics*. 2023;16:294.
530 doi:10.1186/s12920-023-01703-8