Integrative screening identifies functional variants and VNTRs underlying GWAS signals at the 5p15.33 multi-cancer susceptibility locus

Aidan O'Brien[1,2], Hyunkyung Kong[1], Harsh Patel[1], Michelle Ho[1], Minal B. Patel[1], Jun Zhong[1], Mai Xu[1], Brenen W. Papenberg[1], Katelyn E. Connelly[1], Irene Collins[1], Rebecca Hennessey[1], Rohit Thakur[1], Hayley Sowards[1], Karen Funderburk[1], Thong Luong[1], Oscar Florez-Vargas[1], Timothy Myers[3], Ashley Jermusyk[1], Bryan Gorman[4], Wen Luo[5], Kristine Jones[5], Sudipto Das[6], The Pancreatic Cancer Cohort and Case-Control Consortia (PanScan/PanC4), Melanoma Meta-Analysis Consortium, International Lung Cancer Consortium (ILCCO), Qing Lan[7], Nathaniel Rothman[7], James D. McKay[8], Rayjean J. Hung[9], Christopher I. Amos[10], Mark M. Iles[11], Stella Koutros[12], Maria Teresa Landi[4,13], Matthew H. Law[14,15,16], Rachael Z. Stolzenberg-Solomon[17], Brian M Wolpin[18], Manal Hassan[19], Alison P. Klein[20], Samuel O Antwi[21,22], Nick Orr[2], Stephen J. Chanock[23], Sara Lindström[24,25], Jason W. Hoskins[1], Marc-Henri Stern[26], Thorkell Andresson[6], Jianxin Shi[27], Ludmila Prokunina-Olsson[1], Jiyeon Choi[1], Kevin M. Brown[1,28], Laufey T. Amundadottir[1,28]

[1]Laboratory of Translational Genomics, Division of Cancer Epidemiology and Genetics, National Cancer Institute, Rockville, MD, 20850, USA, [2]The Johnston Cancer Research Centre, School of Medicine, Dentistry and Biomedical Sciences, Queen's University Belfast, Belfast, UK, [3]Laboratory of Genetic Susceptibility, Division of Cancer Epidemiology and Genetics, National Cancer Institute, Rockville, MD, 20850, USA, [4]Trans-Divisional Research Program, Division of Cancer Epidemiology and Genetics, National Cancer Institute, Rockville, MD, 20850, USA, [5]Cancer Genomics Research Laboratory, Division of Cancer Epidemiology and Genetics, National Cancer Institute, Rockville, MD, 20850, USA, [6]CCR Protein and Metabolite Characterization Core (PMCC), Frederick National Laboratory for Cancer Research, National Cancer Institute, Frederick, MD, USA, [7]Occupational and Environmental Epidemiology Branch, Division of Cancer Epidemiology and Genetics, National Cancer Institute, Rockville, MD, 20850, USA, [8]International Agency for Research on Cancer, World Health Organization, Lyon, France, [9]Lunenfeld-Tanenbaum Research Institute, Sinai Health, Toronto, Canada, [10]Department of Internal Medicine, University of New Mexico, Albuquerque, NM, [11]Leeds Institute for Data Analytics, University of Leeds, Leeds, UK, [12]Occupational and Environmental Epidemiology Branch, Division of Cancer Epidemiology and Genetics, National Cancer Institute, Rockville, MD, USA, [13]Integrative Tumor Epidemiology Branch, Division of Cancer Epidemiology and Genetics, National Cancer Institute, Rockville, MD, 20850, USA, [14]Statistical Genetics, QIMR Berghofer Medical Research Institute, 300 Herston Road, Brisbane, QLD, 4006, Australia, [15]School of Biomedical Sciences, Faculty of Health, Queensland University of Technology, Brisbane, QLD, 4001, Australia, [16]School of Biomedical Sciences, Faculty of Medicine, University of Queensland, Brisbane, QLD, 4072, Australia, [17]Metabolic Epidemiology Branch, Division of Cancer Epidemiology and Genetics, National Cancer Institute, Rockville, MD, 20850, USA, [18]Department of Medical Oncology, Dana-Farber Harvard Cancer Center, USA, [19]Department of Gastrointestinal Medical Oncology, Division of Cancer Prevention and Population Sciences, MD Anderson Cancer Center, Houston, TX, USA, [20]Departments of Oncology, Pathology and Medicine, Sidney Kimmel Comprehensive Cancer Center, Johns Hopkins School of Medicine, Baltimore, MD, USA, [21]Department of Quantitative Health Sciences, Division of Epidemiology,, Mayo Clinic, Jacksonville, FL, USA, [22]Department of Internal Medicine, Division of Gastroenterology and Hepatology, Mayo Clinic, Jacksonville, FL, USA, [23]Division of Cancer Epidemiology and Genetics, National Cancer Institute, Rockville, MD, 20850, USA, [24]Department of Epidemiology, University of Washington, Seattle, WA, USA, [25]Public Health Sciences Division, Fred Hutchinson Cancer Center, Seattle, WA, USA, [26]Inserm U1339 UMR3666, DNA Repair and Uveal Melanoma (D.R.U.M.), Institut Curie, PSL Research University, Paris, France, [27]Biostatistics Branch, Division of Cancer Epidemiology and Genetics, National Cancer Institute, Rockville, MD, 20850, USA, [28]These authors jointly supervised this work

Abstract

Chromosome 5p15.33 harbors several independent association signals which

demonstrate antagonistic pleiotropy across cancer types, with causal mechanisms

largely unresolved. To identify functional variants and enhancer elements at this locus,

we performed statistical fine-mapping followed by massively parallel reporter assays

(MPRA) and proliferation based CRISPRi screens. This approach identified eight multi-

cancer functional variants (MCFVs) across three GWAS signals. Targeting rs421629

(part of the *CLPTM1L* signal marked by rs465498) with CRISPRi revealed opposing

effects on *TERT* expression in pancreatic versus lung cancer cells, consistent with the

antagonistic pleiotropy observed for this signal. Furthermore, CRISPRi nominated an

intronic *CLPTM1L* variable number tandem repeat (VNTR) as a potent enhancer. Long-

read sequencing established VNTR polymorphisms as potential causal variants for the

rs465498 signal. We showed that Hippo-pathway transcription factors mediate VNTR

enhancer activity in lung and pancreatic cancer cells. Together, these findings indicate

that cancer susceptibility at 5p15.33 may be mediated by both SNPs and VNTRs and

provide an integrated framework for resolving complex pleiotropic loci.

Introduction

While genome-wide association studies (GWAS) have identified hundreds of common risk loci for human cancers, functional characterization of the association signals has lagged behind their identification[1]. Many such loci are thought to be underpinned by causal variants which influence allele-specific *cis*-regulation of nearby target genes[2,3]. Chromosome 5p15.33 is one of the most replicated cancer susceptibility loci, where at least 10 distinct signals have been identified across as many cancer types[4,5]. Successive multi-cancer fine-mapping analyses by Wang and colleagues in 2014[4] and Chen and colleagues in 2021[5] have applied the Association Analysis for SubSETs (ASSET)[6] method to GWAS datasets at the 5p15.33 locus, noting considerable genetic correlation between cancer types. Intriguingly, signals at 5p15.33 demonstrate antagonistic pleiotropy, with risk alleles for one cancer being protective for other cancer types[4,5,7].

At 5p15.33, both *TERT* and *CLPTM1L* are plausible target genes with established roles in tumor development and progression. *TERT* encodes the catalytic subunit of telomerase reverse transcriptase, which maintains telomere length and protects chromosomes from replication-associated erosion. Telomerase activation via upregulation of *TERT* is observed in over 90% of cancers[8–10]. Furthermore, *TERT* represents an established cancer susceptibility gene via both rare and common germline promoter polymorphisms[11–16]. Somatic *TERT* promoter mutations are prognostic indicators for multiple cancers[17–21], and act synergistically with proximal germline polymorphisms (rs2853669) in melanoma[22], bladder cancer[18,23], and glioma[21,24]. Notably, cancers such as pancreatic ductal adenocarcinoma (PDAC) exhibit activated telomerase activity[25] despite lacking canonical *TERT* promoter mutations[26], suggesting alternate *cis*-regulatory activating mechanisms. *CLPTM1L* encodes a transmembrane lipid scramblase which is overexpressed in pancreatic, ovarian, and lung cancers and contributes to RAS-dependent transformation[27], acting as an anti-apoptotic factor upon endoplasmic reticulum stress[28–31]. Recent functional analyses[32,33] and transcriptome-wide association studies[34] further implicate *CLPTM1L* expression as

a potential mediator of risk for uveal melanoma and pancreatic cancer, respectively.

Together, these observations raise the possibility that 5p15.33 risk alleles converge

through expression of *TERT*, *CLPTM1L*, or both, motivating a high-throughput, cross-

cancer functional screen of cancers associated with the 5p15.33 locus using

methodologies sensitive to downstream effects on each gene.

Post-GWAS studies focusing on specific association signals and cancer types have

characterized *cis*-regulatory effects of functional sequence variants on both *TERT* and

CLPTM1L as effector genes[7,12,32,35–40]. More recently, data presented by Florez-Vargas

and colleagues[41] suggests that variable number tandem repeats (VNTRs) may

represent an important class of credible causal variants (CCVs) at 5p15.33[42–44],

indicating that SNP-centric models of GWAS causality may overlook more cryptic

classes of functional variants. With no extensive cross-cancer screen of regulatory

elements and CCVs performed for the larger region to date, we set out to

comprehensively characterize *cis*-regulatory elements and CCVs to better understand

the mechanisms underpinning cancer risk and pleiotropy at this important multi-cancer

susceptibility locus.

To systematically characterize functional variants and *cis*-regulatory elements important

for putative 5p15.33 effector genes, we performed an integrative functional screen at

this locus. First, we fine-mapped GWAS datasets from melanoma, lung, pancreatic, and

bladder cancers and compiled correlated variants from previously published signals to

derive a panel of cross-cancer CCVs. We then implemented a massively parallel

reporter assay (MPRA) to assess allele-specific regulatory activity for CCVs in cell lines

representative of the four fine-mapped cancers. Next, we agnostically identified *cis*-

regulatory elements using a tiled CRISPR-inhibition (CRISPRi) proliferation screen in

cell lines from the same four cancer types. By integrating MPRA and CRISPRi results,

we derived multi-cancer functional variants (MCFV) which mediate their effect through

cis-regulation of *TERT* and/or *CLPTM1L* across cancer types. In addition, CRISPRi

screening nominated a variable number tandem repeat (VNTR) within intron 9 of

CLPTM1L as a potent enhancer element. Long-read sequencing, followed by imputation

of VNTR alleles into a pancreatic cancer GWAS dataset, revealed that polymorphisms

of this element contribute to the credible set of variants underlying the rs465498 signal

at *CLPTM1L*. Subsequent functional analyses revealed VNTR-mediated regulation of

both *TERT* and *CLPTM1L* through Hippo signaling pathway transcription factors.

Collectively, our work applies and extends a generalizable framework for integrative

high-throughput functional screens to assess causal regulatory mechanisms at complex

GWAS loci. By expanding our scope to repeat polymorphisms, we demonstrate that

VNTRs and SNPs may jointly underlie cancer susceptibility signals through *cis-*

regulation of both *TERT* and *CLPTM1L*. In doing so, our work bridges a gap between

statistical genetics and functional biology, providing new insights into the molecular

basis of cancer heritability at one of the most recurrent and complex cancer

susceptibility loci identified through GWAS.

Results

Fine-mapping reveals multiple credible sets at 5p15.33

To comprehensively compile credible causal variants (CCVs) for massively parallel reporter assay (MPRA) analysis, we performed ancestry-specific fine-mapping of GWAS summary statistics from four 5p15.33-implicated cancer types in populations of European ancestry using SuSiE[45] to derive 95% credible causal sets (PDAC: 9,040 cases and 12,496 control subjects[46], melanoma: 36,760 cases and 375,188 controls[47], bladder cancer: 13,790 cases and 343,502 controls[48] and lung adenocarcinoma: 29,266 cases and 56,450 controls[49]). We also assessed a lung adenocarcinoma GWAS of 21,658 cases and 150,676 controls of East Asian[50] (EAS) ancestry (**Table 1, Fig. 1, Supplementary Figs. 1&2)**. These credible sets were denoted by their lead variant (highest posterior inclusion probability, PIP) and assigned to individual pleiotropic ASSET signals reported by Chen and colleagues[5] based on pairwise LD between credible set lead variants and ASSET tag variants ($r^2 \geq 0.6$, 1000 Genomes Project Phase 3, 1kGP p3[51], **Fig. 2, Methods**).

Across studies, SuSiE resolved 11 cancer-specific credible sets (ranging from 1–19 variants; **Table 1, Supplementary Fig. 2**), with 10/11 credible sets recapitulating previously reported ASSET signals (**Table 1**). Lung EUR yielded four credible sets, consistent with previously reported signals[49]. Lung EAS, by contrast, yielded a single credible set marked by rs2736100. Fine-mapping for bladder cancer resulted in a single credible set mapping to the signal marked by rs10069690. Additionally, SuSiE fine-mapping nominated a melanoma signal marked by rs2853667 that was not captured by ASSET (**Table 1**). The rs465498-tagged signal mapping across *CLPTM1L,* notably yielding a larger number of CCVs relative to other signals, was supported in PDAC, melanoma, and lung EUR GWAS, with antagonistic pleiotropy consistent with previous analyses (**Supplementary Fig. 2f**). SuSiE did not return a credible set for several previously reported association signals (PDAC rs2736098[52], lung EAS rs1316780 and rs62332591[50]; **Supplementary Fig. 2.**). In two instances the best-matching ASSET

signal was supported only by modest LD with these SNPs (PDAC rs2735940 with

rs2853677, r^2 = 0.62; melanoma rs2736100 with rs7705526, r^2 = 0.51; **Table 1**).

To ensure comprehensive coverage for our MPRA design, we supplemented ASSET

signals and cancer-specific CCVs with ancestry-matched proxy variants from 1kGP p3

(r^2 ≥ 0.6) to derive a final MPRA panel of 116 5p15.33 CCVs (**Fig. 1a; Supplementary**

Fig. 1, Supplementary Table 1). Given considerable cross-correlation of SuSiE

credible sets (**Supplementary Fig. 3**), subsequent functional annotation of CCVs was

stratified by ASSET signal, with SuSiE results used supplementarily to ensure

comprehensive functional annotation across pleiotropic signals.

MPRA identifies functional *cis*-regulatory CCVs

To assess allele-specific *cis*-regulatory activity for each of the 116 fine-mapped CCVs,

we utilized an MPRA strategy based on previously established workflows[53–56] across

eight cancer cell lines representative of the four fine-mapped cancers (pancreas: PANC-

1 and MIA PaCa-2; melanoma: UACC903 and UACC1113; bladder: 5637 and UM-UC-

3; lung: A549 and NCI-H520). After quality control (see **Methods, Supplementary Fig.**

4), 61.21% of tested CCVs demonstrated a significant allelic effect (False Discovery

Rate, FDR ≤ 0.01) in at least one cell line (**Supplementary Table 2**), although effect

sizes were generally modest (median allelic $\log_2$ fold change for significant CCVs was

0.067, **Supplementary Fig. 5**). We therefore prioritized CCVs using additional

stringency metrics in a step-wise manner: (i) active element status for the 145 bp

sequence harboring each variant determined by divergence of either allele from a

distribution of scrambled controls (n = 79 of 109 CCVs passing MPRA design and QC;

see **Methods; Supplementary Fig. 6**) (ii) FDR ≤ 0.01 for allelic effects (n = 71), (iii)

absolute allelic Δ RNA/DNA TPM ≥ 15% (n = 13). Variants meeting all three criteria in at

least one cell line were designated MPRA-hits (n = 13; **Table 2**). Effect sizes were

concordant with independent MPRA experiments of overlapping CCVs in UACC903

cells by Choi and colleagues[55] (Pearson's r = 0.88, n = 5), and lung cancer cell lines by

Long and colleagues[54] (A549 r = 0.84, NCI-H520 r = 0.82, n = 17). MPRA-functional

CCVs mapped to five ASSET signals (marked by rs465498: n = 7 CCVs; rs11414507:

2; rs7705526: 1; rs7726159: 2; rs451360: 1; **Supplementary Table 3).** We annotated

CCVs with functional scores from RegulomeDB[57], FORGEdb[58], and assessed

disruption/creation of transcription factor binding sites with motifbreakR[59]

(**Supplementary Table 4**), but did not observe robust enrichment of MPRA hits among

higher-scoring or motif-disrupting candidates (see **Supplementary Results**),

underscoring the value of direct functional screening at this locus.

A tiled CRISPRi screen at 5p15.33 supported MPRA-hits and revealed additional regulatory elements

Because episomal MPRA constructs are not integrated into the endogenous chromatin

context, we complemented this approach with an agnostic, tiled CRISPRi proliferation

screen of 25,778 sgRNAs across this locus (chr5:1,233,287-1,380,000, GRCh37)

ensuring robust coverage of CCVs across the sgRNA pool (**Supplementary Table 5,**

Supplementary Results). We used representative cell lines from the same four cancer

types engineered to stably express dCas9-KRAB-ZIM3, retaining seven of the eight

lines described above and substituting TCCSUP for 5637 for bladder cancer (see

Methods). After confirming both *TERT* and *CLPTM1L* as proliferation-essential genes

in these cell lines (consistent with data from DepMap[60], knockdown data not presented),

CRISPRi screens were carried out over 25 cumulative cell doublings with sampling at

T_0, T_1 (~12.5 cell doublings) and T_2 (~25 doublings). Resultant screen data was

analyzed with MAGeCK[61], demonstrating robust depletion of sgRNAs targeting the

promoters of genes selected for proliferation-essentiality and cancer-type susceptibility

status, confirming screen efficacy (**Methods, Supplementary Tables 6-7**). We next

used CRISPR-SURF[62] to deconvolve signals from 5p15.33-tiled sgRNAs across

datapoints into 2,718 proliferation-essential elements at FDR ≤ 0.05 (median length 470

bp; **Supplementary Table 8**), including strong signals intersecting the *TERT* and

CLPTM1L promoters. This nominated a large fraction of the locus as proliferation-

essential (median ~40–50% per cell line per timepoint; **Supplementary Table 9**),

whereas both melanoma lines showed substantially reduced sensitivity (6.1–17.6% of

the locus across timepoints).

To nominate CCVs with the strongest functional support across the four cancers, we

integrated MPRA hits with CRISPRi screening data stratified by pleiotropic ASSET

signals[5] **(Fig. 2)**. We defined CCVs as *functional variants* for a given cancer type upon

concurrent observation of (i) MPRA-hit status and (ii) colocalization with a CRISPR-

SURF-identified regulatory element (FDR ≤ 0.05) in at least one representative cell line

for a given cancer type (**Fig. 2**). We then defined variants as *multi-cancer functional*

variants (MCFVs) upon annotation as functional variants in more than one cancer type,

nominating eight MCFVs. Stratifying MCFVs by corresponding ASSET signal (pairwise

$r^2 \geq 0.6$; 1kGP p3 EUR; **Fig. 2**), the rs465498 signal in *CLPTM1L* again accounted for

the largest set, yielding four MCFVs: rs421629 ($r^2 = 0.98$ with rs465498), rs11133729 (r^2

$= 0.99$), rs31487 ($r^2 = 0.97$), and rs37005 ($r^2 = 0.75$). Notably, rs421629, rs11133729,

and rs31487 were included in the PDAC SuSiE credible set, and rs31487 additionally

mapped to the melanoma credible set (**Fig. 2**). Three MCFVs were detected within

intron 2 of *TERT* (rs7705526 signal: rs4449583 ($r^2 = 0.78$); rs11414507 signal:

rs74682426 ($r^2 = 0.98$) and rs56023411 ($r^2 = 0.62$); **Fig. 2**). Neither of these MCFVs

were captured in SuSiE credible sets but were associated with individual cancer types

(rs4449583: $P_{melanoma} = 2.04 \times 10^{-15}$, $P_{lung\_EUR} = 1.49 \times 10^{-20}$, $P_{lung\_EAS} = 9.82 \times 10^{-43}$;

rs56023411: $P_{melanoma} = 1.27 \times 10^{-17}$). Another *TERT* intron 2 MCFV, rs62332591 was

not correlated to any 2021 ASSET signal[5]) at $r^2 \geq 0.6$, but was most highly correlated to

signal rs2853677 from the 2014 ASSET analysis[4]($r^2 = 0.49$).

Having nominated a panel of MCFVs, we next used SNP-centric individual CRISPRi

guides to link functional variants to their effector gene(s) (**Fig. 3a-b**). Guided by

consistent antagonistic pleiotropy between PDAC and lung cancer in published ASSET

analyses[4,5], we performed CRISPRi to target the respective regions surrounding

MCFVs in representative cell lines (PANC-1 for PDAC; A549 for lung cancer) and

assessed effects on *TERT* and *CLPTM1L* expression by TaqMan RT-qPCR analysis

(**Fig. 3c**; **Methods**). Targeting MCFVs assigned to both the rs7705526 (blue, **Fig. 3**)

and rs11414507 (red, **Fig. 3**) ASSET signals in *TERT*'s intron 2 robustly decreased

TERT expression in tested cells, with weak or no effects on *CLPTM1L*. By contrast,

three MCFVs assigned to the rs465498 signal (purple, **Fig. 3**), comprising CCVs intronic

and upstream of *CLPTM1L*, produced more modest and heterogeneous transcriptional

effects. Perturbation of the respective regions surrounding MCFVs rs31487 and

rs37005 reduced *TERT* expression (with rs37005 also decreasing *CLPTM1L* in A549

cells), whereas perturbation of the region containing rs421629 (within *CLPTM1L* intron

16) yielded opposing *TERT* effects between cell types (increased in PANC-1 but

decreased in A549 for a subset of sgRNAs) (**Fig. 3c**).

A variable number tandem repeat (VNTR) polymorphism in *CLPTM1L* is part of the signal in *CLPTM1L* tagged by rs465498

SNP rs11133729 was identified as a MCFV mapping to the ASSET signal tagged by

rs465498 (r^2 = 0.99, **Fig. 4a**). Located in intron 9 of *CLPTM1L,* this region was

annotated by prominent enhancer/promoter marks across ENCODE cell lines (**Fig. 4a)**

and was nominated as proliferation-essential in 7/8 cell lines (**Fig. 4b**). This SNP maps

within the bounds of a variable number tandem repeat (VNTR) (2,728 bp:

chr5:1,326,295-1,329,022, GRCh38, denoted VNTR1), itself located 371 bp upstream of

a second VNTR (669 bp, denoted VNTR2, chr5:1,329,508-1,330,176, GRCh38), with

both VNTRs comprising distinct 29 bp consensus repeat units (RU) (**Fig. 4c).** Given

emergent appreciation of repeat polymorphisms as underlying drivers of GWAS

association signals[41,63–65], we next aimed to determine whether *CLPMT1L* intron 9

VNTR polymorphisms might represent CCVs at 5p15.33. To achieve this, we utilized a

joint genotyping strategy to infer read-phased VNTR and SNP haplotypes from PacBio

amplicon reads within *CLPTM1L* intron 9 across 484 1kGP EUR samples as well as

DNA samples from 940 patients diagnosed with PDAC and 870 controls from the

PanScan I[66] and II[67] GWAS phases (see **Methods, Supplementary Tables 10-12**). In

the PDAC GWAS cohort, we resolved 459 distinct VNTR1 alleles spanning lengths of

2,149–4,560 bp, with a broadly trimodal frequency distribution (peaks at 2,586 bp, AF =

0.07; 2,789 bp, AF = 0.31; 2,934 bp, AF = 0.31; **Fig. 4d**), and VNTR2 alleles spanning

527–1,830 bp with a bimodal distribution (peaks at 730 bp, AF = 0.42; 933 bp, AF =

0.36; **Fig. 4d**).

Genetic association tests for VNTRs have often grouped alleles as long vs. short to

overcome the challenges in genotyping with traditional sequencing and PCR-based

methods[41,44,68]. Given that TRGT outputs full VNTR sequences as alleles, we assessed

VNTR associations with individual risk signals by encoding VNTR genotypes from the

1,810 PanScan samples in four different ways, with increasing levels of aggregation, as

follows (**Fig. 4d**); (i) discrete VNTR alleles (each unique haploid VNTR sequence as its

own allele, VNTR1_allele_n, VNTR2_allele_n, both represented by allele length

followed by unique index number (n)), (ii) aggregate VNTR lengths (collapsing alleles of

identical length regardless of sequence, VNTR1_length, VNTR2_length), (iii) data-

driven length classes defined by applying k-means clustering to observed allele-length

distributions (k selected based on visual inspection of allele-length distributions,

VNTR1_cluster, VNTR2_cluster, yielding k = 4 for VNTR1 (cluster 1: 2,149–2,673 bp;

cluster 2: 2,702–2,872 bp; cluster 3: 2,876–3,224 bp; cluster 4: 3,253–4,560 bp) and k =

3 for VNTR2 (cluster 1: 527–817 bp; cluster 2: 846–1,020 bp; cluster 3: 1,049–1,803 bp)

and (iv) a binary long/short (VNTR1_LS, VNTR2_LS) encoding obtained by collapsing

adjacent length classes within each VNTR (VNTR1: short = clusters 1–2, long = clusters

3–4; VNTR2: short = clusters 1–2, long = clusters 2-3)). We found VNTR1_LS long

encodings were linked to the PDAC risk haplotype marked by the rs465498 ASSET

signal (rs31490-A, **Fig. 4e**; pairwise LD between VNTR alleles and rs465498; VNTR1

long: r^2 = 0.83, D′ = 0.99; VNTR2 long: r^2 = 0.70, D′ = 0.99), which motivated

subsequent analyses to determine whether VNTR alleles comprise credible causal

variants for this signal.

VNTR alleles associate with PDAC risk

PacBio-derived VNTR alleles were first tested for association with PDAC risk in 1,810

directly sequenced PanScan I and II DNA samples (940 PDAC and 870 controls, see

Methods). Multiple VNTR1 encodings were associated with PDAC (**Supplementary**

Table 13). Among specific VNTR1 alleles, a VNTR1 2,934 bp allele (2934_30) exhibited

the most significant association (VNTR1_allele 2934_30; AF = 0.14, OR = 1.52, *P* =

3.25×10⁻⁵) when compared to all other alleles. Aggregate VNTR1 long alleles also

demonstrated significant association with PDAC (VNTR1_LS long; AF = 0.52, OR =

1.22, *P* = 2.54×10⁻³), with VNTR1-cluster 3 demonstrating comparable results

(VNTR1_cluster 3; AF = 0.44, OR = 1.20, *P* = 7.73×10⁻³). By contrast, both common

and infrequent VNTR2 allele encodings yielded only nominally significant associations,

including VNTR2-long (VNTR2_LS long; AF = 0.55, OR = 1.15, P = 0.05) and a VNTR2

962-bp allele (VNTR2_allele 962; AF = 0.006, OR = 1.50, P = 0.01).

To increase statistical power, we imputed VNTR genotypes into a larger PDAC GWAS

dataset. Using PacBio-derived VNTR encodings and SNP calls from the 1,810 PanScan

I & II samples described above, we imputed each of the four VNTR genotype encodings

(**Fig. 4d**) as distinct multiallelic variants into the remaining samples from the

PanScan/PanC4 PDAC meta-GWAS[66,67] (including an additional 8,066 cases and

11,208 controls; **Methods**). After imputation, we filtered VNTR and SNP alleles (MAF ≥

0.01, imputation quality score R2 ≥ 0.3) prior to application of the same logistic model

applied to directly genotyped samples and subsequent meta-analysis of each cohort. In

agreement with the original GWAS, SNP rs31490 exhibited the most significant

association[46] (**Fig. 5b;** OR = 1.20, P = 2.33x10^{-16}). Results for imputed VNTR alleles

largely replicated the analysis of directly genotyped samples **(Supplementary Table**

14). Of the four VNTR encodings, VNTR1 cluster 3 emerged as the most significant

VNTR association (OR = 1.20, P = 8.86x10^{-15}). We also replicated significant effects for

long/short VNTR1 alleles (OR = 1.20, P = 4.00x10^{-15}). Of the discrete VNTR1 alleles,

the 2,934 bp sequence (2934_30) which previously comprised the most significant

VNTR association in the pilot analysis retained genome-wide significance (OR = 1.25, P

= 1.19x10^{-9}). Collective VNTR2 lengths of 933 bp were the only significant (at P ≤ 5x10^{-8})

VNTR2 encoding (OR = 1.17, P = 5.36x10^{-11}).

Conditioning the association analysis on VNTR1 cluster 3 by including corresponding

VNTR encoding dosages as a fixed covariate resulted in marked attenuation of the

signal for lead SNP rs31490 ($P_{VNTR1\_cluster3}$ = 9.30x10^{-3}). Conditioning on VNTR2 lengths

of 933 bp resulted in more modest attenuation of the rs31490 signal (P_{VNTR2_933bp} =

4.13x10^{-6}). Conditioning the analysis on rs31490 likewise attenuated the VNTR1 cluster

3 association ($P_{rs31490}$ = 0.18). Alternatively, conditioning on SNPs known to tag PDAC

signals confirmed linkage with the rs31490 haplotype (P-values for VNTR1 cluster 3

conditioned on PDAC signal SNPs: $P_{conditional\_rs451360}$ = 1.37x10^{-5}, $P_{conditional\_rs2735940}$ =

3.62x10^{-9}, $P_{conditional\_rs2853677}$ = 2.79x10^{-7}, **Supplementary Table 14**).

VNTR1 is a potent enhancer at the 5p15.33 locus

Given strong evidence that VNTR1 polymorphisms comprise plausible CCVs as part of the rs31490/rs465498-tagging haplotype (**Fig. 4d**), we next sought to characterize the biological mechanism underlying VNTR1-enhancer activity. *In-silico* analysis of VNTR1-transcription factor binding sites using TRAP[69] revealed significant enrichment for TEA Domain Family Member 1 (TEF-1) motifs ($P_{nominal} < 5.0$x10^{-4}, **Fig. 5a**, **Supplementary Table 15**). TEF-1, encoded by *TEAD1,* is a key transcription factor member of the Hippo signaling pathway involved in regulation of apoptosis and proliferation[70–73].

Similar to our approach for small sequence variants, we individually assessed VNTR1 using CRISPRi and individual luciferase reporter assays to assess functionality. CRISPRi repression of VNTR1 reduced expression of both *TERT* and *CLPTM1L* relative to negative control sgRNA in PANC-1 and A549 cells (**Fig. 5c**). VNTR1-sgRNA-A, which directly targets the region harboring rs11133729 in VNTR1 , reduced *TERT* (fold-change = 0.19, $P = 6.69$×10^{-5}) and *CLPTM1L* (fold-change = 0.58, $P = 1.30$×10^{-4}) expression in A549 cells, and similarly decreased *TERT* (fold-change = 0.59, $P = 2.78$×10^{-4}) and *CLPTM1L* (fold-change = 0.72, $P = 1.43$×10^{-2}) expression in PANC-1 cells. Luciferase reporter assays of VNTR1 alleles representative of each of the cluster-genotype encodings confirmed potent enhancer activity in both cell lines (median fold change vs. negative control: PANC-1 = 74; A549 = 609; **Fig. 5d**). Enhancer strength varied across constructs representative of VNTR1 cluster encodings: relative to the shortest tested allele (2,586 bp; assigned to cluster 1), a 2,933 bp sequence assigned to VNTR1 cluster 3 demonstrated higher activity in PANC-1 (fold-change = 1.28, $P = 2.69$×10^{-4}) and A549 (fold-change = 1.20, $P = 2.23$×10^{-2}) cells.

Having established VNTR1 as a potent enhancer for *TERT* and *CLPTM1L*, we proceeded to characterize the transcription factors mediating this effect. We performed DNA pulldown experiments coupled with mass spectrometry by incubating biotinylated VNTR1 repeat units with MIA PaCa-2 nuclear lysate. This analysis identified robust binding of TEF-1 peptides to VNTR1 sequences (**Fig. 5f, Supplementary Table 16**). Notably, competitor experiments led to a striking depletion of TEF-1 (**Fig. 5g**), indicating

highly sequence-specific protein-DNA interactions. This finding was validated by oligo

pulldown experiments followed by western blot showing robust binding of TEF-1

(*TEAD1*) as well as additional Hippo pathway family members TEF-3 (*TEAD4*) and WW

domain-containing transcription regulator protein 1 (*WWTR1*; also named *TAZ*) to the

VNTR1 repeat motif (**Fig. 5h**). Finally, we sought to determine to what degree VNTR1-

luciferase activity was attributable to Hippo-pathway transcription factor (TF) signaling.

Disruption of TEF-YAP complexes with verteporfin, a small molecule inhibitor of TEF-

YAP binding[74,75], markedly reduced VNTR1-driven reporter activity across constructs in

both cell lines (**Fig. 5e**), indicating that VNTR1 enhancer function depends on Hippo-

pathway transcriptional complexes. Collectively, these data support VNTR1

polymorphisms as MCFVs at 5p15.33 with regulatory implications for both *TERT* and

CLPTM1L.

Discussion

Here, we provide a comprehensive dissection of functional *cis*-regulatory CCVs at

5p15.33. By identifying functional variants which mediate causal effects on both target

gene expression and cell proliferation, our work furthers understanding of the

mechanisms underpinning susceptibility for multiple cancers. We derived an integrated

functional map for 116 5p15.33 CCVs and identified eight MCFVs across three distinct

GWAS signals, demonstrating allele preferential *cis*-regulatory effects that localized to

proliferation-essential regulatory elements across cancer cell lines from four cancer

types. Additionally, we build on findings from transcriptome-wide association studies[34]

and post-GWAS functional analyses which support the role of *CLPTM1L* as a possible

target gene for the GWAS signals at 5p15.33.

Antagonistic pleiotropy between cancers has long been a confounding factor at

5p15.33. Context-dependent telomerase biology has been hypothesized as a possible

mediator of such effects, although direct analysis of telomere length across cancer

types has yielded inconclusive results[76–84]. Our analysis marks a major step towards

understanding how these differences may be mediated at a *cis*-regulatory level. For all

eight MCFVs we identified, MPRA allelic effects demonstrated consistent directionalities

across cancer types, suggesting that differences in phenotypic outcome may largely be

driven by effects of downstream target genes and pathways, epigenetic factors, or

potentially by interactions with environmental factors. Our tiled CRISPRi screen

supported this model, with MPRA-functional CCVs intersecting elements nominated as

putative enhancers. Focused interrogation of these variants with CRISPRi proved

largely consistent with this model except for rs421629. Our findings suggest that this

SNP occupies a possible context-dependent regulatory element whose net effect on

TERT may switch between repressive and activating effects depending on the cell type.

An alternative, non-mutually exclusive possibility is that perturbation of this element

reconfigures the local regulatory circuit, such that inhibiting rs421629-dependent activity

unmasks or engages distinct enhancers or promoters in a cell type-specific manner.

This reversal provides a plausible *cis*-regulatory mechanism for antagonistic pleiotropy

at 5p15.33. Motif analysis indicated that the PDAC/melanoma risk allele (rs421629-G)

may introduce high affinity binding motifs for NFATC2, raising the possibility that allele-

specific TF recruitment, combined with lineage-specific cofactor and chromatin

environment, may toggle the element between enhancer- and repressor-like states.

While additional work is required, our data highlights how cell type-specific wiring of a

single *cis*-regulatory element may contribute to antagonistic pleiotropy at 5p15.33.

Our analysis identified a functional VNTR polymorphism as an additional CCV for the

signal at *CLPTM1L* marked by rs465498, where both VNTR and SNP alleles may jointly

underpin the observed association. We characterized VNTR1 as a potent enhancer of

both *TERT* and *CLPTM1L* with activity driven by Hippo pathway TFs. Notably given

antagonistic pleiotropy for this signal, Hippo pathway factors have been implicated in

both oncogenic and tumor suppressive mechanisms across cancer types[85]. We also

note four MCFVs in intron 2 of *TERT* intersecting repeat elements, corresponding to

several independent association signals. Our analysis supports recent studies which

posit VNTRs as a considerable source of missing heritability[86,87]. While post-GWAS

analyses are largely focused on the identification of causal SNPs and their relationships

to target genes, there are several examples of GWAS signals tagging more complex

variants (e.g., in *TERT*[41], *CTRB2*[88], *EGR2*[65]). Repeat elements are pervasive throughout

the genome, with a substantial body of evidence supporting their role as *cis*-regulatory

elements[89–91]. Recent studies have estimated VNTR polymorphisms to underly

prominent glaucoma and colorectal cancer GWAS signals[63]. Moreover, studies have

estimated 11.4% of structural variants and simple tandem repeats to be in LD with

SNPs associated with 701 GWAS traits[86]. Identification of cryptic functional alleles is

particularly pertinent at complex loci such as 5p15.33, where multiple independent

signals and context-dependent regulatory elements confound deeper understanding of

cancer susceptibility. To accelerate functional characterization at such loci, it is

imperative that repeat polymorphisms are included into GWAS imputation reference

panels given recent advances in long-read sequencing technologies.

While this work provides a comprehensive map of cross-cancer regulatory elements at

5p15.33, our approach is not without limitations. While this locus has been implicated in

numerous common cancers, we specifically fine-mapped and functionally interrogated

four representative cancer types. While we assessed 11 pleiotropic signals, functional

screening of only four cancer types may have limited our sensitivity to identify MCFVs

for some signals. Furthermore, all experiments were performed in cancer cell lines,

which do not fully recapitulate the physiological and epigenetic context in which

germline variants act. Future studies should aim to investigate the effects of MCFVs in

primary cells and 3D cell culture models to better understand the mechanisms of

pleiotropy at 5p15.33. In addition, lung cancer GWAS across ancestries have shown

that the signal tagged by rs2736100 is specific to lung adenocarcinoma (EAS), whereas

the signal tagged by rs6554758 is more strongly associated with squamous cell

carcinoma[49]. MCFVs assigned to each signal were supported by evidence from both

adenocarcinoma (A549) and squamous cell carcinoma (NCI-H520) models; however,

subsequent studies should examine how these functional alleles act across lung cancer

subtypes. More broadly, future work should also endeavor to test functional alleles in

contexts that model both genetic and environmental stressors. Because MCFV and

VNTR characterization assays were conducted in *TERT* promoter wild type PANC-1

and A549 cells[7,92], it would also be appropriate to assay the effect of these

polymorphisms in the context of somatically-mutated *TERT* promoters, such as in

UACC903 or UACC1113 melanoma cell lines. Lastly, our CRISPRi screens interrogated

regulatory element function without directly resolving allelic effects; lenti-MPRA or

CRISPR base-editing approaches could quantify allele-specific transcriptional

consequences within native chromatin contexts in models not amenable to transient

transfection.

In summary, our model identifies multiple potential causal variants across the region,

including several linked to the same signals. In addition, this work implicates rs421629

as part of a regulatory element with seemingly dual functions on target gene expression,

highlighting *cis*-regulatory mechanisms as a potential mediatory of antagonistic

pleiotropy at 5p15. Finally, we demonstrate that VNTR polymorphisms contribute to the

CLPTM1L signal marked by rs465498, suggesting contributions from both simple and

complex polymorphisms, as well as demonstrating a potentially broader role of repeat

polymorphism at complex susceptibility loci.

Table 1: Credible sets identified via fine-mapping at 5p15.33 across four cancer types

GWAS Dataset	Most significant 5p15.33 SNP							SuSiE credible sets		Chen et al. 2021 SNP (r^2 1kGP EUR)
	rsID	chr5 Position (GRCh37)	chr5 Position (GRCh38)	Effect Allele	Non-Effect Allele	Odds Ratio	P-Value	CS-tagging CCV	nSNPs	
PDAC[46]	rs31490	1344458	1344343	G	A	0.84	6.02×10^{-17}	rs31490	19	rs465498 (0.96)
								rs2735940	2	rs2853677 (0.62)
Melanoma[47]	rs13178866	1323212	1323097	C	T	0.88	1.10×10^{-43}	rs13178866	11	rs465498 (0.98)
								rs2736100	1	rs7705526 (0.51)
								rs2853667	1	-
Lung EUR[49]	rs7705526	1285974	1285859	G	A	1.25	3.80×10^{-35}	rs7705526	1	rs7705526
								rs6554758	2	rs465498 (0.77)
								rs2853677	1	rs2853677
								rs35033501	2	rs35033501
Lung EAS[50]	rs2736100	1286516	1286401	C	A	0.75	7.92×10^{-58}	rs2736100	1	rs7705526 (0.52, 0.859 EAS)
Bladder[48]	rs2242652	1280028	1279913	A	G	1.08	4.06×10^{-15}	rs10069690	1	rs10069690

rsID corresponds to the most significant 5p15.33 variant for each of the fine-mapped GWAS datasets. SuSiE fine-mapping of 5p15.33-implicated cancer GWAS (PDAC (9,040 cases, 12,496 controls), melanoma (36,760 cases, 375,188 controls), lung cancer (EUR; 29,266 cases, 56,450 controls, EAS; 21,658 cases, 150,676 controls) and bladder cancer (13,790 cases, 343,502 controls).) resulted in 11 credible sets. All but rs2853667 (melanoma) were confidently assigned to a signal-tagging SNP from Chen and colleagues' 2021 ASSET analysis[5]. Considerable overlap between signals was observed (**Supplementary Fig. 3**) with GWAS of varying sample sizes and imputation strategies yielding credible sets of varying sizes. Full fine-mapping results are presented in **Supplementary Table 1**.

431 **Table 2: 5p15.33 MPRA results and functional annotations**

	MPRA-Hit CCVs		Functional Annotations			[†]MPRA Results				
rsID	chr5 Position (GRCh38)	REF>ALT	[‡]Regulome	[‡]FORGE	[f]motifbreakR	PDAC	Melanoma	Lung	Bladder	[*]Number of MPRA positive cell lines
rs10054203	1279849	G>C	4	5	1	↑	↑	↑↑	-	5
rs4449583	1284020	C>T	5	5	7	↑	↑	↑↑	-	5
rs74682426	1289860	A>C	4	3	10	↑	↑	↑↑	-	5
rs62332591	1290204	T>G	1f	6	1	↓↓	↓	↓↓	↓	8
rs56023411	1291625	G>T	4	6	12	↑	↑	↑↑	-	5
rs2736102	1302029	T>C	1f	5	1	-	↓	-	↓	3
rs13170453	1317366	A>G	1b	7	0	-	↓	↓	↓	4
rs421629	1320021	G>A	1b	7	2	↓↓	↓↓	↓	↓	7
rs11133729	1326986	C>G	2b	8	5	↓	-	↓↓	↓	5
rs452384	1330725	T>C	1d	9	1	-	↑	-	↓	2
rs31487	1340986	G>C	1b	5	0	↓↓	↓↓	↓↓	↓↓	7
rs31490	1344343	G>A	1f	9	0	-	-	↓	-	1
rs37005	1356335	C>T	1b	8	10	↓↓	↓	↓↓	-	6

432 [†]↑↑ = ALT allele vs REF strong effect (both cell lines for each cancer type meet criteria: FDR ≤0.01, FC ≥ 15%, Active element), ↑ = Moderate ALT

433 effect (1 cell line meets criteria/cancer), ↓↓ = REF allele strong effect (both cell lines meet criteria), ↓ = REF allele moderate effect (1 cell line

434 meets criteria). [‡]CCV functional scores from RegulomeDB (GRCh38) and FORGEdb. RegulomeDB: integrates eQTL/QTL, chromatin accessibility,

435 footprinting, TF ChIP-seq and motif evidence to assign an inverse rank (1a strongest → 7 weakest); FORGEdb: 0–10 additive evidence scores

436 summing support from eQTL, ABC 3D contacts, DNase hotspots, histone marks, TF motifs, and allele-specific TF occupancy (higher = stronger

437 regulatory support). [f]number of 'strong' transcription factor binding motif events implicated with CCVs as determined by motifbreakR. *Number of

438 cell lines across all cancer CCV where variant was an MPRA-hit.

439 **_Fig. 1_**

Fig. 1: Integrative screening strategy for identification of 5p15.33 multi-cancer functional variants.

a, To collate credible causal variants (CCVs) at 5p15.33 for MPRA, we initially compiled signal-tagging lead variants from two multi-cancer fine-mapping studies (Chen et al. 2021[5]; Wang et al. 2014[4]) along with credible set-tagging variants resulting from SuSiE fine-mapping of GWAS summary statistics from four 5p15.33-implicated cancers. We carried forward lead/tagging variants, additional SuSiE credible-set CCVs, and ancestry-matched LD-correlated variants (1kGP Phase 3; $r^2 \geq 0.6$ for downstream analyses, total 116 CCVs). **b,** Fine mapped CCVs were subject to MPRA to characterize allelic effects on transcriptional output across eight cell lines representative of the four fine mapped cancer types. **c,** Additionally, we assayed across the larger locus to nominate *cis*-regulatory elements using a tiled CRISPRi proliferation screen across eight cell lines spanning four cancer types. **d,** We then integrated results across cancer types to nominate eight multi-cancer functional variants (MCFVs) across three distinct signals.

Fig. 2

Fig. 2: Functional screening reveals multi-cancer functional variants (MCFVs)
across 5p15.33.

a, CCVs were stratified by pairwise LD $r^2 \geq 0.6$ relative to signal-tagging variants

reported by Chen *et al.* 2021[5]. ASSET signals were annotated with a cancer type if

SuSiE fine-mapping yielded a credible set for a given cancer corresponding to that

signal using the same LD threshold. Arrows assigned to each credible set indicate the

relative direction of effect for each fine-mapped cancer ($\uparrow$ risk increasing, $\downarrow$ protective)

(top panel). **b,** CCVs were annotated with RegulomeDB and FORGEdb scores and

motifbreakR predictions of transcription factor motif disruption. MotifbreakR results were

further annotated for concordant TF-ChIP-seq data from the ReMap2022 database[93]

(motifbreakR + ChIP) (middle panel, full annotations in **Supplementary Table 3**). **c,** For

each CCV, binary indicators were assigned per cell line based on MPRA allelic activity

(FDR-significant effect, $\geq 15\%$ absolute fold change, and active-element status) and

overlap with proliferation-essential elements from CRISPRi. CCVs were nominated as

screening hits when MPRA-functional status was supported by CRISPRi within cell lines

from the same cancer type (green filled circles). CCVs were further categorized as

multi-cancer functional variants (MCFVs) when screening hit status was observed in

more than one cancer type (green filled connected circles with black outline; rsIDs

highlighted in red, bottom panels). Functional support was strongest for CCVs mapping

to the rs465498 signal (four MCFVs) over *CLPTM1L* and the rs11414507 signals (two

MCFVs) over *TERT*. rs62332591 was also nominated as a MCFV but was not

correlated at $r^2 \geq 0.6$ to an ASSET signal (and not shown in this figure).

Fig. 3

Fig. 3: MCFV-targeted CRISPRi nominates TERT and CLPTM1L as effector genes.

a, We identified eight MCFVs (rs11133729 assayed as part of subsequent VNTR-

focused analysis) across the four cancer types as MCFVs corresponding to three 2021

ASSET signals (LD r^2 ≥ 0.6 EUR to ASSET signal lead SNP rs7705526 in purple,

rs35226131 in yellow and rs465498 in red, MCFV rs62332591 did not meet LD

correlation threshold to ASSET signal-tagging SNP*). We chose to highlight screening

results and perform subsequent experiments for seven MCFVs in PANC-1 PDAC cells

and A549 lung cancer cells due to antagonistic pleiotropy across signals between these

two cancer types. **b,** MPRA results for MCFVs revealed significant allelic effects on

transcriptional output. Significance indicators reflect P-Value derived from linear model

comparing aggregate normalized RNA/DNA TPM ratios from 35 unique tag sequences

assigned to both reference (blue) and alternate alleles (red). **c,** While our CRISPRi

screening strategy aimed to nominate proliferation-essential elements agnostic to the

location of 5p15.33 CCVs, CCV-focused analysis of MCFV-harboring loci ($\pm$ 100 bp

centered on each CCV) revealed significant depletion of sgRNAs targeting respective

loci in PANC-1 and A549 cells (red dots indicate FDR ≥ 0.05 depletion of sgRNA with y-

axis showing $\log_2$ fold change as determined by CRISPR-SURF following 25 cumulative

screen doublings). **d,** MCFV-targeted CRISPRi using three distinct sgRNAs (x-axis;

sgRNAs A, B and C) reduces *TERT* and *CLPTM1L* mRNA expression relative to a non-

targeting control sgRNA (y-axis; experimental sgRNA $2^{-\Delta\Delta Ct}$ normalized to $2^{-\Delta\Delta Ct}$ of non-

targeting sgRNA, * indicates significance under two-sided t-test). Variants attributable to

the rs465498 ASSET signal across *CLPTM1L* demonstrated more modest effects

relative to *TERT*, with rs421629 showing a striking effect on *TERT* expression across

cell lines, with significant upregulation observed in PANC-1 cells upon epigenetic

silencing contrasted to downregulation when using the same sgRNAs in A549,

suggesting the presence of a context-dependent regulatory element. Color of text of

SNPs above panel A indicates the signal each SNP is part of (in accordance with color

scheme in **Fig. 2**)

Fig. 4

Fig. 4: Tiled CRISPRi proliferation screen nominated VNTRs represent novel
CCVs for PDAC.

a, Functional screening of the 5p15.33 locus nominated three MCFVs intronic to

CLPTM1L. One such MCFV, rs11133729, resides in intron nine of *CLPTM1L*, a region

enriched for enhancer-associated histone marks, DNAse hypersensitivity marks and

transcription factor ChIP-Seq peaks across seven ENCODE cell lines as demonstrated

from the UCSC genome browser's ENCODE regulation track (GRCh37). **b**, Analysis of

tiled CRISPRi screening data of the 5p15.33 locus using CRISPR-SURF nominated a

considerable proportion of *CLPTM1L* as proliferation-essential, including prominent

depletion of sgRNAs within intron 9. Aligned tracks display proliferation-essential

elements from CRISPR-SURF colored by magnitude of CRISPR-SURF depletion FDR.

For each cell line (colored by representative cancer type), screening data was analyzed

across three timepoints (T_0: immediately following puromycin selection and recovery, T_1:

~ 12.5 cumulative screen doublings, T_2: ~ 25 cumulative doublings).**c,** Intron 9 is

notable for the presence of two variable number tandem repeat (VNTR) elements.

CRISPRi screening across tested cell lines nominated both VNTRs as proliferation-

essential elements. As VNTRs are known to be highly polymorphic, we utilized PacBio

long-read amplicon sequencing to genotype both VNTRs in 1kGP and PDAC GWAS

samples. **d,** VNTR genotyping in 1,810 PDAC cases and controls revealed trimodal

(VNTR1) and bimodal (VNTR2) allele length distributions. (VNTR allele count plotted on

y-axis with corresponding VNTR length on x-axis) Bar plots colored as follows;

aggregate VNTR lengths colored by corresponding length, distinct alleles by observed

frequency, clusters by k-means cluster index, long/short by binary indicator). **e,** Read-

based phasing of VNTR and SNP alleles in 484 1kGP EUR samples demonstrated co-

segregation of the rs465498 PDAC risk haplotype (top panel) with longer VNTR1 alleles

(clusters, bottom panel).

Fig. 5

Fig. 5: VNTR1 is a Hippo-dependent regulator of TERT and CLPTM1L.

a, Motifscope plot of VNTR1 as comprised of tandemly repeating 29bp repeats (most

frequently observed repeat units are color-coded and shown at bottom left) which

themselves are enriched for TEF1 binding sites (canonical TEF1 binding motifs shown

in blue and cream, bottom right). **b,** Imputation of VNTR alleles into full GWAS (9,040

PDACs, 12,496 controls) revealed strong association for VNTR alleles (y-axis; -log10 P,

red: $P \leq 5\times10^{-8}$; logistic regression of SNP/VNTR dosage adjusted for age, sex sample

site and principle components) **c,** VNTR1-targeted CRISPRi using three distinct

sgRNAs (x-axis; sgRNAs A, B and C) reduces *TERT* and *CLPTM1L* expression (y-axis;

experimental sgRNA $2^{-\Delta\Delta Ct}$ normalized to $2^{-\Delta\Delta Ct}$ of non-targeting sgRNA, * indicates

significance under two-sided t-test) **d,** Dual luciferase reporter assays incorporating

cluster-representative VNTR1 sequences demonstrated strong VNTR1 enhancer

activity in PANC-1 and A549 cells; cluster 3 alleles show significantly higher luciferase

activity as compared to cluster 1 (Welch's t-test, raw firefly/renilla shown due to intensity

of signal (**Methods**)). **e,** Small molecule inhibition of TEF1/YAP binding negates VNTR1

enhancer activity, indicating Hippo pathway dependence (y-axis; firefly/renilla

luminescence normalized to negative control plasmid, **Methods**). **f,** DNA pulldown/mass

spectrometry comparing peptide intensity following incubation of Mia PaCa-2 nuclear

lysate with biotinylated DNA probes containing consensus VNTR1 repeat unit and

addition of 50-fold excess untagged competitor probe reveals VNTR1-specific TEF1

binding (x-axis; $\log_2$ unlabeled/labelled peptide intensity, y-axis: -$\log_{10} P$, one-sided *t*-

test); **g**, Bar plot of TEF1-specific specific mass-spectrometry results, 10 and 50-fold

excess unlabeled competitor negate TEF1 intensity (independent-samples t-test). **h,**

Western blot following DNA oligo pulldown experiments confirms VNTR1-specific TEF1

binding. Binding is only seen for VNTR1-shifted probe (sequence of RU offset by 15 bp)

to accommodate TF motif in RU junction. Similar binding was observed for Hippo

pathway factors TEF-3 (gene name: *TEAD4*) and WW domain-containing transcription

regulator protein 1 (gene name: *WWTR1*; also named *TAZ*). Nonspecific binding to

beads and probes was seen for YY1 and SP1. "s/n" indicates flowthrough protein levels

after the pulldown

Supplementary Fig. 1

Supplementary Fig. 1: Credible causal variant (CCV) inclusion criteria for MPRA testing.

Two studies have comprehensively identified common risk variants at 5p15.33, which are associated with the risk of several cancer types. Wang et al. in 2014 used the Association analysis based on SubSETs (ASSET) method of conditional and joint meta-analysis of GWAS summary statistics to identify six distinct signals at the locus, which was expanded upon by Chen et al. in 2021, identifying an additional four independent signals. These papers served as a starting point for CCV compilation. These signals were orthogonally supplemented by the application of the Bayesian fine-mapping tool Sum of Single Effects (SuSiE) to individually identify independent signals from GWAS summary statistics for PDAC, lung cancer, bladder cancer and melanoma. GWAS for lung adenocarcinoma were separately analyzed for European and East Asian ancestries. We supplemented ASSET-derived variants and variants assigned to credible sets by identifying signal-tagging ASSET variants along with those variants with the highest PIP for a given credible set and subsequently including correlated proxies with LD $r^2 \geq 0.6$ to derive a final panel of 116 CCVs for inclusion in the MPRA pool.

Supplementary Fig. 2

Supplementary Fig. 2: Fine-mapping of 5p15.33 implicated cancer GWAS.

GWAS signals at the 5p15.33 locus in fine mapped cancers with resultant SuSiE

credible sets are shown with posterior inclusion probabilities (PIPs) for each variant

shown below for; **a,** PDAC, **b,** melanoma, **c,** bladder cancer, **d,** European (EUR) lung

cancer, and **e,** East-Asian (EAS) lung cancer. **f,** Odds ratios and 95% confidence

intervals across all fine-mapped cancers, showing antagonistic pleiotropy between

PDAC, melanoma, bladder cancer and lung cancer EUR/EAS GWAS for one of the

signals at chr5p15.33 (ASSET signal marked by rs465498).

Supplementary Fig. 3

Supplementary Fig. 3: Cross-correlation of ASSET and SuSiE-nominated 5p15.33
signals.

UpSet plot showing intersections between SuSiE fine-mapping credible sets and

ASSET signal-tagging SNPs identified across five cancer GWAS. Variants were

grouped based on linkage disequilibrium (LD $r^2 \geq 0.6$ 1kGP EUR). Horizontal bars (left)

indicate the total number of variants in each credible set or reported signal. Vertical bars

(top) show the size of intersections between sets, with connected dots (matrix below)

indicating which credible sets share variants. The most common intersection (n=44

variants) represents a core set of variants shared across multiple cancer types. Cancer

type abbreviations: PDAC, pancreatic ductal adenocarcinoma; MEL, melanoma; BLAD,

bladder cancer; LUNG_EUR, lung cancer in European ancestry; LUNG_EAS, lung

cancer in East Asian ancestry; Chen (Chen et al. ASSET 2021); Wang (Wang et al.

ASSET 2014). Results demonstrate extensive sharing of credible causal variants across

cancer types, with some cancer-specific signals, consistent with pleiotropic regulatory

mechanisms at this locus.

Supplementary Fig. 4

Supplementary Fig. 4: Inter-replicate correlation of MPRA experiments.

Pairwise correlation matrices showing inter-replicate concordance for MPRA expression

measurements ($\log_{10}$ RNA TPM vs. $\log_{10}$ RNA TPM) in four biological replicates across

eight cancer cell lines. Each panel displays a hexbin scatter plot with Pearson

correlation coefficient (r) for comparisons between replicates (rep1 -rep4). Cell lines

tested: 5637 (bladder cancer), UMUC3 (bladder cancer), A549 (lung adenocarcinoma),

H520 (lung squamous cell carcinoma), UACC903 and UACC1113 (melanoma), MIA

PaCa-2 (PDAC), and PANC-1 (PDAC). Point density is indicated by color intensity (blue

scale). Most pairwise comparisons showed strong correlations (r > 0.96), except for MIA

PaCa-2 replicate 2 (r = 0.017 and r = 0.039, red text) and suggesting technical issues

with MIA PaCa-2 replicate 2 which was removed from downstream analyses. High

correlations across remaining replicates demonstrate robust and reproducible MPRA

measurements for functional variant characterization.

Supplementary Fig. 5

Supplementary Fig. 5: MPRA results for 5p15.33 CCVs revealed variants exhibiting
allele preferential activity across eight cell lines.

Volcano plots showing allelic effects of MPRA-tested variants in; **a,** PANC-1, **b,** MIA

PaCa-2, **c,** UACC903, **d,** UACC1113, **e,** A549, **f,** NCI-H520, **g,** UMUC3, and **h,** 5637

cell lines. X-axis shows $\log_2$ fold-change in expression between alternative (ALT) and

reference (REF) alleles; y-axis shows statistical significance (-$\log_{10}$ adjusted p-value).

Red points indicate variants with significant allelic activity differences (MPRA-functional

variants); gray points and dashed horizontal line indicates non-significant variants (FDR

< 0.01). Selected variants of interest are labeled with their rsID numbers. Multiple

variants demonstrate significant allelic effects across cancer types, with some showing

cell-type-specific regulatory activity. The magnitude and direction of allelic effects vary

across cell lines, reflecting tissue-specific regulatory mechanisms.

Supplementary Fig. 6

643 *Supplementary Fig. 6: Scrambled control constructs assign MPRA-functional*
644 *CCVs to active elements*

645 Ranked distribution of $\log_2$ RNA/DNA TPM values for all tested constructs across eight

646 cancer cell lines: MIA PaCa-2, PANC-1, UACC903, UACC1113, A549, NCI-H520,

647 UMUC3, and 5637/HTB9. Constructs are ranked from highest to lowest expression and

648 colored by type and strand orientation: alternative allele forward strand (alt_fwd, pink),

649 alternative allele reverse strand (alt_rev, orange), reference allele forward strand

650 (ref_fwd, green), reference allele reverse strand (ref_rev, cyan), scrambled control

651 forward strand (scramble_fwd, blue), and scrambled control reverse strand

652 (scramble_rev, purple). Dashed blue lines indicate the first quartile (Q1, lower) and third

653 quartile (Q3, upper) of scrambled control expression within each cell line; dotted red

654 lines indicate Q1-IQR (lower suppressor threshold) and Q3+IQR (upper activator

655 threshold). Alleles with mean expression above Q3 were classified as putative

656 activators; those below Q1 were classified as putative suppressors. CCVs meeting

657 criteria for MPRA-functionality (FDR < 0.01, |fold-change| ≥ 15%, and activator or

658 suppressor classification) demonstrate activity substantially divergent from

659 transcriptionally inert scrambled controls, indicating genuine regulatory function.

Methods

Cell culture

UACC903 and UACC1113 melanoma cell lines were obtained from the University of
Arizona Cancer Center (Phoenix, AZ). All other cells were obtained from American Type
Culture Collection (ATCC; Manassas, VA) and maintained in standard culture conditions
at 37 °C in a humidified atmosphere with 5% CO_2. All cell culture media were obtained
from Thermo Fisher Scientific, Waltham, MA. MIA PaCa-2 (CRM-CRL-1420) cells were
cultured in Dulbecco's Modified Eagle's Medium (DMEM) high glucose supplemented
with 10% fetal bovine serum (FBS) and 2.5% horse serum. PANC-1 (CRL-1469),
UMUC3 and A549 cells were maintained in DMEM high glucose with 10% FBS. NCI-
H520 (HTB-182), UACC903, UACC1113, 5637 (HBT9) and TCCSUP (HTB-5) cells
were cultured in RPMI-1640 with 10% FBS. HEK293T (CRL-3216) cells, used for
lentiviral production, were grown in high-glucose DMEM supplemented with 10% FBS,
1% glutamine, and 1% sodium pyruvate. All cell lines were routinely monitored for
expected morphology and growth rates in addition to regularly being screened for
mycoplasma contamination (Lonza MycoAlert Mycoplasma Detection Kit: LT07-318)
which consistently tested negative. Authentication was performed using a short tandem
repeat (STR) profiling panel (Identifiler kit, Life Technologies, Carlsbad, CA), and
resulting profiles were compared against publicly available reference datasets from
ATCC and DSMZ (https://www.dsmz.de/). All cell lines demonstrated correct STR
identity based on database records.

Fine-mapping and MPRA variant selection

To derive credible sets, we fine mapped previously published GWAS for PDAC[46] (9,040
cases, 12,496 controls), melanoma[47] (36,760 cases, 375,188 controls), lung[49,50] (EUR;
29,266 cases, 56,450 controls, EAS; 21,658 cases, 150,676 controls) and bladder
cancer[48] (13,790 cases, 343,502 controls). We filtered summary statistics[46–50] to include
all available variants within ±500kb of the most significant 5p15.33 SNP for each cancer

type. We then used MungeSummStats[94] (v1.18.0) to lift genomic coordinates over to

GRCh38 and standardized effect alleles to the minor allele (**Supplementary Table 1**).

We then utilized the Sum of Single Effects (SuSiE) method[9] (v0.14.2) susie_rss()

function under default parameters to derive credible sets (CSs) with L = 10 (up to 10

possible causal variants per credible set). LD matrices for SuSiE were generated using

PLINK[95] (v2.0.0-a.7) from the 1000 Genomes Project Phase 3 (1kGP v3) dataset[51],

using genotypes from unrelated East Asian ancestry samples (EAS) for East Asian lung

cancer summary statistics and unrelated European ancestry (EUR) samples for the

remaining four GWAS. For our MPRA, we included CCVs from each of the identified

SuSiE credible sets in addition to the following correlated variants (1kGP v3 EUR for

PDAC, melanoma, bladder and lung EUR, EAS for lung EAS), using an LD threshold r^2

≥ 0.6 as a compromise between inclusivity and specificity:

1. LD r^2 ≥ 0.6 relative to the SNP with the highest posterior inclusion probability

(PIP) for an identified SuSiE credible set.

2. LD r^2 ≥ 0.6 relative to any signal-tagging SNP identified by Chen and

colleagues in 2021.[5]

3. LD r^2 ≥ 0.6 relative to any signal-tagging SNP identified by Wang and

colleagues in 2014.[4]

This approach resulted in 116 5p15.33 CCVs which were included in the MPRA panel

(**Supplementary Figure 1**). In addition to these variants, we also included 44 additional

variants from previous MPRAs and luciferase reporter assays as positive controls

(**Supplementary Table 3**).

Epigenetic annotation of CCVs

CCVs were annotated with functional scoring metrics from RegulomeDB[57] (v2.2) and

FORGEdb[58] using the LDlink online tool[96]. CCVs were further annotated for predicted

disruptions or creations of transcription factor binding motifs using motifbreakR[59].

motifbreakR utilized TF models from HOCOMOCO v11. We used the information

content method and required motif matches to meet a maximum p-value of 10^{-4}. Strong

motifbreakR effects were defined as a difference between reference and alternate

alleles scores greater than 0.7. We further annotated motifbreakR results with

colocalization of matching TF ChIP-Seq data from the ReMap22[93] database.

MPRA library design

For each CCV, we designed oligonucleotides comprised of 145 bases of contextual

sequence (hg38) centered on either the effect or non-effect GWAS alleles, in both

forward and reverse orientations with a unique 12-mer barcode separated by

contiguous KpnI and XbaI restriction sites (structured oligo sequence: 5'-

ACTGGCCGCTTCACTG-145 bases-GGTACCTCTAGA-12 base tag-

AGATCGGAAGAGCGTCG-3') (Agilent Technologies, Santa Clara, CA). Barcode

sequences were preselected from the publicly available FREE barcodes database[12] for

balanced GC content, minimal homopolymer runs, and reduced hairpin propensity.

Additionally, we filtered barcodes to exclude any known mammalian miRNA seed

sequences using miRbase[13]. For each variant, we designed six constructs (reference,

alternate, and core 20 bp randomized in both forward and reverse orientations), with 35

unique barcodes per construct. Seven variants (rs11278847, rs11414507, rs3215401,

rs34218850, rs28379291, rs3030832, rs57516180) failed sequence design due to the

presence of KpnI and XbaI restriction sites within their contextual sequence.

MPRA plasmid pool construction, transfection and sequencing

To clone MPRA oligonucleotide pools into the pMPRA2 delivery vector (Addgene,

Watertown, MA, plasmid #49350), 10 fmol of reconstituted oligonucleotides were first

amplified by emulsion PCR (ePCR) using Herculase II Fusion Polymerase (Agilent) with

primers containing SfiI restriction sites. ePCR was performed with the Micellula DNA

Emulsion & Purification Kit (EURx/CHIMERx, Milwaukee, WI). The amplified

oligonucleotide pools were then ligated to Illumina adapter sequences using the KAPA

HyperPrep Kit (KAPA Biosystems, Wilmington, MA) and quantified with the KAPA qPCR

assay. Tag recovery rates were assessed via Illumina MiSeq v3 (150 bp, paired-end

reads; Illumina, San Diego, CA). ePCR products were digested with SfiI (New England

Biolabs, Ipswich, MA) and ligated into the complementary SfiI site of the pMPRA1

vector (Addgene plasmid #49349) using Electroligase (New England Biolabs, Ipswich,

MA) at a 1:25 vector-to-insert molar ratio. The ligation product was digested with KpnI

and XbaI (New England Biolabs, Ipswich, MA) between the 145 bp element and 12 bp

tag sequence to enable ligation of a *luc2* open reading frame (ORF) cloned from the

corresponding restriction sites of the pMPRA2 vector. The final ligation product was

electrotransformed into ONESHOT electrocompetent cells (Thermo Fisher Scientific),

ensuring that each construct was represented in at least ten independent

transformations. The cloned library intended for transfection was validated as a single

band on an agarose gel following KpnI digestion.

We performed MPRA transient transfections in eight cancer cell lines: PANC-1, MIA

PaCa-2, UACC903, UACC1113, A549, H520, UMUC3, and 5637 (We used 5637

bladder cancer cells for the MPRA (rather than TCCSUP for CRISPRi) because 5637

cells are amenable to transient transfection). Each cell line was transfected in biological

quadruplicate, with replicate cultures plated on separate days. For each transfection,

the number of transfected cells exceeded the library complexity by more than 100-fold

to ensure comprehensive representation of all sequences. 14 µg of MPRA plasmid was

transfected to cell cultures using Lipofectamine 3000 (Thermo Fisher Scientific)

according to the manufacturer's protocol. Cell densities were optimized for each line to

maximize transfection efficiency, which was verified by a parallel GFP transfection

assay and fluorescence microscopy (no less than ~ 50% transfection efficiency). Cells

were harvested 48 hours post-transfection for RNA isolation.

Total RNA was extracted with the RNeasy Kit (QIAGEN, Hilden, Germany) and

enriched for mRNA using the Poly(A)Purist MAG Kit (Thermo Fisher Scientific).

Complementary DNA (cDNA) was synthesized using SuperScript III Reverse

Transcriptase (Thermo Fisher Scientific). Unique 12-mer barcode sequences were

amplified from cDNA with Q5 High-Fidelity DNA Polymerase (New England Biolabs)

using primers containing Illumina TruSeq adapter sequences. Tag sequence libraries

were prepared from both input DNA and cDNA using identical workflows. Sequencing

was performed on a NovaSeq 6000 SP flow cell (Illumina, San Diego, CA) with 100 bp,

dual-indexed, single-end reads at NCI's Cancer Genomics Research Laboratory. For

each cell line, sequencing depth ranged from 150 to 300 million reads per sample,

providing sufficient coverage for downstream analyses (**Supplementary Table 17**).

MPRA data analyses

Using FASTQ files generated from transfected plasmid DNA and RNA transcript

sequencing, we quantified reads that perfectly matched each 12 bp tag sequence, along

with downstream regions encompassing the XbaI recognition site and 20 bp of the 3′

luc2 sequence. For each transfection, tag counts per million (TPM) were calculated by

dividing the raw tag count by the total number of sequence-matching tag counts per

million reads. To infer the transcriptional activity of each element, a single pseudocount

was added to every TPM value and the RNA/DNA ratio for each tag was then computed

by dividing its RNA TPM by the corresponding DNA TPM. For each transfection, the

median ratio across all tags was calculated, and adjusted ratios were obtained by

dividing each tag's ratio by this median to ensure standardized scaling across

replicates. Adjusted ratios were subsequently $\log_2$-transformed to derive activity levels

associated with each tested tag sequence.

Across all eight cell lines, at least 97.05% of barcode sequences were detected in both

input DNA libraries and RNA samples. Reproducibility between transfections was

assessed using Pearson correlation. After removal of a single outlier replicate per cell

line, transcriptional activity levels showed high correlation across transfection replicates

(median Pearson's r; PANC-1 = 0.455, MIA PaCa-2 = 0.236, UACC903 = 0.852,

UACC1113 = 0.903, A549 = 0.999, NCI-H520 = 0.956, UMUC3 = 0.981, 5637 = 0.964;

n = 4 transfections per cell line;). We restricted analysis to barcodes with ≥100 reads in

either RNA or DNA libraries, retaining 89.65% of designed tags. To evaluate the

transcriptional impact of alternate (ALT) versus reference (REF) alleles, we applied a

standard linear regression model adjusting for strand orientation (forward or reverse) as

a binary covariate and biological replicate as a categorical covariate. Model significance

was assessed using the robust sandwich-type variance estimate (Wald test) to account

for potential heteroskedasticity in measurement error, as implemented in the R package

sandwich. P-values for each linear model were corrected for multiple testing using the

Benjamini–Hochberg procedure, applying a false discovery rate (FDR) threshold ≤ 0.01

to identify variants exhibiting significant transcriptional differences between alleles and

scrambled controls.

To further assess whether tested elements displayed divergent activity relative to

transcriptionally inert controls, the mean $\log_2$(RNA/DNA TPM) ratio for each allele was

compared to the distribution of $\log_2$(RNA/DNA TPM) ratios from all scrambled control

constructs within the same cell line. For each cell line, the interquartile range (IQR) of

scrambled control activity was determined: alleles with mean $\log_2$(RNA/DNA TPM)

ratios above the third quartile (Q3) were classified as putative activators, whereas those

below the first quartile (Q1) were classified as putative suppressors (**Supplementary**

Fig. 6). CCVs exhibiting an FDR <0.01 allelic effect, an absolute allelic fold change

≥15% and classification as either activator or suppressor were defined as MPRA-

functional for a given cell line.

Tiled CRISPR interference (CRISPRi) proliferation screen design

sgRNAs targeting chr5:1,233,287–1,380,000 bp (hg19) were downloaded from the

UCSC Genome Browser "CRISPR Targets" track. We also incorporated 992 inert and

validated negative control sgRNAs in addition to sgRNAs targeting a panel of gene

promoters selected based on susceptibility gene status for a given cancer type or noted

as proliferation-essential across DepMap CRISPRi screening datasets[60,97]. CRISPRi

oligonucleotides were synthesized using the SurePrint Oligonucleotide Synthesis

Platform (Agilent) in the following configuration: 5′-AGGCACTTGCTCGTACGACG-

CGTCTCACACCG-[sgRNA20nt]-GTTTCGAGACG-ATGTGGGCCCGGCACCTTAA-3′.

A total of 27,340 sgRNAs were included in the final library, of which 25,778 tiled across

the 5p15.33 region. For the 116 CCVs identified in fine-mapping, there was a median of

14 sgRNAs mapping to within ±50bp of each CCV, with one CCV (rs62651531) failing

to have any within this window. rs2853672 had the most sgRNAs mapping within this

window, with 39 (**Supplementary Table 5**).

CRISPRi sgRNA pool generation

sgRNA oligonucleotides were PCR-amplified using flanking primer sites: Forward – 5′ AGGCACTTGCTCGTACGACG 3′; Reverse – 5′ ATGTGGGCCCGGCACCTTAA 3′. Amplified products were cloned into the pXPR_050 vector (Addgene #96925) using the same emulsion PCR strategy described above for MPRA. ePCR products were purified with a QIAquick PCR Purification Kit (QIAGEN). The pXPR_050 backbone was digested with Esp3I (NEB), gel-extracted to isolate the linearized vector, and ligated to the sgRNA ePCR amplicons using Electroligase (NEB). The ligation product was electroporated into ONESHOT electrocompetent cells in six independent reactions. After a one-hour recovery at 37 °C in S.O.C. medium (Invitrogen; 300 rpm), serial dilutions were plated to yield > 100× colony-forming units (CFUs) per sgRNA in the pool. Bulk cultures were incubated overnight (37 °C, agitation at 300 rpm, 16 h) to minimize selective expansion of individual clones. Plasmids were isolated via six parallel endo-free maxi preps (QIAGEN), pooled, quantified by Qubit High Sensitivity DNA Assay and NanoDrop 8000 (Invitrogen, Waltham, MA), and PCR-amplified. Resultant amplicons were then sequenced for QC (Illumina MiSeq v3, 100 cycles, single-end). FASTQ files were analyzed for sgRNA recovery and quality metrics. >99 % of input sgRNAs were recovered, confirming library complexity sufficient for lentiviral packaging.

Lentiviral Packaging

HEK293T cells were plated and grown to ~70 % confluency in six T175 flasks in virus production medium (high-glucose DMEM media supplemented with 10% FBS, 1% glutamine, and 1% sodium pyruvate). Transfection mixtures were prepared using lipofectamine 3000 (Invitrogen) to the manufacturer's protocol, incubated 15 min at room temperature, and added dropwise to cells. After 6–8 h at 37°C, media was replaced with nutrient-rich DMEM. At 48 hours post-transfection (Day 3), viral supernatants were collected, centrifuged to remove debris, and filtered (0.45 μm). To concentrate virus, 2× PEG precipitation buffer (Millipore Sigma, Burlington, MA) was added and incubated overnight at 4 °C. On Day 4, viral particles were pelleted by

centrifugation, resuspended in 1 mL aliquots of 1× PBS (Quality Biological,

Gaithersburg, MD), and stored at −80 °C pending titration and transduction. Viral titer

was quantified using Lenti-X GoStix Plus (Takara Bio, Kusatsu, Japan). 20 μL of viral

sample were mixed with 80 μL of chase buffer, incubated 10 min at room temperature,

and assessed using the GoStix Plus iOS app. The control (C) and test (T) band intensity

ratio was mapped to batch-specific calibration curves via QR code, yielding an estimate

of p24 concentration and infectious titer.

CRISPRi screen transduction, cell culture and sequencing

CRISPRi screening experiments were performed across eight cell lines (MIA PaCa-2,

PANC-1, UACC903, UACC1113, A549, H520, UMUC3, and TCCSUP) which had been

transduced with lentivirus for dCas9-KRAB-ZIM3 and cultured from clonal subdivisions.

CRISPRi experiments were transduced at a multiplicity of infection (MOI) of 0.3 to

ensure single sgRNA integration per cell. Transductions were conducted at a scale

sufficient to maintain ~500× representation of each sgRNA throughout the screen.

Three biological replicates were performed with transductions performed over

consecutive days. For each replicate, counted cells were resuspended with lentiviral

supernatant and 8 μg/mL polybrene and plated at 1×10^7 cells per 15-cm dish. After 24

h, cultures were washed and selected with 10 μg/mL puromycin until all cells in no-virus

controls were eliminated (approximately 24–72 h). Post-selection, CRISPRi populations

were expanded to generate sufficient material for both the initial time point (T_0) and

continued passaging until subsequent harvests at T_1 (~12.5 doublings) and T_2 (25

doublings).

Genomic DNA (gDNA) was isolated using the Quick-DNA Genomic DNA Kit (Zymo,

Irvine, CA). For each biological replicate, gDNA was harvested at the initial post-

selection time point and again at approximately 12.5 and 25 population doublings. The

full quantity of gDNA from each sample was used as input for the first round of sgRNA

amplification (PCR1), distributing reactions such that each contained 1 μg gDNA to

preserve library complexity. PCR1 products were purified using a 1:1 Ampure XP bead

cleanup (Beckman Coulter, Brea, CA). A second amplification (PCR2) was performed

using the same number of reactions as PCR1 to incorporate sample-specific barcodes

and full Illumina adapter sequences. Each 20 µL PCR2 reaction contained 15 ng of

PCR1 product and PCR2-specific primers and was run for five cycles under the same

cycling parameters used for PCR1. Reactions were pooled and purified using the

QIAquick PCR Purification Kit (QIAGEN). Library size (324 bp) was verified using a

BioAnalyzer High Sensitivity DNA assay (Agilent). A 30 µL aliquot of each prepared

library was submitted to the NCI Cancer Genomics Research Laboratory (Rockville,

MD, USA) for sequencing on an Illumina NovaSeq SP flow cell.

CRISPRi screen quality control and data analyses

Sequencing data from the tiled CRISPRi proliferation screen spanning 5p15.33 (eight

cell lines; biological triplicates; T0 immediately after puromycin recovery, T1 at ~12.5

cumulative doublings, T2 at ~25 doublings) were processed from raw FASTQ files using

the standard MAGeCK-VISPR workflow (v0.5.6) to quantify sgRNA counts for the input

plasmid, baseline, and post-proliferation timepoints and to perform library and assay

QC. Across plasmid controls, we observed high read mapping efficiency (72.28–

73.25%), near-uniform sgRNA representation (median Gini index 0.052; range 0.0488–

0.055), and minimal dropout (mean reads/sgRNA 859.48; mean missing sgRNAs

155.69; **Supplementary Table 6**), whereas cell-line libraries showed progressive

selection-associated skew (Gini increasing from T0 to T2; **Supplementary Table 6**).

For each sample, MAGeCK was used to compute sgRNA log2 fold changes (LFCs) and

gene-level statistics with multiple-testing correction (FDR), including positive-control

evaluation using 95 proliferation-essential genes (**Supplementary Tables 7 and 9**) and

targeted aggregation of sgRNAs assigned to the *TERT* and *CLPTM1L* promoter regions

(*TERT;* chr5:1,294,628–1,298,988: GH05J001293 and *CLPTM1L*: chr5:1,343,491–

1,346,621: GH05J001343; **Supplementary Table 5**). Downstream analysis was

restricted to comparisons between individual timepoints and plasmid counts. Fine-

mapped CCVs were linked to perturbations by intersecting sgRNAs within ±50 bp of

each CCV (median 14 sgRNAs/CCV; **Supplementary Table 5**) and testing CCV-

assigned sgRNA sets for depletion or enrichment (**Supplementary Tables 18&19**).

Finally, proliferation-essential regulatory elements were inferred from sgRNA LFCs

using CRISPR-SURF[62](v1.0) to deconvolve localized peaks of depletion at FDR ≤ 0.05;

datasets in which deconvolution failed to converge (specified in **Supplementary**

Results) were excluded, and locus-level patterns were visually confirmed in IGV by

concordant read coverage and dropout profiles across replicates and cell lines.

Epigenetic silencing of SNP and VNTR loci by targeted CRISPRi

Individual sgRNAs corresponding targeting selected CCVs and both VNTRs were

cloned into the pXPR_050 CRISPRi expession vector. For each sgRNA,

complementary oligonucleotides containing the 20-bp spacer sequence and BsmBI-

compatible overhangs (Forward: 5′-CACCG[$N_{(20)}$]-3′; Reverse: 5′-AAAC[$N_{(20)}$]C-3′)

were synthesized (IDT) and annealed using a standard thermocycler-based annealing

protocol. pXPR_050 was linearized by BsmBI (NEB) digestion and gel-purified prior to

ligation. Annealed oligonucleotides were ligated into the digested backbone using T4

DNA ligase (NEB). Ligations were transformed into *E. coli* TOP10 chemically competent

cells (Thermo Fisher Scientific) and plated on LB-ampicillin agar plates (Quality

Biological). Single colonies were expanded overnight in selective LB medium, and

plasmid DNA was isolated using the ZymoPURE Plasmid Miniprep kit (Zymo). To

confirm correct sgRNA insertion, plasmid DNA from each clone was barcoded and

sequenced using the Oxford Nanopore Rapid Sequencing and Barcoding Kit on a single

Flongle flow cell (Oxford Nanopore, Oxford UK). Sequence-verified clones were

expanded in 50mL cultures and purified using the EndoFree Maxiprep kit (QIAGEN) for

downstream experiments.

Lentiviral particles encoding individual sgRNAs were generated by transient transfection

of HEK293T cells. Briefly, 4×10^6 HEK293T cells were plated in 10cm^2 dishes in high-

glucose DMEM supplemented with 10% FBS, pyruvate, and glutamine (Quality

Biological). Cells were transfected the following day with the sgRNA expression

construct, psPAX2 packaging plasmid (Addgene #12260), and MD2.G envelope

plasmid (Addgene #12259) using Lipofectamine 3000 (Thermo Fisher Scientific) in

accordance with the manufacturer's recommendations. Media was replaced 24 h post-

transfection. Viral supernatant was collected at 48 h, clarified by low-speed

centrifugation, filtered (0.45 μm), and concentrated by overnight PEG precipitation at 4

°C. Viral pellets were resuspended in sterile PBS and stored at −80°C. For sgRNA

delivery, 1×10^5 PANC-1 or A549 cells stably expressing dCas9-KRAB-ZIM3 were

seeded in 6-well plates in selective media containing blasticidin and polybrene. 25μl of

concentrated virus was added to each well, and plates were centrifuged at 300 × g for 1

h to enhance transduction efficiency, followed by overnight incubation at 37 °C. Media

was replaced the next day with blasticidin- and puromycin-containing growth medium,

and cultured for 96 hours before harvesting total RNA.

Total RNA was extracted using a QIAcube connect instrument with the RNeasy mini kit

(QIAGEN). 1 μg of total RNA was converted to cDNA with SuperScript III reverse

transcriptase (Thermo Fisher Scientific). RT–qPCR was performed using a QuantStudio

7 Flex system. *TERT* and *CLPTM1L* expression relative to negative control sgRNAs

was assessed using TaqMan assays (Thermo Fisher Scientific) (*TERT:*

Hs00972656_m1 *CLPTM1L:* Hs00363947_m1) with Ct values normalized to *HPRT1*

values from the same well (Hs99999909_m1): An unpaired t-test with Welch's

correction was used to test for statistical significance.

Targeted PacBio Amplicon Sequencing of *CLPTM1L* Intron 9

Genomic DNA samples for HapMap trios (CEU panel for CEPH Utah residents with

ancestry from Northern and Western Europe, n = 72) and 1000 Genomes Project

Europeans (Toscani in Italia (TSI); n = 114, Finnish in Finland (FIN); n = 100, British

from England and Scotland (GBR); n = 100; Iberian Populations in Spain (IBS); n =100)

were purchased from the Coriell Institute of Medical Research Institute (Camden, NJ).

Genomic DNA samples for pancreatic cancer cases and controls were sourced from the

first two phases of the PanScan GWAS [20,21].

Amplicons spanning chr5:1,325,380–1,330,639 (GRCh38) were generated using

barcoded M13 primers and the PacBio SMRTbell prep kit v3.0 with minor protocol

modifications. PCR cycling consisted of 30 cycles of 98 °C denaturation (10 s) and

combined annealing/extension (6 min), followed by a 10 min final extension at 72 °C.

Amplicons were purified with AMPure beads, quantified by PicoGreen (Invitrogen),

barcoded with an additional 10 PCR cycles, pooled, and sequenced on a Sequel II

SMRT Cell. HiFi reads were processed using SMRT Analysis for circular consensus

sequence (CCS) generation and demultiplexed using Lima.

979 Genotyping of VNTRs and SNPs from HiFi reads

HiFi amplicon reads were aligned to GRCh38 using pbmm2 with a minimum read-length

threshold of 4,700 bp. Aligned HiFI reads were then processed with PacBio Amplicon

Analysis[98] (pbAA). Reads were first clustered to the target region (amplicon ±100 bp;

chr5:1325280–1330739) using pbAA cluster, and haplotype tags (HP) were assigned to

each read using pbAA bampaint. For each sample, the two modal clusters

(corresponding to diploid haplotype sequences) were retained, and reads from minor

artefactual clusters were excluded. Consensus sequences for each haplotype were

reconstructed, realigned to GRCh38, and used as input for tandem repeat calling with

TRGT[99] and SNP genotyping with DeepVariant/GLnexus[100] followed by read-based

phasing of VNTRs and SNPs using HiPhase[101].

We assessed genotyping concordance across 484 1kGP EUR samples between

PacBio-derived SNP calls and orthogonal short read sequence calls[51]. For 17 SNPs

passing quality thresholds (MAF$_{1kGP}$ ≥ 0.01; HWE P$_{1kGP}$ ≥ 0.05), minor allele frequencies

(MAF) derived from PacBio SNP calls were highly concordant with those from 1kGP

short-read WGS calls (Pearson's r = 0.99; **Supplementary Table 10**), with VNTR

genotypes also conforming to Mendelian segregation in 24 HapMap CEU trios

(**Supplementary Table 11**).

For TRGT VNTR genotypes, we derived four complementary allele-encoding

approaches:

1. Discrete VNTR alleles were maintained as distinct sequences denoted by

1000   their length and assigned a unique index number (<LEN_n >).

1001   2. Distinct VNTR alleles described above were grouped together based on

1002   length irrespective of their sequence composition (<LEN>)

3. VNTR lengths and their associated allele counts were used to assign lengths based on k-means clustering (4 clusters for VNTR1, 3 for VNTR2), (<Cluster_n>).
4. Binary long/short VNTR genotypes were derived by combining (<L/S>).

CEU trios and EUR samples were analyzed jointly. Of 503 sequenced samples, 481 were matched to individuals in the 1000 Genomes Phase 3 high-coverage WGS dataset. Mendelian transmission was evaluated across 24 CEU trios using both VNTR1 and VNTR2 consensus alleles with all trios demonstrating complete Mendelian concordance. To further validate genotyping accuracy, variant calls for 31 dbSNP 155-annotated variants were compared with orthogonal 1000 Genomes short-read genotypes[51]. PacBio SNP genotypes demonstrated negligible differences in MAF (Median ΔMAF= 0.015, range = 0.003-0.228) between PacBio and 1000G WGS genotypes **(Supplementary Table 11),** indicating that accompanying VNTR calls were reflective of true genotypes**.**

VNTR imputation and association analysis

To impute VNTR alleles into the broader PanScan GWAS[46], we generated a reference panel designed to only impute VNTR alleles whilst leveraging phase information from both PacBio-genotyped SNPs and accurately-imputed variants from the same samples. Using data from the PanScan 940 PDAC cases and 870 controls subjected to amplicon sequencing, we compiled a multisample VCF file comprised of all VNTR genotypes (each genotyping method represented as a distinct multiallelic row to be imputed as a separate variant) along with corresponding SNP calls that met MAF > 1%, call rate = 1, and Hardy–Weinberg equilibrium $P > 10^{-5}$ (11 SNPs total). Next, we took imputed dosages from the same sample set which had been imputed using the TOPMed imputation server, excluding any variants outside of chr5:0-6Mb, MAF < 1%, imputation R2 < 0.9 and any overlap with SNPs already genotyped by PacBio. VCFs were concatenated into a single file, sorted and then phased using SHAPEIT4.2[102], leveraging HiPhase-derived phase set tags (PS) as a haplotype scaffold to preserve physical phasing information. Recipient PanScan genotypes were prepared similarly to

the 1,810-reference panel cohort, filtering for position, MAF and imputation R2. To

accommodate downstream imputation and association frameworks, multiallelic VNTR

sites were decomposed to biallelic variants using bcftools[103]. VNTR alleles were then

imputed into three cohorts of recipient samples (PanScan I&II (additional 2,585 cases,

2,823 controls)[66,67], PanScan III (1,597 cases, 4,799 controls)[52], and PanC4 (from the

Pancreatic Cancer Case-Control Consortium[46], dbGaP Study Accession:

phs000648.v2.p1; 3,884 cases, 3586 controls; approval to Laufey T. Amundadottir))

using Beagle[104] (v5.4) whilst retaining hard called VNTR alleles in the 1,810 reference

samples for downstream regression. Downstream analysis of linkage disequilibrium and

logistic regression and conditional analyses were performed using PLINK[95] (v2.0.0-a.7).

For logistic regression, we analyzed each cohort independently, including age, sex,

sample collection site and eigenvectors from principal components analysis as fixed

covariates as performed in the original meta-analysis[46]. We then meta-analyzed each

cohort using PLINK (v1.9).

Analysis of VNTR- transcription factor binding

1046

To assess VNTR-enriched transcription factor binding sites, we obtained VNTR1

sequence in FASTA format from the UCSC genome browser simple repeats track

(GRCh38). VNTR1 was then analyzed with the TRAP[69] (Transcription Factor Affinity

Prediction) online tool. TRAP provides an enrichment measure indicative of whether a

transcription factor binds the VNTR sequence more strongly than expected relative to

human promoters curated by TRANSFAC (**Supplementary Table 15**). We

subsequently visualized VNTR1 RU composition and TEAD motif enrichment using

motifscope[105].

To validate *in-silico* TF binding for VNTR1, we designed biotinylated double-stranded

oligonucleotides corresponding to the VNTR1 reference repeat unit (RU) and a +15-bp

shifted RU were synthesized (IDT; **Supplementary Table 20**). Protein–DNA pulldown

and mass spectrometry analyses were performed at the National Cancer Institute's

Protein Characterization Laboratory (Frederick, MD). Oligonucleotides were annealed in

2× binding and wash buffer (10 mM Tris pH 7.5, 1 mM EDTA, 2 M NaCl) and incubated

with Neutravidin beads for immobilization. For each pulldown, 200 pmol of duplex oligo

was incubated with 400 µg of nuclear extract prepared from MIA PaCa-2 cells in IP

buffer (50 mM Tris pH 8.0, 150 mM NaCl, 0.1% NP-40, 1 mM $MgCl_2$, 100 µM $ZnSO_4$, 1

mM DTT, 5% glycerol, protease inhibitors). Competition controls included 50× and 100×

excess of unlabeled duplex DNA to assess binding specificity. Following a 1hour

incubation with oligo-conjugated beads, samples were washed in IP buffer,

resuspended in SDS loading buffer, and submitted for mass spectrometry. Peptide

identification and quantification followed established core facility pipelines. Relative

peptide abundance across competitor and non-competitor conditions was compared

using a one-sided *t*-test to determine significant displacement of DNA-associated

proteins.

VNTR repeat unit (RU) protein binding analysis by DNA pulldown

1072

Biotinylated double-stranded DNA probes were generated by annealing complementary

single-stranded oligonucleotides (100 µM each strand, 3 µl per strand, 300 pmol total,

IDT, **Supplementary Table 20**) comprising sequences representative of the 1 reference

RU and 15bp shifted sequence as used for global proteomics. Annealed probes were

immobilized on NeutrAvidin agarose beads (30 µl) and incubated with nuclear protein

extracts (100 µg) prepared using the NE-PER Nuclear and Cytoplasmic Extraction Kit

(Thermo Fisher Scientific, Waltham, MA). Following incubation, protein-DNA-bead

complexes were pelleted by centrifugation, and both the supernatant and bead-bound

fractions were collected for analysis.

Proteins were eluted from the beads by heating in sample buffer. Input nuclear extract

(20 µg), supernatant fractions (20 µg), and bead-bound proteins were resolved on

Criterion XT 3-8% Tris-Acetate precast gels (Bio-Rad #3450129, Hercules, CA) using

XT Tricine SDS running buffer (Bio-Rad #1610790) optimized for separation of 40-400

kDa proteins. Proteins were transferred to 0.45 µm pore PVDF membranes (Thermo

Fisher Scientific #88518) using standard wet transfer methods. Membranes were

stained with 0.1% Ponceau S (G Biosciences #786-575, VWR #89167-800) to verify

transfer efficiency.

Membranes were blocked in 2% bovine serum albumin (BSA) in TBST (10 mM Tris-HCl
pH 7.5, 150 mM NaCl, 0.05% Tween-20) and incubated overnight at 4°C with primary
antibodies diluted 1:1,000 in 2% BSA/TBST: rabbit polyclonal anti-SP1 (Abcam
ab13370), rabbit polyclonal anti-TAZ (Abcam ab110239), rabbit monoclonal anti-TEAD1
(Abcam ab133533), mouse monoclonal anti-TEAD1 (BD Transduction 610923), rabbit
polyclonal anti-TEAD4 (Thermo Fisher Scientific 12418-1-AP), mouse monoclonal anti-
TEAD4 (Santa Cruz sc-101184), rabbit monoclonal anti-YAP1 (Abcam ab52771), and
rabbit monoclonal anti-YY1 (Abcam ab109228). Following TBST washes, membranes
were incubated for 1 h at room temperature with HRP-conjugated secondary antibodies
diluted 1:2,000 in 2% BSA/TBST: donkey anti-rabbit HRP (Abcam ab205722) or donkey
anti-mouse HRP (Abcam ab7061). Chemiluminescence was detected using
SuperSignal West Femto Maximum Sensitivity Substrate (Thermo Fisher Scientific) and
imaged on a ChemiDoc imaging system (Bio-Rad).

VNTR luciferase assays

To assess VNTR1 transcriptional activity, VNTR1 alleles from samples from the 1000
Genomes European Ancestry Set were amplified using custom PCR primers
(**Supplementary Table 20)**, allowing for Gateway cloning. HapMap samples HG00171
(2,586 bp), GM20759 (2,789 bp), GM20818 (2,933 bp) and GM20787 (3,566 bp) were
selected to represent both the extremes of allelic length distribution and the two modal
lengths. A *negative* control construct was generated by amplifying a 2.7 kb region in
intron 9 of *TERT* which was not annotated by any epigenetic marks in ENCODE or any
CRISPR-SURF peaks from the screen. PCR, Gateway cloning and sequence
verification were performed at the National Cancer Institute's Protein Expression
Laboratory (Frederick, MD, USA) using a protocol described by Wall et al.[26]

PANC-1 and A549 cell lines were seeded in 48-well plates at densities of 2×10^4 and 4
$\times 10^4$ cells per well, respectively. Cells were transfected the following day with firefly
luciferase reporter constructs using Lipofectamine 3000 (Invitrogen), together with 250
ng of the corresponding control plasmid and 2.5 ng of a Renilla luciferase plasmid
(pGL4.74). Each construct was assayed in biological triplicate with six technical

replicates per experiment. Transfection media was replaced after 6 h with complete

growth medium, and cells were lysed 24 h later in passive lysis buffer (Promega

Madison, WI). Firefly and Renilla luciferase activities were quantified using the Dual-

Luciferase Reporter Assay System on a GloMax Explorer plate reader (Promega)

following the manufacturer's protocol. For Hippo pathway inhibition studies, reporter

assays were performed in the presence of 0.5, or 1.0 μM verteporfin (Millipore-Sigma),

alongside a DMSO vehicle control. Firefly luciferase values were normalized to Renilla

luciferase for each well, averaged across technical replicates, and expressed relative to

the activity of the negative control plasmid. Data were pooled across experiments, and

mean values with standard deviations were plotted using ggplot2. Statistical

comparisons between constructs comprised a pairwise two-sided unpaired Welch's t-

test.

1131   Reagents and oligonucleotides

1132   All reagents, oligonucleotides, plasmids, antibodies, commercial kits, and DNA samples

1133   used in this study, including supplier information and catalog numbers, are listed in

1134   **Supplementary Table 20**.

Supplementary Results

MPRA functional score enrichment

Several databases based on publicly available epigenomics data exist for the functional annotation of GWAS SNPs purporting to aide in triaging of functional variants from correlated proxies. To investigate the predictive power of these tools, we assessed whether annotation scores were enriched for MPRA-validated functional variants. For each set of MPRA variants (CCVs and positive controls), we assayed SNPs stratified by cell line and in a combined model and performed Fisher's exact tests to determine whether specific RegulomeDB and FORGEdb scores were associated with MPRA hit status. We evaluated multiple definitions of functional significance: (i) statistical significance alone (FDR ≤ 0.01), (ii) statistical significance with meaningful effect size (FDR ≤ 0.01 and absolute fold change ≥ 15%), (iii) active regulatory element status (activator or suppressor designation in either allele), and (iv) the intersection of all criteria. Annotative scores were found to have limited predictive capacity in for MPRA status, with the only significant associations observed between active element status and FORGEdb scores of 8 in A549 (P_{adj} = 0.0257) regulomeDB scores of 1b in PANC-1 cells (P_{adj} = 0.0469). These findings suggest that annotative scores should be used on a cautiously across individual loci and be used as a complement to comprehensive functional screening.

CRISPRi Screening Validation

Sequencing data from the tiled CRISPRi proliferation screen of 5p15.33 across eight cell lines were processed using MAGeCK-VISPR[106] (v0.5.6) for sgRNA-level and positive control QC. FASTQ files corresponding to the plasmid library, baseline (T_0), and post-proliferation (T_1 and T_2) timepoints were analyzed using the standard MAGeCK-VISPR workflow to assess library quality, read mapping, and assay efficacy. Across all plasmid control libraries, high mapping efficiency (72.28%–73.25%) and uniform sgRNA representation were observed, with Gini indices ranging from 0.05 to 0.06, consistent

with even distribution of sgRNA counts. In contrast, cell-line derived libraries exhibited

selection-mediated variability through timepoints, reflected by progressive elevation of

Gini indices—from 0.05–0.10 at T_0 to 0.13 at T_1 and up to 0.17 at T_2.

To evaluate assay sensitivity and reproducibility, depletion and enrichment statistics

were derived from the normalized sgRNA counts. Enrichment of small P-values was

noted across positive-control proliferation genes (**Supplementary Table 7**), including

KRAS, MYC, and *CCND1,* indicative of proliferation-mediated of associated sgRNAs.

Consistent and significant depletion was observed for sgRNAs targeting the *TERT* and

CLPTM1L promoter regions, defined as chr5:1,294,628–1,298,988 (GeneHancer ID:

GH05J001293) and chr5:1,343,491–1,346,621 (GeneHancer ID: GH05J001343),

respectively, encompassing 915 and 632 intersecting sgRNAs. These loci exhibited

strong depletion signals across multiple cell lines, consistent with their established roles

in supporting proliferative capacity (**Supplementary Table 5**).

Inference of proliferation-essential regulatory elements from the tiling data was

performed using CRISPR-SURF[62] (v1.0) which applies a deconvolution-based approach

to aggregate sgRNA-level effects into distinct regions of proliferation-essentiality.

CRISPR SURF deconvolution successfully resolved high-confidence peaks of depletion

overlapping the *TERT* and *CLPTM1L* promoters in six of eight cell lines. Convolution

failed to converge for MP2 (T_1), UACC903 (all timepoints), and H520 (T_1–T_2) datasets,

which were excluded from downstream integration analyses. Visual inspection of

aligned read coverage and sgRNA depletion profiles in Integrative Genomics Viewer

(IGV) confirmed reproducible patterns of promoter-specific dropout across cell lines.

Together, these analyses demonstrated that the CRISPRi proliferation screens

achieved robust coverage, reproducible sgRNA representation, and consistent detection

of proliferation-essential loci, including canonical oncogenes and core promoter

elements within the 5p15.33 region.

For the 116 CCVs identified in fine mapping, there was a median of 14 sgRNAs

intersecting within ± 50bp of each CCV, with one CCV (rs62651531) failing to have any

within this window. rs2853672 had the most intersecting sgRNAs, with 39

(**Supplementary Table 5**). We annotated 915 and 632 of the tiling sgRNAs as

intersecting the *TERT* and *CLPMT1L* promoters, respectively (Methods). We

transduced lentiviral packaged sgRNAs into the same cell lines used for MPRA which

were clonally engineered to stably express dCas9-KRAB-ZIM3 (**Supplementary Table**

4). Screens were performed in biological triplicate for 25 cumulative cell doublings and

harvested at an initial time point (T_0, immediately following puromycin selection and

initial recovery), T_1 (~12.5 cumulative screen doublings), and T_2 (25 screen doublings).

Following sequencing of sgRNAs from harvested time points and the primary plasmid,

we assessed library quality, read mapping, and assay efficacy using the output of

MAGeCK. Sequencing libraries from input plasmid and cell line demonstrated high

mapping efficiency, even sgRNA read distribution and minimal sgRNA dropout (Average

reads/sgRNA = 859.48, average missing sgRNAs = 155.69, **Supplementary Table 6**).

Gini indices, a measure of sgRNA read distribution across a sample, increased

progressively through plasmid (median Gini = 0.052, range = 0.0488-0.055) and cell line

timepoints from (range T_0 = 0.065, 0.0524-0.175, T_1 = 0.079, 0.055-0.1548, and T_2 =

0.085, 0.052-.1743), indicative of CRISPRi-mediated negative selection of sgRNAs over

time.

We used MAGeCK to calculate $\log_2$ fold changes for each timepoint relative to the

plasmid in addition to its preceding timepoint and measured inter-replicate Pearson

correlation to assess reproducibility (**Supplementary Fig. 6**). Across all comparisons

(three replicate pairs per experiment), the inter-replicate Pearson correlation coefficients

ranged from –0.42 to 1.00, with a median of 0.46 (IQR 0.20–0.81), indicating that while

some comparisons show very strong agreement between replicates, others are

considerably noisier. Reproducibility was highest for the plasmid control samples. For

plasmid experiments the mean inter-replicate correlation was 0.78 (median 0.86), and

~76% of replicate pairs had $r > 0.7$. In contrast, cell line derived time-point contrasts

were less consistent: $T_1\_vs\_T_0$ / $T_2\_vs\_T_0$ comparisons across all cell lines had a

median r of 0.20. $T_2\_vs\_T_1$ comparisons were lower still (median 0.05), with several

replicate pairs near zero or weakly negative. There was also a clear trend with time: at

T0, replicate correlations were generally high (mean 0.68, median 0.82), decreasing at

T_1, (median $r = 0.54$) and further at T_2, with a median $r = 0.27$. By cell line, PANC1, TCC,

UMUC3, A549 and MP2 showed the best overall replicate concordance (median r range

across timepoints = 0.54–0.80), whereas UACC903, H520 and UACC1113 were more

variable (median r = 0.15–0.29). With this in mind, we restricted downstream analysis

only to comparisons between sgRNA counts derived from cell line time points and their

corresponding count in the input plasmid.

To assess the efficacy of CRISPRi systems, we assessed knockdown of 95 genes

selected for proliferation-essentiality and cancer-type susceptibility status using

MAGeCK. Across all timepoints and cell lines, we observed a median of 39 knocked

down genes at FDR ≤ 0.05 (41.05%, range 5-60, **Supplementary Tables 7 and 9**).

Depleted genes varied considerably between cell lines and cancer types, suggestive of

divergent physiology. Ranking each gene within a timepoint in ascending order of

depletion FDR, we found top ranked associations for canonical oncogenes such as

KRAS and *CCND1* in most cell lines across time points (**Supplementary Table 7**).

Depletion testing of both genes in UACC903 melanoma cell lines was consistent with

subsequent confirmation of progressive reduction in dCas9-KRAB-ZIM3 expression

(**Supplementary Fig. 7**). We noted no significant knockdown of the canonical tumor

suppressor *TP53*, concordant with the depletion-based design of our screen.

To further validate our experimental design, we assigned sgRNAs from the tiling

component of the pool to either the *TERT* or *CLPTM1L* promoter and tested both genes

using MAGeCK Analysis of *TERT* promoter targeting-sgRNAs demonstrated striking cell

line-specific effects, suggestive of both physiological differences and experimental

limitations. NCI-H520 lung cancer cells showed the most dramatic depletion of *TERT*

sgRNAs, with $\log_2$ fold changes progressing from -0.028 at T_0 to -1.297 at T_2 (FDR =

3.30×10^{-5}), indicating strong growth dependency on the *TERT* promoter. A549 cells

exhibited more modest but consistent depletion of *TERT* promoter sgRNAs across

timepoints ($\log_2$ fold change = -0.062, FDR = 2.55×10^{-4} at T_2). Among melanoma lines,

UACC1113 demonstrated significant growth dependency on the *TERT* promoter ($\log_2$

fold change= -0.217, FDR = 1.90×10^{-5} at T_2), while UACC903 showed minimal effects

across all timepoints (maximum 7 total depleted positive control genes). Bladder cancer

line 5637 transitioned from modest enrichment at T_0 (LFC = 0.030) to significant

depletion by T_2 (LFC = -0.082, FDR = 3.30×10^{-5}), whereas UMUC3 maintained

consistent low-level enrichment (LFC ~0.04) with marginal significance. Pancreatic lines

showed variable patterns: MIA PaCa-2 exhibited minimal fold changes but significant

FDRs, while PANC-1 progressed from non-significant enrichment at T0 to moderate

depletion at later timepoints (LFC = -0.080, FDR = 2.10×10^{-5} at T2). *CLPTM1L*

promoter silencing generally paralleled *TERT* promoter effects but with attenuated

magnitude. NCI-H520 again displayed the strongest dependency (LFC = -0.703, FDR =

4.32×10^{-4} at T_2), followed by UACC1113 (LFC = -0.167, FDR = 1.14×10^{-3} at T_2).

Notably, several samples showed significant *CLPTM1L* promoter effects at early

timepoints: UACC1113 at T_0 (FDR = 4.32×10^{-4}) and MIA PaCa-2 at T_0 (FDR =

4.50×10^{-5}), suggesting immediate growth consequences upon *CLPTM1L* promoter

perturbation in these contexts. A549 showed modest depletion (LFC = -0.059 at T2),

while UMUC3 and PANC-1 displayed weak or non-significant effects (**Supplementary**

Table 8).

We then went on to apply MAGeCK to 5p15.33 CCVs, assigning sgRNAs to CCVs by

intersecting them with $\pm$ 50 bp each fine mapped CCV. SNP-level analysis identified 5

to 54 significantly depleted variants per sample, with depletion counts generally tracking

overall screen sensitivity (**Supplementary Table 9).** NCI-H520 yielded the highest SNP

hits at T_2 (n=54), coinciding with maximal promoter effects and suggesting coordinated

regulatory dependencies across the locus. 5637 and A549 maintained consistent SNP

depletion across time points while melanoma line UACC903 showed minimal regulatory

variant sensitivity (n=5 across all timepoints), consistent with its weak overall selection.

PANC-1 demonstrated a notable increase in depleted SNPs from T_0 (n=10) to T_1/T_2

(n=35), suggesting that SNP-mediated *cis*-regulatory effects may take a longer period to

manifest in phenotypic divergence. While our we primarily hypothesized that functional

SNPs would occupy loci which would be epigenetically silenced upon CRISPRi

inhibition, we observed six SNPs for which sgRNAs were significantly enriched

(**Supplementary Table 10).**

Next, we used CRISPR-SURF[8] to deconvolve sgRNA $\log_2$ fold changes for each

timepoint into discrete regions of putative functional regulatory activity. Unlike gene-

centric or SNP-focused analyses, CRISPR-SURF operates in an unbiased, genome-

agnostic manner, identifying functional regulatory regions based solely on local

enrichment of depleted sgRNAs without prior assumptions about annotated features or

genetic markers. This approach enabled discovery of both promoter-proximal and distal

regulatory elements, as well as unannotated functional sequences across the tiled

5p15.33 region (**Supplementary Fig. 9**). CRISPR-SURF deconvolution failed for MIA

PaCa-2 T_1 and NCI-H520 T_1, and T_2.and UACC903 T_0. Of the remaining datapoints, we

identified a total of 2,718 proliferation-essential elements at FDR ≤ 0.05, with a median

length of 470 bp.

Acknowledgments

This research was supported in part by the Intramural Research Program of the

National Institutes of Health (NIH). The contributions of the NIH authors are considered

Works of the United States Government. The findings and conclusions presented in this

paper are those of the authors and do not necessarily reflect the views of the NIH or the

U.S. Department of Health and Human Services

The authors acknowledge the research contributions of the NCI Cancer Genomics

Research Laboratory for their expertise, execution, and support of this research in the

areas of project planning, wet laboratory processing of specimens, and bioinformatics

analysis of generated data. We thank Dr. Raj Chari, Director, Genome Modification

Core, Frederick National Laboratory for Cancer Research, National Cancer Institute,

Frederick, MD, for assistance with CRISPR guide design. We also thank Dr. Dominic

Esposito, Director, Protein Expression Laboratory (PEL) Frederick National Laboratory

for Cancer Research, National Cancer Institute, Frederick, MD, for assistance in cloning

of VNTR luciferase constructs.

This work utilized the computational resources of the NIH HPC Biowulf cluster

(https://hpc.nih.gov).

SL was supported by CA297773 (NIH).

The Pancreatic Cancer Cohort Consortium and the Pancreatic Cancer Case-

Control Consortium (PanScan/PanC4) Acknowledgements and Funding

statements:

American Cancer Society. The authors express sincere appreciation to all Cancer

Prevention Study-II participants, and to each member of the study and biospecimen

management group. The authors would like to acknowledge the contribution to this

study from central cancer registries supported through the Centers for Disease Control

and Prevention's National Program of Cancer Registries and cancer registries

supported by the National Cancer Institute's Surveillance Epidemiology and End Results

Program. The American Cancer Society funds the creation, maintenance, and updating

of the Cancer Prevention Study-II cohort.

The **ATBC Study** was supported by the Intramural Research Program, Division of

Cancer Epidemiology and Genetics of the U.S. National Cancer Institute (NCI), National

Institutes of Health.

Cancer incidence data for **CLUE** were provided by the Maryland Cancer Registry,

Center for Cancer Surveillance and Control, Department of Health and Mental Hygiene,

201 W. Preston Street, Room 400, Baltimore, MD 21201,

http://phpa.dhmh.maryland.gov/cancer, 410-767-4055. We acknowledge the State of

Maryland, the Maryland Cigarette Restitution Fund, and the National Program of Cancer

Registries of the Centers for Disease Control and Prevention for the funds that support

the collection and availability of the cancer registry data." We thank all the CLUE

participants.

EPIC-Europe Funding Statement and Disclaimer:

EPIC Financial support: The coordination of EPIC-Europe is financially supported by

International Agency for Research on Cancer (IARC) and also by the Department of

Epidemiology and Biostatistics, School of Public Health, Imperial College London which

has additional infrastructure support provided by the NIHR Imperial Biomedical

Research Centre (BRC).

The national cohorts are supported by: Danish Cancer Society (Denmark);Ligue

Nationale Contre le Cancer, Institut Gustave Roussy, Mutuelle Générale de l'Education

Nationale (MGEN), Institut National de la Santé et de la Recherche Médicale

(INSERM), French National Research Agency (ANR, reference ANR-10-COHO-0006),

French Ministry for Higher Education (subsidy 2102918823, 2103236497, and

2103586016) (France); German Cancer Aid, German Cancer Research Center (DKFZ),

German Institute of Human Nutrition Potsdam-Rehbruecke (DIfE), Federal Ministry of

Research, Technology and Space (BMFTR) (Germany); Associazione Italiana per la

Ricerca sul Cancro-AIRC-Italy, Italian Ministry of Health, Italian Ministry of University

and Research (MUR), Compagnia di San Paolo (Italy); Dutch Ministry of Public Health,

Welfare and Sports (VWS), the Netherlands Organisation for Health Research and

Development (ZonMW), World Cancer Research Fund (WCRF), (The Netherlands); UiT

The Arctic University of Norway; Health Research Fund (FIS) - Instituto de Salud Carlos

III (ISCIII), Regional Governments of Andalucía, Asturias, Basque Country, Murcia and

Navarra, and the Catalan Institute of Oncology - ICO (Spain); Swedish Cancer Society,

Swedish Research Council of Region Skåne and Region Västerbotten (Sweden);

Cancer Research UK (C864/A14136 to EPIC-Norfolk; C8221/A29017 to EPIC-Oxford),

Medical Research Council (MR/N003284/1, MC-UU_12015/1 and MC_UU_00006/1 to

EPIC-Norfolk (DOI 10.22025/2019.10.105.00004); MR/Y013662/1 to EPIC-Oxford)

(United Kingdom). Previous support has come from "Europe against Cancer"

Programme of the European Commission (DG SANCO).

EPIC Disclaimer: Where authors are identified as personnel of the International Agency

for Research on Cancer / World Health Organization, the authors alone are responsible

for the views expressed in this article and they do not necessarily represent the

decisions, policy or views of the International Agency for Research on Cancer / World

Health Organization.

The Health Professionals Follow-up Study is supported by NIH grant UM1
CA167552. from the National Cancer Institute, Bethesda, MD USA.

The work at **Johns Hopkins University** was supported by the NCI Grants
P50CA062924 and R01CA154823. Additional support was provided by the Lustgarten
Foundation, Susan Wojcicki and Dennis Troper and the Sol Goldman Pancreas Cancer
Research Center.

The **Multiethnic Cohort Study (MEC)** is supported by NCI grant U01CA164973.

The **MD Anderson case control study** was supported by NIH R01 CA98380.

The **Melbourne Collaborative Cohort Study (MCCS)** cohort recruitment was funded
by VicHealth and Cancer Council Victoria. The MCCS was further augmented by
Australian National Health and Medical Research Council grants 209057, 396414 and
1074383 and by infrastructure provided by Cancer Council Victoria. Cases and their
vital status were ascertained through the Victorian Cancer Registry and the Australian
Institute of Health and Welfare, including the National Death Index and the Australian
Cancer Database.

Work at **Memorial Sloan Kettering Cancer Center** was funded by the center grant
P30CA008748 (Selwybn Vickers).

The **NYU-WHI** study was funded by NIH R01 CA098661, UM1 CA182934 and center
grants P30 CA016087 and P30 ES000260.

The **Nurses' Health Study** is supported by NIH grants UM1 CA186107, P01 CA87969,
and R01 CA49449 from the National Cancer Institute, Bethesda, MD USA.

The **Physicians' Health Study** was supported by research grants CA-097193, CA-
34944, CA-40360, HL-26490, and HL-34595 from the National Institutes of Health,
Bethesda, MD USA

The **Queensland Pancreatic Cancer Study** was supported by a grant from the
National Health and Medical Research Council of Australia (NHMRC) (Grant number
442302). RE Neale is supported by a NHMRC Senior Research Fellowship (#1060183).

The **SELECT** study is supported by National Institutes of Health grant award number
UG1CA189974 (CD Blanke), and U01 CA182883 (CM Tangen/IM Thompson). The
authors thank the site investigators and staff and, most importantly, the participants who
donated their time to this trial.

Data collection at the **University of Toronto** was supported by the CIHR Canada
Research Chair and from the Canadian Cancer Society Research Institute.

The **WHI** program is funded by the National Heart, Lung, and Blood Institute, National
Institutes of Health, U.S. Department of Health and Human Services through contracts
75N92021D00001, 75N92021D00002, 75N92021D00003, 75N92021D00004,
75N92021D00005.The authors thank the WHI investigators and staff for their
dedication, and the study participants for making the program possible. A full listing of
WHI investigators can be found at: https://www-whi-org.s3.us-west-
2.amazonaws.com/wp-content/uploads/WHI-Investigator-Short-List.pdf.

The **Harvard Women's Health Study** funding was as follows: CA047988, CA182913,
HL043851, HL080467 and HL099355.

The **Yale (CT) pancreas cancer study** is supported by National Cancer Institute at the
U.S. National Institutes of Health, grant 5R01CA098870. The cooperation of 30
Connecticut hospitals, including Stamford Hospital, in allowing patient access, is
gratefully acknowledged. The Connecticut Pancreas Cancer Study was approved by the
State of Connecticut Department of Public Health Human Investigation Committee.
Certain data used in that study were obtained from the Connecticut Tumor Registry in
the Connecticut Department of Public Health. The authors assume full responsibility for
analyses and interpretation of these data.

Consortia authors from the Pancreatic Cancer Cohort Consortium and Pancreatic
Cancer Case-Control Consortium (PanScan/PanC4)

Jun Zhong[1], Demetrius Albanes[2], Gabriella Andreotti[3], Samuel O. Antwi[4], Alan A Arslan[5], William R. Bamlet[6], Laura Beane-Freeman[3], Sonja I Berndt[3], Paige M. Bracci[7], Daniele Campa[8], Federico Canzian[9], Stephen J Chanock[10], Yu Chen[11], Charles C Chung[12], Mengmeng Du[13], A. Heather Eliassen[14,15,16], Steven Gallinger[17], J. Michael Gaziano[18,19], Jeanine Genkinger[20], Michael Goggins[21], Phyllis J Goodman[22], Christopher A Haiman[23], Manal Hassan[24], Belynda Hicks[12], Rayjean J. Hung[25], Amy Hutchinson[12], Verena Katzke[26], Manolis Kogevinas[27,28], Charles Kooperberg[29], Martijn P Kolijn[30,31], Peter Kraft[32], Robert C. Kurtz[33], I-Min Lee[34,14], Loic LeMarchand[35], Núria Malats[36], Roger Milne[37,38,39], Steven C Moore[2], Rachel E. Neale[40], Kimmie Ng[41], Ann L. Oberg[6], Irene Orlow[42], Alpa V Patel[43], Ulrike Peters[44], Miquel Porta[45], Kari G. Rabe[46], Francisco X Real[47,48], Harvey A. Risch[49], Nathaniel Rothman[10], Paul Scheet[50], Howard D Sesso[18,14], Veronica W Setiawan[51], Xiao-Ou Shu[52], Debra Silverman[10], Mingyang Song[14,53,54], Meir J Stampfer[14,15,16], Melissa C Southey[37,38,39], Connie Thurman[14], Geoffrey S Tobias[10], Therese Truong[55], Caroline Um[43], Kala Visvanathan[56,57], Nicolas Wentzensen[10], Jean Wactawski-Wende[58], Herbert Yu[59], Chen Yuan[41], Wei Zheng[52], Brian M Wolpin[41], Rachael Z Stolzenberg-Solomon[10], Alison P. Klein[60,61], Laufey T Amundadottir[1]

[1]Laboratory of Translational Genomics, Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA, [2]Metabolic Epidemiology Branch, Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA, [3]Occupational and Environmental Epidemiology Branch, Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA, [4]Department of Quantitative Health Sciences, Mayo Clinic College of Medicine, Jacksonville, FL, USA, [5]Departments of Obstetrics and Gynecology and Population Health, NYU Grossman School of Medicine, NYU Perlmutter Comprehensive Cancer Center, New York, NY, USA, [6]Department of Quantitative Health Sciences, Mayo Clinic College of Medicine, Rochester, MN, USA, [7]Department of Epidemiology and Biostatistics, University of California San Francisco, San Francisco, CA, USA, [8]Unit of Genetics., Department of Biology, University of Pisa, Pisa, Italy, [9]Genomic Epidemiology Group, German Cancer Research Center (DKFZ), Heidelberg, Germany, [10]Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA, [11]Department of Population Health, NYU Grossman School of Medicine, NYU Perlmutter Comprehensive Cancer Center, New York, NY, USA, [12]Cancer Genomics Research Laboratory, Frederick National Lab for Cancer Research, Frederick, MD, USA, [13]Department of Epidemiology and Biostatistics, Memorial Sloan Kettering Cancer Center,, New York, NY, USA, [14]Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA, USA, [15]Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, USA, [16]Department of Nutrition, Harvard T. H. Chan School of Public Health, Boston, MA, USA, [17]Lunenfeld-Tanenbaum Research Institute, Sinai Health System and University of Toronto,, Toronto, Ontario, Canada, [18]Division of Preventive Medicine, Brigham and Women's Hospital, Boston, MA, USA, [19]Division of Aging, Brigham and Women's Hospital, Boston, MA, USA, [20]Department of Epidemiology, Columbia University, New York, NY, USA, [21]Department of Pathology, Johns Hopkins School of Medicine, Baltimore, MD, USA, [22]SWOG Statistical Center, Fred Hutchinson Cancer Center, Seattle, WA, USA, [23]Department of Preventive Medicine, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA, [24]Department of Epidemiology, Cancer Prevention, The University of Texas MD Anderson Cancer Center, Houston, TX, USA, [25]Prosserman Centre for Population Health Research, Lunenfeld-Tanenbaum Research Institute, Sinai Health System, Toronto, Ontario, Canada, [26]Division of Cancer Epidemiology, German Cancer Research Center (DKFZ), Heidelberg, Germany, [27]ISGlobal, Centre for Research in Environmental Epidemiology (CREAL), Barcelona, Spain, [28]Hospital del Mar Institute of Medical Research (IMIM), Universitat Autònoma de Barcelona, Barcelona, Spain, [29]Division of Public Health Sciences, Fred Hutchinson Cancer Center, Seattle, WA, USA, [30]Division of Environmental Epidemiology and Veterinary Public Health, Institute for Risk Assessment Sciences, Utrecht University, Utrecht, The Netherlands, [31]Julius Global Health, The Julius Center for Health Sciences and Primary Care, University Medical Center, [32]Trans-Divisional Research Program (TDRP), Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA, [33]Gastroenterology, Hepatology, and Nutrition Service, Memorial Sloan Kettering Cancer Center, New York, NY, USA, [34]Division of Preventive Medicine, Department of Medicine, Brigham and Women's Hospital, Boston, MA, USA, [35]Population Sciences in the Pacific Program, Cancer Epidemiology, University of Hawaii Cancer Center, Honolulu, HI, USA, [36]Genetic and Molecular Epidemiology Group, Spanish National Cancer Research Center (CNIO), Madrid, Spain, [37]Cancer Epidemiology Division, Cancer Council Victoria, Melbourne, VIC, Australia, [38]Centre for Epidemiology and

Biostatistics, Melbourne School of Population and Global Health, The University of Melbourne, Melbourne, VIC,
Australia, [39]Precision Medicine, School of Clinical Sciences at Monash Health, Monash University, Clayton, VIC,
Australia, [40]Department of Population Health, QIMR Berghofer Medical Research Institute, Queensland, Australia,
[41]Department of Medical Oncology, Dana-Farber Cancer Institute, Harvard Medical School, Harvard University,
Boston, MA, USA, [42]Department of Epidemiology and Biostatistics, Memorial Sloan Kettering Cancer Center, New
York, NY, USA, [43]Department of Population Science, American Cancer Society, Atlanta, GA, USA, [44]Division of
Public Health Sciences, Fred Hutchinson Cancer Center, Seattle, WA, USA, [45]Hospital del Mar Institute of Medical
Research (IMIM), Centro de Investigacion Biomedica en Red de Epidemiologia y Salud Publica (CIBERESP),
Barcelona, Spain, [46]Department of Quantitative Health Sciences Research, Mayo Clinic College of Medicine,
Rochester, MN, USA, [47]Epithelial Carcinogenesis Group, Tumor Biology Programme, Spanish National Cancer
Research Center (CNIO), Madrid, Spain, [48]Department of Medicine and Life Sciences, Universitat Pompeu Fabra,
Barcelona, Spain, [49]Department of Chronic Disease Epidemiology, Yale School of Public Health, New Haven, CT,
USA, [50]Dept of Epidemiology, The University of Texas MD Anderson Cancer Center, Houston, TX, USA,
[51]Department of Population and Public Health Sciences, Keck School of Medicine, University of Southern California,
Los Angeles, CA, USA, [52]Division of Epidemiology, Department of Medicine, Vanderbilt Epidemiology Center,
Vanderbilt-Ingram Cancer Center, Vanderbilt University School of Medicine, Nashville, TN, USA, [53]Department of
Nutrition, Harvard T.H. Chan School of Public Health, Boston, MA, USA, [54]Clinical and Translational Epidemiology
Unit and Division of Gastroenterology, Massachusetts General Hospital and Harvard Medical School, Boston, MA,
USA, [55]Paris-Saclay University, UVSQ, Inserm, Gustave Roussy, CESP, Villejuif, France, [56]Department of
Epidemiology, Johns Hopkins School of Public Health, Baltimore, MD, USA, [57]Department of Oncology, Sidney
Kimmel Comprehensive Cancer Center, Johns Hopkins School of Medicine, Baltimore, MD, USA, [58]Department of
Epidemiology and Environmental Health, University of Buffalo, Buffalo, NY, USA, [59]Epidemiology Program, University
of Hawaii Cancer Center, Honolulu, HI, USA, [60]Department of Oncology, Sidney Kimmel Comprehensive Cancer
Center, Johns Hopkins School of Medicine, Baltimore, MD, USA, [61]Department of Medicine, Division of
Gastroenterology & Hepatology, Johns Hopkins School of Medicine, Baltimore, MD, USA.

1495

References

1. Alsheikh, A. J. *et al.* The landscape of GWAS validation; systematic review identifying 309 validated non-coding variants across 130 human diseases. *BMC Med Genomics* **15**, 74 (2022).

2. Lappalainen, T. *et al.* Transcriptome and genome sequencing uncovers functional variation in humans. *Nature* **501**, 506–511 (2013).

3. Aguet, F. *et al.* Genetic effects on gene expression across human tissues. *Nature* **550**, 204–213 (2017).

4. Wang, Z. *et al.* Imputation and subset-based association analysis across different cancer types identifies multiple independent risk loci in the TERT-CLPTM1L region on chromosome 5p15.33. *Human Molecular Genetics* **23**, 6616–6633 (2014).

5. Chen, H. *et al.* Large-scale cross-cancer fine-mapping of the 5p15.33 region reveals multiple independent signals. *Human Genetics and Genomics Advances* **2**, 100041 (2021).

6. Bhattacharjee, S. *et al.* A Subset-Based Approach Improves Power and Interpretation for the Combined Analysis of Genetic Association Studies of Heterogeneous Traits. *The American Journal of Human Genetics* **90**, 821–835 (2012).

7.   Fang, J. *et al.* Functional characterization of a multi-cancer risk locus on

chr5p15.33 reveals regulation of TERT by ZNF148. *Nature Communications* **8**, 1–

15 (2017).

8.   Kim, N. W. *et al.* Specific Association of Human Telomerase Activity with Immortal

Cells and Cancer. *Science* **266**, 2011–2015 (1994).

9.   Castelo-Branco, P. *et al.* Methylation of the TERT promoter and risk stratification of

childhood brain tumours: an integrative genomic and molecular study. *The Lancet*

Oncology **14**, 534–542 (2013).

10.   Shay, J. W. & Bacchetti, S. A survey of telomerase activity in human cancer.

European Journal of Cancer **33**, 787–791 (1997).

11.   Yoo, S. S. *et al. TERT* Polymorphism rs2853669 Influences on Lung Cancer Risk

in the Korean Population. *J Korean Med Sci* **30**, 1423 (2015).

12.   Bojesen, S. E. *et al.* Multiple independent variants at the TERT locus are

associated with telomere length and risks of breast and ovarian cancer. *Nat Genet*

45, 371–384 (2013).

13.   Beesley, J. *et al.* Functional Polymorphisms in the TERT Promoter Are Associated

with Risk of Serous Epithelial Ovarian and Breast Cancers. *PLOS ONE* **6**, e24987

(2011).

14.   Walsh, K. M. *et al.* Variants near TERT and TERC influencing telomere length are

associated with high-grade glioma risk. *Nat Genet* **46**, 731–735 (2014).

15. Mosrati, M. A. *et al.* TERT promoter mutations and polymorphisms as prognostic

factors in primary glioblastoma. *Oncotarget* **6**, 16663–16673 (2015).

16. Harland, M. *et al.* Germline TERT promoter mutations are rare in familial

melanoma. *Familial Cancer* **15**, 139–144 (2016).

17. George, J. R. *et al.* Association of TERT Promoter Mutation, But Not BRAF

Mutation, With Increased Mortality in PTC. *J Clin Endocrinol Metab* **100**, E1550–

E1559 (2015).

18. Rachakonda, P. S. *et al.* TERT promoter mutations in bladder cancer affect patient

survival and disease recurrence through modification by a common polymorphism.

Proceedings of the National Academy of Sciences **110**, 17426–17431 (2013).

19. Pópulo, H. *et al.* TERT promoter mutations in skin cancer: the effects of sun

exposure and X-irradiation. *J Invest Dermatol* **134**, 2251–2257 (2014).

20. Griewank, K. G. *et al.* TERT Promoter Mutation Status as an Independent

Prognostic Factor in Cutaneous Melanoma. *J Natl Cancer Inst* **106**, dju246 (2014).

21. Spiegl-Kreinecker, S. *et al.* Prognostic quality of activating TERT promoter

mutations in glioblastoma: interaction with the rs2853669 polymorphism and

patient age at diagnosis. *Neuro Oncol* **17**, 1231–1240 (2015).

22. Horn, S. *et al.* TERT promoter mutations in familial and sporadic melanoma.

Science **339**, 959–961 (2013).

23.　Hosen, I. *et al.* Mutations in TERT promoter and FGFR3 and telomere length in

bladder cancer. *International Journal of Cancer* **137**, 1621–1629 (2015).

24.　Labussière, M. *et al.* TERT promoter mutations in gliomas, genetic associations

and clinico-pathological correlations. *Br J Cancer* **111**, 2024–2032 (2014).

25.　Hiyama, E. *et al.* Telomerase activity is detected in pancreatic cancer but not in

benign tumors. *Cancer Res* **57**, 326–331 (1997).

26.　Bell, R. J. A. *et al.* Understanding TERT Promoter Mutations: a Common Path to

Immortality. *Mol Cancer Res* **14**, 315–323 (2016).

27.　James, M. A., Vikis, H. G., Tate, E., Rymaszewski, A. L. & You, M.

CRR9/CLPTM1L Regulates Cell Survival Signaling and Is Required for Ras

Transformation and Lung Tumorigenesis. *Cancer Res* **74**, 1116–1127 (2014).

28.　Clarke, W. R., Amundadottir, L. & James, M. A. CLPTM1L/CRR9 ectodomain

interaction with GRP78 at the cell surface signals for survival and chemoresistance

upon ER stress in pancreatic adenocarcinoma cells. *International Journal of*

Cancer **144**, 1367–1378 (2019).

29.　Jia, J. *et al.* CLPTM1L Promotes Growth and Enhances Aneuploidy in Pancreatic

Cancer Cells. *Cancer Res* **74**, 2785–2795 (2014).

30.　Parashar, D. *et al.* Targeted biologic inhibition of both tumor cell-intrinsic and

intercellular CLPTM1L/CRR9-mediated chemotherapeutic drug resistance. *npj*

Precis. Onc. **5**, 16 (2021).

31. James, M. A. *et al.* Functional Characterization of CLPTM1L as a Lung Cancer

Risk Candidate Gene in the 5p15.33 Locus. *PLOS ONE* **7**, e36116 (2012).

32. Derrien, A.-C. *et al.* Functional characterization of 5p15.33 risk locus in uveal

melanoma reveals rs452384 as a functional variant and NKX2.4 as an allele-

specific interactor. *Am J Hum Genet* **109**, 2196–2209 (2022).

33. Chang, Y.-H. *et al.* Isoform-level analyses of 6 cancers uncover extensive genetic

risk mechanisms undetected at the gene-level. *Br J Cancer* **133**, 874–885 (2025).

34. Zhong, J. *et al.* A Transcriptome-Wide Association Study Identifies Novel

Candidate Susceptibility Genes for Pancreatic Cancer. *JNCI: Journal of the*

National Cancer Institute https://doi.org/10.1093/jnci/djz246 (2020)

doi:10.1093/jnci/djz246.

35. Yang, Y.-C. *et al.* rs401681 and rs402710 confer lung cancer susceptibility by

regulating TERT expression instead of CLPTM1L in East Asian populations.

Carcinogenesis **39**, 1216–1221 (2018).

36. Wei, R. *et al.* TERT Polymorphism rs2736100-C Is Associated with EGFR

Mutation-Positive Non-Small Cell Lung Cancer. *Clin Cancer Res* **21**, 5173–5180

(2015).

37. Helbig, S. *et al.* Functional dissection of breast cancer risk-associated TERT

promoter variants. *Oncotarget* **8**, 67203–67217 (2017).

38. Ge, M. *et al.* Functional evaluation of TERT-CLPTM1L genetic variants associated

with susceptibility of papillary thyroid carcinoma. *Sci Rep* **6**, 26037 (2016).

39. Zhou, L. *et al.* The identification of two regulatory ESCC susceptibility genetic

variants in the TERT-CLPTM1L loci. *Oncotarget* **7**, 5495–5506 (2015).

40. Hsiung, C. A. *et al.* The 5p15.33 Locus Is Associated with Risk of Lung

Adenocarcinoma in Never-Smoking Females in Asia. *PLoS Genet* **6**, e1001051

(2010).

41. Florez-Vargas, O. *et al.* Genetic regulation of TERT splicing affects cancer risk by

altering cellular longevity and replicative potential. *Nat Commun* **16**, 1676 (2025).

42. Hofer, P. *et al.* MNS16A tandem repeats minisatellite of human telomerase gene: a

risk factor for colorectal cancer. *Carcinogenesis* **32**, 866–871 (2011).

43. Wang, L. *et al.* Association of a functional tandem repeats in the downstream of

human telomerase gene and lung cancer. *Oncogene* **22**, 7123–7129 (2003).

44. Yuan, H. *et al.* MNS16A polymorphism of the TERT gene on cancer risk: a

systematic review and meta-analysis. *Nucleosides, Nucleotides & Nucleic Acids*

42, 1004–1018 (2023).

45. Zou, Y., Carbonetto, P., Wang, G. & Stephens, M. Fine-mapping from summary

data with the "Sum of Single Effects" model. *PLOS Genetics* **18**, e1010299 (2022).

46. Klein, A. P. *et al.* Genome-wide meta-analysis identifies five new susceptibility loci

for pancreatic cancer. *Nat Commun* **9**, 556 (2018).

47. Landi, M. T. *et al.* Genome-wide association meta-analyses combining multiple risk

phenotypes provide insights into the genetic architecture of cutaneous melanoma

susceptibility. *Nat Genet* **52**, 494–504 (2020).

48. Koutros, S. *et al.* Genome-wide Association Study of Bladder Cancer Reveals New

Biological and Translational Insights. *Eur Urol* **84**, 127–137 (2023).

49. McKay, J. D. *et al.* Large-scale association analysis identifies new lung cancer

susceptibility loci and heterogeneity in genetic susceptibility across histological

subtypes. *Nat Genet* **49**, 1126–1132 (2017).

50. Shi, J. *et al.* Genome-wide association study of lung adenocarcinoma in East Asia

and comparison with a European population. *Nat Commun* **14**, 3043 (2023).

51. Byrska-Bishop, M. *et al.* High-coverage whole-genome sequencing of the

expanded 1000 Genomes Project cohort including 602 trios. *Cell* **185**, 3426-

3440.e19 (2022).

52. Wolpin, B. M. *et al.* Genome-wide association study identifies multiple susceptibility

loci for pancreatic cancer. *Nat Genet* **46**, 994–1000 (2014).

53. Zhong, J. *et al.* Large-scale multiomic analysis identifies non-coding somatic driver

mutations and nominates ZFP36L2 as a driver gene for pancreatic ductal

adenocarcinoma. *Gut* https://doi.org/10.1136/gutjnl-2025-335152 (2025)

doi:10.1136/gutjnl-2025-335152.

54. Long, E. *et al.* High-throughput characterization of functional variants highlights

heterogeneity and polygenicity underlying lung cancer susceptibility. *Am J Hum*

Genet **111**, 1405–1419 (2024).

55. Choi, J. *et al.* Massively parallel reporter assays of melanoma risk variants identify

MX2 as a gene promoting melanoma. *Nat Commun* **11**, 2718 (2020).

56. Colli, L. M. *et al.* Altered regulation of DPF3, a member of the SWI/SNF complexes,

underlies the 14q24 renal cancer susceptibility locus. *The American Journal of*

Human Genetics **108**, 1590–1610 (2021).

57. Boyle, A. P. *et al.* Annotation of functional variation in personal genomes using

RegulomeDB. *Genome Res.* **22**, 1790–1797 (2012).

58. Breeze, C. E. *et al.* FORGEdb: a tool for identifying candidate functional variants

and uncovering target genes and mechanisms for complex diseases. *Genome Biol*

25, 3 (2024).

59. Coetzee, S. G., Coetzee, G. A. & Hazelett, D. J. motifbreakR: an R/Bioconductor

package for predicting variant effects at transcription factor binding sites.

Bioinformatics **31**, 3847–3849 (2015).

60. Tsherniak, A. *et al.* Defining a Cancer Dependency Map. *Cell* **170**, 564-576.e16

(2017).

61. Li, W. *et al.* MAGeCK enables robust identification of essential genes from

genome-scale CRISPR/Cas9 knockout screens. *Genome Biol* **15**, 554 (2014).

62. Hsu, J. Y. *et al.* CRISPR-SURF: discovering regulatory elements by deconvolution

of CRISPR tiling screen data. *Nat Methods* **15**, 992–993 (2018).

63. Mukamel, R. E. *et al.* Repeat polymorphisms underlie top genetic risk loci for

glaucoma and colorectal cancer. *Cell* **186**, 3659-3673.e23 (2023).

64. Lee, O. W. *et al.* Targeted long-read sequencing of the Ewing sarcoma 6p25.1

susceptibility locus identifies germline-somatic interactions with EWSR1-FLI1

binding. *The American Journal of Human Genetics* **110**, 427–441 (2023).

65. Grünewald, T. G. P. *et al.* Chimeric EWSR1-FLI1 regulates the Ewing sarcoma

susceptibility gene EGR2 via a GGAA microsatellite. *Nat Genet* **47**, 1073–1078

(2015).

66. Amundadottir, L. *et al.* Genome-wide association study identifies variants in the

ABO locus associated with susceptibility to pancreatic cancer. *Nat Genet* **41**, 986–

990 (2009).

67. Petersen, G. M. *et al.* A genome-wide association study identifies pancreatic

cancer susceptibility loci on chromosomes 13q22.1, 1q32.1 and 5p15.33. *Nat*

Genet **42**, 224–228 (2010).

68. Zagouri, F. *et al.* HTERT MNS16A polymorphism in breast cancer: a case–control

study. *Mol Biol Rep* **39**, 10859–10863 (2012).

69. Thomas-Chollier, M. *et al.* Transcription factor binding predictions using TRAP for

the analysis of ChIP-seq data and regulatory SNPs. *Nat Protoc* **6**, 1860–1869

(2011).

70. Yu, F.-X., Zhao, B. & Guan, K.-L. Hippo Pathway in Organ Size Control, Tissue

Homeostasis, and Cancer. *Cell* **163**, 811–828 (2015).

71. Landin Malt, A. *et al.* Alteration of TEAD1 Expression Levels Confers Apoptotic

Resistance through the Transcriptional Up-Regulation of Livin. *PLoS ONE* **7**,

e45498 (2012).

72. Zhao, B., Li, L., Lei, Q. & Guan, K.-L. The Hippo–YAP pathway in organ size

control and tumorigenesis: an updated version. *Genes Dev.* **24**, 862–874 (2010).

73. Landin Malt, A., Georges, A., Silber, J., Zider, A. & Flagiello, D. Interaction with the

Yes-associated protein (YAP) allows TEAD1 to positively regulate NAIP

expression. *FEBS Letters* **587**, 3216–3223 (2013).

74. Liu-Chittenden, Y. *et al.* Genetic and pharmacological disruption of the TEAD–YAP

complex suppresses the oncogenic activity of YAP. *Genes Dev.* **26**, 1300–1305

(2012).

75. Szeto, S. G. *et al.* YAP/TAZ Are Mechanoregulators of TGF-β-Smad Signaling and

Renal Fibrogenesis. *J Am Soc Nephrol* **27**, 3117–3128 (2016).

76. Weischer, M. *et al.* Short Telomere Length, Cancer Survival, and Cancer Risk in

47102 Individuals. *J Natl Cancer Inst* **105**, 459–468 (2013).

77. Pooley, K. A. *et al.* A genome-wide association scan (GWAS) for mean telomere

length within the COGS project: identified loci show little association with hormone-

related cancer risk. *Hum Mol Genet* **22**, 5056–5064 (2013).

78. Wentzensen, I. M., Mirabello, L., Pfeiffer, R. M. & Savage, S. A. The association of

telomere length and cancer: a meta-analysis. *Cancer Epidemiol Biomarkers Prev*

20, 1238–1250 (2011).

79. Campa, D. *et al.* Leukocyte Telomere Length in Relation to Pancreatic Cancer

Risk: A Prospective Study. *Cancer Epidemiol Biomarkers Prev* **23**, 2447–2454

(2014).

80. Lynch, S. M. *et al.* A prospective analysis of telomere length and pancreatic cancer

in the alpha-tocopherol beta-carotene cancer (ATBC) prevention study.

International Journal of Cancer **133**, 2672–2680 (2013).

81. Nan, H. *et al.* Shorter Telomeres Associate with a Reduced Risk of Melanoma

Development. *Cancer Res* **71**, 6758–6763 (2011).

82. Iles, M. M. *et al.* The Effect on Melanoma Risk of Genes Previously Associated

With Telomere Length. *J Natl Cancer Inst* **106**, dju267 (2014).

83. Seow, W. J. *et al.* Telomere Length in White Blood Cell DNA and Lung Cancer: A

Pooled Analysis of Three Prospective Cohorts. *Cancer Res* **74**, 4090–4098 (2014).

84.   Antwi, S. O. *et al.* Leukocyte Telomere Length and Its Interaction with Germline

1708   Variation in Telomere-Related Genes in Relation to Pancreatic Adenocarcinoma

1709   Risk. *Cancer Epidemiol Biomarkers Prev* **29**, 1492–1500 (2020).

1710   85.   Pearson, J. D. *et al.* Binary pan-cancer classes with distinct vulnerabilities defined

1711   by pro- or anti-cancer YAP/TEAD activity. *Cancer Cell* **39**, 1115-1134.e12 (2021).

1712   86.   Jakubosky, D. *et al.* Properties of structural variants and short tandem repeats

1713   associated with gene expression and complex traits. *Nat Commun* **11**, 2927

1714   (2020).

1715   87.   Bhati, M., Mapel, X. M., Lloret-Villas, A. & Pausch, H. Structural variants and short

1716   tandem repeats impact gene expression and splicing in bovine testis tissue.

1717   *Genetics* **225**, iyad161 (2023).

1718   88.   Jermusyk, A. *et al.* A 584 bp deletion in CTRB2 inhibits chymotrypsin B2 activity

1719   and secretion and confers risk of pancreatic cancer. *Am J Hum Genet* **108**, 1852–

1720   1865 (2021).

1721   89.   Bakhtiari, M. *et al.* Variable number tandem repeats mediate the expression of

1722   proximal genes. *Nat Commun* **12**, 2075 (2021).

1723   90.   Garg, P. *et al.* Pervasive *cis* effects of variation in copy number of large tandem

1724   repeats on local DNA methylation and gene expression. *The American Journal of*

1725   *Human Genetics* **108**, 809–824 (2021).

91.   Gymrek, M. *et al.* Abundant contribution of short tandem repeats to gene

expression variation in humans. *Nat Genet* **48**, 22–29 (2016).

92.   Vinagre, J. *et al.* Frequency of TERT promoter mutations in human cancers. *Nat*

Commun **4**, 2185 (2013).

93.   Coetzee, S. G. & Hazelett, D. J. motifbreakR v2: expanded variant analysis

including indels and integrated evidence from transcription factor binding

databases. *Bioinformatics Advances* **4**, vbae162 (2024).

94.   Murphy, A. E., Schilder, B. M. & Skene, N. G. MungeSumstats: a Bioconductor

package for the standardization and quality control of many GWAS summary

statistics. *Bioinformatics* **37**, 4593–4596 (2021).

95.   Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and

richer datasets. *Gigascience* **4**, s13742-015-0047–8 (2015).

96.   Machiela, M. J. & Chanock, S. J. LDlink: a web-based application for exploring

population-specific haplotype structure and linking correlated alleles of possible

functional variants: Fig. 1. *Bioinformatics* **31**, 3555–3557 (2015).

97.   Sanson, K. R. *et al.* Optimized libraries for CRISPR-Cas9 genetic screens with

multiple modalities. *Nat Commun* **9**, 5416 (2018).

98.   PacificBiosciences/pbAA. PacBio (2025).

99.   Dolzhenko, E. *et al.* Characterization and visualization of tandem repeats at

genome scale. *Nat Biotechnol* **42**, 1606–1614 (2024).

100. Poplin, R. *et al.* A universal SNP and small-indel variant caller using deep neural

networks. *Nat Biotechnol* **36**, 983–987 (2018).

101. Holt, J. M. *et al.* HiPhase: jointly phasing small, structural, and tandem repeat

variants from HiFi sequencing. *Bioinformatics* **40**, btae042 (2024).

102. Delaneau, O., Zagury, J.-F., Robinson, M. R., Marchini, J. L. & Dermitzakis, E. T.

Accurate, scalable and integrative haplotype estimation. *Nat Commun* **10**, 5436

(2019).

103. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *GigaScience* **10**,

giab008 (2021).

104. Browning, B. L., Zhou, Y. & Browning, S. R. A One-Penny Imputed Genome from

Next-Generation Reference Panels. *The American Journal of Human Genetics*

103, 338–348 (2018).

105. Zhang, Y. *et al.* Multisample motif discovery and visualization for tandem repeats.

Genome Res. **35**, 850–862 (2025).

106. Li, W. *et al.* Quality control, modeling, and visualization of CRISPR screens with

MAGeCK-VISPR. *Genome Biology* **16**, 281 (2015).

1762