1   Wildlife hosts predict the distribution of reported coccidioidomycosis in the western United States
2

3   **Authors**
4   Jacob Sussman[1], Katrina M. Derieg[2], Kevin D. Perry[3], Ashlyn Adakai[4], Raniah Corrian[5], Cory Merow[6],
5   Simon C. Brewer[7], Katharine S. Walter[8,9]
6
7   **Affiliations**
8
[1] Department of Biostatistics and Bioinformatics, Duke University School of Medicine, Durham, NC,
USA
[2] Philip L. Wright Zoological Museum, University of Montana, Missoula, MT, USA
[3] Department of Atmospheric Sciences, University of Utah, Salt Lake City, UT, USA
[4] Department of Engineering, University of Arizona, Tucson, AZ, USA
[5] Department of Psychology, Clark Atlanta University, Atlanta, GA, USA
[6] Eversource Energy Center and Department of Ecology and Evolutionary Biology, University of
Connecticut, Storrs, CT, USA
[7] School of Environment, Society and Sustainability, University of Utah, Salt Lake City, UT, USA
[8] Division of Epidemiology, University of Utah, Salt Lake City, UT, USA
[9] Natural History Museum of Utah, Salt Lake City, UT, USA

20
21
Abstract

Global environmental change is reshaping human exposure to zoonotic and environmentally acquired
pathogens, yet predicting disease risk remains challenging. High-resolution risk maps typically rely on
human case data and environmental correlates, often overlooking ecological processes such as wildlife
reservoirs. We evaluated whether mammalian reservoir distributions improve prediction of
coccidioidomycosis (Valley fever), an emerging, environmentally-acquired fungal disease with a poorly
characterized range. Using county-level coccidioidomycosis notification data from the Centers for
Disease Control and Prevention, we developed a hierarchical Bayesian model of county-level endemicity,
defined as >=10 cases per 100,000 population. We incorporated climatic, environmental, and vegetation
covariates, a state-level reporting effect, and species distribution models for 22 mammalian species
previously identified as *Coccidioides* reservoirs. We found that the number of endemic mammalian
reservoirs in a county was the strongest predictor of coccidioidomycosis endemicity, with each standard
deviation increase in reservoir species richness associated with substantially higher odds of endemicity
(log-odds ratio = 1.702; 95% CI: 1.060-2.419). In contrast, maximum vapor pressure deficit, soil
moisture, and land cover were not independently associated with endemicity after accounting for reservoir
distributions. State-level reporting effects revealed substantial heterogeneity, and comparison of models
with and without reporting effects identified regions likely to be endemic but underreported, including
parts of Nevada, Utah, New Mexico, Texas, and Colorado. Our results establish reservoir diversity as a
central predictor of zoonotic fungal disease risk and demonstrate a transferable framework for
distinguishing between ecological drivers of infection from surveillance bias to improve disease risk
mapping and identify areas of potential underreporting.

43
44 **Keywords**
45 Coccidioidomycosis, Valley fever, zoonotic pathogen, spatial model
46
47

48 **Funding**
49 Burroughs Wellcome Climate and Health Interdisciplinary Grant

50
51 **Significance statement**

52
Valley fever, or coccidioidomycosis, is an emerging fungal disease caused by the pathogen *Coccidioides*,
with a geographic range that is not well understood. Maps of Valley fever risk primarily rely on human
case data, overlooking potentially ecological influences like wildlife. This study demonstrates that the
ranges of mammalian reservoirs of *Coccidioides* help predict the distribution of human infection risk. By
integrating species distribution models with epidemiological data, this study also identifies reporting gaps
in the Western United States where *Coccidioides* may be endemic but human cases are underreported.
Our findings highlight areas where unknown Valley fever risk may be present and offers a framework for
anticipating disease risk as environmental shifts occur.

Introduction

Zoonotic and environmentally acquired pathogens are responsible for a large proportion of human infections and are among those most sensitive to global change[1,2]. Unlike for many directly transmitted human diseases, vaccines are often unavailable and public health measures must focus on preventing zoonotic and environmental exposures. Yet we have only a limited understanding of the environmental distributions of many pathogens due to the prohibitive costs of systematic surveys, and must rely on reported human case data, which is sparse and biased.

Coccidioidomycosis, or Valley fever, a fungal disease caused by inhalation of *Coccidioides immitis* and *C. posadasii* spores, is one such disease with poorly defined infection risk. Systematic environmental sampling for *Coccidioides* is infeasible. Therefore, skin testing studies conducted in the 1950s[3,4], along with contemporary reported cases of disease remain the basis for the Center of Disease Control and Prevention's current Valley fever risk maps[5]. Improved risk maps are urgently needed because reported cases of the disease in the United States have dramatically increased in recent years, from 2,271 in 1998 to 21,037 in 2023[8]. Poverty, outdoor occupations such as farming and construction, and incarceration in endemic areas exacerbate risk, compounding inequities in healthcare access [6].

Although climatic and soil characteristics correlate with fungal range, they are insufficient predictors of risk of disease[7]. Further, while soil has long been considered the primary reservoir of *Coccidioides spp.*, there is increasing evidence for the endozoan hypothesis, that zoonotic hosts maintain *Coccidioides spp.* and seed the soil with infectious spores when they die[8]. Biodiversity data is increasingly available[9,10], and host distributions have been successfully incorporated into risk mapping for several zoonotic diseases, including Lassa fever, plague, and hantaviruses, where the presence or abundance of small mammal reservoirs strongly predicts human spillover. Building on this precedent, we tested whether small mammal distributions help explain the endemicity of *Coccidioides*, thereby providing a novel ecological layer for mapping Valley fever risk.

Coccidioidomycosis case reports are subject to substantial biases, making it difficult to infer the true distribution of endemicity using standard modeling approaches. Many infections are never reported because symptoms are mild or resemble other respiratory diseases, and because of low levels of population and physician awareness and lack of access to healthcare. The true incidence has been estimated to be up to thirty times higher than reported cases[11,12]. Reporting also varies widely across the western United States. Although coccidioidomycosis is a nationally notifiable disease with a standardized case definition, states are not obligated to submit reports to the Centers for Disease Control and Prevention. As a result, several endemic states, including Texas, do not report cases[13]. Beyond reporting biases, infection risk for environmentally acquired pathogens such as *Coccidioides* is spatially autocorrelated, further complicating inference with conventional regression approaches.

Integrated Nested Laplace Approximation (INLA) approaches enable efficient Bayesian spatial modelling of spatially autocorrelated data to generate maps of infection risk[10,14]. INLA models have been successfully applied to map other environmentally acquired and zoonotic pathogens, including leptospirosis[15], schistosomiasis[16], dengue[17], and Zika virus[17]. This approach is therefore well suited to generate updated, data-driven maps of coccidioidomycosis endemicity while rigorously testing ecological hypotheses such as the role of rodent reservoirs.

Here, we developed an INLA model to predict coccidioidomycosis endemicity in the western U.S. We tested whether mammalian reservoir distributions predict coccidioidomycosis endemicity, while also assessing environmental and reporting factors. Our approach refines coccidioidomycosis risk maps and provides a generalizable framework for mapping environmentally acquired pathogens.

Methods

Human case data

We obtained Valley fever human case data reported to the National Notifiable Diseases Surveillance System through a data use agreement with the Centers for Disease Control and Prevention

(CDC). Cases are reported from 1998 to 2022 by states. In this study, we use the data reported for the year 2022. The data reports a monthly case frequency grouped by state and county Federal Information Processing Standards code. Additionally, the reporting month is identified as the month of disease onset or diagnosis. We summed the total cases reported in each county for each month to obtain the total number of Valley fever cases reported in each U.S. county in 2022. While coccidioidomycosis is a Nationally Notifiable Disease, reporting varies by state, we used the CDC list of reportable diseases to identify states that report coccidioidomycosis[18].

To align with prediction goals specified by the CDC, we define counties as endemic if incidence is ≥10 cases per 100,000 people, and treated endemicity as a binary outcome. While we are primarily interested in results for counties west of the Mississippi (the "western" U.S.), to avoid introducing artificial spatial boundaries, we include all U.S. counties in our analysis. Regions outside of the spatial neighborhood domain like Alaska, Hawaii, and other territories were not included in analysis.

County data

We obtained county shapefiles from U.S. census data and accessed through the R package tidycensus[19]. Using the same package, the 2022 county population size was obtained from the American Community Survey[20].

Bioclimatic covariates

We obtained bioclimatic data for covariates that have been associated with or that we predicted were associated with *Coccidioides* presence[7,21–24]. We obtained monthly values of maximum temperature, mean temperature, minimum temperature, precipitation, minimum vapor pressure deficit (VPD) and maximum VPD from the PRISM datasets and extracted mean county raster values averaging over all months in 2022 using the R package prism[25,26]. Climate factors like temperature and precipitation are commonly used in models of *Coccidioides* endemicity due to their influence on fungal growth.[7]

We obtained modeled soil properties at a 30-second spatial resolution, and a depth of 0-5 cm from SoilGRIDS, accessed through the package geodata[27,28]. We accessed soil temperature and soil moisture from the 2022 NLDAS-NOAH products from NASA[29]. Soil sampling studies suggest that *Coccidioides* relies on soil as an exclusive source of nutrients, and therefore soil parameters should be considered when building distribution models.[23]

We accessed land cover data from the Multi-Resolution Land Characteristics Consortium database provided by the U.S. Geological Survey.[29] We assigned each county a single value, the "dominant cover type," by identifying the most common land cover category raster value in each county. When the dominant land cover type was water, we assigned the next most common cover type. We extracted mean county values for all covariates and attached all data to a simple features (sf) object in R[31]. While the role of land cover and agriculture on *Coccidioides* presence remains unclear, we include land cover covariates in our modeling to strengthen the diversity of ecological factors used outside of mammal distributions.

Mammal distributions

To test whether reservoir distributions were associated with disease endemicity, we created a map that included the distributions for any mammal previously reported to be infected with *Coccidioides*. First, we compiled all previous studies documenting *Coccidioides* infection through culture, post-mortem serology, or ITS2 amplicon sequencing of mammal lung tissue [32–34] (Supplementary Table 1). We then generated species distribution models (SDMs) for each species, following previously published methods[35]. For each SDM, we converted continuous model outputs into a binary outcome by considering any raster values greater than or equal to 6 to mean that the mammal species is present. We summed the number of species with ranges intersecting each county to obtain a total number of reservoir hosts occurring in each county.

Covariate selection

We excluded covariates that were highly correlated (>80% correlation). Candidate models were
compared using the Widely Applicable Information Criterion (WAIC) and the Deviance Information
Criterion (DIC). Candidate models with different covariate subsets were compared using WAIC to
identify the final specification. In our final model, we included population, mean vegetation, mean soil
moisture, mean soil bulk density (BDOD), maximum VPD, and a simplified grouping of land cover
classes, and number of potential reservoirs.

INLA modeling
We modeled county level endemicity using a Bayesian spatial model fitted with integrated nested
Laplace Approximation (INLA v 25.6.7) for areal data. The outcome, a binary indicator of endemicity,
was modeled with a binomial likelihood and logit link. To capture county-level spatial autocorrelation we
used the BYM2 parametrization of the Besag-York-Mollie model[36], which describes unexplained county-
level variation as the mixture of a spatially-structured intrinsic conditional autoregressive component (u)
and an unstructured independent and identically distributed (i.i.d.) component (v):

180
$$b = 1/\sqrt{\tau}(\sqrt{(1-\phi)}v + \sqrt{\phi}u)$$

The spatial structure was defined using a county-adjacency matrix based on Queen's case
contiguity. We specified a penalized complexity (PC) prior for the spatial hyperparameters[37]. The prior on
the precision (τ) was given as $P\left(\frac{1}{\sqrt{\tau}} > 1\right) = 0.01$, and the prior on the mixing parameter (ϕ) was given as
$P(\phi > 0.5) = 2/3$ [38]. The PC prior penalizes deviation from a base model with no spatial dependence.
In addition to county spatial effects, we included an i.i.d. state-level random effect to account for
unobserved state-specific heterogeneity not captured by the county structure. To characterize the
underlying spatial signal in the outcome, we first fit a spatial-only baseline model that included no
covariates. The baseline model allows us to evaluate how much of the variation in endemicity is
attributable purely to spatial dependence.
We then fit a full covariate model with both county and state-level spatial effects. Fixed effects
included population, species presence count, mean vegetation, mean soil moisture, mean soil bulk density,
VPD maximum and a simplified land-cover category. We standardized continuous predictors prior to
modeling. Models were fit with INLA, and we describe model fit using deviance information criterion
(DIC) and Watanabe-Akaike information criterion (WAIC). We obtained the county-level probability of
endemicity from the full INLA model by extracting the mean of the posterior fitted values for each county
and assigning this value as the probability of endemicity.
To isolate the environmental component of the model, after fitting we extracted the fixed-effect
estimates and constructed a design matrix containing only the environmental covariates. We computed the
linear predictor corresponding to these fixed effects. The environmental-only prediction reflects the
contribution of environmental factors to endemicity independent of county or state spatial effects.

All code is publicly available on GitHub:
https://github.com/sussmanjacob/vf_distribution_modeling.

Human subjects
The University of Utah Institutional Review Board determined this study was Non-Human
Subjects Research (IRB_00177442).

Results
In 2022, 17,562 cases of coccidioidomycosis were reported in the United States (Fig. 1a).
Endemic counties were concentrated in the southwest, and incidence was highest in Kern County,
California (265 cases per 100,000) and Pinal County, Arizona (193 cases per 100,000). Forty nine of
1,501 counties in our study area met CDC criteria for endemicity (incidence >=10 per 100,000) (Fig. 1b).
While most endemic counties are in Arizona (15 counties) and the Central Valley of California (17),
Utah, New Mexico, and Nevada also report endemic regions. We also observed counties meeting the

CDC's criteria for endemicity in Minnesota and Missouri, further east than the expected range of
Coccidioides.
To explore spatial variation in human case reports, we fit a spatial INLA model with no
covariates which accounts for spatial autocorrelation in case reports and serves as a baseline for
evaluating the effect of ecological covariates. The spatial-only model showed substantial county-level
variation, indicating structured spatial heterogeneity in endemicity that environmental covariates might
explain..
To assess whether the distribution *Coccidioides* reservoirs is associated with reported
coccidioidomycosis, we first mapped the distributions of all previously described confirmed small
mammal reservoirs of *Coccidioides* (Fig. 2). Counties with high incidence of reported disease also have
the highest number of potential reservoir species (Fig. 2). Fifteen counties in Arizona, and 19 counties in
California, have more than ten endemic small mammal reservoirs. *Coccidioides* reservoirs are also found
in parts of Utah, Colorado, New Mexico, and Texas.
We then fit a spatial INLA model to predict county-level coccidioidomycosis endemicity
including the number of locally endemic *Coccidioides* reservoir species, VPD, and mean soil moisture
among others. Model predictions of coccidioidomycosis endemicity was largely consistent with reported
cases, with endemic counties concentrated in Arizona and the Central Valley, California (Fig. 4a). Thirty-
one counties had >50% probability of endemicity, clustered in Arizona and California's Central Valley.
The highest predicted probability was 96% in Maricopa County, Arizona.
Presence of *Coccidioides* reservoir species was the strongest predictor of county endemicity (log-
odds ratio per SD = 1.702; 95% CI: 1.060-2.419) (Fig. 3a). Other environmental variables, including
vegetation, soil moisture, soil bulk density, maximum temperature, and land cover category were not
significantly associated with endemicity (Fig. 3a). We performed additional sensitivity analysis across
endemicity thresholds to examine the robustness of the positive association between reservoir presence
and endemicity. Reservoir presence remained significant when the endemicity threshold was reduced to 5
cases per 100,000 (LOR: 1.217; 95% CI: 0.703-1.746) and increased to 15 cases per 100,000 (LOR:
1.674; 95% CI: 0.792-2.461).
We compared multiple covariate specifications in order to examine the additional value of
including reservoir range covariates in the model using WAIC. The model including mammal reservoir
distributions achieved the lowest WAIC among covariate models (WAIC = 280.82), outperforming the
model without reservoir predictors (change in WAIC = 32.71). This provides strong evidence that
reservoir species presence contributes predictive power beyond other included ecological covariates.
State-level random effects captured differences in reporting probability not explained by
ecological covariates (Fig. 3b). The Arizona random effect was significantly higher than zero (3.435, 95%
CI: 0.977-6.411), indicating that cases were more likely to be reported here than expected by a model
where reporting was constant across the Western US. South Dakota (4.07, 95% CI: 1.60-7.61) and
Minnesota (3.358, 95% CI: 1.152-5.993) also had mean state random effects above zero (Fig. 3b). States
with lower random effect values included New Mexico (-1.457, 95% CI: -3.486-0.450), Colorado (-1.955,
95% CI: -4.828-0.565), and Texas (-2.325, 95% CI: -5.107-0.229), though these did not differ
significantly from zero.
The state-level random effect exhibited greater variation (SD = 1.38) than the county-level spatial
effect (SD = 0.05), indicating state-level heterogeneity is primarily responsible for residual variation after
accounting for environmental covariates. This is likely attributable to differences in reporting across
states. The BYM2 parameter (φ = 0.29; 95% CI: 0.01–0.90) suggests predominantly unstructured
variation, though the wide credible interval indicates substantial uncertainty about the balance of spatial
structure. In our model unexplained county-level factors affecting endemicity do not exhibit strong spatial
patterns beyond what is explained by environmental covariates.
Our hierarchical model structure also enabled us to make model predictions based solely on
environmental covariates, excluding population, county spatial effects, and state reporting effects, which
we interpreted as predictions of environmental endemicity of *Coccidioides*, regardless of human case
reporting (Fig. 4b). The probability of environmental endemicity is, like human cases, concentrated in the

Southwestern U.S. However, this model predicts a higher probability of environmental endemicity in
southern Nevada, southern and eastern Utah, and southern New Mexico, and western Texas.
We calculated the difference between predicted environmental endemicity and observed human
case endemicity (Fig. 4c). Negative differences indicate counties where the fungus is predicted to be
present but human cases are under-reported. Arizona and California's Central Valley and Bay Area
exhibited above-average reporting rates. Reporting rates are lower than the mean in southern Nevada,
southern and eastern Utah, southern New Mexico, western Texas, and southwestern Colorado (Fig. 4c).
There is substantial uncertainty in the disparity between predicted environmental and human case
endemicity, as shown by the standard deviation of the difference in predicted probabilities (Fig. 4d). The
greatest uncertainty occurred in northern Arizona and southern Utah.
277
278 **Discussion**
279
We found that the presence of small mammal reservoirs of *Coccidioides* was the most important
predictor of coccidioidomycosis endemicity. We also observed substantial differences in state-level
reporting, suggesting that surveillance infrastructure influences the apparent distribution of disease.
Our findings have important public health implications. First, the observed discrepancies between
environmental suitability and reported cases highlight gaps in surveillance and case recognition. States
with low relative reporting rates or where coccidioidomycosis is not reportable may be underestimating
their true disease burden, limiting their ability to allocate resources and raise awareness of
coccidioidomycosis risk among the public and healthcare providers. Second, the strong relationship
between small mammal reservoirs and endemicity supports the endozoan hypothesis, which suggests that
burrowing mammals harbor *Coccidioides* and contribute to its persistence in soil[8]. This suggests that
mammal distributions provide important information for both understanding the current range of
Coccidioides and predicting changes in its distribution under shifting climate and land use patterns. One
mechanism through which reservoir richness increases cases may be that in areas of high species richness,
both specialist and generalist species are present. Generalists are more likely to be commensal with
humans and spread disease. Generalist species opportunistically occupy burrows made by other species.
In an area with low reservoir richness, multiple types of burrows are less likely to be occupied, potentially
decreasing *Coccidioides* transmission[39].
Our results build upon prior models of the distribution of coccidioidomycosis. Previous work has
focused heavily on climatic and soil covariates. Prior modeling approaches have adopted occupancy
models to account for imperfect reporting[21]. Our approach builds on an occupancy modeling framework
by predicting risk effectively in unsampled areas. By integrating ecological information on potential
reservoirs with human surveillance data, our models advance the capacity to disentangle environmental
and reporting factors underlying of observed endemicity. Although environmental predictors such as soil
moisture and temperature are important for fungal survival, their correlation with small mammal
distributions complicates inference.
Our study has several possible limitations. Reported human case data are biased by under-
diagnosis, under-reporting, and variation in testing practices across states, which may lead to
misclassification of endemic counties. Information on mammalian reservoirs is also incomplete; for
example, the apparent scarcity of reservoir species in Oregon and Washington could reflect biological
reality or simply a lack of systematic screening of mammals in those regions. Incomplete sampling of
small mammals across their range can lead to uncertain range maps Furthermore, many of the covariates
included in our models are interdependent: small mammal ranges are shaped by environmental
conditions, making it difficult to determine causality. Lastly, our study considers the distribution of
Coccidioides in a single year. Temporal and seasonal changes in our covariates and Valley fever case
reporting may add additional information to explain the role of mammal reservoirs in *Coccidioides*
endemicity and may help generate more stable predictions of future Valley fever risk. Expanding our
modeling approach to include a temporal component is an important future direction.

317   Enhanced surveillance systems in under-reporting states will be critical to improve recognition of
Valley fever and to better define its true range. Future research should prioritize systematic surveys of
small mammal populations across the western United States to clarify which species are competent
reservoirs of *Coccidioides*. In addition, preserved samples in natural history collections and
biorepositories could provide additional insight into the historical spread of *Coccidioides*[40,41]. Finally,
longitudinal field studies and genomic approaches that link mammalian ecology, fungal persistence, and
human infections are needed to establish causal pathways and refine predictive models.
324   Together, our findings underscore the importance of considering both ecological and surveillance
325   in understanding the distribution of coccidioidomycosis. By incorporating small mammal reservoirs into
326   models of endemicity, we provide stronger evidence for the endozoan hypothesis and highlight critical
327   gaps in public health reporting. These results set the stage for more targeted ecological and
328   epidemiological studies to inform prevention and control of Valley fever.
329
330

Acknowledgments

This work was supported by a Burroughs Wellcome Fund (Climate and Health Interdisciplinary Award to KSW, KD, and KP).

Competing Interest Statement

The authors declare no competing interest.

References

Allen T, Murray KA, Zambrana-Torrelio C, *et al.* Global hotspots and correlates of emerging zoonotic diseases. *Nature Communications 2017 8:1* 2017; **8**: 1–10.

Jones KE, Patel NG, Levy MA, *et al.* Global trends in emerging infectious diseases. *Nature 2008 451:7181* 2008; **451**: 990–3.

Palmer CE, Edwards PQ, Allfather WE. Characteristics of skin reactions to coccidioidin and histoplasmin, with evidence of an unidentified source of sensitization. *Am J Epidemiol* 1957; **66**: 196–213.

Edwards PQ, Palmer CE. Prevalence of sensitivity to coccidioidin, with special reference to specific and nonspecific reactions to coccidioidin and to histoplasmin. *Dis Chest* 1957; **31**: 35–60.

Centers for Disease Control and Prevention. Valley Fever Maps. 2020.

Rosenstein N, Emery KW, Werner S Ben, *et al.* Risk factors for severe pulmonary and disseminated coccidioidomycosis: Kern County, California, 1995-1996. *Clinical Infectious Diseases* 2001; **32**: 708–15.

Gorris ME, Treseder KK, Zender CS, Randerson JT. Expansion of Coccidioidomycosis Endemic Regions in the United States in Response to Climate Change. *Geohealth* 2019; **3**: 308–27.

Taylor JW, Barker BM. The endozoan, small-mammal reservoir hypothesis and the life cycle of Coccidioides species. *Med Mycol* 2019; **57**: S16–20.

GBIF.org. GBIF Home Page. 2023. https://www.gbif.org/ (accessed Aug 14, 2023).

Rue H, Martino S, Chopin N. Approximate Bayesian inference for latent Gaussian models by using integrated nested Laplace approximations. *J R Stat Soc Series B Stat Methodol* 2009; **71**: 319–92.

Centers for Disease Control and Prevention (CDC). Valley Fever (Coccidioidomycosis) Statistics. 2022. https://www.cdc.gov/fungal/diseases/coccidioidomycosis/statistics.html#one.

Valdivia L, Nix D, Wright M, *et al.* Coccidioidomycosis as a Common Cause of Community-acquired Pneumonia - Volume 12, Number 6—June 2006 - Emerging Infectious Diseases journal - CDC. *Emerg Infect Dis* 2006; **12**: 958–62.

Gorris ME, Ardon-Dryer K, Campuzano A, *et al.* Advocating for Coccidioidomycosis to Be a Reportable Disease Nationwide in the United States and Encouraging Disease Surveillance across North and South America. *Journal of Fungi* 2023; **9**: 83.

Lindgren F, Rue H, Lindström J. An explicit link between Gaussian fields and Gaussian Markov random fields: the stochastic partial differential equation approach. *J R Stat Soc Series B Stat Methodol* 2011; **73**: 423–98.

Baquero OS, Machado G. Spatiotemporal dynamics and risk factors for human Leptospirosis in Brazil. *Sci Rep* 2018; **8**. DOI:10.1038/S41598-018-33381-3.

Hu Y, Ward MP, Xia C, *et al.* Monitoring schistosomiasis risk in East China over space and time using a Bayesian hierarchical modeling approach OPEN. 2016. DOI:10.1038/srep24173.

Martínez-Bello DA, López-Quílez A, Torres Prieto A. Spatio-Temporal Modeling of Zika and Dengue Infections within Colombia. *International Journal of Environmental Research and Public Health 2018, Vol 15, Page 1376* 2018; **15**: 1376.

Centers for Disease Control and Prevention. Reportable Fungal Diseases by State. 2024. https://www.cdc.gov/fungal/php/case-reporting/index.html (accessed Oct 1, 2025).

Walker K, Herman M. tidycensus: Load US Census Boundary and Attribute Data as 'tidyverse' and 'sf'-Ready Data Frames. 2025. https://walker-data.com/tidycensus/ (accessed Oct 1, 2025).

US Census Bureau. American Community Survey. 2024 https://www.census.gov/programs-surveys/acs/ (accessed Oct 2, 2025).

Hepler SA, Kaufeld KA, Kline D, Greene A, Gorris ME. Estimating coccidioidomycosis endemicity while accounting for imperfect detection using spatio-temporal occupancy modeling. *Am J Epidemiol* 2025; **194**: 56–63.

22 Barker BM, Litvintseva AP, Riquelme M, Vargas-Gastélum L. Coccidioides ecology and
genomics. Med. Mycol. 2019; **57**: S21–9.
23 Chow NA, Kangiser D, Gade L, *et al.* Factors Influencing Distribution of Coccidioides immitis in
Soil, Washington State, 2016. *mSphere* 2021; **6**. DOI:10.1128/MSPHERE.00598-
21/ASSET/20386D35-913C-4022-848A-
49E0992AA47A/ASSETS/IMAGES/MEDIUM/MSPHERE.00598-21-F004.GIF.
24 Wagner R, Montoya L, Head JR, Campo S, Remais J, Taylor JW. Coccidioides undetected in soils
from agricultural land and uncorrelated with time or the greater soil fungal community on
undeveloped land. *PLoS Pathog* 2023; **19**. DOI:10.1371/journal.ppat.1011391.
25 PRISM Group OSU. PRISM Data. 2014. https://prism.oregonstate.edu/ (accessed Oct 1, 2025).
26 Edmund H, Bell K. prism: Access Data from the Oregon State Prism Climate Project . CRAN:
Contributed Packages. 2025; published online March 25.
DOI:10.32614/CRAN.PACKAGE.PRISM.
27 Hijmans RJ. Access Geographic Data [R package geodata version 0.6-6]. *CRAN: Contributed*
Packages 2025; published online Sept 30. DOI:10.32614/CRAN.PACKAGE.GEODATA.
28 Poggio L, De Sousa LM, Batjes NH, *et al.* SoilGrids 2.0: Producing soil information for the globe
with quantified spatial uncertainty. *SOIL* 2021; **7**: 217–40.
29 Rodell M, Houser PR, Jambor U, *et al.* The Global Land Data Assimilation System. *Bull Am*
Meteorol Soc 2004; **85**: 381–94.
30 Multi-Resolution Land Characteristics Consortium. Land Cover (CONUS) Data. 2024
https://www.mrlc.gov/data?f%5B0%5D=category%3ALand%20Cover (accessed Oct 2, 2025).
31 Pebesma E, Bivand R. Spatial Data Science. Chapman and Hall/CRC, 2023
DOI:10.1201/9780429459016.
32 del Rocío Reyes-Montes M, Pérez-Huitrón MA, Ocaña-Monroy JL, *et al.* The habitat of
Coccidioides spp. and the role of animals as reservoirs and disseminators in nature. BMC Infect.
Dis. 2016; **16**. DOI:10.1186/s12879-016-1902-7.
33 Emmons CW. Coccidioidomycosis in Wild Rodents. A Method of Determining the Extent of
Endemic Areas. 1943 https://about.jstor.org/terms.
34 Salazar-Hamm PS, Montoya KN, Montoya L, *et al.* Breathing can be dangerous: Opportunistic
fungal pathogens and the diverse community of the small mammal lung mycobiome. *Frontiers in*
Fungal Biology 2022; **3**. DOI:10.3389/ffunb.2022.996574.
35 Carlson CJ, Albery GF, Merow C, *et al.* Climate change increases cross-species viral transmission
risk. *Nature* 2022; **607**: 555–62.
36 Besag J, York J, Mollié A. Bayesian image restoration, with two applications in spatial statistics.
Ann Inst Stat Math 1991; **43**: 1–20.
37 Simpson D, Rue H, Riebler A, Martins TG, Sørbye SH. Penalising Model Component
Complexity: A Principled, Practical Approach to Constructing Priors. *https://doi.org/101214/16-*
STS576 2017; **32**: 1–28.
38 Moraga P. Spatial Statistics for Data Science: Theory and Practice with R. 2023
DOI:10.1201/9781032641522.
39 Kraft JP, Stapp P. Movements and burrow use by northern grasshopper mice as a possible
mechanism of plague spread in prairie dog colonies. *J Mammal* 2013; **94**. DOI:10.1644/12-
MAMM-A-197.1.
40 Paul DL, Thompson CW, Arroyo L, *et al.* Harnessing natural history collections for collaborative
pandemic preparedness. *Bioscience* 2025. DOI:10.1093/biosci/biaf035.
41 Thompson CW, Phelps KL, Allard MW, *et al.* Preserve a voucher specimen! the critical need for
integrating natural history collections in infectious disease studies. *mBio* 2021; **12**: 1–20.
434

Figure 1: Coccidiomycosis incidence per 100,000 individuals at the county level in 2022 (A), and counties which met the endemicity definition of 10 or more cases per 100,000 in 2022 (B). Valley fever case data was provided by the CDC. Does not report means the county is in a state where Coccidiomycosis is not reported to the CDC.

Figure 2: Number of potential *Coccidioides* reservoir mammal species predicted to be present in each western U.S. county, based on species distribution model overlap.

Figure 3: Log-odds ratio means for model fixed effects and 95% C.I. (A), and posterior mean of model state I.I.D. effect (B).

Figure 4: Modeled probability of endemicity of reported Coccidioidomycosis disease from the full model (A), modeled probability of environmental endemicity using only environmental covariates (B), the difference between the two levels (C), and the standard deviation of the predicted difference in the probability of endemicity (D).