# Peak Alpha Frequency as a Neural Marker of Postoperative Pain Outcomes in Spinal Fusion Surgery

Amélie Grandjean[1], Fares Komboz[2], Tatjana Chacon[2], Lukas Weiser[3], Wolfgang Lehman[3], Angelina Nazarenus[2], Dorothee Mielke[2,4], Veit Rohde[2], Ali Mazaheri[1.]* and Tammam Abboud[2] *

1 School of Psychology, Centre for Human Brain Health, University of Birmingham, United Kingdom
2 Department of Neurosurgery, 3, Department of Orthopedic Surgery, University Medical Center Göttingen, Germany
4 Department of Neurosurgery, University Medical Center Augsburg, Germany

* Tammam Abboud and Ali Mazaheri contributed equally to this work.

**Abstract**

**Objective**

Postoperative pain outcomes following spinal fusion surgery remain difficult to predict, as structural and surgical indicators alone offer limited insight into who will experience meaningful relief. A substantial proportion of patients continue to report persistent pain after surgery, underscoring the need for objective markers that can help identify those at risk of poor recovery. Peak alpha frequency (PAF) has emerged as a promising trait-like neural signature of pain sensitivity in experimental models, where individuals with slower PAF tend to exhibit heightened pain sensitivity. Yet despite this link, its ability to forecast longer-term postoperative pain trajectories remains unclear.

**Methods**

Seventeen adults undergoing cervical or lumbar fusion surgery were included. Resting-state, eyes-closed EEG was recorded preoperatively and at multiple visits after surgery. PAF was extracted from central electrodes using the centre-of-mass method. Pain intensity was assessed longitudinally on standardised self-report pain scales. Associations between PAF measures and postoperative pain change were examined using correlation analyses, and receiver operating characteristic (ROC) analyses evaluated discrimination of pain responders ($\geq$50% improvement).

**Results**

Preoperative peak alpha frequency (PAF) was positively associated with longer-term pain reduction at the 3-month follow-up, but showed no consistent relationship with early postoperative pain. Across pain measures, a consistent pattern emerged across the Brief Pain Inventory (BPI), visual analogue scale (VAS), and numerical rating scale (NRS), but not the verbal rating scale (VRS) or Short-Form McGill (SF-MPQ). At the 3-month follow-up, associations reached statistical significance for BPI-Worst ($\rho = 0.67$, p = 0.017), and BPI-Average Pain ($\rho = 0.62$, p = 0.033). VAS and NRS showed moderate-to-strong effects that approached significance in non-parametric analyses and were significant for VAS when treated as an approximately interval measure (*Pearson* r = 0.63, p = 0.022).

ROC analyses using BPI-Worst pain improvement demonstrated good discriminative ability of preoperative PAF for identifying treatment responders at 3 months (*AUC = 0.84; 95% CI: 0.61-1.00*), with high specificity and moderate sensitivity at the Youden-optimal threshold of 10.11 Hz. By contrast, changes in PAF over time were not reliably related to changes in pain scores, suggesting that PAF functions more as a stable, trait-like predictor than a dynamic biomarker in this context.

**Conclusion**

This study demonstrates the feasibility and potential clinical value of preoperative EEG for characterising individual differences in postoperative pain recovery following spinal fusion. The results identify faster preoperative PAF as a stable neural signal that captures meaningful variability in longer-term pain reduction, with convergent support across multiple patient-reported measures. While replication in a larger cohort is required, these findings establish a clear foundation for evaluating PAF as a candidate neurophysiological marker to inform preoperative risk profiling and potentially personalised perioperative pain-management strategies in spinal fusion patients.

## 1. Introduction

Spine pain is one of the most prevalent musculoskeletal sources of pain globally (Wu *et al.*, 2020). Back pain is reported to affect 60-80% of all working adults at some point in their lives (Ehrlich, 2003) and is a leading cause of functional limitation and absence from work (Driscoll *et al.*, 2014). Both low back pain and neck pain are increasing in prevalence and combined are one of the largest contributors to duration of time living with a disability across countries and age ranges (Vos *et al.*, 2016; Hurwitz *et al.*, 2018). Beyond physical disability, chronic back and neck pain is also associated with poorer mental health (Demyttenaere *et al.*, 2007), highlighting the multifaceted impact of spine pain and the importance of understanding how to accurately diagnose and treat it.

**The clinical problem: mismatch between structural measures and symptoms**

Spinal surgery is an established treatment for patients with neck or back pain caused by structural pathology such as stenosis, disc degeneration, or instability when conservative management fails (Scheufler, Dohmen and Vougioukas, 2007; Weinstein *et al.*, 2008; Phillips *et al.*, 2013; Middendorp *et al.*, 2017). Spinal fusion aims to stabilise affected segments and relieve neural compression to reduce pain (Mason *et al.*, 2014; Liang *et al.*, 2024)(Reisener *et al.*, 2020).

Surgical decision-making is typically based on a combination of symptoms, neurological findings, and imaging evidence of degenerative changes (Chou *et al.*, 2007; Weinstein *et al.*, 2008; Hurwitz *et al.*, 2018). Despite substantial advances in imaging and surgical techniques (Mason *et al.*, 2014; Liang *et al.*, 2024) structural correction does not reliably translate into pain relief. Degenerative changes are frequently observed in asymptomatic individuals, while symptomatic patients may show limited explanatory imaging findings (Tonosu *et al.*, 2016; Kasch *et al.*, 2022). This mismatch between structural measures and pain complicates patient selection and outcome prediction in spine surgery.

**Outcome variability: not all patients benefit from surgery**

The discordance between structural pathology and symptoms extends to postoperative outcomes, with a substantial proportion of patients failing to achieve meaningful pain relief after spine surgery (Finkelstein and Schwartz, 2019). MRI-based measures of disease severity do not reliably predict pain or long-term improvement, and surgical success is therefore primarily defined by patient-reported outcome measures rather than radiographic findings (Copay *et al.*, 2008; Dworkin *et al.*, 2008; Weber *et al.*, 2016).

Approximately 15-40% of patients experience insufficient improvement despite technically successful surgery (Trief, Grant and Fredrickson, 2000). While psychological and quality-of-life factors are associated with outcomes, they provide limited predictive accuracy (Celestin, Edwards and Jamison, 2009; Abbott, Tyni-Lenné and Hedlund, 2010). Heterogeneous pain mechanisms and variable recovery trajectories further complicate outcome prediction and patient selection (Tonosu *et al.*, 2016; Cooper *et al.*, 2018).

**Importance of determining objective biomarkers**

Imaging findings and psychological factors, while central to surgical planning, have not been shown to reliably predict postoperative pain outcomes following spinal fusion surgery (Celestin, Edwards and Jamison, 2009; Weber *et al.*, 2016). Moreover, relying too heavily on psychological explanations can be poorly received by patients and may fail to reflect the biological mechanisms that contribute to pain persistence (Newton *et al.*, 2013). This highlights the need for objective biomarkers capable of distinguishing peripheral pathology from centrally mediated pain processes and improving individual pain-profiling approaches (Woolf, 2011).

A growing body of research has explored molecular, inflammatory, biomechanical, and integrative biomarkers of spine pain, including cytokines, inflammatory markers, immune phenotypes, and movement-based measures (Rannou *et al.*, 2007; Shamji *et al.*, 2010; Rathod *et al.*, 2014; Laumet *et al.*, 2019; Vo *et al.*, 2023). However, despite methodological advances and the use of data-driven approaches, no biomarker has yet demonstrated sufficient predictive accuracy for routine clinical use.

**Pain and the brain**

Peripheral and biomechanical measures alone cannot explain the marked interindividual variability in pain experienced after comparable structural injury. Brain-based biomarkers offer direct insight into central pain modulation and the neural mechanisms that construct pain perception (Melzack and Katz, 2004; Tan, Oswald and Kuner, 2021). Cortical oscillations in the alpha frequency range (7-14 Hz) measured using electroencephalography (EEG) and magnetoencephalography (MEG) are closely linked to sensory gating and cortical excitability, processes consistently implicated in pain processing and vulnerability (Klimesch, Sauseng and Hanslmayr, 2007; Jensen and Mazaheri, 2010; Ploner, Sorg and Gross, 2017; Kisler *et al.*, 2020; Kim and Davis, 2021). This has positioned EEG-derived measures as promising candidates for objective pain biomarkers, although clinical translation remains limited (Ahmad and Barkana, 2025).

The peak alpha frequency (PAF) - the dominant frequency within the alpha band is a trait-like measure with high test-retest reliability that reflects fundamental properties of brain function (Grandy *et al.*, 2013; Popov *et al.*, 2023). Slower PAF has been observed across several chronic pain populations and is associated with increased pain sensitivity and longer pain duration (Sarnthein *et al.*, 2006; de Vries *et al.*, 2013; Lim *et al.*, 2016; Furman *et al.*, 2020). Importantly, baseline PAF measured before pain exposure predicts both experimental and prolonged musculoskeletal pain, suggesting that PAF may reflect a pre-existing vulnerability rather than a consequence of pain (Seminowicz *et al.*, 2018; Mazaheri, Seminowicz and Furman, 2022).

Clinically, a slower PAF may indicate that pain processing is more centralised rather than peripherally driven, and therefore less likely to improve through structural surgical intervention. Supporting feasibility, preoperative PAF has been shown to predict acute postoperative pain following thoracic surgery; a context where the outcome of interest is pain *emerging after* the procedure (Millard *et al.*,

2022). In contrast, spinal fusion surgery aims to *reduce pre-existing pain through structural correction*, and it is unknown whether PAF can predict pain improvement in this setting.

**Aims and hypotheses**

This study investigates PAF as a biomarker of pain outcomes in patients undergoing cervical or lumbar fusion surgery. Specifically, we examined PAF as (1) a predictive biomarker, by testing whether preoperative PAF is associated with postoperative pain improvement, and (2) a dynamic biomarker, by assessing whether longitudinal changes in PAF across postoperative visits are related to changes in pain.

Based on evidence that PAF reflects a stable, trait-like index of pain sensitivity and pain-relevant neural dynamics, we tested the following hypotheses:

1. Predictive hypothesis: Slower preoperative PAF is associated with smaller postoperative reductions in pain.
2. Dynamic hypothesis: Increases in PAF over postoperative follow-up are associated with greater improvements in pain, reflecting recovery of pain-relevant brain dynamics.

## 2. Materials and Methods

### 2.2 Study design and population

This is a prospective, observational clinical study that was approved by the local Ethics Committee of Universitätsmedizin Göttingen (approval number: 11/6/23) and registered in the German Clinical Trials Register (DRKS00032164).

Only patients with a planned instrumented spinal fusion procedure for a degenerative disease of the cervical or lumbar spine were eligible for inclusion. Patient recruitment was conducted at the Departments of Neurosurgery and Orthopaedics, University Medical Center Göttingen.

Inclusion criteria were: (1) indication for surgical treatment of a degenerative spine disease requiring an instrumented fusion procedure, (2) age between 18 and 80 years, and (3) ability to understand the nature, significance, and scope of the study and to provide written informed consent.

Exclusion criteria included: severe internal medical or infectious disease, severe psychiatric illness, and current abuse of drugs, alcohol, or medication.

### Surgical procedure

Included patients underwent instrumented fusion of the cervical or lumbar spine, with or without additional decompression. The indication for spinal instrumentation was based on the presence of micro- or macroinstability of the affected spinal segment, as determined by clinical and radiological assessment. Patients treated with isolated decompression procedures or functional pain interventions were not included in this study.

### 2.3 Participant assessments

Participants were assessed at four different time points. Visit 1 data was collected 72 to 24 hours preoperatively, visit 2 data were collected 20 to 24 hours postoperatively, visit 3 at discharge and visit 4 was at 3-month follow-up. All study procedures are presented in Figure 1.

**Figure 1**: A flow diagram including the study visits and procedures.

**EEG recordings**

Resting-state EEG was recorded at three time points: *preoperatively* (72-24 hours before surgery), *at hospital discharge*, and at the *3 month follow-up visit.*

Participants were seated and instructed to remain relaxed with their eyes closed throughout the recording. EEG data were acquired using ISIS Xpert® and NeuroExplorer 2019 (V. 5.1) software (Inomed Medizintechnik GmbH, Emmendingen, Germany) with eight scalp electrodes positioned according to the international 10-20 system. Signals were recorded at a sampling rate of 20,000 Hz with online filtering set to a high-pass filter 0.50 Hz, low-pass filter 70 Hz. Electrode impedances were maintained below 15kΩ threshold for all channels. The duration of each resting-state recording was approximately 10 minutes.

**EEG preprocessing**

The data were preprocessed using the EEGLAB toolbox (Delorme and Makeig, 2004). Continuous recordings were first downsampled to 500 Hz, and the initial and final 3 seconds were removed to avoid edge artefacts. Line noise was removed using a 45-55 Hz notch filter, after which a 0.3 Hz high-pass and 40 Hz low-pass filter were applied. Each recording was then visually inspected to remove overly noisy segments of data due to movement. Across all sessions, the mean duration of rejected data per recording was 9.63 seconds (range: 0 - 72.55 seconds). A single channel was only removed from two recordings; otherwise, all present electrodes were retained.  Finally, the datasets were re-referenced to the average reference and the online reference channel Cz was added back into the data.

**PAF**

FieldTrip toolbox was used for frequency decomposition using the ft_freqanalysis function (Oostenveld *et al.*, 2011). The data were segmented into 5-second epochs, and power spectra computed for each individual epoch using a multitaper Fourier transform with a Hanning taper, yielding power estimates on a per-trial basis. Spectral estimates were calculated over a frequency range of 2-30 Hz.

PAF was estimated for each epoch using the Centre of Mass (CoM) method also called the centre of gravity method) which calculates the weighted average frequency within the predefined alpha band. This approach has been used in previous pain research (e.g. (Furman *et al.*, 2018; Millard *et al.*, 2022) and is advantageous because it captures shifts in the *distribution* of alpha power rather than relying on a single spectral peak. Peak-picking can be unstable across epochs, as the maximum bin is susceptible to noise and may fluctuate substantially from trial to trial.

 We defined as CoM as follows

$$\text{CoM} = \frac{\sum_{i=1}^{n} f_i * a_i}{\sum_{i=1}^{n} a_i}$$

Where $f_i$ is the ith frequency bin within the predetermined frequency range of interest of 7-14 Hz. n represents the 36 frequency bins ((14-7)/0.2 = 35, +1 for inclusion of end points), and $a_i$ is the amplitude at the given $f_i$. This is to say, each frequency within the 7-14 Hz range is multiplied by its power, the values are summed, then divided by the total power in the band, producing a weighted average that reflects the frequency at which alpha power is most concentrated.

PAF was calculated at the central region of interest comprising Cz, C3, and C4 due to alpha activity in sensorimotor regions predicting pain sensitivity in prolonged musculoskeletal pain (Furman *et al.*, 2018; Seminowicz *et al.*, 2018). Epoch-level central PAF values were then averaged over trials and central electrodes for a PAF estimate at each visit.

**Instruments for the assessment of pain**

Although multiple validated pain measures were collected at each visit (outlined below), primary analyses focused on the Brief Pain Inventory (BPI) (Cleeland and Ryan, 1994). Unlike unidimensional intensity scales, the BPI provides a standardised, clinically meaningful assessment of both pain severity and its functional impact, which is particularly relevant in surgical populations where improvement is typically defined by reductions in pain and disability. Using the BPI as the primary outcome therefore offers a more comprehensive and interpretable measure of recovery while the additional scales support convergent validation of pain reports.

Pain assessments were conducted at all four study visits using established unidimensional pain scales:

*Brief Pain Inventory (BPI)*

The BPI is a validated questionnaire assessing pain severity and its impact on daily functioning. Analyses focused on the four pain severity items current pain, and worst, least, and average pain over the past 24 hours. The BPI is widely used in both clinical and research setting, and the worst pain item is of particular relevance to PAF-pain associations in surgical populations (Millard *et al.*, 2022).

*Visual Analogue Scale (VAS)*

Pain intensity was measured using a 100-mm VAS, a validated instruments widely used in clinical pain assessment (Huskisson, 1974; McCormack, Horne and Sheather, 1988).

*Numerical Rating Scale (NRS)*

Participants also completed an 11-point NRS (0-10), a commonly used measure of pain, including in spine surgery research (Jensen, Karoly and Braver, 1986; Khor *et al.*, 2018). Although the NRS often corresponds with the VAS scores (Hjermstad *et al.*, 2011), some research has shown the two scales are not interchangeable across all clinical populations (Hartrick, Kovan and Shapiro, 2003).

*Verbal Rating Scale (VRS)*

The VRS assesses pain using categorical descriptors ('no pain', 'mild pain', 'moderate pain', 'serious/severe pain', and 'worst possible pain'). While some work shows the VAS and VRS may not be interchangeable in spine surgery populations, others demonstrate they are correlated in an orthopaedic clinical setting (Briggs and Closs, 1999; Matamalas *et al.*, 2010).

*Short-Form McGill Pain Questionnaire (SF-MPQ)*

The SF-MPQ is a widely used multidimensional measure of pain experience with good reliability and validity across clinical pain populations (Melzack, 1987). It consists of 15 pain descriptors capturing sensory (11 items) and affective (4 items) dimensions of pain, each rated on a four-point intensity scale ranging from none to severe. Ratings can be summarized into sensory and affective subscale scores, as well as a total pain score reflecting overall pain quality.

**Quality of life**

Health-related quality of life was assessed using the EQ-5D-5L questionnaire (Janssen *et al.*, 2013) at the preoperative visit and at 3-month follow-up. Inclusion of quality of life measure aligns with IMMPACT recommendations, which emphasise outcomes in chronic pain research should extend beyond intensity and consider broader aspects of recovery (Dworkin *et al.*, 2005).

**2.4 Sample Size Justification**

Sample size considerations were informed by the prior study of Millard et al (2022), who reported a strong association between preoperative PAF and postoperative pain intensity in a cohort of 16 thoracicsurgery patients ($p < 0.02$). Although the exploratory nature and feasibility constraints of the present study limit the precision of a formal power calculation, the substantial effect size reported in Millard et al (2022) suggests that a sample of similar magnitude would be sufficient to detect an effect of comparable strength. We therefore set a minimum target enrolment of 16 patients, acknowledging that this sample size is intended to detect large effects and to provide preliminary evidence rather than definitive inferential conclusions.

**2.5 Statistical Analysis**

**Correlation analyses**

Given evidence that 3-month patient-reported outcomes are the strongest and most consistent predictors of 12 month pain, disability, and satisfaction in spine surgery (Rundell *et al.*, 2020), the change from preoperative assessment to 3-month follow-up is considered clinically most meaningful. Our primary correlation analyses therefore examine pain change across this interval. While pain ratings are widely considered to have ordinal measurement properties, the VAS is sometimes considered interval (Begum and Hossain, 2019), however methodological work indicated the VAS values should be interpreted as ordinal (Heller, Manuguerra and Chow, 2016). Similarly, the NRS is frequently treated as interval in

clinical research, although evidence shows that the numerical increments are not consistent across the scale (Hjermstad *et al.*, 2011). The VRS provides categorical information only.

Because the patient-reported pain measures (BPI, VAS, NRS, VRS) are ordinal or commonly treated as approximately interval scaled, we prespecified Spearman's rank correlations ($\rho$) as the primary tests for all PAF-pain associations. Pearson's r was performed only as a sensitivity analysis and is reported in the Supplementary Materials. This approach mitigates distributional assumptions and outlier sensitivity in our small sample while allowing comparison with prior clinical literature that sometimes treats VAS/NRS as interval.

Direction vs magnitude of change. We examined both directional $\Delta$PAF (signed change between visits; e.g., between pre-op and discharge, and pre-op and follow-up) and the magnitude of change in PAF (absolute $|\Delta PAF|$). Directional $\Delta$PAF analyses were conducted with Spearman's $\rho$ (primary) and Pearson's r (sensitivity), whereas magnitude ($|\Delta PAF|$) was analyzed analogously and flagged as exploratory. Full specifications are detailed in the Supplementary Methods

**Receiver Operating Characteristic (ROC) analyses**

To examine the potential predictive value of PAF, ROC analyses were performed to determine the ability of preoperative PAF to distinguish treatment responders vs non-responders. Responders were defined based on change in pain outcomes between the preoperative visit and the 3-month follow-up, reflecting the primary interval of interest for postoperative pain recovery. Analyses were performed using both a 30% improvement in pain and a 50% improvement as measured using the BPI-Worst, VAS, and NRS. These thresholds are based on IMMPACT consensus recommendations, clinically meaningful improvement in chronic pain conditions refers to $\geq$50% reduction in pain intensity as a "substantial" improvement, and a $\geq$30% as "moderately important" improvement (Dworkin *et al.*, 2008). Area under the curve (AUC) values with 95% confidence intervals were computed, with AUC $\geq$ 0.70 interpreted as fair discrimination and AUC $\geq$ 0.80 as good discrimination (Carter *et al.*, 2016).

**Quality of life**

EQ-5D-5L responses were converted into a single index value using the German valuation set, following standard scoring procedures for deriving health-related quality-of-life utilities. Changes in EQ-5D-5L from V1 to V4 were examined using non-parametric correlations with preoperative V1 PAF.

## 3. Results

### Patient characteristics

Between October 2023 and February 2025 seventeen adults undergoing spinal fusion were included in the study (mean age 59.5 ± 10.9, range: 37-75 years). The cohort comprised 10 females (60.4 ± 10.1, range: 44-73 years) and 7 males (58.3 ± 12.7, 37-75 years; Table 1). Fifteen participants underwent lumbar fusion, while two underwent cervical fusion. Of the cervical fusion cases, one participant was male and one was female.

Five patients had previously undergone decompression of the affected spinal segments; no patient had a history of fusion surgery. Eleven patients underwent single-level fusion, whereas the remaining patients underwent multilevel fusion. All surgical procedures were uneventful, and postoperative CT imaging demonstrated correct implant positioning.

**Table 1**: Mean ± standard deviations of baseline characteristics.

| Demographics | n | Mean (SD) | Min-Max |
|---|---|---|---|
| **Age** | 17 | 59.5 (10.9) | 37-75 |
| **Sex** | 17 | Male = 7 (41.2%) Female = 10 (58.8%) | - |

### Change in pain ratings from preoperative visit to three-month follow-up

Pain ratings showed a clear and consistent reduction from the preoperative visit to the 3-month follow-up across all measures. Scores on the BPI (Worst, Least, Average, and Current pain) decreased by approximately **63-71%**, and similar improvements were observed on both the VAS (−61%) and NRS (−63%). By 3-month follow-up, several participants reported no pain, as reflected by minimum values of 0 across multiple scales. These reductions were also evident between patient discharge and 3 months, indicating continued improvement over the postoperative recovery period (Figure 2). Overall, the data demonstrate a substantial decline in self-reported pain levels by 3 months after surgery.

**Figure 2:** Pain trajectories using VAS across the 4 visits

*Note. Individual pain scores as measured using the VAS across time points, with black line to indicate mean.*

Pain scores consistently decreased from pre-op to discharge and pre-op to 3-month follow-up on the Brief Pain Inventory (BPI) subscales of worst, least, and average pain over the past 24 hours, and current pain (Table 2). On the BPI Worst subscale, mean pain scores reduced by 18 points by discharge, and by 36.7 points by 3-month follow-up. On average, pain scores improved from pre-op to follow-up on the Visual Analogue Scale (VAS), Numerical Rating Scale (NRS), and Verbal Rating Scale (VRS) (Tables 2-4). Mean baseline VAS was 63/100 and NRS 6.5/10 (Table 3). Mean reductions were 17.5 (VAS) and 1.6 (NRS) points by discharge, and 32.3 (VAS) and 3.6 (NRS) points by follow-up. On the VRS, most patients showed improvement by follow-up, with 8/15 improving, 5 unchanged, and none worsening (Table 4). For analysis and visualisation, VRS was coded numerically (0 = No pain, 1 = Mild, 2 = Moderate, 3 = Severe, 4 = Worst).

All pain scores were strongly intercorrelated at pre-op ($\rho$ range = 0.62 - 0.98) indicating a substantial overlap across scales. At follow-up, intercorrelations were more heterogeneous ($\rho$ range = 0.16 - 0.99). This reduction was primarily driven by weak associations involving the SF-MPQ affective-subscale. When this subscale was excluded, correlations at follow-up remained moderate to strong ($\rho$ range = 0.52-0.99), with BPI-Least showing the weakest association with other scales (See Supplementary Tables S4-S5). This pattern is not unexpected, given BPI-Least captures the minimum experience pain rather than typical or peak intensity. A larger separation between BPI-Least and other pain scales likely reflects greater variability in pain across time, whereas a smaller separation reflects more constant pain.

**Table 2:** Mean ± standard deviations on BPI at 4 time points

| BPI Scale | Pre-op | | Discharge | | Follow-up | |
|---|---|---|---|---|---|---|
| | n | Mean (SD) | n | Mean (SD) | n | Mean (SD) |
| Worst | 17 | 62.4 (31.9) | 15 | 43.3 (32.0) | 15 | 22.7 (31.0) |
| Least | 17 | 38.8 (32.2) | 15 | 30.7 (30.6) | 15 | 11.3 (17.7) |
| Average | 17 | 50.0 (28.1) | 15 | 38.7 (30.7) | 15 | 16.7 (23.2) |
| Current | 16 | 45.0 (30.6) | 15 | 36.0 (29.0) | 15 | 16.7 (24.7) |

**Table 3:** Mean ± standard deviations on VAS and NRS at 4 time points

| Pain Measure | Pre-op | | 24h Post-op | | Discharge | | Follow-up | |
|---|---|---|---|---|---|---|---|---|
| | n | Mean (SD) | n | Mean (SD) | n | Mean (SD) | n | Mean (SD) |
| VAS | 15 | 63.3 (18.6 ) | 16 | 70.0 (31.0) | 15 | 43.9 (22.5) | 15 | 25.0 (27.5) |
| NRS | 15 | 6.5 (1.7) | 16 | 6.9 (2.9) | 15 | 4.7 (2.2) | 15 | 2.4 (2.7) |

**Table 4:** Mean ± standard deviations on VRS at 3 time points

| VRS Timepoint | n | Mode | Improved | No Change | Worsened |
|---|---|---|---|---|---|
| Pre-op | 15 | Moderate (46.7%) | - | - | - |
| 24h Post-op | 16 | Moderate (31.2%) | n = 2 | n = 8 | n = 4 |
| Discharge | 15 | Moderate (60%) | n = 6 | n = 7 | n = 0 |
| Follow-up | 15 | No Pain (40%) | n = 8 | n = 5 | n = 0 |

### 3.1 Relationship between preoperative PAF and long-term change in pain ratings

By the 3-month follow-up, higher preoperative PAF was significantly associated with greater reductions in pain on several BPI subscales. Specifically, significant correlations were observed for BPI-Worst ($\rho = 0.67$, $p = 0.017$) and BPI-Average ($\rho = 0.62$, $p = 0.033$). BPI-Current showed a similar pattern, with a nonsignificant trend ($\rho = 0.49$, $p = 0.122$). BPI-Least demonstrated the weakest and nonsignificant trend ($\rho = 0.28$, $p = 0.410$). Sensitivity analyses using Pearson correlations supported the overall pattern, though not all effects reached significance, likely due to the modest sample size (see Supplementary Materials).

Because patients reporting no pain prior to surgery could not show measurable improvement, they were excluded from the ΔBPI analyses. Three participants met this criterion, each reporting zero pain at baseline and at all subsequent visits, indicating a stable absence of pain rather than nonresponse. Analyses including these participants are provided in Supplementary Results Section 2.4 (Table S13).

For the VAS, NRS, and VRS, higher preoperative PAF tended to track with greater pain reduction at 3-month follow-up across all three scales. Associations were strongest for VAS ($\rho$ = 0.54, p = 0.055) and were consistent but below the threshold for significance for NRS ($\rho$ = 0.48, p = 0.096). Baseline PAF was not associated with change in VRS score ($\rho$ = 0.42, p = 0.153).

**Figure 3**: Associations between preoperative PAF and change in pain from preoperative visit to 3-month follow-up

*Note. Scatterplots show the relationship between preoperative PAF (x-axis) and change in pain from Pre-op to Follow-up on the BPI-Worst, BPI-Average, NRS, VAS, and VRS (left to right). * indicates a statistically significant association (p < 0.05).*

Preoperative PAF did not predict change on the SF-MPQ. Across total, sensory, and affective scores, correlations for both pre-op to discharge and pre-op to follow-up were small and nonsignificant ($\rho$ = 0.190 to 0.386; p = 0.155 to 0.498). Descriptive trajectories (pre-op, discharge, and follow-up) and full correlation outputs are provided in Supplementary Table S9 (descriptives) and Supplementary Table S10 (correlations).

**3.2 Relationship between preoperative PAF and short-term change in pain ratings**

In contrast, only BPI-Least showed a significant relationship from pre-op to discharge ($\rho$ = 0.62, p = 0.041). All correlations for BPI-Worst, BPI-Average, and BPI-Current were small and nonsignificant during this interval, indicating that PAF was not related to short-term changes in pain ($\rho$ = -0.12, p = 0.709; $\rho$ = -0.11, p = 0.739; $\rho$ = 0.02, p = 0.946, respectively). Correlations were also weak and nonsignificant across all measures; VAS ($\rho$ = 0.17, p = 0.579), NRS ($\rho$ = 0.07, p = 0.828), and VRS ($\rho$ = 0.25, p = 0.415).

Exploratory correlations were conducted for BPI-Impairment and perceived analgesic relief. A significant negative association was observed between baseline PAF and change in BPI-Impairment from pre-op to discharge ($\rho$ = -0.79, p = 0.006). The association was not present at follow-up ($\rho$ = 0.16, p = 0.619), indicating that this relationship was transient and specific to the early recovery phase. For perceived analgesic relief, all participants were retained in the analysis, as a baseline value of zero reflects absence of pain relief rather than absence of pain, and therefore represents a meaningful clinical

state from which improvement can occur. No relationship between preoperative PAF and change in analgesic relief was observed at discharge ($\rho$ = -0.46, p = 0.114) or follow-up ($\rho$ = 0.09, p = 0.773).

**3.3 Sensitivity and specificity of preoperative alpha in predicting outcome category: responder vs non-responder**

Prior to conducting the ROC analyses, we examined group-averaged preoperative power spectra for participants classified as responders and non-responders on the BPI-Worst subscale ($\geq$50% improvement from pre-op to follow-up). This subscale was selected because it showed the strongest association with $\Delta$Pain in earlier analyses. Visual inspection revealed a clear rightward shift of the alpha peak in responders, indicating a faster dominant alpha rhythm relative to non-responders (Figure 4). Consistent with this, the mean preoperative PAF was higher in responders (10.436 Hz) than in non-responders (9.738 Hz), suggesting meaningful spectral separation between outcome groups. These qualitative observations provide context for the formal classification results reported below.

For ROC analyses, 12 patients had usable BPI-Worst data. Participants who reported zero baseline pain were excluded from percentage-based responder analysis to avoid misclassifying individuals with persistently zero pain as non-responders when no improvement was possible.

Using a $\geq$50% improvement threshold, preoperative PAF demonstrated good discrimination of responders on the BPI-Worst subscale (AUC = 0.84; 95% CI: 0.607-1.00). The Youden optimal cut-off was 10.11 Hz, yielding a sensitivity of 0.75 and a specificity of 1.00 (Figure 5).

Thirteen patients had complete data for VAS analyses. Preoperative PAF also showed good discriminative performance on the VAS (AUC = 0.81; 95% CI: 0.561-1.00), again with high specificity (1.00) and moderate sensitivity (0.571). The optimal cut-off for VAS was 10.37 Hz. Notably, all individuals with preoperative PAF values above this threshold exhibited clinically meaningful improvement on both BPI-Worst and VAS, resulting in zero false-positive classifications.

Applying a more liberal $\geq$30% improvement threshold did not change responder classification for either BPI-Worst or VAS, and therefore produced identical ROC metrics. The NRS showed nearly identical results to the VAS; full NRS ROC outputs are presented in the Supplementary Material

**Figure 4**: Preoperative Spectra: Responders vs Non-Responders

*Note. EEG spectra of central channels at preoperative visit averaged over two groups; participants who showed above 50% improvement in pain scores on the BPI-Worst (responders, blue) and patients who reported below 50% improvement on the BPI-Worst (non-responders, green).*

**Figure 5:** ROC curve for responders vs non-responders using the BPI-Worst and VAS

*Note. ROC curve for preoperative PAF predicting 50% improvement from pre-op to follow-up on BPI-Worst and VAS.*

To determine whether preoperative PAF discriminated responders from non-responders above chance (AUC = 0.5), we evaluated the statistical significance of each ROC curve. For the BPI-Worst scale, discrimination approached, but did not reach conventional significance (p ≈ .07), despite a relatively high AUC (0.84; 95% CI: 0.61-1.00). The VAS showed similar borderline evidence (p ≈ .07-.08),

whereas no significant discrimination was observed for the NRS (p = .27). These results suggest potentially meaningful predictive value but also reflect limited power due to sample size and binary outcome classification. Full statistical procedures and supporting analyses are provided in the Supplementary Methods.

**3.4 Association between change in PAF and improvements in pain**

Directional change in PAF ($\Delta$PAF) from pre-op→discharge and pre-op→follow-up was not associated with change in pain on the VAS, NRS, or BPI-Worst (all ρ's non-significant). Cross-visit analyses ($\Delta$PAF pre-op→discharge vs $\Delta$Pain pre-op→follow-up was likewise null. Exploratorily, the magnitude of change (|$\Delta$PAF|) showed test-dependent patterns at pre-op→follow-up (Pearson significant for VAS/NRS and BPI-Average; Spearman non-significant), and should be interpreted cautiously. Full statistics are reported in Supplementary Tables S15-S16 (directional $\Delta$PAF) and Supplementary Tables S17-S18 (|$\Delta$PAF|), with scatterplots in Supplementary Figure S7.

**3.5 Relationship between PAF and concurrent pain ratings**

PAF did not correlate with concurrent pain at any visit (pre-op, discharge, follow-up) on VAS, NRS, VRS, or BPI subscales. Complete visit-wise correlation matrices are provided in Supplementary Table S19.

**3.6 Change in quality of life**

Quality of life scores (EQ-5D-5L index and EQ-VAS) generally improved from pre-op to follow-up. Preoperative PAF showed no association with change in EQ-5D-5L index and a non-significant trend with change in EQ-VAS. Participant-level trajectories and statistics are in Supplementary Table S20.

**3.7 Hospital Anxiety and Depression Questionnaire**

HADS-Anxiety and HADS-Depression decreased from pre-op to follow-up. Preoperative PAF showed a trend level association with change in anxiety (non-significant) and no association with change in depression. Full results appear in Supplementary Table S21.

**3.8 Opioid medication**

Opioid medication was not analysed for outcome associations; individual PAF values and qualitative medication changes per visit are summarized in Supplementary Table S22.

## 4. Discussion

This prospective observational study examined whether peak alpha frequency (PAF) from brief, pre-operative resting-state EEG can serve as a predictive (trait-like) biomarker of longer-term pain improvement after cervical or lumbar spinal fusion, and whether changes in PAF over time track changes in pain (dynamic marker). We observed that patients with faster baseline PAF were more likely to show meaningful pain reduction by 3 months, a pattern that was consistent across standard pain instruments. In contrast, PAF did not relate to early postoperative pain nor to multidimensional pain quality measures, and within-person changes in PAF did not reliably mirror changes in pain over time. Together, these findings position PAF as a stable, pre-surgical signal that helps characterise longer-term pain improvement rather than a dynamic biomarker for short-term pain or complex pain qualities.

**Long-term pain reduction is preferentially predicted by preoperative PAF**

In support of the primary hypothesis, preoperative PAF was correlated with a reduction in pain at the 3-month follow-up. Change in pain was calculated by subtracting follow-up pain scores from preoperative pain scores; higher positive ΔPain scores indicated greater pain reduction. Faster preoperative PAF was positively associated with greater reductions in worst and average pain, particularly on the BPI Worst, aligning with previous work showing that PAF relates most strongly to peak pain intensity (Millard et al, 2022). ROC analyses support the predictive value of preoperative PAF. Using a responder definition based on ≥50% reduction in pain, faster baseline PAF demonstrated high specificity for identifying responders, indicating that individuals with a faster preoperative PAF were unlikely to be misclassified as responders when they did not in fact improve. Taken together, these findings suggest that PAF may serve as a probabilistic neurophysiological marker; individuals with faster PAF are more likely to show clinically meaningful pain improvement, though PAF alone does not deterministically predict postoperative outcomes.

We found that preoperative PAF did not relate to short-term postoperative pain changes (pre-op to discharge) on any measure with the exception of BPI-Least, indicating that baseline PAF is not strongly linked to early pain trajectories. Pain scores at discharge reflect the early recovery period, in which acute surgical injury, analgesic use, and inflammatory processes may dominate pain intensity, possibly influencing change in peak and average pain. In contrast, BPI-Least may be capturing the capacity to achieve periods of relative pain relief, which could be more sensitive to PAF as a trait-like difference in cortical inhibitory processes. As the acute postoperative recovery period subsides, these trait neurophysiological differences become broader across pain measures, as seen by the associations with change in pain at follow-up across various pain measures including peak pain captured by BPI-Worst.

A significant negative association was observed between baseline PAF and change in BPI-Impairment from the preoperative visit to discharge. This association was evident both when participants with zero

baseline BPI-impairment were retained and when they were excluded, with larger effect sizes observed following exclusion. The association was not present at 3-month follow-up, indicating that this relationship was transient and specific to the early recovery phase. The direction of this effect contrasts with the positive association observed between baseline PAF and changes in pain intensity measures, suggesting that PAF may relate differently to early functional interference than to pain perception itself. One possible explanation is that individuals with faster baseline PAF may resume activity earlier in the postoperative period, resulting in greater perceived functional interference despite reductions in pain. Alternatively, this finding may reflect differing psychological or behavioural components captured by the BPI-impairment scale.

**Weak and exploratory associations between PAF shifts and pain outcomes**

We did not find evidence that ΔPAF tracked ΔPain scores across visits. However, when examining absolute change from pre-op to follow-up, we found a weak association between |ΔPAF| and ΔPain using Pearson's correlations. Although exploratory and dependent on the statistical test, this raises the possibility that individuals whose PAF shifts are more malleable over time may show greater improvements in pain.

PAF variability relating to change in pain is conceptually compatible with work suggesting different alpha generators may relate to pain sensitivity and coping strategies (Furman *et al.*, 2024). Furman and colleagues propose that while the slower alpha generator may relate to heightened pain sensitivity (Nir *et al.*, 2012), the faster oscillator relates to a better tolerance for pain (Furman *et al.*, 2020) and that attenuation of the faster generator relates to efficient pain management. It is plausible that a "flexible" PAF may reflect a form of neural adaptability that supports better treatment response.

ΔPAF and |ΔPAF| from pre-op to discharge did not correlate with ΔPain from pre-op to follow-up, suggesting that the change in PAF from preoperative to postoperative cannot yet be considered a reliable marker for tracking postoperative recovery over a longer-term timescale. The exploratory trend suggests that further work is warranted to examine the utility of dynamic changes in PAF in understanding individual variability in pain outcomes.

**Individual variability in pain trajectories**

In the current study, we observed substantial variability in postoperative pain trajectories. Pain scores generally trended down from preoperative to postoperative visits, although not all participants followed the same pattern. These findings were expected given that not all patients benefit from spine surgery (Trief, Grant and Fredrickson, 2000). Most participants showed some improvement, but the amount of change differed depending on the pain scale. On the VAS, two participants had higher pain scores at 3-month follow-up compared to pre-op, whereas on the NRS only one of these showed an increase. These findings therefore support the view that the VAS and NRS are not perfectly interchangeable and support the use of multiple tools to assess pain (Hartrick, Kovan and Shapiro, 2003). On the VRS, nobody showed an increase, but six of fifteen showed no change. Only one person reported an increase in pain

on the BPI-Worst. The inter-individual variability in pain reduction trajectories directly motivates the investigation of why comparable surgical profiles can result in divergent pain outcomes.

Individual differences in PAF and PAF trajectories throughout the visits were also observed. Some participants displayed a reduction in PAF at the time of discharge, but their PAF returned to its baseline frequency at 3 months. The change at discharge may reflect increased postoperative pain. Although some evidence suggests PAF increases in acute experimental pain (Nir *et al.*, 2012), the present findings align with work demonstrating PAF slows in response to newly introduced sustained pain, which may better characterise the postoperative period (Furman *et al.*, 2024). PAF returning to a similar baseline for almost all participants supports the view that PAF represents a stable neurophysiological characteristic (Grandy *et al.*, 2013).

**Possible mechanisms underlying the PAF-pain relationship**

Several mechanisms may account for why higher preoperative PAF, and to a lesser extent long-term changes in PAF relate to postoperative pain improvement. First, the associations were observed at central electrodes, consistent with the role of alpha oscillations in regulating inhibition within sensorimotor networks. Alpha rhythms could suppress somatosensory and motor representations irrelevant to current goal-related networks (Jensen and Mazaheri, 2010), and slower PAF has been linked to reduced inhibitory flexibility and heightened pain sensitivity (Brumagne *et al.*, 2019). From this perspective, individuals with slower PAF may have fewer neurophysiological resources to downregulate pain signals once tissue damage has been addressed, potentially hindering recovery.

A second interpretation centres on thalamocortical function. Slower PAF has been proposed to reflect disrupted thalamic relay or gating processes, which may contribute to amplification and maintenance of pain (Walton, Dubois and Llinás, 2010; Karcz *et al.*, 2024). Individuals with faster baseline PAF may therefore exhibit more efficient thalamocortical processing, making them more likely to benefit from surgical correction of the peripheral pain source. Conversely, slower PAF could indicate greater central dysregulation, reducing the degree to which structural intervention translates into subjective pain relief.

This framework may also explain the exploratory observation that greater long-term modifiability of PAF relates weakly to improved pain outcomes. A more "flexible" PAF over the postoperative period may reflect thalamocortical circuits that adapt more effectively during recovery, aligning with accounts that different alpha generators relate to pain sensitivity versus coping capacities (Furman *et al.*, 2024). However, given the modest and statistically fragile effects, this interpretation remains tentative.

Finally, PAF may index broader cognitive-affective traits rather than pain-specific physiology. Previous work has linked PAF to anxiety, arousal, and attentional control (Knyazev, Savostyanov and Levin, 2004; Barry *et al.*, 2020; Zhang *et al.*, 2021); factors known to influence postoperative trajectories (Adogwa *et al.*, 2016). Under this view, slower PAF may reflect a cognitive-affective profile that confers vulnerability to poorer recovery, whereas faster PAF may indicate greater resilience.

Collectively, these accounts suggest that PAF may capture a neurophysiological or cognitive profile that shapes an individual's likelihood of benefiting from surgery, though the underlying mechanisms remain to be clarified.

**Summary and implications**

This study provides preliminary evidence that PAF may capture individual differences in central pain processing that relate to postoperative improvement following spinal fusion surgery. In this cohort, faster preoperative PAF was probabilistically associated with greater long-term reductions in pain, supporting that PAF may index vulnerability to centrally maintained pain rather than reflecting pain intensity.

An important strength of PAF as a candidate biomarker is its clinical feasibility. We have demonstrated that PAF can be acquired rapidly and with minimal burden to patients, which makes it ideal for clinical translation. In addition, brain rhythms can be modulated. For example, neurofeedback, audio-visual stimulation, and non-invasive brain stimulation have been shown to influence oscillations, including alpha (Helfrich *et al.*, 2014; Notbohm, Kurths and Herrmann, 2016; Vučković *et al.*, 2019). PAF may therefore not only represent a predictive marker, but also a target for future treatments aiming to improve postoperative outcomes.

**Limitations and future directions**

The sample size was modest, and not all statistical associations reached significance. Future studies with larger sample sizes are necessary to confirm whether PAF reliably differentiates patients who will see meaningful improvements in pain scores from those who do not. One methodological consideration is that the VAS and NRS are sometimes considered ordinal, so a larger dataset would allow for continuous ordinal regression (Heller, Manuguerra and Chow, 2016). These modelling approaches would lead to a more rigorous assessment of the predictive value of PAF. In addition, preoperative pain may influence postoperative recovery (Wilder-Smith, 2007) and should be taken into consideration in future larger studies. PAF should not be viewed as an independent indicator, but is best thought of as a potential informative component within a multimodal framework that integrates clinical, psychological, and neurophysiological measures to improve prediction of treatment response (Hodges *et al.*, 2025). Taken together, the present findings offer an encouraging proof-of-concept that PAF may serve as a meaningful neural marker of postoperative pain recovery after spinal fusion surgery.

**Acknowledgements**

Ali Mazaheri is supported by funding from the Wellcome Leap Untangling Addiction Program.

References

Abbott, A.D., Tyni-Lenné, R. and Hedlund, R. (2010) 'The influence of psychological factors on pre-operative levels of pain intensity, disability and health-related quality of life in lumbar spinal fusion surgery patients', *Physiotherapy*, 96(3), pp. 213–221. Available at: https://doi.org/10.1016/j.physio.2009.11.013.

Adogwa, O. *et al.* (2016) 'Pretreatment of Anxiety Before Cervical Spine Surgery Improves Clinical Outcomes: A Prospective, Single-Institution Experience', *World Neurosurgery*, 88, pp. 625–630. Available at: https://doi.org/10.1016/j.wneu.2015.11.014.

Ahmad, B. and Barkana, B.D. (2025) 'Pain and the Brain: A Systematic Review of Methods, EEG Biomarkers, Limitations, and Future Directions', *Neurology International*, 17(4), p. 46. Available at: https://doi.org/10.3390/neurolint17040046.

Barry, R.J. *et al.* (2020) 'Natural alpha frequency components in resting EEG and their relation to arousal', *Clinical Neurophysiology*, 131(1), pp. 205–212. Available at: https://doi.org/10.1016/j.clinph.2019.10.018.

Begum, M.R. and Hossain, M.A. (2019) 'Validity and reliability of visual analogue Scale (vas) for pain measurement', *Journal of Medical Case Reports and Reviews*, 2(11). Available at: https://jmcrr.info/index.php/jmcrr/article/view/44 (Accessed: 10 December 2025).

Briggs, M. and Closs, J.S. (1999) 'A Descriptive Study of the Use of Visual Analogue Scales and Verbal Rating Scales for the Assessment of Postoperative Pain in Orthopedic Patients', *Journal of Pain and Symptom Management*, 18(6), pp. 438–446. Available at: https://doi.org/10.1016/S0885-3924(99)00092-5.

Brumagne, S. *et al.* (2019) 'Neuroplasticity of Sensorimotor Control in Low Back Pain', *Journal of Orthopaedic & Sports Physical Therapy*, 49(6), pp. 402–414. Available at: https://doi.org/10.2519/jospt.2019.8489.

Carter, J.V. *et al.* (2016) 'ROC-ing along: Evaluation and interpretation of receiver operating characteristic curves', *Surgery*, 159(6), pp. 1638–1645. Available at: https://doi.org/10.1016/j.surg.2015.12.029.

Celestin, J., Edwards, R.R. and Jamison, R.N. (2009) 'Pretreatment Psychosocial Variables as Predictors of Outcomes Following Lumbar Surgery and Spinal Cord Stimulation: A Systematic Review and Literature Synthesis', *Pain Medicine*, 10(4), pp. 639–653. Available at: https://doi.org/10.1111/j.1526-4637.2009.00632.x.

Chou, R. *et al.* (2007) 'Diagnosis and Treatment of Low Back Pain: A Joint Clinical Practice Guideline from the American College of Physicians and the American Pain Society', *Annals of Internal Medicine*, Volume 147(Number 7), pp. 441–524.

Cleeland, C.S. and Ryan, K.M. (1994) 'Pain assessment: global use of the Brief Pain Inventory', *Annals of the Academy of Medicine, Singapore*, 23(2), pp. 129–138.

Cooper, M.E. *et al.* (2018) 'Heterogeneity of Reporting Outcomes in the Spine Surgery Literature', *Clinical Spine Surgery*, 31(4), p. E221. Available at: https://doi.org/10.1097/BSD.0000000000000578.

Copay, A.G. *et al.* (2008) 'Minimum clinically important difference in lumbar spine surgery patients: a choice of methods using the Oswestry Disability Index, Medical Outcomes Study questionnaire Short Form 36, and Pain Scales', *The Spine Journal*, 8(6), pp. 968–974. Available at: https://doi.org/10.1016/j.spinee.2007.11.006.

Delorme, A. and Makeig, S. (2004) 'EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis', *Journal of Neuroscience Methods*, 134(1), pp. 9–21. Available at: https://doi.org/10.1016/j.jneumeth.2003.10.009.

Demyttenaere, K. *et al.* (2007) 'Mental disorders among persons with chronic back or neck pain: Results from the world mental health surveys', *PAIN*, 129(3), pp. 332–342. Available at: https://doi.org/10.1016/j.pain.2007.01.022.

Driscoll, T. *et al.* (2014) 'The global burden of occupationally related low back pain: estimates from the Global Burden of Disease 2010 study', *Annals of the Rheumatic Diseases*, 73(6), pp. 975–981. Available at: https://doi.org/10.1136/annrheumdis-2013-204631.

Dworkin, R.H. *et al.* (2005) 'Core outcome measures for chronic pain clinical trials: IMMPACT recommendations', *Pain*, 113(1), pp. 9–19. Available at: https://doi.org/10.1016/j.pain.2004.09.012.

Dworkin, R.H. *et al.* (2008) 'Interpreting the Clinical Importance of Treatment Outcomes in Chronic Pain Clinical Trials: IMMPACT Recommendations', *The Journal of Pain*, 9(2), pp. 105–121. Available at: https://doi.org/10.1016/j.jpain.2007.09.005.

Ehrlich, G.E. (2003) 'Low back pain', *Bulletin of the World Health Organization*, 81(9), pp. 671–676.

Finkelstein, J.A. and Schwartz, C.E. (2019) 'Patient-reported outcomes in spine surgery: past, current, and future directions: JNSPG 75th Anniversary Invited Review Article', *Journal of Neurosurgery: Spine*, 31(2), pp. 155–164. Available at: https://doi.org/10.3171/2019.1.SPINE18770.

Furman, A.J. *et al.* (2018) 'Cerebral peak alpha frequency predicts individual differences in pain sensitivity', *NeuroImage*, 167, pp. 203–210. Available at: https://doi.org/10.1016/j.neuroimage.2017.11.042.

Furman, A.J. *et al.* (2020) 'Sensorimotor Peak Alpha Frequency Is a Reliable Biomarker of Prolonged Pain Sensitivity', *Cerebral Cortex*, 30(12), pp. 6069–6082. Available at: https://doi.org/10.1093/cercor/bhaa124.

Furman, A.J. *et al.* (2024) 'Prolonged Pain Reliably Slows Peak Alpha Frequency by Reducing Fast Alpha Power', *eLife*, 13. Available at: https://doi.org/10.7554/eLife.102096.1.

Grandy, T.H. *et al.* (2013) 'Peak individual alpha frequency qualifies as a stable neurophysiological trait marker in healthy younger and older adults', *Psychophysiology*, 50(6), pp. 570–582. Available at: https://doi.org/10.1111/psyp.12043.

Hartrick, C.T., Kovan, J.P. and Shapiro, S. (2003) 'The Numeric Rating Scale for Clinical Pain Measurement: A Ratio Measure?', *Pain Practice*, 3(4), pp. 310–316. Available at: https://doi.org/10.1111/j.1530-7085.2003.03034.x.

Helfrich, R.F. *et al.* (2014) 'Entrainment of Brain Oscillations by Transcranial Alternating Current Stimulation', *Current Biology*, 24(3), pp. 333–339. Available at: https://doi.org/10.1016/j.cub.2013.12.041.

Heller, G.Z., Manuguerra, M. and Chow, R. (2016) 'How to analyze the Visual Analogue Scale: Myths, truths and clinical relevance', *Scandinavian Journal of Pain*, 13(1), pp. 67–75. Available at: https://doi.org/10.1016/j.sjpain.2016.06.012.

Hjermstad, M.J. *et al.* (2011) 'Studies Comparing Numerical Rating Scales, Verbal Rating Scales, and Visual Analogue Scales for Assessment of Pain Intensity in Adults: A Systematic Literature Review', *Journal of Pain and Symptom Management*, 41(6), pp. 1073–1093. Available at: https://doi.org/10.1016/j.jpainsymman.2010.08.016.

Hodges, P.W. *et al.* (2025) 'Development and application of predictive clinical biomarkers for low back pain care: recommendations from the ISSLS phenotype/precision spine focus group', *European Spine Journal*, 34(4), pp. 1309–1318. Available at: https://doi.org/10.1007/s00586-025-08720-4.

Hurwitz, E.L. *et al.* (2018) 'The Global Spine Care Initiative: a summary of the global burden of low back and neck pain studies', *European Spine Journal*, 27(6), pp. 796–801. Available at: https://doi.org/10.1007/s00586-017-5432-9.

Huskisson, E.C. (1974) 'MEASUREMENT OF PAIN', *The Lancet*, 304(7889), pp. 1127–1131. Available at: https://doi.org/10.1016/S0140-6736(74)90884-8.

Janssen, M.F. *et al.* (2013) 'Measurement properties of the EQ-5D-5L compared to the EQ-5D-3L across eight patient groups: a multi-country study', *Quality of Life Research*, 22(7), pp. 1717–1727. Available at: https://doi.org/10.1007/s11136-012-0322-4.

Jensen, M.P., Karoly, P. and Braver, S. (1986) 'The measurement of clinical pain intensity: a comparison of six methods', *PAIN*, 27(1), p. 117. Available at: https://doi.org/10.1016/0304-3959(86)90228-9.

Jensen, O. and Mazaheri, A. (2010) 'Shaping Functional Architecture by Oscillatory Alpha Activity: Gating by Inhibition', *Frontiers in Human Neuroscience*, 4. Available at: https://www.frontiersin.org/articles/10.3389/fnhum.2010.00186 (Accessed: 17 November 2023).

Karcz, M. *et al.* (2024) 'Pathophysiology of Pain and Mechanisms of Neuromodulation: A Narrative Review (A Neuron Project)', *Journal of Pain Research*, 17, pp. 3757–3790. Available at: https://doi.org/10.2147/JPR.S475351.

Kasch, R. *et al.* (2022) 'Association of Lumbar MRI Findings with Current and Future Back Pain in a Population-based Cohort Study', *Spine*, 47(3), p. 201. Available at: https://doi.org/10.1097/BRS.0000000000004198.

Khor, S. *et al.* (2018) 'Development and Validation of a Prediction Model for Pain and Functional Outcomes After Lumbar Spine Surgery', *JAMA Surgery*, 153(7), pp. 634–642. Available at: https://doi.org/10.1001/jamasurg.2018.0072.

Kim, J.A. and Davis, K.D. (2021) 'Neural Oscillations: Understanding a Neural Code of Pain', *The Neuroscientist: A Review Journal Bringing Neurobiology, Neurology and Psychiatry*, 27(5), pp. 544–570. Available at: https://doi.org/10.1177/1073858420958629.

Kisler, L.B. *et al.* (2020) 'Abnormal alpha band power in the dynamic pain connectome is a marker of chronic pain with a neuropathic component', *NeuroImage. Clinical*, 26, p. 102241. Available at: https://doi.org/10.1016/j.nicl.2020.102241.

Klimesch, W., Sauseng, P. and Hanslmayr, S. (2007) 'EEG alpha oscillations: The inhibition–timing hypothesis', *Brain Research Reviews*, 53(1), pp. 63–88. Available at: https://doi.org/10.1016/j.brainresrev.2006.06.003.

Knyazev, G.G., Savostyanov, A.N. and Levin, E.A. (2004) 'Alpha oscillations as a correlate of trait anxiety', *International Journal of Psychophysiology*, 53(2), pp. 147–160. Available at: https://doi.org/10.1016/j.ijpsycho.2004.03.001.

Laumet, G. *et al.* (2019) 'T Cells as an Emerging Target for Chronic Pain Therapy', *Frontiers in Molecular Neuroscience*, 12. Available at: https://doi.org/10.3389/fnmol.2019.00216.

Liang, W. *et al.* (2024) 'Current advancements in therapeutic approaches in orthopedic surgery: a review of recent trends', *Frontiers in Bioengineering and Biotechnology*, 12. Available at: https://doi.org/10.3389/fbioe.2024.1328997.

Lim, M. *et al.* (2016) 'Increased Low- and High-Frequency Oscillatory Activity in the Prefrontal Cortex of Fibromyalgia Patients', *Frontiers in Human Neuroscience*, 10. Available at: https://doi.org/10.3389/fnhum.2016.00111.

Mason, A. *et al.* (2014) 'The accuracy of pedicle screw placement using intraoperative image guidance systems: A systematic review', *Journal of Neurosurgery: Spine*, 20(2), pp. 196–203. Available at: https://doi.org/10.3171/2013.11.SPINE13413.

Matamalas, A. *et al.* (2010) 'The Visual Analog Scale and a Five-Item Verbal Rating Scale Are Not Interchangeable for Back Pain Assessment in Lumbar Spine Disorders', *Spine*, 35(21), p. E1115. Available at: https://doi.org/10.1097/BRS.0b013e3181e7b315.

Mazaheri, A., Seminowicz, D.A. and Furman, A.J. (2022) 'Peak alpha frequency as a candidate biomarker of pain sensitivity: the importance of distinguishing slow from slowing', *NeuroImage*, 262, p. 119560. Available at: https://doi.org/10.1016/j.neuroimage.2022.119560.

McCormack, H.M., Horne, D.J. de L. and Sheather, S. (1988) 'Clinical applications of visual analogue scales: a critical review', *Psychological Medicine*, 18(4), pp. 1007–1019. Available at: https://doi.org/10.1017/S0033291700009934.

Melzack, R. (1987) 'The short-form McGill Pain Questionnaire', *Pain*, 30(2), pp. 191–197. Available at: https://doi.org/10.1016/0304-3959(87)91074-8.

Melzack, R. and Katz, J. (2004) 'The Gate Control Theory: Reaching for the Brain', *Pain*. Psychology Press.

Middendorp, M. *et al.* (2017) 'Association between intervertebral disc degeneration and the Oswestry Disability Index', *Journal of Back and Musculoskeletal Rehabilitation*, 30(4), pp. 819–823. Available at: https://doi.org/10.3233/BMR-150516.

Millard, S.K. *et al.* (2022) 'Predicting postoperative pain in lung cancer patients using preoperative peak alpha frequency', *British Journal of Anaesthesia*, 128(6), pp. e346–e348. Available at: https://doi.org/10.1016/j.bja.2022.03.006.

Newton, B.J. *et al.* (2013) 'A Narrative Review of the Impact of Disbelief in Chronic Pain', *Pain Management Nursing*, 14(3), pp. 161–171. Available at: https://doi.org/10.1016/j.pmn.2010.09.001.

Nir, R.-R. *et al.* (2012) 'Tonic pain and continuous EEG: Prediction of subjective pain perception by alpha-1 power during stimulation and at rest', *Clinical Neurophysiology*, 123(3), pp. 605–612. Available at: https://doi.org/10.1016/j.clinph.2011.08.006.

Notbohm, A., Kurths, J. and Herrmann, C.S. (2016) 'Modification of Brain Oscillations via Rhythmic Light Stimulation Provides Evidence for Entrainment but Not for Superposition of Event-Related Responses', *Frontiers in Human Neuroscience*, 10. Available at: https://doi.org/10.3389/fnhum.2016.00010.

Oostenveld, R. *et al.* (2011) 'FieldTrip: open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data', *Computational Intelligence and Neuroscience*, 2011, p. 1:1-1:9. Available at: https://doi.org/10.1155/2011/156869.

Phillips, F.M. *et al.* (2013) 'Lumbar Spine Fusion for Chronic Low Back Pain Due to Degenerative Disc Disease: A Systematic Review', *Spine*, 38(7), p. E409. Available at: https://doi.org/10.1097/BRS.0b013e3182877f11.

Ploner, M., Sorg, C. and Gross, J. (2017) 'Brain Rhythms of Pain', *Trends in Cognitive Sciences*, 21(2), pp. 100–110. Available at: https://doi.org/10.1016/j.tics.2016.12.001.

Popov, T. *et al.* (2023) 'Test–retest reliability of resting-state EEG in young and older adults', *Psychophysiology*, 60(7), p. e14268. Available at: https://doi.org/10.1111/psyp.14268.

Rannou, F. *et al.* (2007) 'High-sensitivity C-reactive protein in chronic low back pain with vertebral end-plate modic signal changes', *Arthritis Care & Research*, 57(7), pp. 1311–1315. Available at: https://doi.org/10.1002/art.22985.

Rathod, T.N. *et al.* (2014) 'High sensitive C-reactive protein-Effective tool in determining postoperative recovery in lumbar disc disease', *Indian Journal of Orthopaedics*, 48(4), pp. 354–359. Available at: https://doi.org/10.4103/0019-5413.136216.

Reisener, M.-J. *et al.* (2020) 'Trends in lumbar spinal fusion—a literature review', *Journal of Spine Surgery*, 6(4), pp. 752–761. Available at: https://doi.org/10.21037/jss-20-492.

Rundell, S.D. *et al.* (2020) 'Adding 3-month patient data improves prognostic models of 12-month disability, pain, and satisfaction after specific lumbar spine surgical procedures: development and validation of a prediction model', *The Spine Journal*, 20(4), pp. 600–613. Available at: https://doi.org/10.1016/j.spinee.2019.12.010.

Sarnthein, J. *et al.* (2006) 'Increased EEG power and slowed dominant frequency in patients with neurogenic pain', *Brain*, 129(1), pp. 55–64. Available at: https://doi.org/10.1093/brain/awh631.

Scheufler, K.-M., Dohmen, H. and Vougioukas, V.I. (2007) 'PERCUTANEOUS TRANSFORAMINAL LUMBAR INTERBODY FUSION FOR THE TREATMENT OF DEGENERATIVE LUMBAR INSTABILITY', *Operative Neurosurgery*, 60(4), p. 203. Available at: https://doi.org/10.1227/01.NEU.0000255388.03088.B7.

Seminowicz, D.A. *et al.* (2018) 'Slow peak alpha frequency and corticomotor depression linked to high pain susceptibility in transition to sustained pain'. bioRxiv, p. 278598. Available at: https://doi.org/10.1101/278598.

Shamji, M.F. *et al.* (2010) 'Proinflammatory cytokine expression profile in degenerated and herniated human intervertebral disc tissues', *Arthritis & Rheumatism*, 62(7), pp. 1974–1982. Available at: https://doi.org/10.1002/art.27444.

Tan, L.L., Oswald, M.J. and Kuner, R. (2021) 'Neurobiology of brain oscillations in acute and chronic pain', *Trends in Neurosciences*, 44(8), pp. 629–642. Available at: https://doi.org/10.1016/j.tins.2021.05.003.

Tonosu, J. *et al.* (2016) 'The relationship between findings on magnetic resonance imaging and previous history of low back pain', *Journal of Pain Research*, 10, pp. 47–52. Available at: https://doi.org/10.2147/JPR.S122380.

Trief, P.M., Grant, W. and Fredrickson, B. (2000) 'A Prospective Study of Psychological Predictors of Lumbar Surgery Outcome', *Spine*, 25(20), p. 2616.

Vo, N.V. *et al.* (2023) 'Toward the Identification of Distinct Phenotypes: Research Protocol for the Low Back Pain Biological, Biomechanical, and Behavioral (LB3P) Cohort Study and the BACPAC Mechanistic Research Center at the University of Pittsburgh', *Pain Medicine*, 24(Supplement_1), pp. S36–S47. Available at: https://doi.org/10.1093/pm/pnad009.

Vos, T. *et al.* (2016) 'Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015', *The Lancet*, 388(10053), pp. 1545–1602. Available at: https://doi.org/10.1016/S0140-6736(16)31678-6.

de Vries, M. *et al.* (2013) 'Altered resting state EEG in chronic pancreatitis patients: toward a marker for chronic pain', *Journal of Pain Research*, 6, pp. 815–824. Available at: https://doi.org/10.2147/JPR.S50919.

Vučković, A. *et al.* (2019) 'EEG Correlates of Self-Managed Neurofeedback Treatment of Central Neuropathic Pain in Chronic Spinal Cord Injury', *Frontiers in Neuroscience*, 13. Available at: https://doi.org/10.3389/fnins.2019.00762.

Walton, K.D., Dubois, M. and Llinás, R.R. (2010) 'Abnormal thalamocortical activity in patients with Complex Regional Pain Syndrome (CRPS) Type I', *PAIN*, 150(1), pp. 41–51. Available at: https://doi.org/10.1016/j.pain.2010.02.023.

Weber, C. *et al.* (2016) 'Is There an Association Between Radiological Severity of Lumbar Spinal Stenosis and Disability, Pain, or Surgical Outcome?: A Multicenter Observational Study', *Spine*, 41(2), p. E78. Available at: https://doi.org/10.1097/BRS.0000000000001166.

Weinstein, J.N. *et al.* (2008) 'Surgical versus Nonsurgical Therapy for Lumbar Spinal Stenosis', *New England Journal of Medicine*, 358(8), pp. 794–810. Available at: https://doi.org/10.1056/NEJMoa0707136.

Wilder-Smith, C.H. (2007) 'Predicting pain and pain responses to opioids', *European Journal of Pain Supplements*, 1(1), pp. 31–37. Available at: https://doi.org/10.1016/S1754-3207(08)60009-7.

Woolf, C.J. (2011) 'Central sensitization: Implications for the diagnosis and treatment of pain', *PAIN*, 152(3, Supplement), pp. S2–S15. Available at: https://doi.org/10.1016/j.pain.2010.09.030.

Wu, A. *et al.* (2020) 'Global low back pain prevalence and years lived with disability from 1990 to 2017: estimates from the Global Burden of Disease Study 2017', *Annals of Translational Medicine*, 8(6), p. 299. Available at: https://doi.org/10.21037/atm.2020.02.175.

Zhang, Y. *et al.* (2021) 'Relationship between individual alpha peak frequency and attentional performance in a multiple object tracking task among ice-hockey players', *PLOS ONE*, 16(5), p. e0251443. Available at: https://doi.org/10.1371/journal.pone.0251443.