Neuroinvasive Human Parvovirus 4 is associated with increased mortality in

children: a multiyear retrospective observational study

Deb Purna Keya (MSc)[1]*, Apurba Rajib Malaker (MSc)[1]*, Naito Kanon (MSc)[1], Arif

Mohammad Tanmoy (PhD)[1], Saffat Reaz (BSc)[1], Preonath Chondrow Dev (MSc)[1],

Hafizur Rahman (BSc)[1], Lubana Tanvia (MSc)[1], Anisur Rahman (MSc)[1], Afroza Akter

Tanni (MSc)[1], Dipu Chandra Das (MSc)[1], Anannya Barman Jui (MBBS)[1], Mirza Md.

Ziaul Islam (MD)[2], Reaz Mobarak (FRCP)[3], Shamsun Nahar (MBBS)[1], Cristina Tato

(PhD)[4], ASM Nawshad Uddin Ahmed (FCPS)[1], Farhad Imam (PhD)[5], Joseph L DeRisi

(PhD)[4,6#], Samir K Saha (PhD)[1#], Yogesh Hooda (PhD)[1#], Senjuti Saha (PhD)[1#$]

[1]Child Health Research Foundation, Dhaka, Bangladesh

[2]Bangladesh Institute of Child Health, Bangladesh Shishu Hospital & Institute, Dhaka,

Bangladesh

[3]Department of Epidemiology & Research, Bangladesh Shishu Hospital & Institute,

Dhaka, Bangladesh

[4]Chan Zuckerberg Biohub, San Francisco, California, USA

[5]Gates Foundation, Seattle, WA, USA

[6]Department of Biochemistry and Biophysics, University of California, San Francisco,

California, USA

*authors with equal contribution

[#]senior authors

24
[$]Correspondence to

Senjuti Saha, PhD, MPH

Deputy Executive Director

Child Health Research Foundation

Dhaka Bangladesh

Telephone: +880 1711646038

senjutisaha@chrfbd.org

Abstract

Background: Infections of the central nervous system (CNS) in children remain a major cause of mortality and long-term disability globally, particularly in low- and middle-income countries (LMICs), where a high proportion of cases lack an identified pathogen. Sporadically, human parvovirus 4 (PARV4) has been detected in a small number of cerebrospinal fluid (CSF) from children with CNS infections, but its pathogenic role is unclear. We investigated the prevalence, clinical impact, and genomic characteristics of PARV4 in children with suspected meningitis.

Methods: We retrospectively analyzed CSF samples collected from children with WHO-defined suspected meningitis at the largest pediatric hospital in Bangladesh between 2015-2022. All samples underwent routine diagnostics, including bacterial culture and serological testing. Additional testing for PARV4 and parvovirus B19 was performed using qPCR of samples with >9 white blood cell (WBC)/µl followed by metagenomic sequencing of a subset. Clinical and laboratory data were extracted from patient records. Associations between PARV4 detection and mortality were assessed using logistic regression, adjusting for age, WBC count, and co-infections. Genomic and phylogenetic analyses were conducted on PARV4-positive samples.

Findings: Among 2,793 CSF samples with >9 WBC/µl, 526 (18·8%) were PARV4-positive. The median age of PARV4-positive cases was lower than that of PARV4-negative cases (4 vs 7 months, $p<0·001$). Co-infections were more common among PARV4-positive cases (49·6%) than PARV4-negative cases (16·4%). PARV4 positivity was independently associated with increased in-hospital mortality (adjusted odds ratio 2·09, 95%CI:1·46–2·96; $p<0·001$). Phylogenetic analysis indicated most strains belonged to genotype 2, with two sequences forming a distinct clade.

Interpretation: PARV4 is frequently detected in the CSF of children with suspected meningitis and is associated with increased in-hospital mortality. Its high prevalence,

63 detection early in life, and frequent co-infection with other pathogens highlight the need

64 to investigate PARV4 as an emerging CNS pathogen in LMICs.

Introduction

Central nervous system (CNS) infections including meningitis remains associated with high morbidity and mortality worldwide.[1] Globally, an estimated 10 million meningitis cases and 213,962 deaths occur each year, with 70% of these cases concentrated in low- and middle-income countries (LMICs).[2,3] Survivors often experience long-term neurological sequelae, emphasizing the need for early diagnosis, pathogen-specific treatment, and prevention.[4] Many pathogens including bacteria, viruses, fungi and parasites are known to cause CNS infections. In LMICs, where disease burden is the highest, pathogen detection rates remain particularly low: up to 80–85% of pediatric CNS infection cases may remain unexplained.[5,6] This diagnostic gap undermines treatment, surveillance, and policy development for effective prevention strategies, including vaccine prioritization.

In recent years, the application of multiplex PCR panels and unbiased metagenomic sequencing has expanded our understanding of CNS infection etiology, leading to the detection of emerging or previously unrecognized viruses in cerebrospinal fluid (CSF), such as chikungunya virus, SARS-CoV-2, and parechoviruses.[7–9] These discoveries suggest that additional understudied viruses may contribute to the burden of CNS infections in children, particularly in LMICs where surveillance is limited.

In recent years, human parvovirus 4 (PARV4 or tetraparvovirus) has surfaced as a virus of interest in CNS infections.[10,11] Two independent studies from India reported PARV4 DNA in the CSF of children with suspected CNS infection (4 of 22 cases tested). A previous study from Bangladesh also detected PARV4 RNA in 3 of 25 CSF samples from suspected meningitis cases using unbiased RNA metagenomics.[9] However, these findings, were limited by small sample sizes and lacked statistical association with clinical outcomes, leaving the clinical significance unresolved.

PARV4 is a small, non-enveloped, single-stranded DNA virus belonging to the *Parvoviridae* family. It was first identified in 2005 using sequence-independent PCR in an individual with febrile illness and hepatitis-like symptoms.[12] It is one of only four

parvoviruses known to infect humans, the others being parvovirus B19, human

bocavirus, and adeno-associated viruses.[13] The PARV4 genome is approximately 5·2

kb[14], encoding two major open reading frames (ORFs).

While PARV4 is strongly associated with parenteral transmission, especially among

people who inject drugs or those co-infected with HIV, hepatitis B virus (HBV), or

hepatitis C virus (HCV), seroprevalence studies have also detected PARV4 IgG in

individuals without these exposures, including healthy children, although the potential

for serological cross reactivity to other viruses remains is unknown.[13–17] Vertical

transmission has also been documented in neonates born to IgM-positive mothers, with

evidence of viremia at birth.[18]

Despite multiple reports of detection, the pathogenic role of PARV4 remains unclear.

There is currently no *in vitro* culture system for PARV4, and whether it replicates

autonomously is uncertain.[14] Most infections appear to be asymptomatic, but PARV4

DNA has been detected coincident with respiratory, gastrointestinal, hepatic, and

neurological symptoms.[11,12,19,20] It has also been found in blood, bone marrow, lymphoid

tissue, and liver.[13,21,22]

Overall, whether PARV4 contributes to disease, acts synergistically with other

pathogens, or represents an innocuous passenger is not yet known. Its frequent co-

detection with other viruses and association with young age raise questions about its

tissue tropism, transmission dynamics, and relation to disease severity. In this study, we

assess the prevalence, clinical associations, and genomic characteristics of PARV4 in a

large collection of CSF samples from children with suspected meningitis, collected over

eight years (2015-2022) at the largest pediatric hospital in Bangladesh. We evaluate

associations with age, co-infections, and in-hospital mortality, and analyse genomic

characteristics of circulating PARV4 strains.

Methods

Ethical considerations

The study protocol was approved by the Ethics Review Committee of the Bangladesh

Institute of Child Health. Written informed consent was obtained from the parents or

legal guardians of all participants prior to sample collection. All clinical data were de-

identified prior to analysis.

Study site and sample collection

Cerebrospinal fluid (CSF) samples were collected at Bangladesh Shishu Hospital and

Institute (BSHI), formerly known as Dhaka Shishu Hospital, the largest tertiary pediatric

hospital in the country, with 665 beds. BSHI and Child Health Research Foundation

(CHRF) collaborate to conduct meningitis surveillance, which was part of the invasive

bacterial vaccine-preventable disease (IB-VPD) surveillance.[9]

Lumbar puncture was performed as part of routine diagnostic evaluation for children

presenting with suspected meningitis, at the discretion of the attending physician.

Children were enrolled in the surveillance if their parent or legal guardian provided

written informed consent, they met the WHO-defined case criteria for meningitis (Text

S1; appendix p4) and had a CSF specimen collected as part of their clinical evaluation.

Residual CSF specimens were stored at –80°C for long-term preservation and used for

retrospective analyses in this study.

Sample processing and screening

CSF specimens were cultured using standard microbiological methods and evaluated

for meningitis causing pathogens (Text S2; appendix p4).

CSF samples with >9 WBC/μl, biobanked between 2015 and 2022, were screened for

PARV4. We designed and optimized qPCR assays to detect PARV4 and

Erythroparvovirus (B19) directly from whole CSF (Text S3; appendix p4). 16S Sanger

sequencing was performed on culture-positive samples for which bacterial species

could not be identified through standard microbiological methods (Text S4, appendix

p5).

158 Unbiased metagenomics and amplicon sequencing

Nucleic acid was extracted from 200 µl CSF samples using QIAamp DNA Mini kit

(Qiagen, Germany) for DNA and Quick DNA/RNA Miniprep Plus kit (Zymo Research,

USA) for RNA following the manufacturer's instructions. Using previously optimized

protocols, we used unbiased RNA and DNA metagenomics to detect PARV4 in the CSF

samples and generate high coverage full genomes (Text S5; appendix p6).[9]

Near-complete genomes of PARV4 from unbiased metagenomics were used to

establish a cost-effective amplicon sequencing protocol for PARV4 using

PrimalScheme[23] (Table S1; appendix p16, Figure S1; appendix p9) and following the

protocol described in Text S6; appendix p6. One PARV4-confirmed CSF with a Ct (cycle

threshold)-value <25 was randomly selected from each month between 2015-2022.

Consensus genomes of PARV4 from metagenomics and amplicon sequencing data

were generated using CZ-ID viral consensus genome pipeline v3·5.0, an open source

and cloud-based bioinformatics portal for analyzing NGS data (Text S6; appendix p6).

We used 30 available sequences from Matthew *et al*[24] to develop a genotyping scheme

for PARV4 sequences using custom python scripts (Text S7; appendix p7).

177 Genomic and phylogenetic analysis

To obtain multiple sequence alignment, all near complete genomes were aligned using

MAFFT v7·490. Maximum likelihood tree (MLT) was constructed employing the

generalized time-reversible model and a gamma distribution to model site-specific rate

variation. 100 bootstrap replicates were used to assess MLT statistical support. We

generated an outgroup rooted MLT (using MH215556·1 as the outgroup) with global

sequences and visualized it.

We estimated genome-wide substitutions/year (Text S8; appendix p7, Table S5;

appendix p26) and analyzed the mutation profile of PARV4 (Text S9, appendix p7).

188 Clinical data collection and analysis

Demographic and clinical data were systematically collected from electronically stored
health records for all cases. Descriptive analyses of the variables are presented as
frequencies (percentages) or medians with interquartile ranges (IQR), as most variables
exhibited non-normal distributions. Mann-Whitney U and Pearson chi-square tests were
used to compare demographic and clinical characteristics between PARV4-positive and
PARV4-negative groups. A two-proportion test was conducted to compare mortality
rates between the two groups.
196
197 To further assess the association between PARV4 positivity and mortality, we performed
198 both crude and adjusted logistic regression analyses. The adjusted model accounted for
199 potential confounders, including age, WBC count, and co-infections, to provide a
200 comprehensive understanding of the odds of mortality in PARV4-positive cases. All
201 statistical tests were two-sided, with a significance level set at 5%.
202

203 **Role of the funding source**
204 The funders of the study had no role in study design, data collection, analysis,
205 interpretation, or the decision to submit this manuscript for publication.
206

Results
PARV4 prevalence, demographics, and clinical outcomes
Between 2015 and 2022, a total of 8,238 CSF samples were collected from pediatric
patients presenting with suspected meningitis at BSHI (Figure 1). Among these, 3,235
samples had elevated WBC counts (>9 WBC/µl), of which 2,793 samples were
preserved in the biobank. Samples were evenly distributed over the eight-year period,
averaging 349 stored samples per year, except for 2020 when surveillance was
interrupted due to COVID-19 lockdowns. All 2,793 stored CSF samples were tested for
PARV4 using qPCR, with 526 (18·8%) testing positive for PARV4 DNA. 64·4% of the
samples were collected within 72 hours of hospitalization (Table S2; appendix p19).
Among PARV4-positive samples, Ct values were broadly distributed, with a median of
30·2 (IQR 24·4-33·9), with 12·9% of samples below Ct<20, and 30·0% of samples below
Ct<25 (Figure S2; appendix p10).

Figure 1. Flow diagram of CSF sample collection and PARV4 testing. This figure illustrates
the process of sample collection from children under 5 years of age meeting the case definition
of suspected meningitis at the Bangladesh Shishu Hospital and Institute. Samples were tested
for PARV4, and eligible positive samples underwent genome sequencing.

The PARV4 positivity rate remained consistent over the study years, with the highest
rate in 2021 at 31% and the lowest in 2016 at 11% (Figure S3A; appendix p11). Monthly
prevalence analysis across the eight-year study period showed fluctuations in positivity
rates but there appeared to be no consistent seasonal trend (Figure S3B; appendix
p11). BSHI is the largest tertiary pediatric care hospital in Bangladesh, and cases
included in this study came from different regions from across the country, with 24·7% of
positive cases coming from the Dhaka district, where the hospital is located, and the
remaining from other districts (Figure S4; appendix p12). Some areas had higher
positivity rates than others, such as Barisal, Bhola, and Bogura, but this study was not
designed or powered to assess statistical differences in spatial distribution.
236
The age distribution of PARV4-positive cases showed a peak in early infancy (Table 1,
Figure 2A). The median age of PARV4-positive cases was 4 months (IQR 1-17 months),
significantly younger than PARV4-negative cases whose median age was 7 months
(IQR 2-30, p<0·001). Among the positive cases, 12·9% of the cases were in their first 2
weeks of life, 55·3% were <6 months, and 79·6% were <2 years of age. Sex distribution
was similar in all groups, with approximate 60% of cases being male.
243
Among the 2,793 patients evaluated, respiratory symptoms such as difficulty breathing
(22·1% vs. 11·7%, p<0·001) and fast breathing (12·4% vs. 6·0%, p<0·001) were more
frequent among PARV4-positive cases (Table 1). Poor sucking was also significantly
higher in PARV4-positive cases (20·7% vs. 13·4%, p<0·001). Although neurological
symptoms were common, with convulsions observed in 70·3% of PARV4-positive cases,
this rate did not differ significantly in PARV4-negative cases (70·1%).
250
251 **Table 1. Clinical characteristics of cases (n = 2,793) admitted to study wards by PARV4**
252 **status.** A list of diagnoses included in the "other" category is provided in Table S3; appendix
253 p19). Data are n (%) or median [IQR] unless otherwise specified. p-values were calculated for
254 positive vs. negative groups using the ranksum or chi-square test based on data type.
255 DORB=discharge on risk bond.

	Overall	PARV4	PARV4	p-

		Negative	Positive	value
Number of Patients	**2793**	**2267 (81·2%)**	**526 (18·8%)**	
Sex- Female (%)	1019 (36·5)	819 (36·1)	200 (38·0)	0·416
Age in months (median [IQR])	7 [2·00, 27·00]	7·00 [2·00, 30·00]	4·00 [1·00, 17·00]	<0·001
< 2 weeks	194 (6·9)	126 (5·6)	68 (12·9)	
2 weeks - 3 months	860 (30·8)	687 (30·3)	173 (32·9)	
4 - 6 months	336 (12·0)	269 (11·9)	67 (12·7)	
7 months - 2 years	657 (23·5)	546 (24·1)	111 (21·1)	
2 - 5years	354 (12·7)	304 (13·4)	50 (9·5)	
>5 years	392 (14·0)	335 (14·8)	57 (10·8)	
In- hospital case fatality (%)	155 (5·5)	106 (4·7)	49 (9·3)	<0·001
Length of Stay (median [IQR])	12 [7·00,18·00]	11·00 [7·00, 16·00]	15·00 [10·00, 24·00]	<0·001
Hospital Outcome				0·330
Discharged	2376 (85·1)	1939 (85·5)	437 (83·1)	
Referred	14 (0·5)	13 (0·6)	1 (0·2)	
DORB	248 (8·9)	209 (9·2)	39 (7·4)	
Symptoms (%)				
Fever	2203 (78·9)	1820 (80·3)	383 (72·8)	<0·001
Cough	453 (16·2)	374 (16·5)	79 (15·0)	0·119
Difficulty Breathing	380 (13·6)	264 (11·7)	116 (22·1)	<0·001
Fast Breathing	201 (7·2)	136 (6)	65 (12·4)	<0·001
Convulsion	1959 (70·1)	1589 (70·1)	370 (70·3)	0·950
Vomiting	241 (8·6)	206 (9·1)	35 (6·7)	0·024
Headache	109 (3·9)	98 (4·3)	11 (2·1)	0·006
Diarrhoea	102 (3·7)	87 (3·8)	15 (2·9)	0·014
Lethargy	823 (29·5)	650 (28·7)	173 (32·9)	0·003
Chest In Drawing	108 (3·9)	79 (3·5)	29 (5·5)	0·060
Poor Sucking	412 (14·8)	303 (13·4)	109 (20·7)	<0·001

Diagnosis Cluster (%)				
Neurological Manifestation	1865 (66·8)	1513 (66·7)	352 (66·9)	0·937
Respiratory Manifestation	283 (10·1)	204 (9·0)	79 (15·0)	<0·001
Systemic Infections	313 (11·2)	230 (10·2)	83 (15·8)	<0·001
Febrile Illness	241 (8·6)	223 (10·0)	18 (3·4)	<0·001
Gastrointestinal Manifestation	47 (1·7)	41 (1·8)	6 (1·1)	0·283
Genitourinary Manifestation	20 (0·7)	14 (0·6)	6 (1·1)	0·200
Congenital Cardiovascular Manifestation	92 (3·3)	63 (2·8)	29 (5·5)	0·002
Perinatal Asphyxia	122 (4·4)	78 (3·4)	44 (8·4)	<0·001
Preterm low-birth weight	19 (0·7)	13 (0·6)	6 (1·1)	0·154
Neonatal Jaundice	24 (0·9)	16 (0·7)	8 (1·5)	0·068
Others	986 (35·3)	748 (33·0)	238 (45·2)	<0·001

We also assessed differences in final diagnosis at death or end of hospitalization.

Neurological manifestations were prevalent in both groups, affecting 66·9% of PARV4-

positive cases, similar to 66·7% of PARV4-negative cases. However, respiratory

manifestations and systemic infections were more common in PARV4-positive cases

(15·0% and 15·8%, respectively) compared to PARV4-negative cases (9·0% and 10·2%,

respectively; p < 0·001). The median WBC count for PARV4-positive cases was 50

cells/μl (IQR 20 - 210), which was the same (50 cells/μl) (IQR 20 - 170) for PARV4-

negative cases (Figure S5; appendix p13).

Figure 2. Age distribution and hospital duration of PARV4-positive and PARV4-negative
cases. (A) Age distribution (months) of children with suspected meningitis according to PARV4
status; dashed lines indicate median ages. **(B)** Duration of hospital stay (days) by PARV4
status; dashed lines indicate median duration.

The median length of hospital stay for PARV4-positive cases was significantly higher

than for PARV4-negative cases (15 days vs 11 days, p<0·001; Figure 2B). There were

no significant differences in the proportion of cases discharged, referred to other

hospitals, or cases where patients left against medical advice (Table 1). However, the

in-hospital case fatality rate among PARV4 positive cases was 9·3%, significantly higher

than the 4·7% observed in PARV4-negative (p<0·001).

Additional pathogens identified in PARV4-positive cases

All 2,793 CSF samples underwent bacteriological culture and immunochromatographic

testing for *Streptococcus pneumoniae* (pneumococcus) antigen as part of routine

diagnostics, and these results were available in the clinical records. In addition, in this

study, we conduced qPCR for pneumococcus, *Neisseria meningitidis, Haemophilus*

influenzae, and parvovirus B19 (B19). Among the 526 PARV4-positive cases, 262

(49·8%) were co-infected with at least one other pathogen, a rate significantly higher

than that of the PARV4-negative cases, where in only 374 out of 2,267 (16·5%) cases

another pathogen was detected (Table 2).

B19 was the only other virus tested for using PCR and was commonly detected in

PARV4-positive cases (173 of 526 cases, 32·9%). In comparison, only 157 out of 2,267

PARV4-negative cases (6·9%) were B19-positive. Pneumococcus (the only bacteria

whose presence is tested both through serology and culture) was the second most

frequently detected organism in PARV4-positive cases, found in 62 of 526 cases

(11·8%). This was almost double the detection rate in PARV4-negative cases, where

136 out of 2,267 cases (6·0%) tested positive for *Streptococcus pneumoniae*. Other

bacterial pathogens like *Neisseria meningitidis* and *Haemophilus influenzae* were rarely

detected in either PARV4 positive or negative cases.

Table 2. Distribution of additional pathogens among PARV4-positive and PARV4-negative
suspected meningitis cases. Organisms detected by culture, immunochromatographic test
(ICT), or PCR in cerebrospinal fluid samples stratified by PARV4 status. Data are presented as
number (%). B19 – erythroparvovirus B19, ICT – immunochromatographic test, PCR –
polymerase chain reaction.

	PARV4 Positive (n=526)				PARV4 Negative (n=2267)			
Organism	Total	Culture	ICT	PCR	Total	Culture	ICT	PCR
B19	173 (32·9%)	0	0	173	157 (6·9%)	0	0	157
Streptococcus pneumoniae	62 (11·78%)	2	59	1	136 (6·00%)	5	130	1
Neisseria meningitidis	9 (1·71%)	1	0	8	47 (2·07%)	1	0	46
Haemophilus influenzae	5 (0·95%)	0	0	5	11 (0·44%)	0	0	11
Acinetobacter baumannii	3 (0·57%)	3	0	0	3 (0·13%)	3	0	0
Klebsiella pneumoniae	3 (0·57%)	3	0	0	3 (0·13%)	3	0	0
Escherichia coli	3 (0·57%)	3	0	0	2 (0·09%)	2	0	0
Elizabethkingia anopheles	2 (0·38%)	2	0	0	1 (0·04%)	1	0	0
Acinetobacter baumannii	1 (0·19%)	1	0	0	3 (0·13%)	3	0	0
Flavobacterium meningosepticum	1 (0·19%)	1	0	0	2 (0·09%)	2	0	0
Salmonella enterica serovar Typhimurium	0	0	0	0	3 (0·13%)	3	0	0
Desemzia incerta	0	0	0	0	1 (0·04%)	1	0	0
Achromobacter xylosoxidans	0	0	0	0	2 (0·09%)	2	0	0
Streptococcus agalactiae	0	0	0	0	1 (0·04%)	1	0	0
Acinetobacter lwoffii	0	0	0	0	1 (0·04%)	1	0	0
Providencia alcalifaciens	0	0	0	0	1 (0·04%)	1	0	0

No organism detected	264 (50·4%)	0	0	0	1893 (83·6%)	0	0	0

Association of PARV4 with in-hospital case fatality

We examined the odds of in-hospital mortality associated with PARV4 positivity while adjusting for demographic and clinical factors. Individuals testing positive for PARV4 had 2·09 times higher odds of mortality compared to PARV4-negative individuals (Figure 3). This association remained consistent when adjusted for age (aOR = 2·06) and for WBC count (aOR = 2·05) in separate models. After accounting for co-infection with only B19, the commonly detected co-pathogen, the odds of mortality in PARV4-positive cases remained 2·13 times higher than in the negative group, regardless of B19 co-infection status.

Figure 3. Association between PARV4 positivity and in-hospital case fatality. Odds ratios (ORs) for in-hospital case fatality associated with PARV4 positivity under crude and adjusted models. Models were adjusted for age, other pathogens detected, erythroparvovirus B19, and white blood cell (WBC) count. Points indicate ORs and horizontal lines show 95% CIs; the vertical line at OR=1 indicates no association.

Genomic analysis of PARV4

A total of 73 samples underwent genome sequencing, including 20 sequenced using unbiased metagenomics and 53 using amplicon sequencing. Of these, 70 samples yielded sufficient data for whole genome assembly and further analysis. The coverage

depth across amplicon-based sequences ranged from 1694x- 31248·9x with 98%

samples having coverage depth more than 1694x. In these samples, over 97% of the

genome met thresholds for calling consensus bases. An overview of the consensus

genomes with coverage depth is given in Table S4; appendix p24.

Bayesian analysis of Bangladeshi PARV4 population (n=70) estimated a genome-wide

substitution rate of 0·5911475 substitutions per year (Table S5; appendix p26). This rate

represents the average number of nucleotide changes accumulating in the PARV4

genome annually in Bangladesh. We then analysed the mutation profile of 66 PARV4

genomes which had complete ORF1 and 2 and used a local genome

(CHRF_CSF_0090, accession number: ERS29405649) from 2015 as a reference. Non-

synonymous (NS) mutation profiling revealed region-specific variability in the frequency

of mutations (Figure 4A, Figure S6; appendix p14). ORF1 coding for NS1 exhibited

sparse distribution of NS substitutions. In ORF2, coding for VP1 and VP2 structural

proteins, NS substitutions were concentrated towards the unstructured VP1 region (non-

VP2 region of VP1) with over 70% samples having amino acid substitutions at various

regions between aa 913 and aa 1125. PARV4 showed limited variation along the capsid

forming VP2 with 77·6% of samples showing substitutions in variable regions-II (VR-II)

and 37·3% in VR-VIII.

Phylogenetic analysis of the 70 assembled genomes and 37 publicly available PARV4

sequences showed that most sequences from Bangladesh clustered within genotype 2,

aligning closely with strains previously identified in India (Figure 4B). Two sequences

formed a distinct clade between genotypes 1 and 2, suggesting a potential novel

genotype within Bangladesh (we term genotype 1,2).

Figure 4. Genomic variation and phylogenetic analysis of PARV4. (A) Frequency of non-synonymous mutations across the PARV4 genome. The x-axis shows nucleotide position with annotated coding regions (NS1, VP1 and VP2); y-axis indicates the proportion of isolates with NS mutations at each position. Variable regions (VR-I to VR-IX and the HI loop) within VP2 region are highlighted in blue. **(B)** Maximum-likelihood phylogenetic tree of PARV4 genomes generated in this study together with publicly available global sequences. Outer annotations indicate geographic origin and genotype. The scale bar represents substitutions per site.

Discussion

With significant positive associations between PARV4 positivity, co-infections, and in-hospital case fatality, this study generates evidence that PARV4 is neuroinvasive and could be significant contributor to CNS infections particularly in settings where rates of idiopathic meningitis continue to be high.

We detected PARV4 in 18·8% of CSF samples from suspected meningitis cases. This rate aligns with the two reports from India - one detected PARV4 DNA in 2 of 10 CSF samples from children with acute CNS infection and another detected in 2 of 12.[10,11] Our data also revealed a higher prevalence of PARV4 among children under 1 year of age, with a significantly lower median age of 4 months in PARV4-positive cases compared to PARV4-negative cases (7 months). 13% of the positive cases were in babies younger than 2 weeks. This trend may indicate increased susceptibility to PARV4 infection in early life with potential vertical transmission. In a study from Taiwan, placental transmission of PARV4 was reported in neonates who presented with viremia and were born to IgM-positive mothers, suggesting the virus may be transmitted vertically, in line with our findings.[18]

Almost half of the PARV4-positive cases had at least one additional pathogen detected in the CSF. The most common co-infection was with parvovirus B19 (only virus tested for), a related but distinct member of the same viral family. Parvovirus B19 was detected in a third of PARV4-positive cases compared to only 6·97% in PARV4-negative cases. Several previous studies have also reported strong association of PARV4 with other viruses like HBV, HCV and HIV.[13,16,17] In addition to B19, *Streptococcus pneumoniae*

(SPN) was the second most frequently detected co-infecting pathogen in PARV4-
positive cases. While the immunochromatographic test used for detection of
pneumococcus in the CSF is highly sensitive compared to only culture used for other
bacteria,[25] pneumococcal antigen was present in nearly double the rate in PARV4-
positive cases compared to negative cases. Given that pneumococcus is a leading
cause of bacterial meningitis, co-infection with PARV4 could complicate clinical
management, particularly if PARV4 contributes to atypical presentations or immune
modulation in infected individuals. Although co-infections with *Neisseria meningitidis*
and *Haemophilus influenzae* were less common, overall higher rates of co-infections
may suggest that PARV4 increases susceptibility to other CNS pathogens or causes
secondary infections upon infection with other pathogens that impact clinical outcomes.
398
A key finding of this study is the association between PARV4 positivity and increased in-
hospital mortality. Nearly 10% of PARV4-positive cases died during admission, and
compared with PARV4-negative cases, PARV4-positive children had approximately
twice the odds of death. This association remained after adjustment for age, white blood
cell count, and detection of other pathogens. The relatively low cycle-threshold values
observed, together with prior detection of PARV4 RNA in unbiased metagenomic
studies[9], suggest the likely possibility of active viral replication within the CNS, and
potential neurotropism. Given that infants comprised of the majority of PARV4-positive
cases, the observed mortality underscores the potential clinical relevance of PARV4
detection in this vulnerable population.
409
Beyond the differences in mortality, respiratory manifestations and systemic infections
were more frequently observed among PARV4-positive cases than PARV4-negative
cases. These findings suggest that PARV4-associated illness may have distinct clinical
features compared to other pediatric CNS infections. Detection of PARV4 in respiratory
samples from children with respiratory illness in Ghana and India has also been
reported[19,26], raising the possibility of respiratory involvement and potential respiratory
transmission, similar to B19. However, further studies are needed to clarify transmission
dynamics and clinical pathogenesis.

Bayesian phylogenetic analysis of the Bangladeshi PARV4 genomes revealed a
genome-wide substitution rate of 0·5911475 substitutions per year, consistent with those
observed in related human parvoviruses, which generally exhibit substitution rates
between ~1×10^{-5} and 1×10^{-4} substitutions per site per year.[27] The unique N-terminal
unstructured region of VP1 in PARV4 exhibited the presence of several NS mutations,
suggesting a potential site for positive selection. In B19, this region of VP1 has been
demonstrated to be important during the internalization of the virus in the host cell and
was found to be present on the exterior side of capsid which could potentially enable
interaction with their host for infection and to elicit immune response.[28,29] The role of this
region of VP1 in PARV4 pathogenicity remains unknown.

Phylogenetic analysis of 70 PARV4 genomes showed that most isolates clustered within
genotype 2. This aligns with previous reports showing that the three PARV4 genotypes
classified to date have varying geographic distributions: genotypes 1 and 2 are
prevalent in Europe, North America, and Asia, while genotype 3 is common in sub-
Saharan Africa.[13,17,30] The PARV4 genomes sequenced from CSF samples in India also
belong to genotype 2.[10] Notably, two sequences formed a distinct clade positioned
between genotypes 1 and 2, indicating the circulation of a distinct genotype within
Bangladesh, which we term 1,2.

The findings of this study must be interpreted within the context of several limitations.
First, our study was conducted at a single tertiary care hospital, potentially limiting the
generalizability of the results to other settings in Bangladesh or other LMICs. However,
the large dataset of over 2,000 samples collected over eight years from patients across
the country suggests that our findings are likely to reflect broader trends and are not
isolated from a single institution. Second, the cross-sectional design of the study
restricts our ability to establish causality between PARV4 infection and clinical
outcomes, including mortality. Although logistic regression analysis suggests a strong
association, prospective and longitudinal studies, paired with laboratory studies with ex-
vivo models are needed to confirm the role of PARV4 in disease causality, progression

and outcomes. Finally, our detection methods relied on PCR and metagenomic

sequencing, which, while sensitive, may not fully distinguish between active infection

and chronic viremia. Future studies incorporating viral load quantification or markers of

active infection could help determine whether PARV4 presence directly contributes to

CNS pathology.

While we are yet to understand the causal relation of PARV4 with CNS infection, the

findings reveal a high prevalence of the virus, significant co-infection rates with other

pathogens, and an association with increased mortality in cases of suspected

meningitis. Understanding PARV4's biology, clinical implications, and potential

transmission routes will be essential for informing public health strategies and improving

outcomes for vulnerable pediatric populations.

Contributors

DPK and ARM contributed equally to this work. DPK, ARM, NK, and AMT contributed to

drafting the first version of the manuscript, data analysis, and figure preparation. NK

contributed to analysis, figures, and study design. AMT contributed to study design,

genomics analyses, manuscript review and editing. SR, PCD, HR, LT, AR, AAT, DCD,

ABJ, MZI, RM, SN, ANUA, and FI contributed to data interpretation, manuscript review,

and editing. PCD and AAT contributed to genomic analyses. HR, ABJ, and SN

contributed to surveillance activities and study implementation. DCD contributed to

laboratory work. MZI, RM, and SN contributed to study implementation and clinical

oversight. CT contributed to training and analytical support. JLD, SKS, YH, and SS

conceptualised and designed the study and contributed to data interpretation and

manuscript revision. SS obtained funding and oversaw project administration and

management.

All authors had full access to the data in the study and accept responsibility for the

decision to submit for publication.

Data sharing

The sequence reads of 70 PARV4 genomes are available in the European Nucleotide

Archive (ENA) under the study accession PRJEB88738 / ERP171815. All scripts and

codes used to create the figures can be found in github folder https://github.com/CHRF-

Genomics/PARV4-Analysis

Declaration of interests

All authors declare no competing interests.

Acknowledgements

This study was supported by the Gates Foundation and the Child Health Research

Foundation, Dhaka, Bangladesh. From the Child Health Research Foundation, we

would like to thank Nusrat Alam, Sharmistha Goswami, Md Mobarak Hossain for helping

us with amplicon sequencing set up and Adittya Arefin for assisting with literature

review.

References:

1 McIntyre PB, O'Brien KL, Greenwood B, Beek D van de. Effect of vaccines on
bacterial meningitis worldwide. *The Lancet* 2012; **380**: 1703–11.

2 Roth GA, Abate D, Abate KH, *et al.* Global, regional, and national age-sex-specific
mortality for 282 causes of death in 195 countries and territories, 1980–2017: a
systematic analysis for the Global Burden of Disease Study 2017. *The Lancet* 2018;
392: 1736–88.

3 James SL, Abate D, Abate KH, *et al.* Global, regional, and national incidence,
prevalence, and years lived with disability for 354 diseases and injuries for 195
countries and territories, 1990–2017: a systematic analysis for the Global Burden of
Disease Study 2017. *The Lancet* 2018; **392**: 1789–858.

4 Ramakrishnan M, Ulland AJ, Steinhardt LC, Moïsi JC, Were F, Levine OS. Sequelae
due to bacterial meningitis among African children: a systematic literature review.
BMC Medicine 2009; **7**: 47.

5 Granerod J, Tam CC, Crowcroft NS, Davies NWS, Borchert M, Thomas SL.
Challenge of the unknown: A systematic review of acute encephalitis in non-outbreak
situations. *Neurology* 2010; **75**: 924–32.

6 Kwambana-Adams BA, Liu J, Okoi C, *et al.* Etiology of Pediatric Meningitis in West
Africa Using Molecular Methods in the Era of Conjugate Vaccines against
Pneumococcus, Meningococcus, and Haemophilus influenzae Type b. *Am J Trop*
Med Hyg 2020; **103**: 696–703.

7 Guan Y, Yu C, Fei Y, *et al.* Case Report: Four cases of SARS-CoV-2-associated
Guillain-Barré Syndrome with SARS-CoV-2-positive cerebrospinal fluid detected by
metagenomic next-generation sequencing: a retrospective case series from China.
Front Immunol 2023; **14**: 1258579.

8 Suthar PP, Hughes K, Kadam G, Jhaveri M, Gaddikeri S. Human parechovirus
meningoencephalitis. *SA Journal of Radiology* 2023; **27**: 2589.

9 Saha S, Ramesh A, Kalantar K, *et al.* Unbiased Metagenomic Sequencing for
Pediatric Meningitis in Bangladesh Reveals Neuroinvasive Chikungunya Virus
Outbreak and Other Unrealized Pathogens. *mBio* 2019; **10**: e02877-19.

10   Prakash S, Jain A, Seth A, Singh AK, Jain B. Complete Genome Sequences of
Two Isolates of Human Parvovirus 4 from Patients with Acute Encephalitis Syndrome.
Genome Announc 2015; **3**: e01472-14.

11 Benjamin LA, Lewthwaite P, Vasanthapuram R, *et al.* Human Parvovirus 4 as
Potential Cause of Encephalitis in Children, India. *Emerg Infect Dis* 2011; **17**: 1484–7.

12   Jones MS, Kapoor A, Lukashov VV, Simmonds P, Hecht F, Delwart E. New DNA
viruses identified in patients with acute viral infection syndrome. *J Virol* 2005; **79**:
8230–6.

13   Manning A, Willey SJ, Bell JE, Simmonds P. Comparison of tissue distribution,
persistence, and molecular epidemiology of parvovirus B19 and novel human
parvoviruses PARV4 and human bocavirus. *J Infect Dis* 2007; **195**: 1345–52.

14   Matthews PC, Malik A, Simmons R, Sharp C, Simmonds P, Klenerman P. PARV4:
An Emerging Tetraparvovirus. *PLoS Pathogens* 2014; **10**: e1004036.

15   Sharp CP, Lail A, Donfield S, *et al.* High frequencies of exposure to the novel
human parvovirus, PARV4 in haemophiliacs and injecting drug users detected by a
serological assay for PARV4 antibodies. *J Infect Dis* 2009; **200**: 1119–25.

16   Simmons R, Sharp C, McClure CP, *et al.* Parvovirus 4 Infection and Clinical
Outcome in High-Risk Populations. *The Journal of Infectious Diseases* 2012; **205**:
1816–20.

17   Yu X, Zhang J, Hong L, *et al.* High Prevalence of Human Parvovirus 4 Infection in
HBV and HCV Infected Individuals in Shanghai. *PLoS ONE* 2012; **7**: e29474.

18   Chen M-Y, Yang S-J, Hung C-C. Placental Transmission of Human Parvovirus 4
in Newborns with Hydrops, Taiwan. *Emerging Infectious Diseases* 2011; **17**: 1954.

19   Drexler JF, Reber U, Muth D, *et al.* Human Parvovirus 4 in Nasal and Fecal
Specimens from Children, Ghana. *Emerg Infect Dis* 2012; **18**: 1650–3.

20   Fryer JF, Lucas SB, Padley D, Baylis SA. Parvoviruses PARV4/5 in hepatitis C
virus-infected patient. *Emerg Infect Dis* 2007; **13**: 175–6.

21   Schneider B, Fryer JF, Reber U, *et al.* Persistence of novel human parvovirus
PARV4 in liver tissue of adults. *J Med Virol* 2008; **80**: 345–51.

22   Chen M-Y, Hung C-C, Lee K-L. Detection of human parvovirus 4 viremia in the
follow-up blood samples from seropositive individuals suggests the existence of
persistent viral replication or reactivation of latent viral infection. *Virol J* 2015; **12**: 94.

23   Kent C, Smith AD, Tyson J, *et al.* PrimalScheme: open-source community
resources for low-cost viral genome sequencing. 2024; : 2024·12·20·629611.

24   Matthews PC, Sharp C, Simmonds P, Klenerman P. Human parvovirus 4 'PARV4'
remains elusive despite a decade of study. *F1000Res* 2017; **6**: 82.

25   Hasanuzzaman M, Saha S, Malaker R, *et al.* Comparison of Culture, Antigen
Test, and Polymerase Chain Reaction for Pneumococcal Detection in Cerebrospinal
Fluid of Children. *The Journal of Infectious Diseases* 2021; **224**: S209.

26   Prakash S, Shukla S, Ramakrishna V, Mishra H, Bhagat AK, Jain A. Human
Parvovirus 4: A harmless bystander or a pathogen of severe acute respiratory illness.
Int J Infect Dis 2020; **90**: 21–5.

27   Firth C, Kitchen A, Shapiro B, Suchard MA, Holmes EC, Rambaut A. Using time-
structured data to estimate evolutionary rates of double-stranded DNA viruses. *Mol*
Biol Evol 2010; **27**: 2038–51.

28   Leisi R, Von Nordheim M, Ros C, Kempf C. The VP1u Receptor Restricts
Parvovirus B19 Uptake to Permissive Erythroid Cells. *Viruses* 2016; **8**: 265.

29   Lakshmanan R, Mietzsch M, Jimenez Ybargollin A, *et al.* Capsid Structure of
Aleutian Mink Disease Virus and Human Parvovirus 4: New Faces in the Parvovirus
Family Portrait. *Viruses* 2022; **14**: 2219.

30   Simmonds P, Douglas J, Bestetti G, *et al.* A third genotype of the human
parvovirus PARV4 in sub-Saharan Africa. *Journal of General Virology* 2008; **89**:
2299–302.