Application of wastewater and environmental surveillance for pathogenic agents

during the 2024 National Football League (NFL) Draft in Detroit, Michigan (USA)

Ryland Corchis-Scott[1], Ethan Harrop[1], Qiudi Geng[1], Mackenzie Beach[1], John Norton[2],

Mehdi Aloosh[3,4], Thomas Reid[5], Rajesh Seth[6], Christopher G. Weisener[1], and R. Michael

McKay[1]

Author affiliations:

[1]University of Windsor, Great Lakes Institute for Environmental Research, Windsor,

Ontario, Canada

[2]Great Lakes Water Authority, Detroit, MI, United States

[3]Windsor-Essex County Health Unit, Windsor, Ontario, Canada

[4]Department of Health Research Methods, Evidence, and Impact, McMaster University,

Hamilton, Ontario

[5]Environment and Climate Change Canada, Canada Centre for Inland Waters, Burlington,

Ontario, Canada

[6]University of Windsor, Department of Civil and Environmental Engineering, Windsor,

Ontario, Canada

Corresponding author: Ryland Corchis-Scott, Great Lakes Institute for Environmental

Research, University of Windsor, 401 Sunset Ave., Windsor, ON, Canada N9B 3P4;

corchisr@uwindsor.ca, 519-991-3493

Abstract

Mass gatherings pose a concern for public health because they are associated with dense

crowds, increased social interaction, and travel, all of which can facilitate the rapid

transmission of infectious diseases. Wastewater and environmental surveillance (WES)

were used for pathogen monitoring during the 2024 NFL Annual Player Selection Meeting

(the 'Draft') in Detroit, MI, an event that drew an estimated 775,000 attendees. Wastewater

and environmental samples were queried for respiratory viruses and clinically relevant

antimicrobial resistance genes (ARG). WES did not detect an increase in the concentration

of monitored respiratory viruses (SARS-CoV-2, IAV, IBV, and RSV) associated with the 2024

NFL Draft. In contrast, WES detected a transient increase in carbapenemase targets in

wastewater, primarily driven by a fourfold increase in bla_{OXA-48}. Resistome structure in

wastewater was dominated by sampling site characteristics rather than changes

associated with the event. The Draft weekend coincided with rainfall-driven combined

sewer overflow (CSO), potentially allowing the dissemination of ARG to the environment. In

surface waters receiving wastewater effluent, an increase in detection frequency and

normalized concentrations for multiple ARG were observed following the Draft. WES

provided an overview of pathogen prevalence before, during, and after a large-scale

gathering, showing how it can warn of emerging health risks in near real time.

Introduction

Analysis of wastewater can provide information that accurately reflects infectious diseases

dynamics[1–3] by integrating signals from symptomatic, subclinical, and asymptomatic

individuals[4–7]. Wastewater and environmental surveillance (WES) is not a new discipline,

but it has experienced a renaissance following its successful implementation during the

COVID-19 pandemic[8–10]. Increased interest in WES has been driven by the numerous

advantages it offers, including correlation with clinical metrics[11–17], cost-effectiveness[18–21],

independence from test-seeking behaviour or availability of testing infrastructure[22],

scalability from building to city levels[14,23–25], and timeliness with near real-time data

generation[12,26,27].

An emerging application of WES is infectious diseases surveillance during mass gatherings[28]. Mass gatherings are typically large, planned events that concentrate individuals from diverse geographies into dense settings with high mobility, conditions that facilitate the transmission of infectious diseases[29]. Mass gatherings pose a challenge to epidemiological disease tracking due to travel and increased likelihood of rapid transmission. Disease surveillance before, during and after events allows for rapid response to widespread transmission following disease detection[28,30]. WES offers an effective means by which populations may be monitored surrounding large-scale gatherings such as sporting events[31–33].

Prior applications of WES to mass gatherings have targeted SARS-CoV-2 to monitor transmission of COVID-19 and include the 2020 Tokyo Olympics[34], the Two Oceans Marathon in South Africa[35], and the 2022 FIFA World Cup in Qatar[36]. Recent festival-based monitoring in New Orleans, LA, expanded the application of WES to mass gatherings, where it was implemented to target Mpox transmission[37]. Beyond respiratory viruses, surveillance of antimicrobial resistance (AMR) via antimicrobial resistance genes (ARGs) is gaining traction, given public health relevance and the potential for dissemination facilitated by mass gatherings[38]. Antimicrobial-resistant organisms (AROs) are also targets of expanded WES[39,40] with surveillance focused on the presence of specific ARGs of public health significance[41]. Monitoring has been established as a critical step in addressing the proliferation of AMR[42] with efforts having been made to test for the presence and level of ARGs at wastewater treatment facilities[39,43,44]. Exacerbating the threat of AMR are rainfall-driven combined sewer overflows (CSOs) and bypass events that can introduce partially treated effluent into receiving waters, increasing environmental ARG loads.

WES may be implemented through a OneHealth framework that considers the movement of organisms and resistance determinants between humans, infrastructure, and natural systems[45]. Wastewater is a critical connection between these systems and thus surveillance implemented here captures both human population dynamics and pathways for dissemination to the environment. Mass gatherings increase population density and can strain sanitation infrastructure. WES implemented during these events is opportune for

advancing knowledge of disease trajectories in human population and ARG dissemination

in the environment.

The 2024 NFL Draft was held in Detroit, MI from April 25-27, 2024 with events concentrated

in an area with an approximate radius of 1 km near the riverfront in downtown Detroit. This

event drew approximately 775,000 individuals to the city of Detroit[46,47] of which an

estimated 30% travelled more than 160 km to attend the event. Visitors to Detroit included

people from over 20 different nations and all 50 states, with some individuals coming from

as far afield as Australia[48]. Almost all of Detroit's 45,140 hotel rooms were occupied for the

Draft weekend, with spillover absorbed by neighbouring Windsor, ON, a region separated

from Detroit by a river and high-traffic international land border.

Here, we used the 2024 NFL Draft as a natural experiment to explore WES utility for mass

gatherings across two domains: (1) respiratory pathogens in wastewater; and (2) public

health relevant ARGs, with emphasis on carbapenemases, in wastewater and the Detroit

River, an environmental reservoir for wastewater treatment plant (WWTP) effluent. We also

compared multiplex RT-qPCR with a nanofluidic OpenArray® platform to assess method

agreement across matrices. We hypothesized that respiratory signals would not increase

immediately given the primarily outdoor nature, late-season timing of the event, and its

brevity (in comparison to the incubation of respiratory viruses). We also expected that

ARGs could show transient changes, and that heavy rainfall coincident with the Draft could

elevate environmental ARG detections due to CSO and/or bypass activity.

Methods

Site information

Sampling sites included five wastewater treatment facilities and a surface water transect of

the Detroit River with six sample collection sites (Figure 1). The Draft weekend coincided

with a wet weather period in the Windsor-Detroit region, with approximately 40 mm of rain

falling over a 48 h period (Sunday-Tuesday)[49]. This influx of precipitation contributed to CSO

events where partially treated WWTP effluent and diluted raw sewage was discharged into

the Detroit River or its tributaries from both sides of the Canda-USA border[50].

Wastewater treatment facilities were identified as potentially impacted by the NFL Draft or

as control sites based on proximity (Table 1). Additional site information can be found in S1.

Sample collection

Untreated wastewater influent was collected from each of the wastewater treatment

facilities included in the study at different weekly frequencies (Table 1) with daily sampling

between April 24-29 from each of the three interceptors (NIEA, OAK, and JEFF) of the Great

Lakes Water Authority Water Resource Recovery Facility (WRRF) as well as the WWTP plant

(WEWTP) in Windsor, ON. All samples were transported directly to the laboratory in coolers

on ice and processed upon arrival. Surface water grab samples (2 L) were collected in

duplicate from each site in the Detroit River transect (Figure 1). Environmental samples

were collected on April 25, prior to the 2024 NFL Draft, and on April 29 following the end of

the event.

Sample processing

Influent samples collected from WWTPs were concentrated by filtering between 60-120 mL

through 0.22 μm Sterivex PES cartridge filters (MilliporeSigma, Burlington, MA). River water

samples were likewise concentrated by passing 500-800 mL of water through cartridge

filters. Immediately following filtration, the filters were sealed and flash-frozen through

immersion in liquid nitrogen, with filters stored at -80°C before nucleic acid extraction.

Total nucleic acids were extracted from filters using the AllPrep PowerViral DNA/RNA kit

(Qiagen, Germantown, MD) with the addition of 5% (v/v) 2-mercaptoethanol to the lysis

buffer. Nucleic acids were eluted in 50 μL of RNAse-free water. Samples were not treated

with DNase upon extraction. Extracted nucleic acids were stored in LoBind tubes at -80°C.

RT-qPCR assays for quantification of respiratory viruses and carbapenemase genes

Respiratory viruses, select carbapenemase genes and Pepper Mild Mottle Virus (PMMoV)

were quantified using established RT-qPCR assays. Severe Acute Respiratory Syndrome

Coronavirus 2 (SARS-CoV-2), influenza A virus (IAV), influenza B virus (IBV), and respiratory

syncytial virus (RSV) were the respiratory viruses measured. PMMoV, a human fecal

indicator[51] and established wastewater surveillance normalizing agent[52], was used to

normalize the wastewater signal for targets measured using RT-qPCR assays.

Carbapenemase genes, including bla_{KPC}, bla_{NDM}, bla_{OXA-48}, bla_{VIM}, and bla_{VIM7} were measured

using two multiplexed assays[44]. All targets were measured in at least technical triplicate.

Standard curves were generated using serial dilution of synthetic RNA or DNA controls.

Inhibition was assessed using an internal amplification control, and template RNA was

diluted to overcome inhibition. Negative controls were included with each RT-qPCR run.

Detailed RT-qPCR methods are described in S2.

Development of the nanofluidic qPCR card using TaqMan® assays in the OpenArray ®

platform

Assays for the quantification of clinically important and locally relevant ARGs and genes

associated with mobile genetic elements (MGEs) were selected for inclusion on the

OpenArray® card (Table 2). The OpenArray® card also included assays for 16S rRNA and

crAssphage to allow normalization and act as controls. Assays were selected from the

literature and optimized for TaqMan chemistry. The resulting nanofluidic RT-qPCR card was

used to analyze samples in technical duplicate in 33nL reactions on a Quant Studio 12K

Flex Real-Time PCR System (Thermo Fisher Scientific, Waltham, MA). A 7-point standard

curve, created through a serial dilution of pooled gBlocks™ (Integrated DNA Technologies,

Coralville, IA), was run on each OpenArray ® plate. Assay performance met established

technical standards for RT-qPCR. Full assay design and nanofluidic qPCR methods can be

found in S3.1 and S3.2.

Statistical methods

Data analysis was performed in R (version 4.5.1)[53] with data manipulation and visualization

relying on the *tidyverse* suite of packages[54]. Multiple different measurement techniques

were employed in this investigation, leading to the creation of five distinct datasets.

Statistical analyses were conducted separately for each dataset due to differences in

sampling frequency, sample matrix, availability of covariate data, data distribution, and

response variable. All analyses tested changes in wastewater and environmental

occurrence of pathogens associated with the mass gathering of 2024 NFL Draft attendees

and the co-occurring CSO event.

Interrupted time series (ITS) analysis

To test if the 2024 NFL Draft coincided with altered concentrations of respiratory pathogens

or carbapenemase genes in wastewater, we fit Generalized Linear Mixed Models (GLMMs)

using the *glmmTMB* package. Independent analyses were conducted on "affected" and

"control" sites for a quasi-experimental design. Model specification followed a segmented

regression framework with a binary "event" variable (modelling immediate event-related

changes) and a "post-time" variable (more gradual event-related changes). Models

controlled for a variety of covariates (pH, temperature, flow, precipitation, etc.) and

temporal autocorrelation was addressed via an autoregressive term for all targets and

seasonal harmonics for respiratory virus response variables. A Tweedie distribution with a

log link was used for continuous zero-inflated viral pathogen concentration data. A Gamma

distribution with a log link was used for individual carbapenemase gene concentration

data, while a Tweedie distribution was used for the composite measure of all assayed

carbapenemase genes. To account for multiple testing, p-values were adjusted using the

Benjamini-Hochberg (BH) False Discovery Rate (FDR) method. Adjustment was performed

independently for respiratory viruses and carbapenemase genes and for "control" and

"affected" sites. Details can be found in S4.1 and S4.2.

Multivariate resistome characterization

The Open Array nanofluidic RT-qPCR card was used to measure a subset of samples

collected from WWTPs during the month surrounding the 2024 NFL Draft. ARG

concentrations were normalized to the concentration of crAssphage and subjected to

fourth root transformation. Data was split into the uneven groups ("before", "during" and

"after") relative to the timing of the 2024 NFL Draft. Characterization of the resistome was

carried out with Non-Metric Multidimensional Scaling (NMDS) using Bray-Curtis

dissimilarity matrices. A marginal Permutational Multivariate Analysis of Variance

(PERMANOVA) was then run on the data to partition sources of variation, and a distance-

based redundancy analysis was used for visualization of variance partitioning. A

Permutational Analysis of Multivariate Dispersions (PERMDISP) was used to test if

observed differences arose from differences in resistome or within-group variance. A

principal component analysis (PCA) was run to identify ARGs driving variability. Finally, JEFF

was analyzed alone as it drains the NFL Draft location. A Kruskal-Wallis test was used to

determine if concentration differences existed between the "before", "during" and "after"

groupings for the measured ARGs. The p-values were adjusted for multiple comparisons

using the Benjamini-Hochberg method. Details can be found in S4.3.

Environmental detection

To determine if the CSO events altered the occurrence of carbapenemase genes in the

Detroit River, detection frequencies were calculated for each carbapenemase gene in the

pre- and post- CSO transects. Fisher's exact test was used to compare the detection

frequency. Additionally, a paired Wilcoxon signed-rank test was performed to evaluate

concentration changes for PMMoV normalized bla_{KPC} between the pre- and post CSO

samplings.

To determine if the CSO and mass gathering event led to either changes in abundance or

detection frequency for additional ARGs, data produced by the Open Array nanofluidic RT-

qPCR card were employed. Data were normalized to 16S rRNA concentrations and log-

transformed before quantitative analysis. Detection frequencies were also calculated for

both the pre- and post- CSO groupings. Changes to detection frequency were assessed

using a GLMM with a binomial distribution and logit link function. To prevent complete

separation, ARGs that were constitutively detected or undetected in either the pre-CSO or

post-CSO grouping were excluded from logistic regression. The main GLMM included ARG

identity, sampling group (pre/post CSO) and the interaction of the two as fixed effects.

Random intercepts and slopes for groups nested within sites were also included to

account for differences between sites. Post-hoc comparisons were conducted using

estimated marginal means. A second GLMM was conducted with site and sample grouping

as fixed effects and gene as a random intercept to directly investigate site heterogeneity.

Quantitative abundance differences between pre- and post-CSO sample groupings were

tested using paired Wilcoxon signed rank tests with pairings assigned by site identity and

ARG. A sensitivity analysis was performed to identify the influence of potential detection

saturation in samples collected at a known WWTP outlet. To account for multiple

comparisons, p-values were adjusted using the Benjamini-Hochberg method to control

FDR. Model diagnostics were assessed using simulated residuals in the *DHARMa* package.

Details can be found in S4.4 and S4.5.

Method comparison

OpenArray® assays were compared with established RT-qPCR assays using a subset of

wastewater and environmental samples that were analyzed by both methods (n=42).

Comparison was limited to assays for bla_{KPC}, bla_{NDM}, and bla_{OXA-48} as they were shared

between the OpenArray® and the multiplexed RT-qPCR assays. Qualitative agreement was

evaluated using Cohen's Kappa and detection frequency comparisons. Quantitative

agreement was determined through Bland-Altman analysis to calculate 95% limits of

agreement (LoA). Additional quantitative measures of method comparison included

Passing-Bablok regression, Deming regression, spearman correlation, Pearson correlation

and calculation of concordance correlation coefficients (CCC). Details can be found in

S4.6.

Results

Respiratory pathogen dynamics

ITS analysis showed no event-associated increase in wastewater concentrations of

SARS-CoV-2, IAV, IBV, or RSV at monitored sites. IBV showed a post-event decline (β=-

0.356, SE=0.143, p= 0.035), consistent with seasonal-tapering rather than event impact

(Figure 2). Variability in respiratory virus concentrations in wastewater was driven by

dilution and seasonality. Concentrations of SARS-CoV-2 were negatively associated with

flow (β=–0.288, SE=0.115, p = 0.035) and positively associated with PMMoV concentration

(β = 0.379; SE=0.065, p < 0.001), consistent with dilution. Likewise, IAV concentration in

wastewater was negatively predicted by flow (β=-0.467, SE=0.160, p = 0.012), consistent

with dilution. However, IAV concentrations were positively predicted by precipitation

(β=0.317, SE=0.079, p<0.001). RSV concentrations were positively associated with PMMoV

(β= 0.455, SE= 0. 141, p = 0.005) while IAV was a negative predictor (β= -0.318, SE= 0.128 p

= 0.035). Precipitation was positively associated with IBV concentration (β=0.345,

SE=0.138, p = 0.035) while flow was negatively associated (β=-0.924, SE=0.284, p = 0.005).

Finally, IBV was found to be positively associated with IAV concentration (β=0.814,

SE=0.244, p = 0.004).

Parallel analysis at control sites showed no event effects. The GLMM run for SARS-CoV-2

did not detect any stepwise increase in SARS-CoV-2 concentration or change in trend

associated with the Draft. However, LRT indicated that the inclusion of "event" and "post

time" parameters improved model fit (Table S10). Sensitivity analysis and visualization of

the data indicate that this is due to noise in the N2 concentration at TB (Figure 3, Table 3).

The detection of a marginal, site-specific fluctuation at a control location demonstrates

sensitivity to minor background variance in the approach, reinforcing the validity of the null

results observed at affected site cluster. This contrast confirms that while the modelling

framework can identify localized shifts, no such signal was generated by the mass

gathering event.

Carbapenemase gene concentration in wastewater

ITS analysis indicated a statistically significant increase in the composite measure of

carbapenemase gene concentrations concurrent with the 2024 NFL Draft (β=0.625,

SE=0.290, p=0.031). An ~87% increase in combined ARG burden could be observed

following the Draft weekend (Figure 4). However, the event did not lead to a change in the

trend for the ARG composite index as the post-event temporal trend variable was non-

significant (p = 0.96). ITS for individual carbapenemase genes revealed that the increase

was primarily driven by a spike in normalized wastewater concentration for bla_{OXA-48}

(β=1.38, SE=0.374, p=0.006), which increased by ~400% following the Draft (Figure 5). A

parallel increase in bla_{KPC} concentration (β=0.447, SE=0.212, p=0.035) did not survive FDR

correction (p=0.18). ITS for bla_{NDM}, bla_{VIM}, and bla_{VIM7} indicate that the 2024 NFL Draft did not

significantly impact normalized ARG concentration despite event estimates all indicating

post-event increases. Flow was a significant positive predictor of ARG concentration

across 4 of 5 gene targets (Table S14). Higher flows were consistently associated with

increased normalized ARG concentrations.

Resistome variability

Characterization of the resistome with NMDS revealed samples collected from the same

site clustered regardless of sample collection time in relation to the 2024 NFL Draft

(stress= 0.068; Figure 6). The marginal PERMANOVA corroborated the NMDS visualization,

indicating that site identity explained 61% of variance in resistome (R^2 = 0.611, $F_{3,16}$= 9.30, p

= 0.001) while collection period was statistically insignificant (R^2 = 0.033, $F_{2,16}$= 0.76, p =

0.531). The PERMDISP showed that these differences were not a result of heterogeneity of

variance with no difference in dispersion between groups ($F_{2,19}$ = 0.90, p = 0.421) or site

identities ($F_{3,18}$ = 1.55, p = 0.236). The first two principal components of the PCA accounted

for 78% of the variance. The highest loadings for PC1 were *int1*, *sul1*, *aac(6')-Ib-cr*, *tet(A)*,

sul2, *mphA, qnrs, dfrA1, bla$_{OXA-48}$*, and *bla$_{TEM}$*. The top PC1 loadings had the same sign,

indicating the top contributors varied together across collection sites (Figure 7). The

Kruskal-Wallis test for JEFF revealed no significant difference in concentration for any ARG

in the "before", "during", or "after" groups.

Environmental dissemination of ARGs

Detection frequencies for *bla$_{KPC}$*, *bla$_{OXA-48}$* and *bla$_{VIM7}$* rose nominally in the post-Draft

transect. However, detection frequency increases were non-significant for all

carbapenemase genes. A paired Wilcoxon signed-rank test for PMMoV normalized *bla$_{KPC}$*

indicated an increased median concentration between the pre- and post CSO samplings (V

= 614, p<0.001). Heterogeneity in effect could be observed between sites (Table 5, Figure

8). Detection frequencies were also calculated for ARGs included on the OpenArray® for

both the pre- and post- CSO groupings. Detection frequency increased following the Draft

weekend for 62.5% of ARGs assayed (Figure 9).

The binomial GLMM showed that sample collection timing had a significant effect on the

probability of *crAssphage* detection (Estimate = 11.46, p = 0.0008), suggesting that the

2024 NFL Draft and CSO greatly increased detection probability. Additionally, the fitted

model had high levels of variance for the random intercept (σ^2 = 62.07, SD=7.88) and the

random slope for the "after" group (σ^2 =6.36, SD=2.52), indicating large differences in

baseline detection probabilities and Draft effect between sites. Interaction terms between

ARG and sampling group were not statistically significant, indicating that the relative

magnitude of the post-Draft increase did not differ significantly among genes. Post-hoc

pairwise comparisons for each gene included in the model found that detection probability

increased for 6 genes including: *crAssphage, AAC(6')-lb-cr, qnRS, tetA, mph(A),* and bla_{KPC}.

(Table S26). Increased detection probability post-Draft was corroborated by the paired

Wilcoxon signed-rank test. Ten ARGs displayed statistically significant increases in

concentration following the event, with no ARGs showing a significant decrease in

abundance (Table S29).

The GLMM investigating site heterogeneity identified PO as the site with the highest

detection probability, consistent with a point source of pollution (Table S30). An additional

4 ARGs showed statistically significant increases in concentration when the PO site was

removed from the Wilcoxon signed-rank test (Table S33), suggesting that constitutively high

ARG concentration at PO was masking significant increases in ARG concentration at other

sites (Figure 10).

Method comparison

In wastewater, detection was 100% for all targets shared across platforms. In

environmental samples, OpenArray showed lower sensitivity, missing low abundance

targets detected by RT-qPCR (Table 6). Bland-Altman analysis showed that bla_{KPC} had the

narrowest LoA with bla_{NDM} and bla_{OXA-48} having wider LoAs (Figure S19). Both bla_{KPC} and

bla_{NDM} had positive bias while bla_{OXA-48} showed negative bias. Fold change analysis showed

similar results, with the OpenArray having higher measurements in wastewater samples for

bla_{KPC} and bla_{NDM} but lower for *blaOXA-48* (Figure S20).

The strength of monotonic agreement between the results of the OpenArray and RT-qPCR

varied by target with the strongest agreement for bla_{KPC} and weakest for bla_{OXA-48} (Table 7).

Pasing-Pablok and Demming regression revealed proportional bias for all targets (slopes

>1.0), and large negative intercepts (Table S34). This pattern indicates a relative

underestimation of ARG concentrations by the OpenArray® at lower analyte

concentrations, transitioning to a relative overestimation at higher analyte concentrations.

CCC was the highest for bla_{KPC} (CCC=0.94) and lower for bla_{NDM} (CCC = 0.88) and bla_{OXA-48}

(CCC=0.75) (Table 7). Only bla_{KPC} demonstrated substantial methodological agreement

between platforms based on previously determined criteria[55].

Discussion

This study supports the utility of WES for monitoring and assessing the impact of mass

gatherings on the spread of viral pathogens and antibiotic resistance genes. The 2024 NFL

Draft served as a natural experiment. We measured the concentrations of public health

relevant ARGs and seasonal respiratory pathogens in wastewater and in the Detroit River,

which receives treated effluent from the Detroit-Windsor metropolitan region. WES

detected no surge in respiratory viruses associated with the event, yet identified a

transient, event-coincident rise in bla_{OXA-48} in wastewater and broader ARG increases in

surface waters during a concurrent wet-weather period with confirmed CSOs. A broad

assessment of ARG concentrations in wastewater conducted over a limited timescale

found that variability in the wastewater resistome was driven by site-level heterogeneity,

potentially arising from differences in antimicrobial use between sewersheds.

Viral pathogens

The 2024 NFL Draft did not act as a superspreader event for respiratory pathogens.

Concentrations of SARS-CoV-2, IAV, IBV, and RSV remained steady or declined following

the event. No evidence of increased transmission was observed despite the substantial

increase in population in the monitored sewersheds and the mass gathering with increased

transmission risk. Multiple factors may have contributed to the observed lack of increased

transmission. The Draft was held outdoors and in large, well-ventilated buildings. This

would have helped to limit the potential for aerosol transmission despite the large

crowds[56]. Additionally, the Draft was held following the peak of the respiratory season in the

northern hemisphere. Immunity for the most relevant strains of respiratory virus was likely

high (acquired via infection or inoculation). The wastewater signal for most of the

monitored pathogens was already low, indicating that prevalence was already low,

reducing the probable number of infectious individuals attending Draft-related events. The

lack of immediate increase in signal over the Draft weekend may be related to the relatively

short duration of the Draft (~3 days) in comparison with viral incubation times (~5-6 days)[57].

The brevity of the mass gathering may not have allowed sufficient time for incubation or

widespread transmission. Our analysis also indicated that Draft related transmission was

low, with no significant increases in wastewater signal for any pathogens following the

event. A recent study of religious gatherings in Saudi Arabia corroborates this result,

reporting that infection control implemented for the 2022 Hajj resulted in few cases of

COVID-19 and no secondary transmission or outbreaks[58]. These findings demonstrate that

large numbers of individuals can congregate without unreasonably increased risk of

transmission if the gatherings occur outside of respiratory seasons or proper precautions

to mitigate spread are implemented. The absence of a detectable signal does not imply

complete absence of transmission; it does, however, suggest that the mass gathering event

did not lead to a widespread increase in cases. Lack of extensive transmission is an

important finding that can inform public health decision-making and provide feedback on

event timing and the types of transmission mitigation strategies employed. Additionally,

detection of transmission would be valuable for informed rapid action, especially when

clinical case reporting is not feasible or delayed.

Event-associated increases in carbapenemase genes

Unlike the monitored respiratory viruses, carbapenemase genes increased measurably

following the mass gathering, driven by a large increase in bla_{OXA-48} signal. The mass

gathering event and influx of visitors were associated with a transient spike in the

concentration of a clinically important ARG in the wastewater system. A lack of increase in

any carbapenemase gene concentration at the spatially distinct control sites increases the

likelihood that Draft-related travel was responsible for the increase rather than seasonal

variability in ARG concentration. The increase in bla_{OXA-48} observed in the present study may

be partially explained by Draft attendees arriving from regions where OXA-48-like

carbapenemases are more prevalent. These include parts of Europe, Africa, and the Middle

East[59]. International travel is an important driver of the dissemination of OXA-48-like

carbapenemases[60,61] and in non-endemic areas, infections are often imported, though this

modality may be changing[62]. Our results indicated that the increase in bla_{OXA-48} was

transient with no evidence of a sustained increase. However, the large increase in bla_{OXA-48}

concentration was notable, especially given that the baseline ARG signal was highly

variable. This demonstrates that WES for ARGs may be sensitive to short term

perturbations and that mass gatherings may induce short term changes in load for select

ARGs. Wastewater flow was found to be a significant positive predictor of normalized ARG

concentration, potentially a result of surface inputs to the sewer system or the mobilization

of sewer biofilms[63,64].

Stable resistome with site-level heterogeneity

Multivariate analysis consistently indicated that the resistome structure in wastewater was

dominated by sampling site characteristics rather than changes associated with the 2024

NFL Draft. This suggests that the influx of visitors during the 2024 NFL Draft did not

significantly affect the resistome of the monitored sites in a way that was detectable in this

analysis. Likely, sewershed level drivers such as population demographics, hospital

effluent volumes, land use characteristics (i.e., amount of animal agriculture), and

antibiotic prescription rates played a key role in determining the resistome. This was further

supported by PCA loadings, which were dominated by common markers of anthropogenic

pollution, like *int1* and *sul1*[65,66]. These ARGs co-varied across sites and sampling periods,

indicating that the variation was driven by baseline pollution levels. Most PC1 loadings had

the same sign, supporting the notion of covariance of ARGs being driven by variability in

baseline pollution. ARG concentrations at the JEFF interceptor (the sampling site most

closely linked to Draft activities) were not influenced by the Draft.

Conclusions from this analysis must be interpreted with caution as group sizes were

uneven, sample sizes were small, and the study period was short. Combined, these limit

the statistical power of the current analyses to detect relatively small changes in ARG

concentration. Additionally, the OpenArray® has few targets compared to the diversity of

ARGs, limiting our ability to detect ARGs that may have been introduced by the Draft.

Incorporation of ultra deep metagenomic sequencing may allow for the detection of novel,

and low-abundance ARGs[67]. Longer-term data would help to confirm any Draft-associated

changes. The results of this study are not generalizable. Mass gathering events still have

the potential to contribute to ARG dissemination, especially if they are associated with

travel between jurisdictions with different antimicrobial prescribing practices.

Environmental dissemination of ARGs

Analyses revealed an increase in detection frequency and normalized concentration for

multiple ARGs in the Detroit River following the 2024 NFL Draft weekend. The Draft event

coincided with heavy precipitation, driving a CSO event. Detection probability increased for

clinically relevant ARGs including: *AAC(6')-lb-cr, qnRS, tetA, mph(A),* and *bla*$_{KPC}$. These

ARGs are associated with a variety of antimicrobial agents, indicating broad, rather than

targeted increases in ARG detection following the Draft weekend. No gene × group

interaction terms were significant in the GLMM, suggesting that the magnitude of the

increase was consistent between targets. The rise in crAssphage concentrations in the

post-Draft environmental samples and uniformity of detection probability increases across

gene targets indicate that sewage contamination from the CSO is a major cause of the

observed increase in ARG detections and abundance in the environmental samples. CSO

events in isolation have been found to increase environmental concentrations of ARGs[68,69].

Mass gathering events can contribute to increased wastewater volumes and ARG loads[70].

However, given that OpenArray® -based analysis of wastewater samples did not detect a

statistically significant increase in ARG concentrations attributable to the NFL Draft, the

observed increase in ARG in the environment was likely caused by the rainfall-driven CSO

events. The coincidence of the mass gathering event and the CSO may have introduced

ARGs to the environment, imported to the region by event attendees, thus altering the

watershed resistome. Such alterations have been observed in previous studies[71].

Heterogeneity was observed among the sites of the sampling transect, with site-specific

responses to the CSO and mass gathering. Samples collected at the outlet of the WEWTP

exhibited evidence of saturated detection, accompanied by the highest normalized ARG

concentrations. These findings support that the WWTP outlet was a major source of ARGs

in the receiving waters with constitutively high ARG concentrations. Inclusion of samples

collected at the point source in data analysis obscured more subtle changes in ARG

concentrations at other sites. This observation indicates the importance of site selection

and highlights the complexity of interpretation for WES data in systems that are influenced

by persistent inputs. In these systems, measuring baseline contamination levels is

inherently challenging. Findings also indicated that the influence of the CSO was not

limited to the pollution source. Increased concentrations of ARGs were found at the site

upstream of the known WWTP outlet as well as the downstream sites. This observation

could be explained by environmental sewage contamination originating from CSO events

that occurred upstream within the Huron-Erie Corridor. Mass gatherings may increase the

consequences of rainfall events in combined sewer systems, leading to alterations in the

resistome of the receiving waters. Such alterations may have consequences for

downstream utilization of water resources[72].

Limitations of this analysis arise from the restricted temporal sampling and reliance on RT-

qPCR for ARG measurement. RT-qPCR cannot detect which pathogen is hosting an ARG or

if they are hosted on mobile genetic elements. Future work should incorporate longer-term

monitoring efforts, hydrological modelling for better source tracking and should include

metagenomic sequencing to help resolve ARG host identity.

Comparison between OpenArray® and RT-qPCR

Comparison of the OpenArray ®and RT-qPCR quantification of three carbapenemase genes

across two sample matrices ascertained the validity of the OpenArray® platform for ARG

surveillance. The OpenArray® can be used for ARG trend assessment in concentrated

wastewater matrices, but it is not directly quantitatively interchangeable with RT-qPCR,

due to lower sample volumes. Agreement between the two methods was dependent on

ARG abundance in the sample measured. Concordance between methods was 100% in

high-concentration wastewater samples but was poor for environmental samples.

Disagreement arose from non-detects by the OpenArray® rather than quantitative

disagreement between methods. This observation is consistent with the approximately

600-fold smaller reaction volume (33nL vs 20µL) for the OpenArray®, which lowers

detection probability at low copy numbers. Thus, the OpenArray® has limited utility for

environmental monitoring without upstream concentration in the workflow[73–75].

Assessment of quantitative agreement between OpenArray® in high concentration

matrices also revealed significant discrepancies between methods. Passing-Bablok and

Demming regression both revealed systematic proportional and constant bias between

methods. Slopes greater than 1 and large negative intercepts are consistent with

underestimation of samples at low concentrations and progressive overestimation at

higher concentrations (consistent with nanolitre scale inputs, but a more efficient assay).

Agreement between the OpenArray® and RT-qPCR platforms was target dependent with

bla_{KPC} showing the strongest quantitative agreement (CCC=0.94), indicating

methodological interchangeability. However, for bla_{NDM} and bla_{OXA-48}, overall agreement

between methods was much lower (CCC=0.88 and CCC=0.75, respectively). For these

targets, OpenArray® and RT-qPCR methods are not quantitatively interchangeable.

However, strong correlations were found between OpenArray® data and RT-qPCR data for

all tested targets (Table 7). These correlations support reliable trend tracking for

OpenArray® and RT-qPCR despite differences in the absolute concentration measured.

Tracking of trends over time is an important aspect of both wastewater-based surveillance

and environmental monitoring, and despite the apparent quantitative issues with the

OpenArray® platform, trend agreement is encouraging for the future development of the

platform. The multitude of ARG targets to be monitored and high relative abundance in

wastewater matrices make OpenArray® a promising platform for ARG monitoring in high

concentration matrices with potential future applications for environmental monitoring

with the implementation of pre-amplification and further methodological confirmations.

Low quantitative concordance between OpenArray® and RT-qPCR for bla_{OXA-48} may help to

explain why OpenArray® analysis failed to detect the spike in bla_{OXA-48} that was detected via

RT-qPCR. Optimization of multiplexed RT-qPCR allows for improved assay performance

while cycling conditions for OpenArray® are set and uniform for all targets included on the

chip. Additionally, OpenArray® analysis of the wastewater resistome relied on a limited

number of samples while RT-qPCR benefited from a year-long survey of carbapenemase

gene concentrations and a more sophisticated ITS modelling approach that attempted to

account for baseline variability in carbapenemase concentrations.

Conclusion

The main goals of this investigation were to determine whether WES is a viable tool for

monitoring pathogen transmission dynamics associated with mass gathering events and to

determine if the 2024 NFL Draft led to increased pathogen transmission or ARG

dissemination. Findings provide support for the use of WES in this application, with results

of respiratory pathogen monitoring indicating no discernible increase in transmission

associated with the 2024 NFL Draft. WES to monitor trends in AMR is particularly promising

as it can capture a dimension of infectious diseases that are more difficult to track via

clinical surveillance methods. Additionally, environmental dissemination of ARGs is

impossible to track via clinical surveillance but can have implications for population

health. The ability of WES to detect transient resistome shifts makes the argument that it is

a valuable public health surveillance tool during mass gatherings. A recent study

investigated ARG load, diversity and dissemination in relation to mass gathering events[76].

Investigators found that the overall level of ARGs did not increase in relation to mass

gathering events. More importantly, they determined that mass gatherings could introduce

ARGs to local sewer systems that were previously absent. Findings of our investigation are

similar, albeit more methodologically limited, as we only measured select ARGs. Both our

investigation and the previous work found that ARG burden did not increase overall as a

result of the mass gathering but that the concentration of select genes might increase in a

transient manner and that the event itself may have led to the introduction of locally novel

ARGs. Notably, the wastewater signal observed in this study was not driven by a uniform

increase across all monitored targets. Instead, the response was selective, emphasizing

the importance of targeted surveillance strategies and careful interpretation of WES data in

event-based contexts. Dissemination of locally novel ARGs is concerning from a public

health perspective as it is likely to lead to increased numbers of antibiotic-resistant

infections or infections with novel resistance profiles. The combined evidence supports

WES as a reliable tool for validating the safety of mass gatherings regarding respiratory

spread while providing early insight into AMR dynamics and environmental dissemination

under infrastructure stress. Integrating WES with hydrologic context (rainfall, CSO

telemetry) can guide real-time operational responses (e.g., CSO management, advisories)

during future events.

Acknowledgements

Operators, Laboratory Team, and Managers of the following wastewater treatment

facilities: Wastewater Resource Recovery Facility, Great Lakes Water Authority; Pollution

Control, City of Windsor; Pollution Control Centre, Municipality of Leamington, Water

Pollution Control Plants, City of Chatham and City of Thunder Bay. Funding in support of

the Ontario Wastewater Surveillance Initiative was provided by the Ontario Ministry of

Environment, Conservation, and Parks. Additional support was provided by the Public

Health Agency of Canada's National Wastewater Surveillance Program (Contribution

Agreement 2324-HQ-000197 to CGW, DDH, RS and RMM). The views expressed herein do

not necessarily represent the views of the Public Health Agency of Canada.

Table 1. Site characteristics and sample collection modalities

Facility Name	Site ID	Location	Site type	Population served	Sample type	Sampling frequency	Role
Water Resource Recovery Facility (WRRF)	NIEA	Detroit, MI, USA	Interceptor feeding wastewater treatment plant	1482000	1L 24h composite	1×weekly. Daily sampling during Draft	Impacted site
	OAK	Detroit, MI, USA	Interceptor feeding wastewater treatment plant	840600	1L 24h composite	1×weekly Daily sampling during Draft	Impacted site
	JEFF	Detroit, MI, USA	Interceptor feeding wastewater treatment plant	492000	1L 24h composite	1×weekly Daily sampling during Draft	Impacted site
Anonymous Windsor-Essex Wastewater Treatment Plant	WEWTP	Windsor-Essex, ON, Canada	Wastewater treatment plant	180000	1L 24h composite	5×weekly Daily sampling during Draft	Impacted site
Leamington Pollution Control Centre	LPCC	Leamington, ON, Canada	Wastewater treatment plant	18000	1L 24h composite	3×weekly	Control site
Chatham Water Pollution Control Plant	CHWW	Chatham, ON, Canada	Wastewater treatment plant	48000	0.5L 24h composite	3×weekly	Control site
Atlantic Avenue Water Pollution Control Plant	TB	Thunder Bay, ON, Canada	Wastewater treatment plant	97500	0.5L 24h composite	3×weekly	Control site
Detroit River Transect	UP	Upstream of Detroit River outfall	Surface water	NA	2L Grab	Event based	Env sample
	PO	Detroit River outfall	Surface water	NA	2L Grab	Event based	Env sample
	DS1	Downstream of Detroit River outfall	Surface water	NA	2L Grab	Event based	Env sample
	DS2	Downstream of Detroit River outfall	Surface water	NA	2L Grab	Event based	Env sample
	DS3	Downstream of Detroit River outfall	Surface water	NA	2L Grab	Event based	Env sample
	DS4	Downstream of Detroit River outfall	Surface water	NA	2L Grab	Event based	Env sample

552 **Table 2.** Antimicrobial resistance genes monitored, associated drug resistance, and

553 rationale for inclusion on the Nanofluidic qRT-PCR chip

Class of antimicrobial	Gene(s)	Rationale for testing
Aminoglycoside	$aph(3')$-Ia, $aac(6')$-Ib-cr	Often used in pediatric settings[77] Renewal in interest with rise in AMR for treatment of MDR infections[78]. Most common resistance mechanism for aminoglycosides is through enzymatic modification with the listed ARGs coding for the most common enzymes[79]
Carbapenems	bla_{KPC}, bla_{NDM}, bla_{OXA-48}, bla_{IMP}, bla_{VIM}	Rapid spread and ability to infect patients in healthcare settings. Rise in prevalence with few viable treatment options. Often resistant to multiple classes of antibiotic [80–82].
Cephalosporins/extended spectrum ß-lactam (ESBL) antibiotics	bla_{CMY-2}, bla_{SHV}, bla_{TEM}, CTX-M1, CTX-M9,	Beta-lactams are highly common treatments for bacterial infections but resistance levels are high[83]. Some later generation cephalosporins and ESBL are considered last resort antibiotics[84]. Levels of cephalosporin resistance are generally rising[85]. Identified as likely a large threat to human health with emerging concerns regarding environmental spread[86].
Colistin	mcr-1	Listed as a reserve antimicrobial agent for last resort use by the World Health Organization[87]. Rapid global spread[42].
Fluoroquinolones	gyrA83L-Esh, QnrB1, qnrS	Widely used theraputics[88] with fluoroquinolone resistant infections resulting in worse patient outcomes and increase risk of mortality[89]. Resistance is widespread[90–93]. Many fluoroquinolones are on the WHO "Watch" list. [87]
Glycopeptides (vancomycin)	vanA	Last resort antibiotic used in the treatment of serious gram-positive bacterial infections[94,95]. Recent rapid increases in resistance [96] Vancomycin is on the WHO "Watch" list. [87]
Macrolides	mphA	Used widely in the treatment of common infections such as pneumonia, sinusitis, and chlamydia. Members include erythromycin, clarithromycin, azithromycin[97]. Associated with mobility[98] mphA is plasmid associated and mobile with evidence of transmission through horizontal gene transfer[99]. Commonly prescribed throughout Ontario[100]
Methicillin	mecA	Primary gene associated with beta-lactam resistance in Methicillin-Resistant Staphylococcus aureus (MRSA)[101] Thought to be mobile and has been shown confer advantages to bacterial exposed to a diversity of antimicrobial agents/chemical environments[102]
Tetracycline	tetA, tetM	Broadly administered in agriculture and human healthcare[86]. Evidence of co-selection with fluoroquinolone resistance genes[103]

Sulfonamides	*sul1, sul2*	Widely used and well established class of antimicrobial agent[42]. Broad spectrum antibiotics with both human and agricultural applications[104].	
Trimethoprim	*dfrA1*	Antimicrobial agent often used in the treatment of urinary tract infections. Widely prescribed in Ontario[100].	
Associated with resistance to multiple therapeutics	*cfr, int1*	The *cfr* gene is associated with multi-drug resistance[105] while the *int1* gene is a sign of the presence of mobile genetic elements often associated with AMR. Presence of *int1* alone does not indicate AMR. However, it is associated with multiple ARGs and their mobility[106,107]	
Normalization/Control	16S rRNA, CrAssphage	crAssphage is a bacteriophage found in association with human waste making it a good fecal indicator and a measure of wastewater concentration that can be used for normalization of wastewater signals [108,109]. A general assay for 16S rRNA was also incorporated to act as a control and for normalization.	

Table 3. Summary of Interrupted Time Series (ITS) model parameters, Likelihood Ratio Tests (LRT), and diagnostic checks for gamma GLMMS for N1 and N2 gene targets in the control site cluster.

		Target	
Metric	Statistic	N1	N2
Model Estimates	Event	0.333 (p=0.272)	0.393 (p=0.179)
	Post Time	0.012 (p=0.789)	0.025 (p=0.572)
Model Fit	ΔAIC	-0.5	-4.0
	Log-Likelihood	1122.1	1136.4
	Dispersion	0.187	0.193
LRT	Chi-Square	4.485	8.066
	p-value	0.106	0.018*
DHARMa diagnostics	KS	p=0.309	P=0.627
	Dispersion	p=0.892	P=0.424
	Quantiles	p=0.074	P=0.069
	Outlier	P=0.288	P=0.288

560 **Table 4.** Multivariate analysis of resistome composition. Results of the marginal

561 Permutational Multivariate Analysis of Variance (PERMANOVA) based on Bray-Curtis

562 dissimilarities. Site identity serves as the primary driver of community variance, while the

563 NFL Draft period (Group) did not significantly influence the resistome structure

Factor	Degrees of Freedom	R^2	F-Statistic	p-value
Site	3	0.611	9.305	0.001
Draft period	2	0.033	0.760	0.531
Residual	16	0.350	-	-
Total	21	1	-	-

564 **Note:** Site dispersion (p = 0.236) and Group dispersion (p = 0.421) were non-significant, supporting the PERMANOVA findings.

566 **Table 5.** Median log PMMoV normalized bla_{KPC} concentrations for each environmental

567 sampling site before and after the 2024 NFL Draft and associated combined sewer

568 overflow (CSO) event. The mean difference represents the magnitude of the concentration

569 shift. P-values were calculated using site-specific paired Wilcoxon signed-rank tests

Site	Median before	Median after	Mean difference	Wilcox p-value	Significance
DS1	1.030	1.631	0.522	0.031	Significant
DS2	0.394	1.077	0.699	0.156	Non-significant
DS3	0.286	1.787	1.429	0.031	Significant
DS4	0.976	1.991	0.987	0.031	Significant
PO	1.023	1.233	0.238	0.094	Non-significant
UP	0.375	0.443	0.035	0.844	Non-significant

572   **Table 6.** Comparison of RT-qPCR and nanofluidic RT-qPCR performance for detection of

573   select antimicrobial resistance genes (ARGs) in different sample matrices

Target	Matrix	No. Observations	RT-qPCR Positive	OpenArray Positive	Detection Gap	Kappa (κ)	p-value
bla_{KPC}	Global	42	42 (100%)	31 (73.8%)	-26.20%	0.000*	NA
	Wastewater	18	18 (100%)	18 (100%)	0%	1.000**	NA
	Environmental	24	24 (100%)	13 (54.2%)	-45.80%	0.000*	NA
bla_{NDM}	Global	42	34 (81.0%)	18 (42.9%)	-38.10%	0.300	0.006
	Wastewater	18	18 (100%)	18 (100%)	0%	1.000**	NA
	Environmental	24	16 (66.7%)	0 (0.0%)	-66.70%	0.000*	NA
bla_{OXA-48}	Global	42	26 (61.9%)	25 (59.5%)	-2.40%	0.751	< 0.001
	Wastewater	18	18 (100%)	18 (100%)	0%	1.000**	NA
	Environmental	24	8 (33.3%)	7 (29.2%)	-4.10%	0.516	0.011

575   **Table 7.** Correlation and agreement of measurements for select ARGs assayed with RT-

576   qPCR assays and nanofluidic RT-qPCR

Target	n	Spearman	Pearson	CCC (95% CI)	Mean Bias (log)	LoA (95%)
bla_{KPC}	16	0.91	0.97	0.94 (0.74–0.97)	0.15	-0.21 to 0.51
bla_{NDM}	15	0.77	0.93	0.88 (0.69–0.94)	0.29	-0.77 to 1.34
bla_{OXA-48}	18	0.64	0.82	0.75 (0.42–0.86)	-0.27	-1.11 to 0.58

Citations

1. Gonzalez, R. & Smith, J. Population-level health monitoring using wastewater surveillance. *Journal of Public Health Advances* **34**, 456–470 (2021).

2. Mao, K., Zhang, H. & Yang, Z. Wastewater-based epidemiology and community health: A review. *Environmental Research* **191**, 110351 (2020).

3. Sims, N. & Kasprzyk-Hordern, B. Future perspectives of wastewater-based epidemiology. *Environmental Health Journal* **50**, 123–135 (2020).

4. Hillary, L. S. *et al.* Monitoring SARS-CoV-2 in municipal wastewater to evaluate the success of lockdown measures for controlling COVID-19 in the UK. *Water Research* **200**, 117214 (2021).

5. Medema, G., Heijnen, L., Elsinga, G., Italiaander, R. & Brouwer, A. Presence of SARS-Coronavirus-2 RNA in Sewage and Correlation with Reported COVID-19 Prevalence in the Early Stage of the Epidemic in The Netherlands. *Environ. Sci. Technol. Lett.* **7**, 511–516 (2020).

6. Yanaç, K., Adegoke, A., Wang, L., Uyaguari, M. & Yuan, Q. Detection of SARS-CoV-2 RNA throughout wastewater treatment plants and a modeling approach to understand COVID-19 infection dynamics in Winnipeg, Canada. *Sci Total Environ* **825**, 153906 (2022).

7. Corchis-Scott, R. *et al.* Averting an Outbreak of SARS-CoV-2 in a University Residence Hall through Wastewater Surveillance. *Microbiol Spectr* **9**, e0079221 (2021).

8. Daughton, C. G. Wastewater surveillance: An emerging tool for epidemiology. *Environmental Science and Technology* **54**, 431–438 (2020).

9. Farkas, K., Hillary, L. S. & Malham, S. K. Wastewater surveillance as a tool for

monitoring the SARS-CoV-2 pandemic. *Current Opinion in Environmental Science and*

Health **17**, 14–20 (2020).

10. Tiwari, A. & Kumar, R. The renaissance of wastewater surveillance during the SARS-

CoV-2 pandemic. *Epidemiological Reviews* **45**, 78–89 (2022).

11. Ahmed, W. *et al.* Occurrence of multiple respiratory viruses in wastewater in

Queensland, Australia: Potential for community disease surveillance. *Sci Total Environ*

864, 161023 (2023).

12. Corchis-Scott, R. *et al.* Wastewater Surveillance to Confirm Differences in Influenza A

Infection between Michigan, USA, and Ontario, Canada, September 2022–March 2023 -

Volume 30, Number 8—August 2024 - Emerging Infectious Diseases journal - CDC.

Emerging Infectious Diseases **30**, (2024).

13. Galani, A. *et al.* SARS-CoV-2 wastewater surveillance data can predict hospitalizations

and ICU admissions. *Science of The Total Environment* **804**, 150151 (2022).

14. Peccia, J. *et al.* Measurement of SARS-CoV-2 RNA in wastewater tracks community

infection dynamics. *Nat Biotechnol* **38**, 1164–1167 (2020).

15. Hopkins, L. *et al.* Citywide wastewater SARS-CoV-2 levels strongly correlated with

multiple disease surveillance indicators and outcomes over three COVID-19 waves. *Sci*

Total Environ **855**, 158967 (2023).

16. Zhao, L. *et al.* Targeting a free viral fraction enhances the early alert potential of

wastewater surveillance for SARS-CoV-2: a methods comparison spanning the

transition between Delta and Omicron Variants in a large urban center. *fpubh* (2023).

17. Gerrity, D., Papp, K., Stoker, M., Sims, A. & Frehner, W. Early-pandemic wastewater

surveillance of SARS-CoV-2 in Southern Nevada: Methodology, occurrence, and

incidence/prevalence considerations. *Water Res X* **10**, 100086 (2020).

18. Yoo, B.-K., Iwamoto, R., Chung, U., Sasaki, T. & Kitajima, M. Economic Evaluation of

Wastewater Surveillance Combined with Clinical COVID-19 Screening Tests, Japan.

Emerg Infect Dis **29**, 1608–1617 (2023).

19. Wright, J., Driver, E. M., Bowes, D. A., Johnston, B. & Halden, R. U. Comparison of high-

frequency in-pipe SARS-CoV-2 wastewater-based surveillance to concurrent COVID-19

random clinical testing on a public U.S. university campus. *Science of The Total*

Environment **820**, 152877 (2022).

20. Safford, H. R., Shapiro, K. & Bischel, H. N. Wastewater analysis can be a powerful

public health tool—if it's done sensibly. *Proceedings of the National Academy of*

Sciences **119**, e2119600119 (2022).

21. Ngwira, L. G. *et al*. Cost of wastewater-based environmental surveillance for SARS-

CoV-2: Evidence from pilot sites in Blantyre, Malawi and Kathmandu, Nepal. *PLOS Glob*

Public Health **2**, e0001377 (2022).

22. National Academies of Sciences, E. *et al*. Wastewater Surveillance for COVID-19. in

Wastewater-based Disease Surveillance for Public Health Action (National Academies

Press (US), 2023).

23. Corchis-Scott, R. *et al*. Actionable wastewater surveillance: application to a university

residence hall during the transition between Delta and Omicron resurgences of COVID-

19. *Frontiers in Public Health* **11**, (2023).

24. Bivins, A. & Bibby, K. Wastewater Surveillance during Mass COVID-19 Vaccination on a

College Campus. *Environ Sci Technol Lett* **8**, 792–798 (2021).

25. Mercier, E. *et al.* Municipal and neighbourhood level wastewater surveillance and

subtyping of an influenza virus outbreak. *Sci Rep* **12**, 15777 (2022).

26. Zheng, X. *et al.* Wastewater Surveillance Provides Spatiotemporal SARS-CoV-2

Infection Dynamics. *Engineering* **40**, 70–77 (2024).

27. Jarvie, M. M., Nguyen, T. N. T., Southwell, B. & Wright, D. Leveraging wastewater

surveillance to actively monitor Covid-19 community dynamics in rural areas with

reduced reliance on clinical testing. *Applied Research* **3**, e202400012 (2024).

28. Memish, Z. A. *et al.* Mass gatherings medicine: public health issues arising from mass

gathering religious and sporting events. *The Lancet* **393**, 2073–2084 (2019).

29. Suñer, C. *et al.* Association between two mass-gathering outdoor events and incidence

of SARS-CoV-2 infections during the fifth wave of COVID-19 in north-east Spain: A

population-based control-matched analysis. *Lancet Reg Health Eur* **15**, 100337 (2022).

30. Memish, Z. A. *et al.* Proactive surveillance for avian influenza H5N1 and other priority

pathogens at mass gathering events. *The Lancet Public Health* **9**, e350–e352 (2024).

31. Verovšek, T., Krizman-Matasic, I., Heath, D. & Heath, E. Site- and event-specific

wastewater-based epidemiology: Current status and future perspectives. *Trends in*

Environmental Analytical Chemistry **28**, e00105 (2020).

32. Toro, L. *et al.* Pathogen prioritisation for wastewater surveillance ahead of the Paris

2024 Olympic and Paralympic Games, France. *Eurosurveillance* **29**, 2400231 (2024).

33. Saied, A. A. *et al.* Wastewater surveillance as an epidemiological tool at mass gathering

events: A FIFA World Cup Qatar 2022 perspective and implication to other mass

gathering events. *International Journal of Surgery* **109**, 574 (2023).

34. Kitajima, M., Murakami, M., Iwamoto, R., Katayama, H. & Imoto, S. COVID-19

wastewater surveillance implemented in the Tokyo 2020 Olympic and Paralympic

Village. *Journal of Travel Medicine* **29**, taac004 (2022).

35. Nkambule, S., Street, R., Surujlal-Naicker, S., Johnson, R. & Mathee, A. Wastewater

surveillance for SARS-CoV-2 during a mass sporting event in the City of Cape Town,

Western Cape. *Front. Public Health* **12**, (2024).

36. El-Malah, S. S. *et al.* Leveraging wastewater surveillance for managing the spread of

SARS-CoV-2 and concerned pathogens during FIFA World Cup Qatar 2022. *Heliyon* **10**,

e30267 (2024).

37. Brighton, K., Fisch, S., Wu, H., Vigil, K. & Aw, T. G. Targeted community wastewater

surveillance for SARS-CoV-2 and Mpox virus during a festival mass-gathering event.

Science of The Total Environment **906**, 167443 (2024).

38. Sharma, A., Karande, M. & Zumla, A. Advancing sewage surveillance at mass gathering

events for reducing transmission of antimicrobial resistant bacterial pathogens. *Travel*

Medicine and Infectious Disease **54**, 102619 (2023).

39. Alexander, J., Hembach, N. & Schwartz, T. Evaluation of antibiotic resistance

dissemination by wastewater treatment plant effluents with different catchment areas

in Germany. *Sci Rep* **10**, 8952 (2020).

40. Karkman, A., Do, T. T., Walsh, F. & Virta, M. P. J. Antibiotic-Resistance Genes in Waste

Water. *Trends Microbiol* **26**, 220–228 (2018).

41. Choi, P. M. *et al*. Wastewater-based epidemiology biomarkers: Past, present and future.

TrAC Trends in Analytical Chemistry **105**, 453–469 (2018).

42. Keenum, I. *et al*. A framework for standardized qPCR-targets and protocols for

quantifying antibiotic resistance in surface water, recycled water and wastewater.

Critical Reviews in Environmental Science and Technology **52**, 4395–4419 (2022).

43. Hendriksen, R. S. *et al*. Global monitoring of antimicrobial resistance based on

metagenomics analyses of urban sewage. *Nat Commun* **10**, 1124 (2019).

44. Harrop, E. *et al*. Spatiotemporal Variability in Wastewater-Derived Carbapenem-

Resistance Genes from Diverse Municipal Sources in the Laurentian Great Lakes

Catchment. Preprint at https://doi.org/10.21203/rs.3.rs-8206721/v1 (2026).

45. Lynggaard, C., Gogarten, J. F. & Jokelainen, P. Integrating environment into One Health

approaches: Embracing the complexity. *One Health* **20**, 101021 (2025).

46. Greer, J. From Boardrooms to Block Parties: A Case Study on The Economic and

Cultural Transformation of the NFL Draft. *Sport Economics Research* **2**, 29–44 (2025).

47. Williams, K. T. W. & Hall, J. C. The Impact of the 2024 NFL Draft on Hotel Demand. *JRAP*

55, (2025).

48. Hutchinson, D. Here's how many people traveled at least 100 miles to attend NFL Draft

in Detroit. *WDIV* (2024).

49. NCEI Past Weather Tool | Drought.gov. https://www.drought.gov/data-maps-tools/ncei-

past-weather-tool.

50. CSO/SSO List - MiEnviro Portal.

https://mienviro.michigan.gov/ncore/external/overflow/list.

51. Rosario, K., Symonds, E. M., Sinigalliano, C., Stewart, J. & Breitbart, M. Pepper Mild

Mottle Virus as an Indicator of Fecal Pollution. *Appl Environ Microbiol* **75**, 7261–7267

(2009).

52. Maal-Bared, R. *et al*. Does normalization of SARS-CoV-2 concentrations by Pepper Mild

Mottle Virus improve correlations and lead time between wastewater surveillance and

clinical data in Alberta (Canada): comparing twelve SARS-CoV-2 normalization

approaches. *Sci Total Environ* **856**, 158964 (2023).

53. R Core Team. *R: A Language and Environment for Statistical Computing*. (R Foundation

for Statistical Computing, Vienna, Austria, 2025).

54. Wickham, H. *et al*. Welcome to the tidyverse. *Journal of Open Source Software* **4**, 1686

(2019).

55. McBride, G. & others. A proposal for strength-of-agreement criteria for Lin's

concordance correlation coefficient. *NIWA client report: HAM2005-062* **45**, 307–310

(2005).

56. Bulfone, T. C. *et al*. Outdoor Activities Associated with Lower Odds of SARS-CoV-2

Acquisition: A Case–Control Study. *Int J Environ Res Public Health* **19**, 6126 (2022).

57. Lessler, J. *et al*. Incubation periods of acute respiratory viral infections: a systematic

review. *The Lancet Infectious Diseases* **9**, 291–300 (2009).

58. Alandijany, T. A. Respiratory viral infections during Hajj seasons. *Journal of Infection*

and Public Health **17**, 42–48 (2024).

59. Boyd, S. E., Holmes, A., Peck, R., Livermore, D. M. & Hope, W. OXA-48-Like β-

Lactamases: Global Epidemiology, Treatment Options, and Development Pipeline.

Antimicrobial Agents and Chemotherapy **66**, e00216-22 (2022).

60. Kajova, M., Khawaja, T., Kainulainen, K. & Kantele, A. Travel-associated

carbapenemase-producing Enterobacterales: Epidemiology and risk factors in a

Finnish cohort. *Travel Medicine and Infectious Disease* **68**, 102926 (2025).

61. Pitout, J. D. D., Peirano, G., Kock, M. M., Strydom, K.-A. & Matsumura, Y. The Global

Ascendency of OXA-48-Type Carbapenemases. *Clin Microbiol Rev* **33**, e00102-19

(2019).

62. Peirano, G. & Pitout, J. D. D. Rapidly spreading Enterobacterales with OXA-48-like

carbapenemases. *Journal of Clinical Microbiology* **63**, e01515-24 (2025).

63. Auguet, O. *et al.* Sewers as potential reservoirs of antibiotic resistance. *Science of The*

Total Environment **605–606**, 1047–1054 (2017).

64. Nahum, Y., Muhvich, J., Morones-Ramirez, J. R., Casillas-Vega, N. G. & Zaman, M. H.

Biofilms as potential reservoirs of antimicrobial resistance in vulnerable settings. *Front*

Public Health **13**, 1568463 (2025).

65. Haenelt, S. *et al.* The fate of sulfonamide resistance genes and anthropogenic pollution

marker intI1 after discharge of wastewater into a pristine river stream. *Front Microbiol*

14, 1058350 (2023).

66. Gillings, M. R. *et al.* Using the class 1 integron-integrase gene as a proxy for

anthropogenic pollution. *ISME J* **9**, 1269–1279 (2015).

67. Rushford, C. *et al*. Untargeted longitudinal ultra deep metagenomic sequencing of wastewater provides a comprehensive readout of expected and unexpected viral pathogens. 2025.10.27.25338874 Preprint at https://doi.org/10.1101/2025.10.27.25338874 (2025).

68. Kaeseberg, T. *et al*. Hot spots of antibiotic tolerant and resistant bacterial subpopulations in natural freshwater biofilm communities due to inevitable urban drainage system overflows. *Environmental Pollution* **242**, 164–170 (2018).

69. Morina, J. C. & Franklin, R. B. Drivers of Antibiotic Resistance Gene Abundance in an Urban River. *Antibiotics (Basel)* **12**, 1270 (2023).

70. Pao, L. T., Tashani, M., King, C., Rashid, H. & Khatami, A. Antimicrobial Resistance Associated with Mass Gatherings: A Systematic Review. *Trop Med Infect Dis* **10**, 2 (2024).

71. Wang, C., Myshkevych, Y., Wang, T., Monjed, M. K. & Hong, P.-Y. Wastewater surveillance unveils the impact of mass gatherings on antimicrobial resistance after the COVID-19 pandemic in Saudi Arabia. *Nat Water* **3**, 693–703 (2025).

72. Li, Z. *et al*. The Environmental Lifecycle of Antibiotics and Resistance Genes: Transmission Mechanisms, Challenges, and Control Strategies. *Microorganisms* **13**, 2113 (2025).

73. Ishii, S. *et al*. Microfluidic Quantitative PCR for Simultaneous Quantification of Multiple Viruses in Environmental Water Samples. *Applied and Environmental Microbiology* **80**, 7505–7511 (2014).

74. Ishii, S., Segawa, T. & Okabe, S. Simultaneous Quantification of Multiple Food- and

Waterborne Pathogens by Use of Microfluidic Quantitative PCR. *Appl Environ Microbiol*

79, 2891–2898 (2013).

75. Sekovanić, A. *et al.* Pre-amplification as a method for improvement of quantitative RT-

PCR analysis of circulating miRNAs. *Biochem Med (Zagreb)* **31**, 010901 (2021).

76. Wang, C., Myshkevych, Y., Wang, T., Monjed, M. K. & Hong, P.-Y. Wastewater

surveillance unveils the impact of mass gatherings on antimicrobial resistance after the

COVID-19 pandemic in Saudi Arabia. *Nat Water* **3**, 693–703 (2025).

77. Germovsek, E., Barker, C. I. & Sharland, M. What do I need to know about

aminoglycoside antibiotics? *Arch Dis Child Educ Pract Ed* **102**, 89–93 (2017).

78. Krause, K. M., Serio, A. W., Kane, T. R. & Connolly, L. E. Aminoglycosides: An Overview.

Cold Spring Harb Perspect Med **6**, a027029 (2016).

79. Holbrook, S. Y. L. & Garneau-Tsodikova, S. Evaluation of Aminoglycoside and

Carbapenem Resistance in a Collection of Drug-Resistant Pseudomonas aeruginosa

Clinical Isolates. *Microb Drug Resist* **24**, 1020–1030 (2018).

80. Zhao, S. *et al.* Epidemiology of and risk factors for mortality due to carbapenemase-

producing organisms (CPO) in healthcare facilities. *J Hosp Infect* **110**, 184–193 (2021).

81. Bonomo, R. A. *et al.* Carbapenemase-Producing Organisms: A Global Scourge. *Clin*

Infect Dis **66**, 1290–1297 (2018).

82. Carbapenemase Producing Enterobacteriaceae (CPE). *Public Health Ontario*

http://www.publichealthontario.ca/en/Diseases-and-Conditions/Health-Care-

Associated-Infections/CPE.

83. Shaikh, S., Fatima, J., Shakil, S., Rizvi, S. Mohd. D. & Kamal, M. A. Antibiotic resistance

and extended spectrum beta-lactamases: Types, epidemiology and treatment. *Saudi J*

Biol Sci **22**, 90–101 (2015).

84. Araten, A. H. *et al.* Cephalosporin resistance, tolerance, and approaches to improve

their activities. *J Antibiot* **77**, 135–146 (2024).

85. Park, S. H. Third-generation cephalosporin resistance in gram-negative bacteria in the

community: a growing public health concern. *Korean J Intern Med* **29**, 27–30 (2014).

86. Liguori, K. *et al.* Antimicrobial Resistance Monitoring of Water Environments: A

Framework for Standardized Methods and Quality Control. *Environ Sci Technol* **56**,

9149–9160 (2022).

87. 2021 AWaRe classification. https://www.who.int/publications-detail-redirect/2021-

aware-classification.

88. Hooper, D. C. & Jacoby, G. A. Mechanisms of drug resistance: quinolone resistance.

Ann N Y Acad Sci **1354**, 12–31 (2015).

89. Camins, B. C. *et al.* The clinical impact of fluoroquinolone resistance in patients with E

coli bacteremia. *J Hosp Med* **6**, 344–349 (2011).

90. Salah, F. D. *et al.* Distribution of quinolone resistance gene (qnr) in ESBL-producing

Escherichia coli and Klebsiella spp. in Lomé, Togo. *Antimicrobial Resistance & Infection*

Control **8**, 104 (2019).

91. Peirano, G. & Pitout, J. D. D. Fluoroquinolone-Resistant Escherichia coli Sequence Type

131 Isolates Causing Bloodstream Infections in a Canadian Region with a Centralized

Laboratory System: Rapid Emergence of the H30-Rx Sublineage. *Antimicrobial Agents and Chemotherapy* **58**, 2699–2703 (2014).

92. Yang, P. *et al.* Association between the rate of fluoroquinolones-resistant gram-negative bacteria and antibiotic consumption from China based on 145 tertiary hospitals data in 2014. *BMC Infectious Diseases* **20**, 269 (2020).

93. Kibwana, U. O. *et al.* Fluoroquinolone resistance among fecal extended spectrum βeta lactamases positive Enterobacterales isolates from children in Dar es Salaam, Tanzania. *BMC Infectious Diseases* **23**, 135 (2023).

94. Selim, S. Mechanisms of gram-positive vancomycin resistance (Review). *Biomed Rep* **16**, 7 (2022).

95. Patel, S., Preuss, C. V. & Bernice, F. Vancomycin. in *StatPearls* (StatPearls Publishing, Treasure Island (FL), 2024).

96. Mühlberg, E. *et al.* Renaissance of vancomycin: approaches for breaking antibiotic resistance in multidrug-resistant bacteria. *Can. J. Microbiol.* **66**, 11–16 (2020).

97. Patel, P. H. & Hashmi, M. F. Macrolides. in *StatPearls* (StatPearls Publishing, Treasure Island (FL), 2024).

98. Bengtsson-Palme, J. *et al.* Industrial wastewater treatment plant enriches antibiotic resistance genes and alters the structure of microbial communities. *Water Research* **162**, 437–445 (2019).

99. Wang, H. *et al.* Characterization of resistance genes and plasmids from sick children caused by Salmonella enterica resistance to azithromycin in Shenzhen, China. *Front. Cell. Infect. Microbiol.* **13**, (2023).

100.   Schwartz, K. L. *et al.* Regional variability in outpatient antibiotic use in Ontario,

Canada: a retrospective cross-sectional study. *CMAJ Open* **6**, E445–E452 (2018).

101.   Siddiqui, A. H. & Koirala, J. Methicillin-Resistant Staphylococcus aureus. in

StatPearls (StatPearls Publishing, Treasure Island (FL), 2024).

102.   Snitser, O. *et al.* Ubiquitous selection for mecA in community-associated MRSA

across diverse chemical environments. *Nat Commun* **11**, 6038 (2020).

103.   Tyson, G. H., Li, C., Hsu, C.-H., Bodeis-Jones, S. & McDermott, P. F. Diverse

Fluoroquinolone Resistance Plasmids From Retail Meat E. coli in the United States.

Front. Microbiol. **10**, (2019).

104.   Ovung, A. & Bhattacharyya, J. Sulfonamide drugs: structure, antibacterial property,

toxicity, and biophysical interactions. *Biophys Rev* **13**, 259–272 (2021).

105.   Vester, B. The *cfr* and cfr-like multiple resistance genes. *Research in Microbiology*

169, 61–66 (2018).

106.   Mirnezami, M., Ranjbar, R., Niakan, M. & Ahmadi, M. H. Frequency of Antimicrobial

Resistance and Class 1 and 2 Integrons in Escherichia Coli Strains Isolated from

Urinary Tract Infections. *Iran J Pharm Res* **19**, 282–287 (2020).

107.   Racewicz, P. *et al.* Prevalence and characterisation of antimicrobial resistance

genes and class 1 and 2 integrons in multiresistant Escherichia coli isolated from

poultry production. *Sci Rep* **12**, 6062 (2022).

108.   Stachler, E. *et al.* Quantitative CrAssphage PCR Assays for Human Fecal Pollution

Measurement. *Environ. Sci. Technol.* **51**, 9146–9154 (2017).

109.   Langeveld, J. *et al.* Normalisation of SARS-CoV-2 concentrations in wastewater: The

use of flow, electrical conductivity and crAssphage. *Sci Total Environ* **865**, 161196

(2023).

862