An in silico framework for evaluating PRS-guided prognostic enrichment in clinical trial design

Ruoyi Cai[1,2], Joshua Gillard[1,2], Sherry Yang[3], Samvel B. Gasparyan[4], Ying Lu[1], Lu Tian[1], Ola Vedin[5], Euan A. Ashley[†1,2]*, Manuel A. Rivas[†1]*, Jack W. O'Sullivan[†1,2]*

[1]Department of Biomedical Data Science, Stanford University, Stanford, CA, USA, 94305

[2]Department of Medicine, Stanford University, Stanford, CA, USA, 94305

[3]Department of Bioengineering, Stanford University, Stanford, CA, USA, 94305

[4]Late Stage Development, BioPharmaceuticals Research & Development, AstraZeneca, Boston, MA, USA

[5]Late Stage Development, BioPharmaceuticals Research & Development, AstraZeneca, Gothenburg, Sweden

*Work jointly supervised.

†Corresponding authors: jackos@stanford.edu, euan@stanford.edu, mrivas@stanford.edu

Abstract

Background: Clinical trials are essential for therapeutic development but increasingly face challenges due to imprecise inclusion criteria, leading to low event rates and the need for large sample sizes. This inefficiency makes modern trials costly and time-consuming. Despite the availability of extensive clinical, genomic, and biological data, current trial enrollment strategies do not fully leverage this information. Incorporating genomic information into trial design could enable risk-based participant enrichment by preferentially enrolling individuals with higher disease risk, thereby increasing event rates and improving trial efficiency.

Methods: In this study, we developed an in silico framework for evaluating prognostic enrichment guided by polygenic risk scores (PRS) in clinical trial design using genomic and electronic health record data from large-scale biobanks. Naturally occurring protective genetic variants were used as analogs of therapeutic interventions, with variant carriers treated as 'treatment' arms and non-carriers as 'control' arms. We compared unenriched designs, in which carriers and non-carriers were drawn from the full population, against PRS-enriched designs in which both arms were restricted to participants in the upper 75%, 50%, or 25% of the PRS distribution, respectively. Across these four designs, we quantified disease prevalence, statistical power, sample size requirements, and time-to-event accrual.

Results: We applied this approach to the UK Biobank using three model gene–disease pairs: the protective variant p.Arg46Leu in *PCSK9* for coronary artery disease (CAD), p.Gln175His in *ANGPTL7* for glaucoma, and p.Arg381Gln in *IL23R* for inflammatory bowel disease (IBD). Across all three disease contexts, PRS-enriched designs increased disease prevalence, improved empirical power, and accelerated event accrual relative to unenriched cohorts. At 80% power, restricting enrollment to the upper 25% of the PRS distribution reduced required per-arm sample sizes by approximately 60% for CAD-PCSK9 and 78% for IBD-IL23R. Consistent reductions in time-to-event were also observed across enriched strata, suggesting that PRS-enriched trials could achieve target event counts with both smaller sample sizes and shorter follow-up. However, for glaucoma-ANGPTL7, the most restrictive threshold did not yield additional gains over moderate enrichment, as reduced sample size attenuated the detectable difference between arms. These results highlight the need to balance enrichment for higher-risk participants against retaining a sufficient eligible population, and underscore that optimal PRS thresholds are disease-context dependent.

Conclusions: These findings establish a generalizable, data-driven framework for prospectively evaluating PRS-guided prognostic enrichment prior to trial initiation. In general, PRS-guided study designs lead to improved empirical power, lower required sample sizes, and faster trials. As population-scale genomic data become increasingly available within healthcare systems and biobanks, this framework provides a scalable foundation for integrating genetic risk information into clinical trial design.

Introduction

Despite being the gold standard for evaluating therapeutic interventions, traditional randomized controlled trials (RCTs) are often constrained by low event rates, limiting statistical power to detect clinically relevant treatment effects.[1–4] Currently, only a small fraction of participants may experience the primary endpoint within the study follow-up period. A systematic review of 344 contemporary cardiovascular trials reported a median observed primary endpoint event rate of only 9.0%.[5] In addition, improvements in background medical therapy and preventive care have further reduced disease risk in the general population.[4,6,7] Accruing a sufficient number of clinical events in trials typically requires large sample sizes and prolonged follow-up, substantially increasing trial cost and duration.[4,7–9] Insufficient events lead to null trials, and thus uncertainty whether the medication is beneficial, or if the trial was too small or too short.

To address these challenges, prognostic enrichment strategies have been proposed to improve trial power and efficiency by selectively recruiting individuals with higher disease risk, thereby increasing the expected number of clinical events of interest for a given sample size.[10–15] However, it remains that current trials use no or few variables to identify and include high risk participants. With the advancement of sequencing technologies, genetic data have become powerful resources for risk prediction at the population level. In particular, polygenic risk scores (PRS) combine the effects of many common genetic variants across the genome to capture an individual's inherited predisposition to disease.[16] PRS have demonstrated strong predictive value for a wide range of complex traits and diseases, and can be calculated from birth.[17] By identifying individuals at higher genetic risk, PRS can enrich trial populations for clinical events, thereby increasing statistical power and reducing the required sample size.[18]

Previous studies have highlighted the potential of PRS to stratify disease risk, enhance clinical risk scores, and have suggested their use for prognostic enrichment in clinical trials.[18–24] For example, retrospective analyses of completed cardiovascular outcome trials of statins and PCSK9 inhibitors have consistently demonstrated that participants in high-PRS subgroups have a higher event rate and receive a greater relative and absolute clinical benefit.[19,23,24] A post hoc analysis of the FOURIER trial by Fahed et al. showed that restricting enrollment to individuals in the highest genetic risk strata could theoretically achieve the same level of statistical power with a substantially reduced sample size.[18] More recently, German et al. incorporated PRS into target trial emulations using observational data and showed that restricting enrollment to participants with higher PRS in emulated cohorts could achieve the required number of events with a smaller sample size than the original trials.[20]

While these studies demonstrate that PRS can meaningfully differentiate risk and identify subgroups with higher event rates, they were retrospective analyses embedded within existing or emulated trials. As such, they evaluated how PRS would have stratified risk within previously conducted trials and were therefore inherently tied to specific completed studies, limiting their applicability for proactively designing enrichment strategies for new trials. Furthermore, prior

work generally focused on single diseases within particular trial settings, rather than
systematically comparing PRS-based enrichment across multiple gene–disease contexts or
alternative risk thresholds. As a result, existing work does not establish a generalizable
framework for informing future trial design by quantifying the degree of enrichment achievable
under various PRS thresholds through systematic evaluation of sample size requirements,
statistical power, and expected event accrual.

To address this gap between the theoretical promise of PRS-guided enrichment and its practical
application in prospective trial design, we propose an in silico framework for quantitatively
evaluating PRS-based prognostic enrichment using large-scale biobanks with linked genomic
and electronic health record data (Figure 1). Our approach leverages naturally occurring
protective genetic variants as human genetic analogs of therapeutic inhibition targets, enabling
treatment-like effects to be studied without interventional trials. For example, loss-of-function
variants in *PCSK9*, such as p.Arg46Leu, lower low-density lipoprotein cholesterol and reduce
the risk of coronary artery disease (CAD).[25,26] Such protective loss-of-function variants have also
provided the biological foundation for therapeutic target development, and PCSK9 inhibition is
now a well-established clinical strategy.[27,28] The random distribution of carriers and non-carriers
of these naturally occurring protective variants thus creates a "natural experiment," in which
carriers serve as a genetic proxy for the treatment arm and non-carriers for the control arm.

Using this framework, we stratified participants by PRS quantiles and systematically assessed
four key trial design parameters: disease prevalence in carriers versus non-carriers across PRS
strata, required sample sizes to achieve target statistical power, empirical power under fixed
sample sizes, and prospective event accrual in disease-free cohorts followed over time. Across
multiple disease–gene examples, we found that PRS-based enrichment consistently increased
disease burden and improved power relative to unenriched designs, although the magnitude of
benefit varied by disease context. This approach enables data-driven evaluation of how PRS-
based enrichment increases the power to detect treatment effects leveraging protective variants
with well-established biological effects, operating entirely using existing population data and
feasible prior to trial initiation. Together, these analyses illustrate the potential of human genetics
to inform and optimize clinical trial design.

Figure 1. An in silico framework for evaluating PRS-guided prognostic enrichment in clinical trial
design using protective genetic variants as therapeutic proxies. (A) Large-scale biobanks with linked
genomic and electronic health record data provide the foundation for analysis. (B) Naturally occurring
protective variants define a genetic natural experiment in which carriers serve as proxies for therapeutic
inhibition and non-carriers serve as controls. (C) Within the full cohort, participants are stratified by
polygenic risk score (PRS) thresholds to quantitatively evaluate enrichment across alternative risk cutoffs.
(D) Analyses on disease prevalence, required sample size, statistical power, and event accrual within
PRS-defined subgroups inform key design parameters for prospective clinical trials.

Methods

Study population

The study included White British individuals from the UK Biobank.[29] Individuals were classified as White British based on genetic ethnicity grouping reported in UK Biobank. Participants with missing information on the relevant genetic variants, disease outcomes, or covariates were excluded from downstream analyses.

Definition of carrier status

Whole-exome sequencing data from the UK Biobank were used to define carrier status for protective genetic variants.[29,30] For each gene, individuals carrying at least one copy of the protective variant were classified as variant carriers, while non-carriers served as the comparison group. Within the in silico framework, carriers and non-carriers were treated as genetically defined "treatment" and "control" arms, respectively, reflecting the treatment-like effects of these biologically validated variants.

Definition of disease endpoints

Disease phenotypes were derived from linked hospital inpatient electronic health record data in the UK Biobank and defined using International Classification of Diseases, 10th Revision (ICD-10) diagnosis codes recorded in primary or secondary diagnostic positions.[29] The ICD-10 code definitions of all disease endpoints considered in this study were summarized in Table S1. Disease status was defined as a binary indicator reflecting the presence of at least one relevant ICD-10 diagnosis code recorded during UK Biobank follow-up.

Polygenic risk score

Disease-specific polygenic risk scores (PRS) were obtained from the UK Biobank PRS released
by Thompson et al. (2022).[31,32] Within the White British study population, we defined three PRS
strata, "Top 75%", "Top 50%", and "Top 25%" using the 25th, 50th, and 75th percentiles of
each disease-specific PRS, respectively. Individuals above each percentile threshold q were
assigned to the corresponding "Top $(1 - q)$%" PRS group, and the full study population, labeled
as the "All" group, served as the reference group for comparisons with the enriched groups.

Selection of disease-gene examples for implementing the PRS-guided in silico framework

The workflow used to identify disease–gene examples appropriate for implementing the PRS-
guided prognostic enrichment is summarized in Figure S1. We began with a structured literature
search to identify candidate disease–gene pairs. Eligible pairs were required to involve well-
characterized monogenic variants demonstrating protective effects against the disease of interest,
supported by prior genetic or functional evidence.

For each disease–gene pair with verified biological foundation, we evaluated three empirical
features to determine whether sufficient statistical information is available in the finite dataset to
evaluate PRS-guided enrichment framework. The first feature was the presence of a genetic loss
of function variant in the source population, assessed by estimating the odds ratio comparing
carriers to non-carriers leading to a significant difference in disease occurrence. This was
quantified by examining disease occurrence between carriers (of the variant) vs non-carriers via
the Wald statistic and corresponding p-value. Variants demonstrating a statistically significant
difference (using nominal level of 0.05) in disease risk between carriers and non-carriers in the
full cohort were considered for further evaluation in our PRS-guided trial design framework.

The second feature was whether the PRS meaningfully stratified disease burden. This was
evaluated using the odds ratio per standard deviation of PRS and by examining disease
prevalence across progressively enriched PRS strata. A statistically significant association
between PRS and disease, accompanied by a generally increasing pattern of disease prevalence
across strata, indicates that restriction to higher PRS groups shifts disease risk in the intended
direction.

The third feature assessed whether the difference in disease risk between carriers and non-
carriers remained statistically significant within PRS-enriched strata. Specifically, we evaluated
whether the association observed in the overall population persisted when restricting to
individuals above the 25%, 50%, and 75% thresholds of the PRS distribution using the stratum-
specific carrier odds ratio, its Wald statistic, and corresponding p-value. Because restriction to
higher PRS strata reduces the effective sample size and the number of carrier–case observations,
the variance of the carrier effect estimate may increase substantially after restriction, and the
carrier effect may no longer reach statistical significance in enriched subgroups even if it is
significant in the full cohort. This typically reflects insufficient sample sizes of enriched PRS
strata and suggests that overly aggressive restriction to high PRS strata is unlikely to yield
meaningful efficiency gains.

We summarize the three empirical features described above using a joint information score,
denoted by Ψ, to provide a quantitative summary of the empirical support for applying the PRS-
guided enrichment framework to each candidate disease–gene pair:

$$\Psi = I_{\{p_{full}<\alpha\}} \times ln\,(OR_{PRS})^{\,2} \times Z_{min}^2.$$

This metric combines three components, corresponding to the presence of a carrier-non-carrier
difference in disease prevalence, the magnitude of polygenic risk stratification, and the strength
of the carrier association within enriched PRS strata, respectively. The first component,
$I_{\{p_{full}<\alpha\}}$, is an indicator function equal to one when the p-value associated with the carrier effect
in the full cohort (p_{full}) satisfies the nominal significance threshold α, and zero otherwise. The
second component, $ln\,(OR_{PRS})^{\,2}$, represents the squared log-odds ratio per standard deviation of
PRS estimated in the full cohort, capturing the magnitude of the association between PRS and
disease prevalence. In the third term, Z_{min} is the minimum Wald Z-statistic for testing the carrier
effect across the evaluated PRS-enriched strata, which reflects the smallest stratum-specific
signal to noise ratio for the carrier effect under increasing PRS restriction.

We note that these features focus on whether the available sample size and event counts allow
stable estimation of the protective variant effect under PRS restriction, rather than on therapeutic
importance or magnitude of biological effects. The latter considerations mainly relate to the
operational implications and translational impact but do not determine whether the framework
can be assessed with adequate statistical power in the observed data.

Evaluation of PRS-guided prognostic enrichment

We first examine disease prevalence among protective carriers and non-carriers across PRS-
defined subgroups (top 25th, 50th, 75th). Within each PRS-defined stratum, disease prevalence
was estimated separately for carriers and non-carriers, and compared.

We next estimated sample size requirements and statistical power based on disease prevalence
across PRS-enriched groups to quantify how enrichment by PRS influences trial design
parameters. To estimate disease prevalence in each arm for sample size calculations, we fit a
logistic regression model of disease outcome on PRS, carrier status of the protective variant, and
their interaction (PRS × carrier), adjusting for age, sex, and the first three genetic principal
components. For each observed individual, the carrier disease probability was estimated from the
fitted logistic regression model assuming the individual is a carrier, while maintaining their
observed PRS, age, sex and genetic principal components unchanged. Then for each PRS-
defined subgroup, we obtained the average carrier disease probabilities across individuals in the
subgroup as the average carrier disease prevalence for the subgroup. Similarly, we calculated the
average non-carrier disease prevalence for each PRS-defined subgroup. These population-level
disease prevalences were then used for sample size calculation in PRS-based enrichment study
design, whose target population is a PRS-defined subgroup.

Sample sizes were calculated based on standard two-proportions Z test at a two-sided 5%
significance level across power targets ranging from 60% to 85%. Specifically, the required
sample size per group was given by

$$n = \frac{(Z_{0.975} + Z_{\mathrm{power}})^2 [p_1(1-p_1) + p_0(1-p_0)]}{(p_1 - p_0)^2},$$

where Z_α is the α-th quantile of a standard normal, and p_1 and p_0 are carrier and non-carrier
disease prevalences for the PRS-defined subgroup, respectively. To further assess the
recruitment feasibility and operational implications of PRS-based enrichment, we additionally
estimated the required screening sample size and trial cost using a streamlined cost framework
incorporating participant screening, PRS ascertainment, and trial execution costs based on the
calculated sample size. Details of the screening and cost assumptions and calculations are
provided in the Supplementary Methods.

Finally, we conducted simulations to empirically quantify statistical power achievable under
different combinations of sample size and PRS-based enrichment. Within each PRS-defined
subgroup, we constructed synthetic datasets with 100, 200, 500, and 1000 individuals per arm
(carriers vs non-carriers) by sampling the corresponding number of individuals with replacement
from carriers and non-carriers, separately. Each generated dataset was analyzed using logistic
regression with disease status as the binary outcome and carrier status as the independent
variable of the primary interest adjusting for age, sex, and three genetic principal components.
This process was repeated 1,000 times for each given combination of sample size and PRS-based
enrichment, and the empirical power was calculated as the proportion of replicates in which the
carrier effect reached nominal statistical significance. To be conservative, when fewer than five
disease cases were observed in either group within a resampled cohort, we recorded the
comparison as a non-rejection without formally testing the carrier effect to avoid unstable
estimates arising from extremely small numbers of events.

Simulation of prospective event accumulation under PRS-guided enrichment

The PRS-based enrichment can not only increase the overall disease prevalence but also reduce
the length of follow-up time to observe a sufficient number of disease events. To quantify how
PRS-guided enrichment influences the waiting time to incidence disease, we further analyzed
individuals who were free of the relevant disease at baseline, defined as the date of UK Biobank
enrollment. Participants were followed prospectively from baseline until incident disease or
censoring. Incident disease was defined as the first recorded diagnosis after baseline, and
censoring was defined at 10 years of follow-up.

Restricted mean survival time (RMST), defined as the area under the survival curve up to a
prespecified truncation time, was used to summarize average event-free time over the follow-up
horizon. Within each PRS-enriched subgroup and for each specific target sample size, we
generated a synthetic dataset by sampling the corresponding number of individuals with
replacement from the baseline disease-free cohort. Then, the RMST up to 10-years was

estimated based on the synthetic data. This process was repeated 1,000 times and the empirical
distribution of estimated RMST was obtained. We then compared the distribution of RMST
across PRS-enriched groups to evaluate how PRS-based recruitment influences the rate of
prospective event accrual.

Results

Selection of candidate disease-gene pairs

Through our structured literature review, we first identified loss of function variants that confer
natural protection against disease. The identified examples included *PCSK9* p.Arg46Leu for
coronary artery disease (CAD-PCSK9),[25–28] *ANGPTL7* p.Gln175His for glaucoma (glaucoma-
ANGPTL7),[33–36] *IL23R* p.Arg381Gln for inflammatory bowel disease (IBD-IL23R),[37–40] *IL6R*
p.Asp358Ala for rheumatoid arthritis (RA-IL6R),[41–44] *SLC30A8* p.Arg138Ter for type 2 diabetes
(T2D-SLC30A8),[45–48] and *APOC3* p.Ala43Thr for coronary artery disease (CAD-APOC3).[49–52]

We then calculated the joint information score for each disease-gene pair to determine if it met
our criteria for inclusion in our framework. To do this, first we obtained GRCh38 genomic
coordinates for the protective variant in each disease-gene pair, as well as the ICD-10 code
definition of the corresponding disease endpoint are summarized in Table S1. Second, we used
the UK Biobank PRS release to obtain PRS for coronary artery disease, glaucoma, type 2
diabetes, and rheumatoid arthritis. Because inflammatory bowel disease was defined in our study
as a composite endpoint including Crohn's disease and ulcerative colitis, and no unified IBD
PRS was available, we constructed an IBD PRS proxy by taking the maximum of the CD PRS
and UC PRS for each individual. This definition aligns the genetic risk stratification with the
composite disease endpoint and prioritizes sensitivity to genetic risk for either IBD subtype.

These data allowed us to calculate joint information scores for each disease-gene pair, which are
presented in Figure 2 and detailed in Table S2. Three disease–gene pairs meet our criteria for
inclusion (score > 0) and were selected for full in silico enrichment analysis: CAD-PCSK9,
glaucoma-ANGPTL7, and IBD-IL23R. PCSK9-CAD and IL23R-IBD satisfied all selection
criteria. In the full cohort, both showed a statistically significant odds ratio comparing carriers
with non-carriers, as well as a strong association between PRS and disease risk. The carrier
versus non-carrier odds ratio remained detectable across progressively enriched PRS strata.
These features are reflected in high joint information scores of 4.57 for CAD-PCSK9 and 3.11
for IBD-IL23R, supporting the use of these examples for evaluating PRS-guided prognostic
enrichment within our framework.

Glaucoma-ANGPTL7 met baseline statistical criteria and showed meaningful stratification of
disease risk by PRS. However, the difference in disease prevalence between carriers and non-
carriers was attenuated and no longer statistically significant within the most restrictive Top 25%
PRS stratum. Consistent with this pattern, the joint information score was lower at 0.45,

reflecting more limited within-stratum information under aggressive restriction. We retained this
example to illustrate how limited carrier–case observations under stringent enrichment thresholds
can influence the performance of PRS-guided enrichment.

In contrast, RA-IL6R, T2D-SLC30A8, and CAD-APOC3 failed to demonstrate a statistically
significant odds ratio comparing carriers with non-carriers in the full cohort and therefore did not
meet the minimum criteria for enrichment evaluation in our dataset, although protective effects
have been reported in previous studies. For RA-IL6R, the estimated carrier vs. non-carrier OR
was 1.01 (95% CI: [0.97, 1.06]), consistent with the moderate effect size of the common variant
p.Asp358Ala.[42,43] For T2D-SLC30A8 and CAD-APOC3, the wide confidence intervals (T2D-
SLC30A8: OR 1.11 [0.46–2.65]; CAD-APOC3: OR 1.24 [0.79–1.93]) likely reflect limited
statistical power due to small numbers of variant carriers in the study population (66 carriers of
SLC30A8 p.Arg138Ter and 199 carriers of *APOC3* p.Ala43Thr). Correspondingly, the joint
information scores of these examples were all zero, indicating insufficient signal for enrichment
evaluation. These examples were therefore not pursued for further evaluation under PRS-guided
prognostic enrichment.

Figure 2. Evaluation of candidate disease–gene pairs for PRS-guided in silico enrichment. (A) Joint
information score Ψ for each disease–gene example. (B) Carrier versus non-carrier odds ratios (ORs)
with 95% confidence intervals estimated in the full cohort. (C) Odds ratio per standard deviation of PRS
estimated in the full cohort. (D) Heatmap of within-stratum carrier association p-values across
progressively enriched PRS strata (Top 75%, Top 50%, and Top 25%), shown as $-log_{10}(p)$.

Effects of PRS and protective variant carrier status on disease prevalence

We analyzed 393,610 White British participants from the UK Biobank to evaluate the effects of protective genetic variants across PRS quantiles on disease prevalence for the three gene–disease pairs. The CAD–PCSK9 analysis included 41,697 CAD cases and 13,717 carriers of the *PCSK9* p.Arg46Leu variant. The glaucoma–ANGPTL7 analysis included 12,092 glaucoma cases and 6,314 carriers of the *ANGPTL7* p.Gln175His variant. The IBD-IL23R analysis included 6,172 IBD cases (2,436 Crohn's disease and 4,396 ulcerative colitis) and 50,633 carriers of *IL23R* p.Arg381Gln.

Across all three disease contexts, carriers of the protective variants exhibited lower disease prevalence than non-carriers within each PRS stratum (Figure 3). In all examples, disease prevalence increased monotonically with higher PRS quantiles in both carriers and non-carriers, consistent with the expected positive association between polygenic risk and disease burden. At more stringent PRS thresholds, subgroup sample sizes reduced substantially, leading to larger uncertainty and reduced power to detect differences. For the glaucoma–ANGPTL7 example, given only 72 cases among 1,530 carriers in the Top 25% PRS stratum, the difference in disease prevalence between carriers and non-carriers was not statistically significant.

Figure 3. Disease prevalence in carriers versus non-carriers of protective variants across PRS strata for CAD–PCSK9, glaucoma–ANGPTL7, and IBD–IL23R. Numbers denote case counts and total sample sizes in each subgroup. P-values correspond to two-sided two-proportion tests comparing carriers and non-carriers within each stratum.

Logistic regression models adjusting for age, sex, and genetic principal components confirmed that carrier status was associated with lower disease risk across all three gene–disease pairs. For coronary artery disease, the *PCSK9* p.Arg46Leu variant was associated with an odds ratio of 0.878 (p = 0.003). For glaucoma, the *ANGPTL7* p.Gln175His variant was associated with an odds ratio of 0.760 (p = 0.004). For inflammatory bowel disease, the *IL23R* p.Arg381Gln variant was associated with an odds ratio of 0.913 (p = 0.049). In all three examples, PRS was strongly associated with disease risk (CAD PRS: OR = 1.438, p < 1 × 10^{-16}; glaucoma PRS: OR = 1.843, p < 1 × 10^{-16}; IBD PRS: OR = 1.85, p < 1 × 10^{-16}). Complete model estimates are provided in Tables S3–S5, and model-predicted disease prevalence across PRS is shown in Figure S2.

Sample size requirements and operational implications of PRS-enriched designs

Using model-based predictions of disease prevalence within each PRS stratum, we derived the minimum per-arm sample sizes required to achieve 60–85% statistical power under a two-sided 5% significance level across PRS-enriched subgroups (Figure 4). Across all three gene–disease pairs, restricting recruitment to higher PRS strata reduced required sample sizes. However, both the estimated per-arm sample size required in the unenriched population and the magnitude of reduction achieved under PRS enrichment differed substantially across disease contexts.

At 80% power, the CAD–PCSK9 example required 6,099 participants per arm in the full cohort. Restricting recruitment to the top 25% of the PRS distribution reduced this requirement to 2,383 per arm, representing an approximate 60% reduction. For glaucoma–ANGPTL7, the required per-arm sample size at 80% power was 8,404 in the full cohort and 5,853 in the Top 25% PRS group, corresponding to a more modest reduction of approximately 30%. In contrast, the IBD–IL23R example required substantially larger absolute sample sizes due to lower disease prevalence. At 80% power, 51,710 participants per arm were required in the full cohort, compared with 11,089 in the Top 25% PRS group. Despite the larger absolute numbers, this represented the greatest relative reduction (78%) among the three examples.

Projected total trial cost followed similar patterns (Figure S3). Under a general assumption that sequencing is required to obtain PRS, restriction to the Top 25% PRS group reduced projected total cost by 41.8% for CAD–PCSK9 and 69.5% for IBD–IL23R relative to the unenriched design. In glaucoma–ANGPTL7, however, the greatest relative cost reduction (9.3%) was observed at the Top 50% PRS threshold, whereas restriction to the Top 25% PRS group yielded only a 1.0% reduction. In contrast, under a scenario in which genetic data were assumed to be pre-existing and no additional sequencing cost was incurred, projected total trial cost decreased monotonically across PRS strata in all three examples. Restricting recruitment to the Top 25% PRS group reduced projected total cost by 51.7% for CAD–PCSK9, 17.8% for glaucoma–ANGPTL7, and 74.7% for IBD–IL23R.

Figure 4. Estimated per-arm sample size required to achieve target power (60–85%) under PRS-guided enrichment, assuming 1:1 randomization and two-sided 5% significance level. Per-arm sample size requirements were derived from analytical power calculations based on disease prevalence predicted by logistic regression models for carriers and non-carriers of protective variants, adjusting for covariates as described in the Methods. Curves show the per-arm sample size required to achieve target power levels ranging from 60% to 85% across polygenic risk score (PRS) strata.

Empirical power under PRS-guided enrichment

We next evaluated empirical power through bootstrap simulations under fixed per-arm sample sizes of 100, 200, 500, and 1,000 participants (Figure 5). For the CAD–PCSK9 example, empirical power increased consistently with increasing PRS enrichment across all sample sizes, with the highest power consistently observed in the top 25% PRS group. At 1,000 participants per arm, empirical power increased from approximately 20% in the full cohort to 50% in the Top 25% PRS group.

A similar monotonic pattern was observed for the IBD–IL23R example, although absolute power remained modest even at larger sample sizes. At 1,000 participants per arm, empirical power increased from approximately 2–3% in the full cohort to about 5–6% in the Top 25% PRS group, reflecting the lower overall prevalence of IBD in the study population.

In contrast, the glaucoma–ANGPTL7 simulations did not show a consistent increase in empirical power across progressively higher PRS thresholds. While PRS-enriched groups generally outperformed the full cohort, the Top 25% PRS group did not uniformly achieve higher power than the Top 50% or Top 75% groups. At 1,000 participants per arm, empirical power increased from 14.5% in the full cohort to 18.4% and 19.3% in Top 75% and Top 50% PRS groups, respectively, but declined to 15.9% in the Top 25% PRS group. This non-monotonic pattern is consistent with the observation that disease prevalence did not differ significantly between carriers and non-carriers within the Top 25% PRS group due to substantially reduced number of carriers–case observations.

We note that in the IL23R–IBD and glaucoma–ANGPTL7 examples, empirical power fell below 5% in some small-sample scenarios, as our simulation classified comparisons between carriers and non-carriers as nonsignificant when fewer than five disease cases were observed in either resampled group.

Figure 5. Empirical power of simulated trials with 100, 200, 500, or 1000 participants per arm. For each sample-size scenario, the proportion of significant simulations (empirical power) is compared across PRS strata. Pairwise differences between strata are tested using two-sided proportion tests, and significant contrasts after adjusting for multiple testing are annotated on the plot using standard significance codes (*** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$).

 Prospective event accrual under PRS-enriched recruitment

We constructed baseline disease-free cohorts for the CAD–PCSK9, glaucoma–ANGPTL7, and
IBD–IL23R gene–disease examples and followed participants for up to 10 years from UK
Biobank enrollment. Baseline demographic characteristics and follow-up summaries are shown
in Table 1. Age and sex distributions were similar across cohorts, whereas overall event counts
differed substantially by disease context.

Table 1. Baseline characteristics of baseline disease-free cohorts for prospective event simulations.
Demographic characteristics, follow-up duration, and total incident events for CAD–PCSK9, glaucoma–
ANGPTL7, and IBD–IL23R cohorts. All participants were free of the relevant disease at baseline and
followed for up to 10 years.

	CAD-PCSK9	Glaucoma-ANGPTL7	IBD-IL23R
N participants	380678	392276	390820
Age and baseline, mean (SD)	56.7 (8.0)	56.9 (8.0)	56.9 (8.0)
Female, %	44.9	45.9	45.9
Number of events	20122	9759	2485
Median follow-up, years	10.0	10.0	10.0

Across all three examples, distributions of the restricted mean survival time (RMST) consistently
shifted downwards from the full cohort to the top 25% PRS group, indicating earlier
accumulation of events within enriched groups (Figure 6). In the CAD–PCSK9 example, at
1,000 participants per arm, the mean RMST across 1,000 simulation replicates decreased from
9.77 years (95% CI: [9.71, 9.82]) in the unenriched cohort to 9.66 years (95% CI: [9.60, 9.73]) in
the Top 25% PRS group. In the glaucoma–ANGPTL7 example, the mean RMST was 9.94 years
(95% CI: [9.92, 9.97]) in the unenriched cohort and 9.88 years (95% CI: [9.85, 9.92]) in the Top
25% PRS group. In the IBD–IL23R example, the difference was more modest, with mean RMST
of 9.97 years (95% CI: [9.95, 9.99]) in the unenriched cohort and 9.95 years (95% CI: [9.93,
9.98]) in the Top 25% PRS group.

We note that RMST estimates depend on the chosen follow-up horizon and may also be sensitive
to sampling variability. In our setting, the absolute differences in RMST were modest in part
because the number of observed events was limited, particularly in the glaucoma–ANGPTL7 and
IBD–IL23R examples. Variability in RMST estimates also increased in more restrictive PRS
groups due to the reduced number of eligible participants and events within narrower PRS strata.

Figure 6. Restricted mean survival time under PRS-enriched recruitment. Distributions of restricted
mean survival time (RMST) from fixed-size bootstrap simulations across PRS strata (All, Top 75%, Top
50%, Top 25%) and per-arm sample sizes (100, 200, 500, 1,000). RMST was calculated for up to 10 years
of follow-up.

Discussion

In this study, we present an in silico framework for prospectively evaluating how polygenic risk
information can be incorporated into clinical trial design through prognostic enrichment using
large-scale biobank data. By combining naturally occurring protective genetic variants as proxies
for therapeutic interventions with PRS-based stratification of disease risk, we demonstrate how
observed population data can be leveraged to inform key trial design parameters prior to trial
initiation. We systematically reviewed disease–gene examples with established protective variant
effects and assessed whether the available data provided sufficient statistical information to
power the evaluation of PRS-guided enrichment in each setting. Our quantitative framework
integrates population-level estimation of disease prevalence, analytical sample size calculation,
simulation-based evaluation of empirical power and prospective event accrual, enabling
systematic assessment of how PRS enrichment influences sample size requirements, statistical
power, and the rate of event accumulation over follow-up.

Across three distinct disease systems illustrated by CAD–PCSK9, glaucoma–ANGPTL7, and IBD–IL23R use-cases, protective variant carriers consistently exhibited lower disease prevalence within PRS strata, and higher PRS was uniformly associated with increased disease risk. These shared features translated into reduced sample size requirements, lower projected operational costs, improved empirical power, and accelerated event accrual relative to unenriched designs. However, the magnitude and pattern of efficiency gains under PRS enrichment varied across disease contexts. In both the CAD-PCSK9 and IBD-IL23R example, the greatest enrichment benefits were observed in the most restrictive Top 25% PRS stratum. The glaucoma–ANGPTL7 example showed a similar pattern but more modest gains in sample size reduction and cost savings across progressively higher PRS strata, with moderate enrichment at the Top 50% threshold achieving empirical power and projected cost reductions comparable to, or exceeding, those obtained with the Top 25% threshold.

The variation in optimal PRS cutoffs across diseases highlights important operational considerations for implementing PRS-based enrichment in clinical trials. One key consideration is that the optimal enrichment threshold depends on the trade-off between enriching the cohort for higher disease risk and retaining a sufficiently large pool of eligible participants. In the CAD–PCSK9 and IBD–IL23R examples, enrichment to the upper PRS quartile was an efficient design choice, as it concentrated risk while preserving sufficient representation of protective variant carriers. In comparison, in the glaucoma–ANGPTL7 example, more moderate PRS thresholds provided comparable or superior efficiency by avoiding excessive loss of eligible participants. Another practical consideration is that increasingly stringent PRS thresholds require screening larger populations to identify eligible participants. When the cost of running the trial exceeds the cost of screening potential participants, as assumed in our model, reductions in required sample size generally outweigh the additional screening costs, especially when genetic data are already available. However, this trade-off depends on the relative costs of screening and trial execution, which vary across diseases and trial designs. In addition, more stringent enrichment criteria may require longer screening periods to identify eligible participants. Although our RMST analyses indicated a consistent pattern of earlier event occurrence in higher-risk cohorts, the absolute reductions in follow-up time were modest, and whether these gains would offset potentially longer recruitment periods would require further evaluation under realistic assumptions about trial recruitment and implementation. Together, our findings emphasize that PRS enrichment should be implemented through a structured, data-driven framework tailored to disease context, genetic risk distribution, and trial settings, rather than relying on a single uniform threshold.

Beyond methodological innovations, our work also presents broader implications for the clinical research and healthcare community. At present, germline genetic information is not routinely incorporated into clinical trials, despite the rapidly expanding availability of genomic data in population cohorts and healthcare systems. Our results contribute quantitative evidence demonstrating the potential value of human genetic data, including naturally occurring protective

variants and polygenic risk scores, for informing clinical trial design. As such evidence accumulates, clinical trialists and sponsors may increasingly consider PRS-guided enrichment as a strategy to improve trial efficiency. In the meantime, governing bodies, trial registers, and adjudicating agencies may have increasing opportunities to develop guidance for genetically informed clinical trials. Moreover, while our study focuses on PRS and genetic data, the framework developed here is not limited to PRS-based enrichment. Other predictors, including molecular biomarkers, imaging measures, and clinical risk scores, may also serve as enrichment variables for clinical trials. The quantitative framework presented here provides a general approach for evaluating such enrichment strategies and may facilitate further comparison of alternative enrichment variables in future trial design studies.

While our framework provides a structured approach for quantitatively evaluating and optimizing PRS-guided enrichment in clinical trial design, several limitations warrant consideration. First, the implementation of this framework in the present study is constrained using a single population cohort, specifically the White British subset of the UK Biobank. Although UK Biobank is a well-characterized population-based resource, evaluation of enrichment performance is inevitably constrained by the available carrier counts and event numbers within the dataset, particularly for rare variants and low-prevalence diseases. In such settings, reduced statistical information may limit precision and render some gene–disease pairs underpowered for robust evaluation of enrichment effects. Integration of additional large-scale biobanks and meta-analyses across independent cohorts would increase carrier representation and event counts and enable more comprehensive evaluation of enrichment strategies, expanding application of this framework across a broader range of therapeutic targets and disease contexts. Such extension will be essential to further establish the generalizability and robustness of the proposed approach.

Second, PRS-guided enrichment raises considerations regarding the generalizability of trial results. Restricting enrollment to individuals with elevated PRS is intended to increase event rates and improve statistical efficiency, but treatment effects estimated in PRS-enriched populations may not directly generalize to individuals across the full PRS distribution. This limitation is an inherent challenge in trial enrichment strategies that restrict enrollment to a subgroup of the target population. A related consideration is potential labeling implications, as trials conducted in PRS-enriched populations may lead to indications tied to PRS-based eligibility. However, PRS can be viewed as one approach to identifying individuals at elevated disease risk, and the resulting evidence may be broadly informative for high-risk patients beyond those defined strictly by PRS, although the extent of this translation will depend on the clinical and regulatory context. In addition, our results derived from the UK Biobank White British cohort may also not directly generalize to populations with different genetic architectures, linkage disequilibrium patterns, or environmental exposures. In practice, enriched trials may provide an efficient approach for establishing initial evidence of therapeutic efficacy, while

subsequent studies or post-marketing analyses would be necessary to evaluate effectiveness
across broader patient populations.

Finally, our analyses rely on several assumptions related to the use of human genetic variation to
inform therapeutic target evaluation. We assume that naturally occurring protective variants can
serve as proxies for pharmacologic target modulation. Although this approach does not fully
capture the complexity of therapeutic intervention, prior work has shown that human genetic
perturbations often provide informative models of target inhibition and drug response, supporting
the conceptual validity of this strategy.[53–55] Nevertheless, because the protective effects
evaluated here reflect naturally occurring genetic variation, they may underestimate the
magnitude of effect achievable by a therapeutic intervention with stronger or more sustained
target modulation. In addition, the effectiveness of PRS-guided enrichment depends on the
accuracy and transferability of the PRS and the heritability of the target disease. For conditions
with limited genetic contribution or poorly calibrated PRS, the benefits of enrichment may be
attenuated.

In summary, our study demonstrates how population-scale genomic and health record data can
be used to evaluate PRS-guided prognostic enrichment as part of early-stage trial planning.
Rather than relying on retrospective analyses of completed trials, our results quantify how PRS-
based enrichment is expected to influence event burden, statistical power, and operational
considerations before a trial is conducted, while highlighting the difference of enrichment effects
across disease contexts. These findings emphasize that the utility of PRS-guided enrichment is
context dependent and should be assessed with respect to disease-specific prevalence, risk
gradients, and feasibility considerations. Ultimately, this framework offers a scalable foundation
for incorporating genomic risk information into the design of future clinical trials and may be
extended to evaluate other prognostic enrichment variables using similar principles.

Author contributions

538 M.A.R. and J.W.OS. designed the study. R. C. led the analysis and manuscript writing. E.A.,
539 Y.L., J.W.OS., and L.T. contributed to the prognostic trial enrichment design approach. All
540 authors contributed to the writing of the manuscript and figures generation. All authors reviewed
541 the manuscript.

Acknowledgements

This research has been conducted using the UK Biobank Resource under Application Number
24983 and 22282. Based on the information provided in Protocol 44532 the Stanford IRB has
determined that the research does not involve human subjects as defined in 45 CFR 46.102(f) or
21 CFR 50.3(g). All participants in the UK Biobank provided written informed consent (more
information is available at https://www.ukbiobank.ac.uk/2018/02/gdpr/). We thank all the
participants in the UK Biobank study. Some of the computing for this project was performed on
the Sherlock cluster. We would like to thank Stanford University and the Stanford Research

Competing interests

M.A.R. is a co-Founder of Broadwing Bio, and holds equity in Maze Therapeutics, Danger Bio, and Braveheart Bio. J.W.O'S is supported by the AHA Postdoctoral fellowship and ACC postdoctoral fellowship and has had consultancy relationships with Google AI, Google Deepmind, Curie Bio, and Foresite Labs (outside the current work). E.A.A. is a Founder of Personalis, Deepcell, Svexa, Saturnus Bio, and Swift Bio, Founding Advisor of Candela, and Parameter Health, Advisor for Pacific Biosciences, and Non-executive director of AstraZeneca, and Dexcom. O.V. and S.B.G. are shareholders and employed by AstraZeneca.

Funding

R.C. is a postdoctoral fellow supported by a sponsored research agreement between Stanford University and AstraZeneca PLC.

References

1. Hariton, E. & Locascio, J. J. Randomised controlled trials - the gold standard for effectiveness research: Study design: randomised controlled trials. *BJOG* **125**, 1716 (2018).

2. Rauch, B. *et al.* OMEGA, a randomized, placebo-controlled trial to test the effect of highly purified omega-3 fatty acids on top of modern guideline-adjusted therapy after myocardial infarction. *Circulation* **122**, 2152–2159 (2010).

3. Yusuf, S., Collins, R. & Peto, R. Why do we need some large, simple randomized trials? *Stat. Med.* **3**, 409–422 (1984).

4. Solomon, S. D. & Pfeffer, M. A. The future of clinical trials in cardiovascular medicine. *Circulation* **133**, 2662–2670 (2016).

5. Olivier, C. B. *et al.* Accuracy of event rate and effect size estimation in major cardiovascular trials: A systematic review. *JAMA Netw. Open* **7**, e248818 (2024).

6. Shen, L. *et al.* Declining risk of sudden death in heart failure. *N. Engl. J. Med.* **377**, 41–51 (2017).

7. Bowman, L. *et al.* Randomized trials fit for the 21st century: A joint opinion from the European society of cardiology, American heart association, American college of cardiology, and the world heart federation. *Circulation* **147**, 925–929 (2023).

8.   Rauch, G., Rauch, B., Schüler, S. & Kieser, M. Opportunities and challenges of clinical trials in

cardiology using composite primary endpoints. *World J. Cardiol.* **7**, 1–5 (2015).

9.   Davidson, M. H., Hsieh, A., Dicklin, M. R. & Maki, K. C. The imperative to enhance cost-

effectiveness for cardiovascular therapeutic development. *JACC Basic Transl. Sci.* **9**, 1029–1040

(2024).

10. Temple, R. Enrichment of clinical study populations. *Clin. Pharmacol. Ther.* **88**, 774–778 (2010).

11. Office of Communications, Division of Drug Information Center for Drug Evaluation and Research.

Enrichment Strategies for Clinical Trials to Support Determination of Effectiveness of Human Drugs

and Biological Products. (US Food and Drug Administration, 2019).

12. Kerr, K. F. *et al.* Evaluating biomarkers for prognostic enrichment of clinical trials. *Clin. Trials* **14**,

629–638 (2017).

13. Viele, K. & Girard, T. D. Risk, results, and costs: Optimizing clinical trial efficiency through

prognostic enrichment. *Am. J. Respir. Crit. Care Med.* **203**, 671–672 (2021).

14. Jering, K. S. *et al.* Improving clinical trial efficiency using a machine learning-based risk score to

enrich study populations. *Eur. J. Heart Fail.* **24**, 1418–1426 (2022).

15. Scott, J. V., Garnett, C. E., Kanwar, M. K., Stockbridge, N. L. & Benza, R. L. Enrichment benefits

of risk algorithms for Pulmonary Arterial Hypertension clinical trials. *Am. J. Respir. Crit. Care Med.*

203, 726–736 (2021).

16. Torkamani, A., Wineinger, N. E. & Topol, E. J. The personal and clinical utility of polygenic risk

scores. *Nat. Rev. Genet.* **19**, 581–590 (2018).

17. Khera, A. V. *et al.* Genome-wide polygenic scores for common diseases identify individuals with

risk equivalent to monogenic mutations. *Nat. Genet.* **50**, 1219–1224 (2018).

18. Fahed, A. C., Philippakis, A. A. & Khera, A. V. The potential of polygenic scores to improve cost

and efficiency of clinical trials. *Nat. Commun.* **13**, 2922 (2022).

19. Damask, A. *et al.* Patients with high genome-wide polygenic risk scores for coronary artery disease

may receive greater clinical benefit from alirocumab treatment in the ODYSSEY OUTCOMES trial.

Circulation **141**, 624–636 (2020).

20. German, J. *et al.* Incorporating genetic data improves target trial emulations and informs the use of

polygenic scores in randomized controlled trial design. *Nat. Genet.* **57**, 1620–1627 (2025).

21. Pressly, M. A., Schuck, R., Mummaneni, P., Roman, Y. M. & Pacanowski, M. Utilization of

polygenic risk scores in drug development protocols. *Pharmacogenomics* **26**, 75–79 (2025).

22. Fahed, A. C. & Natarajan, P. Clinical applications of polygenic risk score for coronary artery disease

through the life course. *Atherosclerosis* **386**, 117356 (2023).

23. Natarajan, P. *et al.* Polygenic risk score identifies subgroup with higher burden of atherosclerosis

and greater relative benefit from statin therapy in the primary prevention setting. *Circulation* **135**,

2091–2101 (2017).

24. Marston, N. A. *et al.* Predicting benefit from evolocumab therapy in patients with atherosclerotic

disease using a genetic risk score: Results from the FOURIER trial: Results from the FOURIER

trial. *Circulation* **141**, 616–623 (2020).

25. Cohen, J. C., Boerwinkle, E., Mosley, T. H., Jr & Hobbs, H. H. Sequence variations in PCSK9, low

LDL, and protection against coronary heart disease. *N. Engl. J. Med.* **354**, 1264–1272 (2006).

26. Kent, S. T. *et al.* PCSK9 loss-of-function variants, low-density lipoprotein cholesterol, and risk of

coronary heart disease and stroke: Data from 9 studies of blacks and whites: Data from 9 studies of

blacks and whites. *Circ. Cardiovasc. Genet.* **10**, e001632 (2017).

27. Sabatine, M. S. *et al.* Evolocumab and clinical outcomes in patients with cardiovascular disease. *N.*

Engl. J. Med. **376**, 1713–1722 (2017).

28. Schwartz, G. G. *et al.* Alirocumab and cardiovascular outcomes after acute coronary syndrome. *N.*

Engl. J. Med. **379**, 2097–2107 (2018).

29. Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic data. *Nature* **562**,

203–209 (2018).

30. UK Biobank Whole-Genome Sequencing Consortium. Whole-genome sequencing of 490,640 UK

Biobank participants. *Nature* **645**, 692–701 (2025).

31. Thompson, D. J. *et al.* UK Biobank release and systematic evaluation of optimised polygenic risk

scores for 53 diseases and quantitative traits. *bioRxiv* 2022.06.16.22276246 (2022)

doi:10.1101/2022.06.16.22276246.

32. Thompson, D. J. *et al.* A systematic evaluation of the performance and properties of the UK Biobank

Polygenic Risk Score (PRS) Release. *PLoS One* **19**, e0307270 (2024).

33. Tanigawa, Y. *et al.* Rare protein-altering variants in ANGPTL7 lower intraocular pressure and

protect against glaucoma. *PLoS Genet.* **16**, e1008682 (2020).

34. Praveen, K. *et al.* ANGPTL7, a therapeutic target for increased intraocular pressure and glaucoma.

Commun. Biol. **5**, 1051 (2022).

35. Brown, S. F.-K. *et al.* ANGPTL7 and its role in IOP and glaucoma. *Invest. Ophthalmol. Vis. Sci.* **65**,

22 (2024).

36. Zaveri, S. R. *et al.* ANGPTL7 silencing reduces intraocular pressure in genetic and steroid-induced

glaucoma models. *Invest. Ophthalmol. Vis. Sci.* **66**, 836–836 (2025).

37. Duerr, R. H. *et al.* A genome-wide association study identifies IL23R as an inflammatory bowel

disease gene. *Science* **314**, 1461–1463 (2006).

38. Pidasheva, S. *et al.* Functional studies on the IBD susceptibility gene IL23R implicate reduced

receptor function in the protective genetic variant R381Q. *PLoS One* **6**, e25038 (2011).

39. Kaste, R. *et al.* Efficacy and safety of IL-23 inhibitors in the treatment of Crohn's disease and

Ulcerative Colitis: Systemic review and network meta-analysis. *medRxiv* 2025.09.03.25334996

(2025) doi:10.1101/2025.09.03.25334996.

40. Parisio, L. *et al.* Interleukin 23: Pathogenetic involvement and therapeutic target for ulcerative

colitis. *J. Clin. Med.* **14**, 4590 (2025).

41. Eyre, S. *et al.* High-density genetic mapping identifies new susceptibility loci for rheumatoid

arthritis. *Nat. Genet.* **44**, 1336–1340 (2012).

42. Ferreira, R. C. *et al.* Functional IL6R 358Ala allele impairs classical IL-6 receptor signaling and

influences risk of diverse inflammatory diseases. *PLoS Genet.* **9**, e1003444 (2013).

43. Janahiraman, S., Too, C. L., Lee, K. W., Shahril, N. S. & Leong, C. O. Genetic biomarkers as

predictors of response to tocilizumab in rheumatoid arthritis: A systematic review and meta-analysis.

Genes (Basel) **13**, 1284 (2022).

44. Zorc, R. *et al.* A coding single nucleotide polymorphism in the interleukin-6 receptor enhances IL-6

signalling in CD4 T cells and predicts treatment response to tocilizumab in giant cell arteritis. *Ann.*

Rheum. Dis. **84**, 1401–1411 (2025).

45. Flannick, J. *et al.* Loss-of-function mutations in SLC30A8 protect against type 2 diabetes. *Nat.*

Genet. **46**, 357–363 (2014).

46. Rutter, G. A. & Chimienti, F. SLC30A8 mutations in type 2 diabetes. *Diabetologia* **58**, 31–36

(2015).

47. Dwivedi, O. P. *et al.* Loss of ZnT8 function protects against diabetes by enhanced insulin secretion.

Nat. Genet. **51**, 1596–1606 (2019).

48. Lamarche, L. B. *et al.* Complete loss of SLC30A8 in humans improves glucose metabolism and beta

cell function. *Diabetologia* **68**, 2754–2766 (2025).

49. TG and HDL Working Group of the Exome Sequencing Project, National Heart, Lung, and Blood

Institute *et al.* Loss-of-function mutations in APOC3, triglycerides, and coronary disease. *N. Engl. J.*

Med. **371**, 22–31 (2014).

50. Jørgensen, A. B., Frikke-Schmidt, R., Nordestgaard, B. G. & Tybjærg-Hansen, A. Loss-of-function

mutations in APOC3 and risk of ischemic vascular disease. *N. Engl. J. Med.* **371**, 32–41 (2014).

51. Khetarpal, S. A. *et al.* A human APOC3 missense variant and monoclonal antibody accelerate apoC-

III clearance and lower triglyceride-rich lipoprotein levels. *Nat. Med.* **23**, 1086–1094 (2017).

52. Ashraf, T. *et al.* The impact of olezarsen on hypertriglyceridemia in high cardiovascular risk

patients: a systematic review, meta-analysis, and meta-regression. *Ann. Med. Surg. (Lond.)* **87**,

5173–5184 (2025).

53. Plenge, R. M., Scolnick, E. M. & Altshuler, D. Validating therapeutic targets through human

genetics. *Nat. Rev. Drug Discov.* **12**, 581–594 (2013).

54. Nelson, M. R. *et al.* The support of human genetic evidence for approved drug indications. *Nat.*

Genet. **47**, 856–860 (2015).

55. Hingorani, A. D. *et al.* Improving the odds of drug development success through human genomics:

modelling study. *Sci. Rep.* **9**, 18911 (2019).

Supplementary Materials

Supplementary Methods

Cost model for PRS-guided clinical trial enrichment

To quantify the operational implications of polygenic risk score (PRS)–based trial enrichment,
we developed a simplified cost model that translates the PRS-specific sample size requirements
into total projected trial cost. Our goal is to illustrate relative changes in projected cost across
PRS thresholds rather than estimate absolute expenditures, which in practice depend strongly on
disease-specific, assay-specific, and trial-specific factors. Accordingly, we characterize the
relative change in projected cost when restricting enrollment to individuals above a given PRS
threshold compared with an unenriched trial. Our cost model conceptualizes the trial workflow
as three sequential cost components, including general screening, genotyping and PRS
computation, and trial execution for randomized participants, each scaled according to the
number of individuals required at that stage under a given PRS threshold.

For each PRS threshold q, defined as enrolling only individuals in the upper $(1 - q)$ quantile of
the PRS distribution, where the unenriched trial corresponds to $q = 0$, let $n_{arm}(q)$ denote the
analytically derived per-arm sample size required to achieve a given target power. The total
number of randomized participants is:

704
$$N_{rand}(q) = 2n_{arm}(q).$$

705   The number of individuals who must undergo genotyping and PRS assessment would then be

706
$$N_{PRS}(q) = \frac{N_{rand}(q)}{1 - q} = \frac{2n_{arm}(q)}{1 - q}.$$

707   In practice, we might consider only genotyping individuals who pass the inclusion criteria and
708   provide consent during an initial screening stage, during which a proportion p_{exclu} is excluded
709   due to ineligibility or lack of consent. The total number of individuals requiring general
710   screening is therefore

711
$$N_{screen}(q) = \frac{N_{PRS}(q)}{1 - p_{exclu}} = \frac{2n_{arm}(q)}{(1 - q)(1 - p_{exclu})}.$$

712   Suppose all individuals entering the screening incur a base screening cost C_{screen}, and
713   individuals who fail screening additionally incur a smaller incremental failure cost C_{fail}. Then
714   the cost during the screening stage would be:

715
$$Cost_{screen}(q) = N_{screen}(q)C_{screen} + N_{screen}(q)p_{exclu}C_{fail}.$$

716   Assuming the cost of genotyping and PRS calculation is C_{geno}, then the total cost to obtain PRS
717   on individuals who passed the initial screening would be:

$$718 \qquad Cost_{PRS}(q) = N_{PRS}(q)C_{geno} = \frac{2n_{arm}(q)}{(1-q)}C_{geno}.$$

719 Finally, for participants whose PRS exceeding threshold q, assuming the cost of running the trial
720 on each individual is C_{trial}, which includes study visits, drug supply, safety monitoring, and
721 endpoint adjudication, the total cost of trial execution would be:

$$722 \qquad Cost_{trial}(q) = N_{rand}(q)C_{trial} = 2n_{arm}(q)C_{trial}.$$

723 Combining the three components yields the total projected cost for a PRS-enriched design at
724 threshold q:

$$725 \qquad TotalCost(q) = Cost_{screen}(q) + Cost_{PRS}(q) + Cost_{trial}(q)$$

$$726 \qquad = 2n_{arm}(q)[C_{trial} + \frac{C_{geno}}{1-q} + \frac{C_{screen}+p_{exclu}C_{fail}}{(1-q)(1-p_{exclu})}].$$

727 We define $A(q) = C_{trial} + \frac{C_{geno}}{1-q} + \frac{C_{screen}+p_{exclu}C_{fail}}{(1-q)(1-p_{exclu})}$ to represent the average per-randomized-
728 participant cost at threshold q. The percent reduction in total cost of a PRS-enriched trial at
729 threshold q relative to an unenriched trial can then be expressed as:

$$730 \qquad \Delta(q) = \frac{TotalCost(0) - TotalCost(q)}{TotalCost(0)} = 1 - \frac{n_{arm}(q)A(q)}{n_{arm}(0)A(0)}.$$

We produced illustrative numerical examples of $\Delta(q)$ using analytically derived sample size
estimates for the CAD–PCSK9 and glaucoma–ANGPTL7 simulations, assuming $C_{screen} = 500$,
$C_{trial} = 10000$, and $p_{exclu}=0.2$. Recognizing that large population-genomic initiatives are
making genomic data increasingly available, we evaluated two scenarios: one in which per-
person sequencing costs are included ($C_{geno} = 1000$), and another assuming that genetic data
are already available at scale and can be obtained at negligible marginal cost ($C_{geno} = 0$), as
anticipated in future biobank-enabled health systems. Since $TotalCost(q)$ is linear in $n_{arm}(q)$,
percent changes in cost are invariant across power targets. Therefore, we evaluate cost reductions
at a representative target of 80% power, which fully captures the behavior of the model. The
result is shown in Figure S2.

Supplementary Figures and Tables

Figure S1. Workflow for selection of disease–gene pairs to evaluate PRS-guided in silico
enrichment. Candidate disease–gene pairs involving biologically supported monogenic variants
with established protective association to the disease of interest were identified through
structured literature. These pairs were subsequently evaluated in the full cohort for a statistically
supported carrier–non-carrier difference in disease risk and a significant association between
polygenic risk score (PRS) and disease burden. Finally, we assessed whether the carrier–non-
carrier contrast remained statistically supported within progressively enriched PRS strata.
Disease–gene pairs satisfying these criteria were selected for full in silico enrichment analysis.

Figure S2. Predicted disease probabilities by logistic regression. Predicted disease risk was estimated using logistic regression models incorporating PRS, carrier status, and covariates. Curves represent the model-predicted risk for carriers and non-carriers across the continuous PRS distribution, with vertical dashed lines indicating PRS quantile thresholds used to define enrichment strata.

Figure S3. Relative projected trial cost under PRS-enriched recruitment. For each gene–disease example (CAD–PCSK9, glaucoma–ANGPTL7, and IBD–IL23R), bars display the total projected cost of a trial restricted to the top 75%, 50%, or 25% of the PRS distribution, scaled relative to an unenriched trial. The height of each bar corresponds to $\frac{TotalCost(q)}{TotalCost(0)}$. Annotated brackets indicate $\Delta(q)$, the percent reduction in overall cost compared with the full-population design. Bars are partitioned into cost components attributable to screening, sequencing/PRS, and per-participant trial costs, distinguished using different hatch patterns. Results are shown under two assumptions: a current-state scenario in which sequencing is required, and a future-state scenario in which no additional sequencing cost is incurred.

764 **Table S1. Candidate monogenic variants and corresponding disease endpoint definitions.**
765 Disease–variant pairs identified through structured literature review and considered for
766 evaluation under the PRS-guided enrichment framework. Variants are denoted using HGVS
767 protein nomenclature. Disease endpoints were defined using ICD-10 codes as specified.
768 Genomic coordinates correspond to the GRCh38 reference genome assembly.

Variant	Genomic coordinates	Disease	ICD-10 codes for endpoint
PCSK9 p.Arg46Leu	chr1:55039974	CAD	I21 and I25, excluding I25.2
ANGPTL7 p.Gln175His	chr1:11252389	Glaucoma	H40
IL23R p.Arg381Gln	chr1:67705958	IBD	K50, K51
IL6R p.Asp358Ala	chr1:154426970	RA	M05, M06
SLC30A8 p.Arg138Ter	chr8:118253964	T2D	E11
APOC3 p.Ala43Thr	chr11:116701713	CAD	I21 and I25, excluding I25.2

Table S2. Evaluation of disease–gene examples for implementing the PRS-guided in silico enrichment framework. Numbers of cases and participants in the full cohort and among variant carriers are reported. Baseline genetic and PRS signals are assessed in the full cohort using carrier versus non-carrier odds ratios and the odds ratio per standard deviation of PRS. Within-stratum signal is evaluated using the Wald statistic for the carrier contrast in progressively enriched PRS strata. OR: odds ratio. Z_carrier: Wald statistic from logistic regression of disease prevalence on carrier status within the specified PRS stratum.

	CAD (PCSK9)	Glaucoma (ANGPTL)	IBD (IL23R)	RA (IL6R)	T2D (SLC30A8)	CAD (APOC3)
Sample counts						
Full cohort (Cases/N)	$\frac{41{,}697}{393{,}610}$	$\frac{12{,}092}{393{,}610}$	$\frac{6{,}172}{393{,}610}$	$\frac{8{,}254}{393{,}610}$	$\frac{32{,}274}{393{,}610}$	$\frac{41{,}697}{393{,}610}$
Carriers (Cases/N)	$\frac{1{,}236}{13{,}717}$	$\frac{152}{6{,}314}$	$\frac{530}{50{,}633}$	$\frac{5{,}353}{255{,}320}$	$\frac{6}{66}$	$\frac{25}{199}$
Genetic and PRS association in full cohort						
Carrier OR (95% CI)	0.84 (0.80, 0.89)	0.72 (0.61, 0.84)	0.65 (0.60, 0.70)	1.01 (0.97, 1.06)	1.11 (0.46, 2.65)	1.24 (0.79, 1.93)
PRS OR per SD (95% CI)	1.59 (1.58, 1.61)	1.71 (1.68, 1.74)	1.82 (1.78, 1.87)	1.51 (1.48, 1.54)	2.07 (2.04, 2.10)	1.60 (1.58, 1.62)
Carrier effect within PRS strata						
Z_carrier (p-value), Top 75% PRS	5.86 (4.7e-09)	3.52 (4.4e-04)	6.20 (5.6e-10)	0.51 (0.613)	0.03 (0.974)	0.85 (0.394)
Z_carrier (p-value), Top 50% PRS	4.85 (1.3e-06)	2.49 (0.0129)	4.23 (2.4e-05)	0.19 (0.849)	0.30 (0.763)	0.83 (0.408)
Z_carrier (p-value), Top 25% PRS	3.94 (8.0e-05)	1.61 (0.107)	2.56 (0.0105)	1.48 (0.139)	0.02 (0.985)	0.45 (0.651)
Joint information score						
Ψ	4.57	0.453	3.112	0	0	0

Table S3. Logistic regression model of coronary artery disease on PRS and PCSK9 carrier status. Odds ratios (ORs) and 95% confidence intervals are estimated from a logistic regression model including carrier status of the *PCSK9* p.Arg46Leu variant, standardized polygenic risk score (PRS), their interaction term, and covariates (age, sex, and the first three genetic principal components). No statistically significant interactions were observed between PRS and protective variant carrier status.

	OR (95% CI)	p-value
Intercept	0.023 (0.020, 0.026)	< 0.0001
Carrier	0.878 (0.806, 0.956)	0.003
PRS	1.438 (1.416, 1.460)	< 0.0001
Carrier * PRS	0.930 (0.854, 1.012)	0.092
Age	1.083 (1.081, 1.086)	< 0.0001
Sex (Male)	2.321 (2.252, 2.393)	< 0.0001
PC1	0.992 (0.975, 1.006)	0.0004
PC2	0.996 (0.983, 1.002)	0.116
PC3	1.001 (0.991, 1.010)	0.880

Table S4. Logistic regression model of glaucoma on PRS and ANGPTL7 carrier status.
Odds ratios (ORs) and 95% confidence intervals are estimated from a logistic regression model including carrier status of the *ANGPTL7* p.Gln175His variant, standardized glaucoma PRS, their interaction term, and covariates (age, sex, and the first three genetic principal components). No statistically significant interactions were observed between PRS and protective variant carrier status.

	OR (95% CI)	p-value
Intercept	0.020 (0.017, 0.023)	< 0.0001
Carrier	0.759 (0.631, 0.914)	0.004
PRS	1.722 (1.690, 1.755)	< 0.0001
Carrier * PRS	1.028 (0.872, 1.211)	0.741
Age	1.103 (1.100, 1.107)	< 0.0001
Sex (Male)	1.073 (1.034, 1.113)	0.0002
PC1	1.000 (0.988, 1.012)	0.994
PC2	1.005 (0.993, 1.018)	0.402
PC3	0.989 (0.977, 1.000)	0.060

Table S5. Logistic regression model of IBD on PRS and IL23R carrier status. Odds ratios (ORs) and 95% confidence intervals are estimated from a logistic regression model including carrier status of the *IL23R* p.Arg381Gln variant, standardized glaucoma PRS, their interaction term, and covariates (age, sex, and the first three genetic principal components). A weak interaction effect was observed between PRS and protective variant carrier status.

	OR (95% CI)	p-value
Intercept	0.013 (0.010, 0.016)	< 0.0001
Carrier	0.912 (0.833, 0.999)	0.049
PRS	1.843 (1.794, 1.893)	< 0.0001
Carrier * PRS	0.904 (0.826, 0.990)	0.029
Age	1.010 (1.007, 1.013)	< 0.0001
Sex (Male)	1.141 (1.085, 1.200)	< 0.0001
PC1	1.009 (0.993, 1.025)	0.267
PC2	1.010 (0.993, 1.027)	0.268
PC3	0.997 (0.980, 1.013)	0.673