Title: Characterizing Oseltamivir Use among Community-Dwelling Patients Diagnosed with
Influenza Virus Infection, 2023-2025

Authors:

Emily A. McNair, MSPH[1,2]; Jennie H. Kwon, DO, MSCI[3]; Carlos G. Grijalva, MD, MPH[4]; Son
H. McLaren, MD, MS[5]; Jessica E. Biddle, MPH[2]; Samantha Dean, PhD[2]; Lydia Bristol, MPH,
CPH[2]; Elizabeth B. White, PhD[2]; Stephanie A. Fritz, MD, MSCI[3]; Rachel M. Presti, MD, PhD[3];
Caroline A. O'Neil, MA, MPH[3]; Ellen Sano, DO[5]; Celibell Vargas, MD[5]; Jonathan Schmitz,
MD, PhD[4]; Yuwei Zhu, MD, MS[4]; Theresa A. Scott, MS[4]; Stacey House, MD, PhD[3]; H. Keipp
Talbot, MD, MPH[4]; Melissa S. Stockwell, MD, MPH[5]; Alexandra M. Mellis, PhD[2] on behalf of
the RVTN Flu study group

Affiliations:

[1] STI Federal

[2]Centers for Disease Control and Prevention, Atlanta, GA, United States

[3] Washington University in St. Louis, St. Louis, MO, United States

[4] Vanderbilt University Medical Center, Nashville, TN, United States

[5] Columbia University Irving Medical Center, New York, NY, United States

Acknowledgements

Thank you to the RVTNFlu Study Group Members: Kim T. Vu, Regine Burton, Francesca
Yerbic, Lucy Vogt, Carleigh Samuels, Kelly Bono, Olivia Arter, Alyssa Valencia, Elianora
Ovchiyan, Akshay Saluja, Mengesha Teshome, Alexandra Astarita, Sara Hubert, Tina Robinson,
Mary G. Boyle, Alaina L. Schneider, Lisa M. Richardson, Tara Curley, Jamie Mills, Elizabeth
Capasso-Gulve, Sarah Pruitt-Welsh, Sam Kerz, Heena Kahn, Jamie Garner, Judy A. King, Erica
Anderson, Brianna Schibley-Laird, Rebecca Tablada, Onika Abrams, Sydney A. Cornelison,
Dandan Liu, Liqun Wang, Anny L Diaz Perez, Raul A. Silverio Francisco, Ana Valdez de
Romero

Corresponding Author:

Emily McNair

1600 Clifton Rd. NE
Atlanta, GA 30333
United States
zob8@cdc.gov
32

33 **Conflict of Interest Statements:** Author Melissa Stockwell makes the following financial
34 disclosures: subcontract originating from CDC paid to Trustees of Columbia for grants related to
35 influenza vaccination and disease. Service agreement paid to Trustees of Columbia from
36 American Academy of Pediatrics. Author Carlos Grijalva makes the following financial

42 **Article Summary:** We described and analyzed oseltamivir initiation and course completion
43 among influenza-positive outpatients enrolled in a household transmission study, finding low
44 overall initiation and completion. Results highlight differences in antiviral treatment for groups
45 at higher risk for influenza complications.

Abstract:

Background: Oseltamivir is an antiviral medication for influenza that can reduce the duration of
symptoms and may lower the risk of some complications. Recommendations for use of
oseltamivir include in the outpatient setting for individuals at higher risk of developing influenza
complications.

Objectives: To describe oseltamivir initiation and treatment completion among influenza-
positive outpatients and identify factors associated with each.

Methods: In a U.S. outpatient household transmission study, index participants with laboratory-
confirmed influenza provided up to 12 days of detailed information on medication use. We
described oseltamivir initiation among index cases and treatment course completion of ≥ 10
doses among cases who initiated oseltamivir. We used unadjusted and adjusted logistic
regression to identify factors associated with initiation and course completion.

Results: Among 823 enrolled index cases, 324 (39%) initiated oseltamivir treatment. Of 406
persons at higher risk for influenza complications, 172 (42%) initiated treatment. Oseltamivir
initiation was lowest among children aged 2 to < 5 years (19%) compared to all other age groups.
Among 313 cases who initiated oseltamivir, 42% completed the recommended treatment course
of ≥ 10 doses. Among 163 individuals at higher risk of influenza complications, 69 (42%)
completed the recommended treatment course of ≥ 10 doses. Children < 2 years were
significantly less likely to complete treatment compared to adults aged 18-50 years (aOR: 0.21,
95% CI: 0.04, 0.78, p= 0.030); reasons for discontinuation could not be determined.

Conclusions: These findings reveal differences in oseltamivir treatment in an outpatient setting
among groups at higher risk for influenza complications.

Keywords: Influenza, Antiviral, Oseltamivir, Initiation, Course completion

I. Introduction

Influenza contributes to a substantial burden of disease each year and can cause serious illness and complications among those infected [1]. Four antivirals are FDA-approved and currently recommended for treatment of influenza in the United States; oseltamivir is the most widely prescribed [2]. Oseltamivir is FDA approved for the treatment of influenza among individuals aged 14 days and older. The CDC [2] and American Academy of Pediatrics [3] recommend that oseltamivir should be started as soon as possible (< 48 hours of symptom onset) for people at higher risk of influenza complications [4] and recommends providers use clinical judgement for all others. Individuals at higher risk for influenza complications, or those who are hospitalized with influenza, may benefit from treatment initiation more than 48 hours after symptom onset [5]. The recommended course of oseltamivir for treatment of uncomplicated influenza is two doses per day for 5 days or 10 consecutive doses [2]. Among outpatients with mild influenza illnesses, oseltamivir may reduce symptom duration and hospital admission with most evidence suggesting a greater benefit when treatment is started within 48 hours after symptom onset [6-11].

Oseltamivir usage among outpatients varies by age and risk group. [12]. One investigation found that from 2010-2019, 37% of children under 2 years and 34% of children aged 2-5 years at higher risk of influenza complications and with an influenza diagnosis received prescriptions for oseltamivir [13]. A study of influenza-positive adults at higher risk for influenza complications treated in emergency department and urgent care settings enrolled in the VISION network during the 2023-2024 influenza season found that 58% were prescribed antivirals [14]. Once prescribed, taking an incomplete course of oseltamivir may reduce the impact treatment has on the course and severity of disease because current evidence is based on a 5 day or 10 dose course [2]. Currently there are few to no detailed data regarding course completion in the outpatient setting specific to oseltamivir in the United States. During the 2009 influenza H1N1 pandemic, global investigations reported rates of 79-88% for completing a five-day course (10 doses) of oseltamivir [5, 15-17]. However, findings during that period may not be representative of current US outpatient settings, where guidelines and dispensing patterns differ under non-pandemic influenza seasons.

The objective of our study is to provide a description and analysis of oseltamivir initiation and treatment completion in a population of influenza-positive individuals who presented for outpatient care and subsequently enrolled in a household transmission study.

II. Methods

Study design

As part of a case-ascertained household transmission investigation, we identified and enrolled individuals ranging in age from 0 to 96 years with laboratory-confirmed influenza infections. These individuals sought care at non-emergency ambulatory and emergency department settings in Tennessee, New York, and Missouri during the 2023-2024 and 2024-2025 influenza seasons. Index participants were defined as the first individuals within their households to test positive for influenza from a test performed by a health care provider. To be eligible for

inclusion in the study, index participants had to have received a positive influenza test result
within the previous five days, reside in a household with at least one other person, and have no
household members who exhibited clinical symptoms suggestive of influenza on or preceding
the date of symptom onset or the date of illness onset for the index case. Following provision of
written consent, index participants were enrolled on study day 0, together with their household
contacts, and prospectively followed for 7 days. For this analysis, we restricted our study
population to index participants, since all received medical care and had the opportunity to be
prescribed oseltamivir for treatment of influenza.

The study protocol was reviewed and approved as research by the Institutional Review
Board at Washington University in St. Louis School of Medicine, with reliance from Columbia
University Irving Medical Center, Vanderbilt University Medical Center, and CDC and was
conducted consistent with applicable federal law and CDC policy.

Data collection

Index participants self-reported sociodemographic information, including age, sex, and
race and ethnicity at the time of enrollment. Participants also reported the presence of preexisting
health conditions, including heart disease, asthma, cancer, diabetes, kidney, obesity, liver, and
immune conditions. Self-reported influenza vaccination history was reviewed and verified by the
study teams using available medical records, state immunization registries, and data from
vaccine providers. Participants were considered vaccinated for the current season if they received
a verified influenza vaccination or reported both a valid date and known location of vaccination.
The date of vaccination had to occur at least 14 days prior to the onset date of the index case's
illness to be considered vaccinated. Participants were also instructed to self-collect nasal swab
specimens daily during the 7-day follow-up period. Specimens were shipped to the central
RVTN laboratory at Vanderbilt in Tennessee for processing and testing for influenza using the
Hologic Panther Fusion^{TM} quantitative RT-PCR instrument.

Medication and Symptom Diaries

Index participants completed prospective and retrospective surveys designed to collect
daily information on symptoms, medication usage, and medical care encounters. Retrospective
daily surveys were conducted from the day prior to index symptom onset up until enrollment.
After enrollment 7 days were assessed prospectively for a possibility of up to 12 days of data.
Participants reported daily whether they had taken any over the counter and/or prescription
medications including influenza antivirals (oseltamivir, baloxavir, peramivir, and/or zanamivir),
and the number of doses taken for their current illness. In addition to medication use, participants
reported daily any fever, chills, cough, sore throat, runny nose, nasal congestion, wheezing,
shortness of breath, chest tightness, fatigue, muscle or body aches, loss of smell or taste,
headache, abdominal pain, diarrhea, nausea, vomiting or any other symptom.

Analysis

Index participants were included in analysis if they reported either "yes" or "no" to the
question regarding the initiation of oseltamivir on at least 4 daily diaries. If we could not carry

out last observation carried forward for participants missing initiation and dose data, they were
excluded from the analysis. See the supplemental methods for more details. Among all included
influenza-positive index cases, initiation of oseltamivir was defined as having ever taken 1 or
more doses of oseltamivir throughout the course of the study. We described initiation across
participant characteristics including characteristics that classify individuals as higher risk for
influenza complications. In this analysis, higher-risk individuals were defined as people < 5
years (especially age < 2 years) or ≥ 65 years of age or people with heart disease, asthma, cancer,
diabetes, kidney, obesity, liver, and immune conditions [4]. We also looked at the percentage of
individuals at higher risk for influenza complications who initiated oseltamivir after seeking care
more than 48 hours or 2 days after symptom onset because these individuals are recommended to
initiate oseltamivir regardless of time to seek care. Finally, we performed unadjusted logistic
regression using bidirectional Akaike information criterion (AIC) stepwise selection to select the
final covariates for inclusion in our adjusted model for overall initiation. Variables controlled for
in our adjusted models included season, sex, age, study enrollment site, care setting, preexisting
conditions, and influenza vaccination.

To investigate oseltamivir course completion, we used a subset of the population who
initiated oseltamivir. Participants were excluded from our population subset if their oseltamivir
use was likely censored, defined as having reported taking oseltamivir on the first (one day prior
to symptom onset or first baseline diary entry) or last day (last daily diary entry) of data
collection. We defined a complete course of oseltamivir as 10 or more doses of oseltamivir to
reflect clinical guidelines. We described each participant's days of oseltamivir use, total doses of
oseltamivir, dose distribution by age group, and whether their course was complete or
incomplete. We did not ask participants to report reasons for oseltamivir discontinuation,
however, among the subset of individuals who initiated oseltamivir, we conducted a
supplemental analysis comparing symptoms at illness onset, treatment initiation, and treatment
discontinuation between those who completed versus did not complete a full course of
oseltamivir. Finally, we performed unadjusted and adjusted logistic regression for oseltamivir
course completion among those who initiated treatment, controlling for season, sex, age, study
enrollment site, care setting, preexisting conditions and influenza vaccination.

III. Results

Study Cohort

Of 890 index participants enrolled in the 2023-2024 and 2024-2025 seasons, 823 (92%)
met inclusion criteria for the oseltamivir initiation analysis. (see Supplemental Figure 1). Of the
total participants, 46% were enrolled during the 2023-2024 season and 54% were enrolled during
the 2024-2025 season. Half (49%) of participants were at higher risk for influenza complications
(Table 1). A majority of participants were children < 18, with 409/823 (50%) aged 5-17, 101/823
(13%) aged 2 to < 5 and 53/823 (6%) aged < 2. Of the 823 overall participants, 263 (32%) had
preexisting medical conditions. Most (54%) were recruited from non-emergency ambulatory care
settings and 36% had received a seasonal influenza vaccine (Table 1).

Oseltamivir Initiation

Overall, 324 out of 823 participants (39%) reported initiating oseltamivir (Table 1).
Among other influenza antivirals, baloxavir was initiated by 4 participants (one of whom also
reported oseltamivir use) and zanamivir by 1 participant. These participants were retained in the
analysis. Among all participants who are at higher risk for influenza complications, 172/406
(42%) initiated oseltamivir (Table 1). Oseltamivir initiation was higher among children aged < 2
(42%, 22 of 53) compared to children aged 2 to < 5 (19%, 19 of 101). In contrast, adults aged ≥
65 had a higher initiation rate of 67% (16 of 24, Table 1). Among 263 participants with
preexisting conditions (including any of: heart disease, cancer, diabetes, obesity, and various
liver, kidney, lung, and immune conditions), 133 (51%) initiated oseltamivir compared to 34%
with no preexisting conditions (Table 1). Pre-existing conditions were more common among
older adults than children, and frequencies of initiation stratified by both age and underlying
conditions are given in Supplemental Table 1. Across study sites, initiation was highest in
Tennessee (57%), and more (50%) participants screened and enrolled from non-emergency
ambulatory care settings initiated oseltamivir compared to those from emergency care settings
(27%) (Table 1).

Across all index participants, the median duration from symptom onset to seeking care
was 2 days (interquartile range (IQR): 0–3 days) and a majority of participants initiated
oseltamivir on the day they sought care, as detailed in Supplemental Table 2. Initiation of
oseltamivir was more common among the 585 overall participants who sought care within the
first 48 hours after symptom onset (45%) compared to the 237 overall participants who sought
care more than 48 hours after symptom onset (26%) (see Supplemental Table 2 and Table 3).
This was also true among individuals at higher risk for influenza complications recommended to
receive oseltamivir, with a consistent pattern across all age groups and presence of preexisting
medical conditions (see Supplemental Table 3).

Odds of Oseltamivir Initiation

In the unadjusted analysis, participants at higher risk for influenza complications based
on age and preexisting conditions were 1.28 times more likely to initiate oseltamivir, but the
association was not significant ($p > .05$) (Figure 1). Ages 2 to < 5 were associated with decreased
odds of oseltamivir initiation compared to adults aged 18-50 (OR: 0.24, 95% CI: 0.13, 0.42, $p <$
0.001), while the presence of preexisting conditions was associated with increased odds of
oseltamivir initiation (OR: 1.98, 95% CI: 1.47–2.67, $p < 0.001$) compared to those without
preexisting conditions (Figure 1) Adjusted analyses did not include overall higher-risk status but
instead included age and underlying medical conditions as individual covariates. In these
adjusted analyses children aged 2 to < 5 were significantly less likely to initiate oseltamivir than
adults aged 18-50 (aOR: 0.30, 95% CI: 0.15, 0.58, $p < 0.001$) (Figure 1). Additionally,
participants enrolled from Tennessee exhibited higher odds of initiating oseltamivir compared to
those from New York (aOR: 2.41, 95% CI: 1.32, 4.38, $p = 0.008$) (Figure 1).

Oseltamivir Course Completion

258   The number of days of oseltamivir use ranged from 1 day to 11 days and the median
259   number of days of use was 5 for all enrolled index participants (IQR: 4-6 days) (Figure 1, Panel

A). The number of oseltamivir doses taken by all participants ranged from 1 dose to 22 doses and the median number of doses taken was 9 (IQR: 6-10 doses) (Figure 1, Panel B). The median number of oseltamivir doses taken was lowest among participants < 2 years of age (median: 5, IQR: 3-8 doses) (Figure 1, Panel C).

We described oseltamivir course completion among a subset of 313 index participants whose oseltamivir use was not censored by the beginning or end of follow-up (see Supplemental Figure 1). To align with clinical guidelines, we defined completion as receiving ≥ 10 doses of oseltamivir. Overall, 42% of participants completed the recommended course (Table 1). Participants at risk for influenza complications completed a course of ≥ 10 doses at the same rate as the overall population (42%) (Table 1). Completion of a ≥ 10 dose course was lowest among children aged < 2 years (25%) and adults aged 65 years and older (31%) while higher among children aged 2 to < 5 years (56%) (Table 1). Additionally, 45% of individuals with preexisting conditions completed a ≥ 10 dose course (Table 1). A greater proportion of participants who initiated oseltamivir on the day of symptom onset completed a ≥ 10 dose course compared to the proportion of participants who initiated oseltamivir 1-7+ days after symptom onset (Table 1). The proportion of participants reporting symptoms was similar at illness onset, treatment initiation, or treatment discontinuation between individuals who completed versus did not complete a ≥ 10 dose course. (see Supplemental Figure 3).

Odds of Oseltamivir Completion

In unadjusted analysis, participants at higher risk for influenza complications and recommended to receive oseltamivir had 0.95 odds of completing a ≥ 10 dose course compared to those not at higher for influenza complications, but the results were not significant (p > .05). Individuals aged < 2 years had significantly lower odds of completing a ≥ 10 dose course of oseltamivir in the unadjusted model compared to those aged 18-50 years (OR: 0.15, 95% CI: 0.03, 0.48, p= 0.004) and in the adjusted model (aOR: 0.21, 95% CI: 0.04, 0.78, p= 0.030). No additional variables including the prespecified variables for selection in the model, season, sex, age, presence of preexisting conditions, study site, care setting, or influenza vaccination status reached statistical significance in either the unadjusted or adjusted analyses. (Figure 2).

IV. Discussion

This report describes oseltamivir initiation and course completion among people of all ages with laboratory-confirmed influenza infection enrolled in an outpatient household transmission study. Identifying patterns in oseltamivir initiation and course completion enhances our understanding of outpatient antiviral use in the United States. A minority of overall participants (39%) initiated oseltamivir. Among those classified as higher risk for influenza complications, less than half (42%) initiated oseltamivir, compared to 36% of participants not considered higher risk. However, we found that some higher-risk groups are less likely to initiate oseltamivir than others. Only 19% of participants aged 2 to < 5 years initiated oseltamivir while initiation was greater among participants aged ≥ 65 years (67%). We additionally found that overall, fewer than half of people who initiated oseltamivir completed a ≥ 10 dose course, representing the recommended twice daily dosing for 5 days. Individual courses varied, but the lowest median

number of doses taken was among children at higher risk for influenza complications aged < 2
years.

The 2024-2025 influenza season was classified as high severity for all ages, with high
hospitalization rates among children [18], a high number of pediatric deaths [19], and reported
cases of influenza associated encephalopathy (IAE), a rare but serious neurologic influenza
complication, in children [20]. Despite severe influenza seasons, outpatient investigations have
found low initiation of oseltamivir, particularly among children, beyond the present report. One
investigation using outpatient and emergency department prescription claims included in the
IBM Marketscan Commercial Claims and Encounters Database, found that from 2010-2019,
37% of children < 2 of age and 34% of children 2-5 years of age with an influenza diagnosis
received antiviral treatment [13].

Although we did not collect data on prescribing practices, several factors can influence
whether medical professionals prescribe antivirals to groups at higher risk for influenza
complications, including children. These include providers' perceptions of the risks and benefits
for each patient, as well as concerns about potential side effects, which have been reported in
provider surveys [21]. Additional factors may influence whether participants choose to initiate
oseltamivir, such as concerns about potential side effects or a perceived lack of illness or risk for
severe disease [22]. While we did not capture these factors in our analysis, others such as time to
seek care were found to be related to antiviral initiation in our analysis. Data from the Influenza
Vaccine Effectiveness Network also found that early presentation to care was significantly
associated with receipt of an influenza antiviral prescription [23], likely reflecting provider
adherence to guidelines recommending treatment initiation within 48 hours of symptom onset.
We also captured differences in initiation across settings, finding higher initiation among
participants enrolled from Tennessee. Geographic differences may reflect oseltamivir availability
or differences in provider knowledge. Our findings highlight the heterogeneity of oseltamivir
initiation in the outpatient setting, highlighting the difference between current recommendations
and real-world use.

This study is among the first to describe individual courses of oseltamivir in the outpatient
setting, and found that among the individuals that initiated oseltamivir, not all completed a
recommended course of ≥ 10 doses. Fewer than half of participants with preexisting conditions
completed a full course and children < 2 years of age had the lowest rate of course completion.
Although evidence from a systematic review found oseltamivir to be associated with
gastrointestinal side effects, especially for pediatric age groups [24], we did not observe
significant differences in reported gastrointestinal symptoms at illness onset, treatment initiation,
or treatment discontinuation between individuals who completed and those who did not complete
oseltamivir courses. These data could not confirm that gastrointestinal side effects drive early
discontinuation. We also did not observe differences in other reported symptoms and could not
confirm that symptom resolution drove discontinuation. While we did not collect any additional
information from participants relating to reasons for discontinuation, other factors such as
perceptions of antiviral effectiveness and counseling by healthcare providers, specifically
pharmacists, may substantially impact treatment completion [25, 26]. As is currently
recommended, timely initiation of a 5-day treatment course or ≥ 10 doses of oseltamivir

maximizes the benefit of oseltamivir but was not completed by a majority of our study
participants.

Our study is subject to several limitations. First, data on oseltamivir use was self-reported,
which may introduce bias into our study. Second, participants enrolled in the study were not
asked to report whether they received a prescription for an antiviral, so the number of people
receiving a prescription may have been different from the number who ultimately initiated
oseltamivir. Third, precision of our estimates was limited due to small cell sizes, particularly
within pediatric age groups. Additionally, our findings are likely not representative as we only
enrolled participants who sought care from three U.S academic medical centers and had been
given an influenza test and ultimately agreed to participate in a prospective study with intensive
daily follow-up.

This study describes the initiation and completion of a course of 10 or more doses of
oseltamivir among influenza-positive participants presenting for outpatient care across three US
sites. We found low overall rates of both initiation and completion of a $\geq$ 10-dose course of
oseltamivir. Initiation was lowest among pediatric participants, specifically ages 2 to < 5 years,
while course completion was lowest for ages < 2 years. Improved understanding of oseltamivir
initiation and treatment completion for influenza in populations at higher risk for influenza
complications can aid in the evaluation of clinical outcomes, which in turn helps determine
whether there is a need for improving prescribing patterns for outpatients. However, given the
high burden of influenza among patients at higher risk for influenza complications, antiviral
treatment and course completion remain important.

Citations

376
1. CDC. Flu Burden. Available at: https://www.cdc.gov/flu-burden/php/data-vis/index.html.
2. CDC. Influenza Antiviral Medications: Summary for Clinicians. Available at:
https://www.cdc.gov/flu/hcp/antivirals/summary-clinicians.html.
3. Committee on Infectious D. Recommendations for Prevention and Control of Influenza in
Children, 2025-2026: Policy Statement. Pediatrics **2025**; 156(6).
4. CDC. People at Increased Risk for Flu Complications. Available at:
https://www.cdc.gov/flu/highrisk/index.htm.
5. Butler CC, van der Velden AW, Bongard E, et al. Oseltamivir plus usual care versus
usual care for influenza-like illness in primary care: an open-label, pragmatic,
randomised controlled trial. Lancet **2020**; 395(10217): 42-52.
6. Fry AM, Goswami D, Nahar K, et al. Efficacy of oseltamivir treatment started within 5
days of symptom onset to reduce influenza illness duration and virus shedding in an
urban setting in Bangladesh: a randomised placebo-controlled trial. Lancet Infect Dis
2014; 14(2): 109-18.
7. Malosh RE, Martin ET, Heikkinen T, Brooks WA, Whitley RJ, Monto AS. Efficacy and
Safety of Oseltamivir in Children: Systematic Review and Individual Patient Data Meta-
analysis of Randomized Controlled Trials. Clin Infect Dis **2018**; 66(10): 1492-500.
8. Nicholson KG, Aoki FY, Osterhaus AD, et al. Efficacy and safety of oseltamivir in
treatment of acute influenza: a randomised controlled trial. Neuraminidase Inhibitor Flu
Treatment Investigator Group. Lancet **2000**; 355(9218): 1845-50.
9. Whitley RJ, Hayden FG, Reisinger KS, et al. Oral oseltamivir treatment of influenza in
children. Pediatr Infect Dis J **2001**; 20(2): 127-33.
10. Treanor JJ, Hayden FG, Vrooman PS, et al. Efficacy and safety of the oral neuraminidase
inhibitor oseltamivir in treating acute influenza: a randomized controlled trial. US Oral
Neuraminidase Study Group. JAMA **2000**; 283(8): 1016-24.
11. Venkatesan S, Myles PR, Leonardi-Bee J, et al. Impact of Outpatient Neuraminidase
Inhibitor Treatment in Patients Infected With Influenza A(H1N1)pdm09 at High Risk of
Hospitalization: An Individual Participant Data Metaanalysis. Clin Infect Dis **2017**;
64(10): 1328-34.
12. Piedra PA, Schulman KL, Blumentals WA. Effects of oseltamivir on influenza-related
complications in children with chronic medical conditions. Pediatrics **2009**; 124(1): 170-
8.
13. Antoon JW, Sarker J, Abdelaziz A, et al. Trends in Outpatient Influenza Antiviral Use
Among Children and Adolescents in the United States. Pediatrics **2023**; 152(6).
14. Adams K, Garg S, Tartof SY, et al. Patterns in prescribing and dispensing of influenza
antivirals among adults with influenza presenting to urgent care and emergency
department settings, VISION Network, 2023-2024. Clin Infect Dis **2025**.
15. Ushijima K, Higuchi S, Fujimura A. Survey of compliance with oseltamivir phosphate
therapy in Japan. Am J Ther **2009**; 16(1): 8-10.
16. Lonsdale J WR. Antivirals management in Lancashire during the 2009 swine flu
pandemic. Pharm J **2010**; 285.
17. Koliou M, Soteriades ES, Toumasi MM, Demosthenous A, Hadjidemetriou A.
Epidemiological and clinical characteristics of influenza A(H1N1)v infection in children:
The first 45 cases in Cyprus, June - August 2009. Euro Surveill **2009**; 14(33).

18. CDC. Flu Disease Burden: Past Seasons. Available at: https://www.cdc.gov/flu-burden/php/data-vis/past-seasons.html.

19. CDC. FluView Interactive: Influenza-Associated Peadiatric Mortality. Available at: https://gis.cdc.gov/GRASP/Fluview/PedFluDeath.html.

20. Fazal A, Reinhart K, Huang S, et al. Reports of Encephalopathy Among Children with Influenza-Associated Mortality - United States, 2010-11 Through 2024-25 Influenza Seasons. MMWR Morb Mortal Wkly Rep **2025**; 74(6): 91-5.

21. Bassett HK, Rao S, Beck J, et al. Clinician Preferences for Oseltamivir Use in Children With Influenza in the Outpatient Setting. Pediatrics **2025**; 156(3).

22. Singleton JB, Wang D, Louis S, et al. Perceptions and Barriers to Outpatient Antiviral Therapy for COVID-19 and Influenza as Observed by Infectious Disease Specialists in North America: Results of an Emerging Infections Network (EIN) Survey, February 2024. Open Forum Infect Dis **2024**; 11(12): ofae666.

23. Schicker RS, Flannery B, Chung J, et al. Influenza Antiviral Use Among High-Risk Outpatients During Four Recent Influenza Seasons—United States, 2011–2015. Open Forum Infectious Diseases **2016**; 3(suppl_1).

24. Hanula R, Bortolussi-Courval E, Mendel A, Ward BJ, Lee TC, McDonald EG. Evaluation of Oseltamivir Used to Prevent Hospitalization in Outpatients With Influenza: A Systematic Review and Meta-Analysis. JAMA Intern Med **2024**; 184(1): 18-27.

25. Smith LE, D'Antoni D, Jain V, Pearce JM, Weinman J, Rubin GJ. A systematic review of factors affecting intended and actual adherence with antiviral medication as treatment or prophylaxis in seasonal and pandemic flu. Influenza Other Respir Viruses **2016**; 10(6): 462-78.

26. Rajiah K, Sivarasa S, Maharajan MK. Impact of Pharmacists' Interventions and Patients' Decision on Health Outcomes in Terms of Medication Adherence and Quality Use of Medicines among Patients Attending Community Pharmacies: A Systematic Review. Int J Environ Res Public Health **2021**; 18(9).

 Tables and Figures

461   **Table 1.** Characteristics of influenza-positive index participants enrolled in the RVTN, stratified
462   by analytic group **(n (%)).**

		Oseltamivir initiation (N=823)		Oseltamivir completion (N=313)	
	Overall N = 823	No N = 499	Yes[1] N = 324	No (< 10 doses) N = 180	Yes (≥ 10 doses N = 133)
Season					
2023-2024	376 (46%)	256 (68%)	120 (32%)	56 (50%)	56 (50%)
2024-2025	447 (54%)	243 (54%)	204 (46%)	124 (62%)	77 (38%)
Sex[2]					
Female	444 (54%)	278 (63%)	166 (37%)	96 (59%)	67 (41%)
Male	372 (45%)	217 (58%)	155 (42%)	83 (56%)	64 (44%)
Age (years)					
< 2	53 (6%)	31 (58%)	22 (42%)	19 (86%)	3 (14%)
2-4	101 (13%)	82 (81%)	19 (19%)	8 (44%)	10 (56%)
5-17	409 (50%)	272 (67%)	137 (33%)	77 (60%)	51 (40%)
18-50	176 (21%)	89 (51%)	87 (49%)	42 (49%)	44 (51%)
51-64	60 (7%)	17 (28%)	43 (72%)	23 (53%)	20 (47%)
≥ 65	24 (3%)	8 (33%)	16 (67%)	11 (69%)	5 (31%)
Race-Ethnicity					
American Indian and Alaska Native, Non-Hispanic	4 (1%)	2 (50%)	2 (50%)	1 (50%)	1 (50%)
Asian, Non-Hispanic	11 (1%)	4 (36%)	7 (64%)	5 (71%)	2 (29%)
Black, Non-Hispanic	125 (15%)	82 (66%)	43 (34%)	23 (56%)	18 (44%)
Hispanic/Latino	280 (34%)	198 (71%)	82 (29%)	46 (61%)	30 (39%)
Multiple race, Non-Hispanic	8 (1%)	2 (25%)	6 (75%)	4 (67%)	2 (33%)
White, Non-Hispanic	355 (43%)	183 (52%)	172 (48%)	94 (56%)	75 (44%)
Unknown/Refused	13 (2%)	7 (54%)	6 (46%)	4 (67%)	2 (33%)
Missing	27 (3%)	21 (78%)	6 (22%)	3 (50%)	3 (50%)
Site					
New York	289 (35%)	216 (75%)	73 (25%)	44 (66%)	23 (34%)
Tennessee	243 (30%)	105 (43%)	138 (57%)	75 (56%)	60 (44%)
Missouri	291 (35%)	178 (61%)	113 (39%)	61 (55%)	50 (45%)
Index Screening Site					
Non-emergency ambulatory care[3]	443 (54%)	220 (50%)	223 (50%)	123 (56%)	98 (44%)
Emergency care	380 (46%)	279 (73%)	101 (27%)	57 (62%)	35 (38%)
Preexisting medical conditions					

No underlying condition[4]	552 (67%)	364 (66%)	188 (34%)	109 (60%)	73 (40%)
Underlying condition[5]	263 (32%)	130 (49%)	133 (51%)	71 (55%)	57 (45%)
Missing/Unknown	8 (1%)	5 (63%)	3 (37%)	0 (0%)	3 (100%)
At higher risk for influenza complications					
No	411 (50%)	261 (64%)	150 (36%)	83 (57%)	62 (43%)
Yes[6]	406 (49%)	234 (58%)	172 (42%)	97 (58%)	69 (42%)
Missing/Unknown	6 (1%)	4 (67%)	2 (33%)	0 (0%)	2 (100%)
Influenza vaccination status					
Unvaccinated	512 (62%)	335 (65%)	177 (35%)	100 (60%)	68 (40%)
Vaccinated[7]	286 (35%)	149 (52%)	137 (48%)	75 (55%)	61 (45%)
Missing/Unknown	25 (3%)	15 (60%)	10 (40%)	5 (56%)	4 (44%)
Symptom onset to care seeking date (days)					
0 days	82 (10%)	42 (51%)	40 (49%)	21 (55%)	17 (45%)
1-2 days	503 (61%)	280 (56%)	223 (44%)	126 (58%)	92 (42%)
3-4 days	220 (27%)	163 (74%)	57 (26%)	31 (57%)	23 (43%)
5-6 days	18 (2%)	14 (78%)	4 (22%)	2 (67%)	1 (33%)
Symptom onset to oseltamivir initiation (days)					
0 days	29 (4%)	0 (0%)	29 (100%)	11 (41%)	16 (59%)
1-2 days	188 (23%)	0 (0%)	188 (100%)	101 (55%)	84 (45%)
3-4 days	77 (9%)	0 (0%)	77 (100%)	47 (62%)	29 (38%)
5-6 days	21 (3%)	0 (0%)	21 (100%)	15 (79%)	4 (21%)
7+ days	9 (1%)	0 (0%)	9 (100%)	6 (100%)	0 (0%)

[1]Initiation is defined as any participant who took ≥ 1 dose of oseltamivir throughout follow-up.

[2]Overall, 7 out of 823 participants (1%) did not report their sex. Of these, 4 (57%) did not initiate
oseltamivir, while 3 (43%) did. Among those who initiated oseltamivir, 1 (33%) took < 10 doses and 2
(67%) took ≥ 10 doses of oseltamivir.

[3]Ambulatory care settings include walk-in clinics, pharmacies, occupational health, and specialized care
clinics.

[4-5]Underlying conditions include heart disease, asthma, cancer, diabetes, kidney, obesity, liver, and
immune conditions.

[6]At higher risk for influenza complications is defined as anyone < 5 years of age (especially age < 2
years), ≥ 65 years of age, or who have a preexisting condition defined as increasing the risk for influenza
complications [4].

[7]Participants were considered vaccinated for influenza if they received a verified influenza vaccination or
if a valid date and location of vaccination were reported at least 14 days prior to the onset date of the
index case's illness.

Figure 1. Descriptions of oseltamivir initiation and dosages in the outpatient cohort of influenza-
positive index participants enrolled in the Respiratory Viral Transmission Network.

Panels A, B, and C: Histograms and box plot illustrating the use of oseltamivir among participants who initiated treatment, excluding those with likely censored usage. **Panel A:** Histogram displaying the distribution of days of oseltamivir use, with a maximum of 12 days of data collection available. **Panel B:** Histogram depicting the distribution of oseltamivir doses. **Panel C:** Box plots representing the median doses of oseltamivir stratified by age group.

493   **Figure 2.** Unadjusted and adjusted odds ratios of oseltamivir initiation and completion by select
494   characteristics in our outpatient cohort of influenza-positive index participants enrolled in the
495   Respiratory Viral Transmission Network.

Factors evaluated for association with oseltamivir initiation

Model ● Adj. ▲ Unadj.

Factors evaluated for association with oseltamivir course completion

Model ● Adj. ▲ Unadj.

Results of logistic regression for oseltamivir treatment outcomes: initiation (defined as taking at least one dose) and course completion (defined as completing a full course of at least 10 doses). All estimates except for higher-risk status were adjusted for season, sex, age, recruitment site, screening site (care setting), preexisting conditions, and vaccination status. Estimates with confidence intervals overlapping 1 are considered non-significant.