Early-childhood temperament deviations mark psychiatric risk into early adulthood

J. Kopal[1*], N.R. Bakken[1], P. Parekh[1,2], A.A. Shadrin[1,3], P. Jaholkowski[1], L.A.R. Ystaas[1], N. Parker[1], O.B. Smeland[1], E.P. Tissink[1], I.E. Sønderby[1,3,4], K.S. O'Connell[1], O. Frei[1], A.M. Dale[2,5,6], O.A. Andreassen[1,3,*]

[1] Centre for Precision Psychiatry, Division of Mental Health and Addiction, Oslo University Hospital & Institute of Clinical Medicine, University of Oslo, Oslo, Norway

[2] Center for Multimodal Imaging and Genetics, J. Craig Venter Institute, La Jolla, CA, USA

[3] K.G. Jebsen Centre for Neurodevelopmental Disorders, University of Oslo and Oslo University Hospital, Oslo, Norway

[4] Department of Medical Genetics, Oslo University Hospital & University of Oslo, Oslo, Norway

[5] University of California San Diego, La Jolla, CA, USA

[6] Oslo University Hospital, Oslo, Norway

Corresponding authors:
Jakub Kopal, jakub.kopal@medisin.uio.no
Ole A. Andreassen, ole.andreassen@medisin.uio.no

Abstract

Early-childhood temperament is associated with mental health outcomes decades later. Temperament reflects early-emerging individual differences in emotional and behavioral tendencies. These differences are relatively stable across development and shaped by both genetic and environmental influences. However, the consequences of departures from expected developmental trajectories remain largely unexplored. Using data from more than 50,000 children in the Norwegian Mother, Father and Child Cohort Study, we modeled longitudinal temperament trajectories at 1.5, 3, and 5 years of age and quantified deviations from expected development. Multivariate pattern analysis revealed latent dimensions linking these deviations to clinical diagnoses, with ADHD as the most prominent outcome. Time-to-event analysis showed that these dimensions were associated with a higher hazard of ADHD diagnosis across childhood and adolescence. Finally, genetic analyses identified loci jointly associated with temperament trajectories and ADHD, revealing age-dependent genetic effects. Together, these findings show that deviations from temperament trajectories in early childhood capture transdiagnostic vulnerability across development. Early temperament monitoring may thus serve as an indicator of later mental health risk.

Introduction

Well before children speak their first word, they display temperamental tendencies that have long been linked to later emotional and behavioral functioning[1]. By temperament, we here refer to early-appearing individual differences in emotional and behavioral reactivity, shaped by the child's partly biologically rooted tendency to approach and react to the environment[2]. These individual patterns are partly attributable to genetic variation[3,4] and reflect innate predispositions that guide early behavior[5,6]. Temperament is further shaped by neurobiological maturation and environmental experiences, both of which substantially change across childhood[7–9]. Temperament therefore shows partial stability while continuing to evolve[3,10]. This combination of stability and change implies that, despite individual variability, children tend to follow broadly predictable temperament developmental trajectories[7]. When children depart from these expected trajectories, the deviations may serve as early markers of clinical vulnerability[11,12]. However, how deviations from normative trajectories in early childhood relate to later mental health outcomes remains unclear.

One of the most widely used instruments in early-childhood temperament research is the Emotionality, Activity, Shyness, and Sociability Temperament questionnaire (EAS)[13]. The EAS characterizes temperament using four core traits: *emotionality*, reflecting sensitivity and intensity of negative affect; *activity*, indexing the vigor and tempo of motor behavior; *shyness*, referring to inhibition in the presence of unfamiliar people; and *sociability*, reflecting the desire to seek and enjoy social interaction. The validity of these four traits is supported across diverse populations, ages, and measurement approaches[14–16]. Twin and family studies estimate substantial heritability for these traits, typically in the range of 30–60%[3,4]. Evidence suggests that genetic influences on temperament may vary across development[3,17], which underscores the importance of longitudinal approaches. The EAS thus provides a well-established framework for characterizing early temperament and examining its developmental course.

Extensive longitudinal work demonstrates that early temperament predicts a wide range of later developmental, behavioral, and mental health outcomes[18–21]. These predictive links extend well into adulthood, including associations with social functioning more than three decades later [22]. Specific psychiatric and neurodevelopmental disorders are associated with characteristic early temperamental profiles. Children later diagnosed with ADHD often show elevated activity, high negative emotionality, and early regulatory difficulties[23–26]. Anxiety and depressive disorders are linked to early behavioral inhibition and negative emotionality[27–30]. Increasing shyness across childhood has been associated with heightened risk for substance-use disorders[22]. Collectively, these findings suggest that variation in early temperament may signal vulnerability to specific mental health conditions[31,32]. However, prior work has largely focused on static trait levels rather than on how these temperament patterns change over time. Because temperament unfolds dynamically across childhood, single-time measurements may obscure meaningful differences in developmental trajectories. Departures from expected trajectories may thus capture patterns of change that are not reflected in static trait levels.

The availability of large-scale longitudinal cohorts and advances in statistical modeling now enable population-scale analyses of early-childhood temperament development, the identification of clinically relevant deviations from expected trajectories, and the characterization of age-dependent genetic influences. Here, we leverage data from the Norwegian Mother, Father and Child Cohort Study (MoBa), a prospective, genotyped, deeply phenotyped, population-based cohort[33,34]. We use mother-reported EAS measurements collected at 1.5, 3 and 5 years of age for more than 50,000 children, linked to nationwide health registry diagnoses extending from childhood into early adulthood. Using the FEMA-Long framework, a scalable mixed-effects approach designed for longitudinal population cohorts[35,36], we model non-linear temperament trajectories across early childhood and distill each child's trajectory into a single measure of persistent departure from developmental norms. Canonical Partial Least Squares (PLS) analysis then relates these deviations to a spectrum of mental and behavioral disorders. Furthermore, time-to-event modeling examines how early deviations are associated with the risk and timing of psychiatric disorders. Finally, by conducting genome-wide analyses of age-dependent genetic effects, we map how genetic influences on temperament change across development. Together, these analyses provide an integrated developmental framework that links early temperament trajectories and their genetic architecture to later mental health outcomes.

Results

Longitudinal structure of early-childhood temperament

We leveraged data from the MoBa Study, which prospectively follows more than 110,000 children with genetic profiling and comprehensive linkage to health registries. Our aim was to chart how early-childhood temperament trajectories relate to later mental health outcomes. In the present study, we used mother-reported temperament assessments collected when children were approximately 1.5, 3 and 5 years old (Fig. 1a). These assessments were based on the EAS instrument[14], which quantifies four core temperament traits: *emotionality*, *activity*, *shyness*, and *sociability*. Temperament measurements from at least one time point were available for 50,529 children (45,028 at 1.5 years; 36,435 at 3 years; 25,969 at 5 years). These children were linked to data from the Norwegian Patient Registry (NPR), a nationwide registry of specialist-care diagnoses since 2008. Psychiatric diagnoses were defined according to ICD-10 Chapter V (Mental and behavioral disorders; F00–F99). Follow-up for psychiatric diagnoses extended through December 31, 2024. At the end of follow-up, participants were between 16 and 24 years of age, enabling capture of adolescent- and early adult-onset psychiatric diagnoses. Across the full sample of 50,529 children (Fig. 1b), the most prevalent condition was ADHD (F90.0; 3,211 cases), followed by moderate depressive episode (F32.1; 1,390 cases), nonorganic enuresis (F98.0; 1,361 cases), social phobia (F40.1; 1,165 cases) and Asperger syndrome (F84.5; 884 cases). In total, 372 unique ICD-10 codes including subcategories were observed in our sample (Supplementary Data 1).

Each temperament trait was constructed as a composite score summarizing multiple questionnaire items with five response levels. All four traits showed moderate temporal stability across early childhood (Fig. 1c). Correlations ranged from $r = 0.35$ to 0.53 between ages 1.5 and 3 years, and from $r = 0.40$ to 0.55 between ages 3 and 5 years. *Activity* showed the highest stability and *shyness* the lowest across both intervals. Over the longer 1.5-to-5-year interval, correlations remained evident but weaker ($r = 0.27$–0.41; Fig. 1d). Temperament thus retains consistent individual differences while continuing to change.

In addition to temporal stability, we examined cross-trait relationships. The four temperament traits showed weak intercorrelations (Fig. 1e). Averaged across the three timepoints, the strongest relationship was observed between *shyness* and *sociability* (Pearson's $r = -0.36$), whereas *sociability* and *emotionality* showed essentially no linear association ($r = 0.00$). These pairwise relationships were highly consistent across time, varying no more than $r = 0.07$ between any timepoint (Fig. 1f). Collectively, early-childhood temperament exhibited structured, rather stable patterns that form the foundation for the deviation analyses that follow.

Deviations from population-based temperament trajectories

We leveraged the FEMA-Long framework[35,36] to model age-dependent patterns in each temperament trait, accounting for repeated measurements and family structures inherent to population cohorts. The non-linear effect of age was modeled using natural cubic splines with unit height at knots (Fig. 2a; see Methods). In addition to the smooth effect of age, we included sex, genetic principal components and genotyping batch effects. Sex effects varied

across traits, where *activity* showed the largest difference (Supplementary Data 2) since boys
scored approximately 0.16 standard deviations higher than girls ($\beta = -0.16 \pm 0.01$).
In the next step, we generated fitted values for each child using the estimated fixed
effects. Fitted values represent the expected temperament scores based on age, sex, genetic
principal components, and batch effect. As expected, fitted and observed scores values were
only weakly correlated ($r = 0.06$–0.10 across timepoints; Fig. 2b), indicating that most
variation in temperament lies beyond what is captured by the modeled covariates. Next, we
defined temperament deviations as the difference between observed and fitted values (Fig. 2c,
d). A positive deviation indicates that a child scores above expectation for their age and sex, a
negative deviation the opposite. Deviations were then averaged across the three timepoints
for the 18,651 children with complete data to emphasize consistent departures across early
childhood rather than timepoint-specific fluctuations. This yielded four deviation measures
that formed the basis for subsequent clinical analyses.

Latent dimensions connecting temperament deviations and mental health

We examined whether deviations from population-level temperament trajectories in
early childhood were linked to clinical diagnoses until early adulthood. To do so, we applied
canonical PLS, a multivariate framework designed to extract latent dimensions that maximize
shared variance between two sets of variables. In our case, PLS identified linear
combinations of the four temperament deviations and linear combinations of all observed
mental and behavioral disorders diagnoses (266 observed ICD-10 codes including
subcategories) that covaried most strongly. This procedure yielded individual-level PLS
scores that indicate how strongly each child expresses the multivariate pattern represented by
each dimension.
Using permutation testing (Fig. S1), we identified two significant latent dimensions
linking early-childhood temperament deviations to clinical diagnoses in early adulthood. To
interpret each latent dimension, we examined the correlations between the original
temperament deviations and the corresponding latent dimension scores (i.e., PLS scores). The
first dimension reflected a temperament deviation profile marked by elevated *activity* ($r = 0.71$), and *emotionality* ($r = 0.67$), together with lower *sociability* ($r = -0.38$) (Fig. 3a). This
dimension showed its strongest association with ADHD ($r = 0.97$), but was also associated
with Asperger syndrome ($r = 0.24$), Tourette's syndrome ($r = 0.23$), hyperkinetic conduct
disorder ($r = 0.23$), and specific reading disorder ($r = 0.22$).
The second dimension reflected a distinct temperament deviation profile characterized
by elevated *shyness* ($r = 0.70$) and *emotionality* ($r = 0.63$), together with reduced *sociability*
($r = -0.67$) and *activity* ($r = -0.37$) (Fig. 3b). This dimension was associated with increased
risk for Asperger syndrome ($r = 0.72$), social phobia ($r = 0.36$), generalized anxiety disorder
($r = 0.22$), and major depressive disorder ($r = 0.31$). Notably, it showed a negative
relationship with ADHD ($r = -0.27$). Taken together, the two latent dimensions captured
orthogonal, data-driven patterns of temperament deviation from the developmental norm
during childhood, each differently associated with psychiatric outcomes in adolescence and
early adulthood.

182   To assess the robustness of these associations, we repeated the analysis using ICD-10
183 category codes, which aggregate the 266 subcategory codes into 50 broader diagnostic
184 categories (Fig. 3c). Rerunning the PLS model yielded again two significant dimensions with
185 temperament profiles similar to those obtained in the full analysis (Fig. S2). The first latent
186 dimension was dominated by hyperkinetic disorders ($r = 0.86$), the major-category umbrella
187 for all ADHD subtypes, followed by pervasive developmental disorder ($r = 0.39$) and other
188 behavioral and emotional disorders that usually occur in childhood and adolescence ($r =$
189 0.36). The second dimension showed the highest loadings for pervasive developmental
190 disorder ($r = 0.68$), hyperkinetic disorders ($r = -0.48$) and other anxiety disorders ($r = 0.26$).
191 As a sensitivity analysis, we repeated this analysis using diagnoses from both the NPR
192 (specialist healthcare) and the KUHR database (Norwegian Control and Payment of Health
193 Reimbursements Database, see Methods). KUHR captures reimbursed healthcare encounters
194 including primary care consultations. Results were consistent with the NPR-only analysis, as
195 very few additional diagnoses were present in KUHR alone (Fig. S3). All subsequent
196 analyses therefore used NPR diagnoses only. All associations and confidence intervals (CIs)
197 are summarized in Supplementary Data 3. Collectively, these results reinforced the strong
198 links between deviations from temperament trajectories and mental and behavioral diagnoses.

Temperament deviations and the timing of ADHD diagnosis

200   Having identified two diagnostically relevant latent dimensions, we examined their
201 association with the risk and timing of first recorded diagnoses. ADHD was selected for
202 follow-up time-to-event analyses because it showed the most consistent and robust
203 associations across both latent dimensions in the multivariate model (Fig. S4). To quantify
204 this association, we modeled time to first ADHD diagnosis using Cox proportional hazards
205 models with age in months as the underlying time scale. The two latent dimension scores
206 were included as predictors together with relevant covariates (Table S1), allowing us to
207 estimate how variation along each latent dimension related to the hazard of receiving an
208 ADHD diagnosis.

209   Only the first latent dimension showed a significant association with increased hazard
210 of ADHD diagnosis (PLS 1: hazard ratio (HR) = 1.52, 95% CI 1.41–1.65), whereas the
211 second dimension was not significant (PLS 2: HR = 1.05, 95% CI 0.98–1.12; Fig. 4a).
212 Kaplan–Meier survival curves showed a clear increase in cumulative risk with high scores on
213 the first dimension (Fig. 4b). Cumulative risk also differed by sex, with males showing
214 consistently higher risk and the sex difference widening from early childhood onward (Fig.
215 4c).

217   Although the temperament deviation dimensions were not optimized specifically for
218 ADHD diagnosis, the hazard ratio associated with the first latent dimension score appeared to
219 exceed those observed for individual temperament deviations (Fig. 4a). The higher hazard
220 ratio indicates that the latent dimension efficiently captures the multivariate temperament
221 signal relevant to ADHD risk. Given the prominence of ADHD in both latent dimensions,
222 time-to-event analyses initially focused on this diagnosis. For comparison, we applied the

same time-to-event analysis pipeline to Asperger syndrome and social phobia, diagnoses that
were prominent in the second latent dimension. Both latent dimensions were associated with
increased hazard of Asperger syndrome (PLS 1: HR = 1.16, 95% CI 1.01–1.34; PLS 2: HR =
1.60, 95% CI 1.42–1.82). In contrast, only the second latent dimension was associated with
increased hazard of social phobia (HR = 1.19, 95% CI 1.07–1.32; Fig. S5). Notably, females
exhibited markedly higher cumulative risk of social phobia across childhood and adolescence
compared to males (Fig. S5). Together, these findings showed that deviations from
temperament trajectories are associated with both the risk and timing of psychiatric diagnosis.

Genetics of temperament trajectories

The temperament deviations examined above reflect individual differences in how
children depart from population-level developmental norms. Because these deviations capture
variation in the underlying temperament traits, we next examined the genetic architecture of
temperament itself. The FEMA-Long framework decomposes the total phenotypic variance
into individual-level, family-level and additive genetic variance components, each of which
can vary across traits and over time within a trait (Fig. S6). Focusing on the genetic
component revealed variation in its relative contribution across the four traits (Fig. 5a). For
example, the proportion of variance in *activity* attributable to the genetic component
increased from 0.41 at 1.5 years, to 0.50 at 3 years, and then decreased to 0.46 at 5 years.
Overall, these estimates indicate that the relative contribution of additive genetic variance
remains largely stable across early childhood, with only modest fluctuations over time.
To investigate genetic influences on temperament trajectories, we applied the FEMA-
Long framework to perform genome-wide association analyses (GWAS) that estimate both
main and age-dependent single nucleotide polymorphism (SNP) effects. Unlike standard
cross-sectional GWAS, this approach allows genetic effects to vary non-linearly across
development. For each temperament trait, we modeled every SNP with both a main effect and
an interaction with smooth age basis functions (see Methods). Each GWAS thus quantified
the overall genetic effect of each SNP and whether that effect changed across early
childhood. The resulting Q-Q-plots indicated polygenic enrichment (Fig. 5b, Fig. S7), though
few SNPs reached genome-wide significance ($p < 5\times10^{\Box\Box}$, Fig. S8).

The strong phenotypic association between early temperament and ADHD suggested
potential shared genetic architecture. To investigate this possibility, we applied conjunctional
false discovery rate (conjFDR) analysis[37,38] to identify SNPs jointly associated with ADHD
and each temperament trait, considering both main genetic effects and age-dependent effects
(Figs. S9–S10; Table S2). As an illustrative example, three loci with significant main effects
and two loci with significant age-dependent effects were shared between *activity* and ADHD
(Fig. 5c). Comparing the conjFDR significance of main and age-dependent effects revealed
both shared and distinct patterns of genetic association (Fig. 5d). Specifically, lead SNP
rs6484367 on chromosome 11 showed predominantly age-dependent effects, whereas
rs10789442 on chromosome 1 displayed a stronger main effect. Lead SNP rs1343667 on
chromosome 6 showed evidence of both components. These findings highlight the

contribution of age-dependent effects to the shared genetic architecture between temperament
and ADHD.
Across the four temperament traits, *activity* showed the highest number of SNPs
shared with ADHD (Fig. 5e). For lead SNPs with significant age-dependent effects in the
conjFDR analyses, we reconstructed their effect trajectories across early childhood using the
spline-based interaction terms from the fitted FEMA-Long model (Fig. 5f). Several SNPs,
such as rs1368742 (*emotionality*) and rs6484367 (*shyness*), displayed decreasing effect sizes
across age. In contrast, other SNPs (e.g., rs1343667 for *activity* and rs1397128 for *shyness*)
showed effect sizes that increased with age. Notably, these SNPs have no previously reported
associations in the GWAS Catalog[39]. Another SNP with an increasing negative effect size
(rs10789442 for *activity*) was previously associated with self-reported mathematical ability.
Finally, we mapped lead SNPs from the conjFDR analysis to genes based on genomic
position (FUMA; see Methods) and identified eight genes across all temperament traits (Fig.
5g, Table S2). *ST3GAL3* was implicated across four trait–effect combinations (*activity* main
and age-dependent effects, *shyness* main effect, *emotionality* main effect), suggesting a broad
contribution to early temperament variation. Two additional genes, *KDM4A* and *PTPRF*,
were each implicated across three combinations (*activity* main and age-dependent effects,
shyness main effect). The remaining genes (*CADM2*, *CDH8*, *SLC6A9*, *ZNF462*, and *RP11-*
508N12.4) were each associated with two or fewer trait–effect combinations. GWAS Catalog
queries indicated that seven of the eight genes have previously been associated with
cognitive, educational, psychiatric, or neurodevelopmental phenotypes, including educational
attainment, socioeconomic status, ADHD, autism spectrum disorder, and intelligence (Fig.
S11). *RP11-508N12.4*, which was not represented in the GWAS Catalog, has been linked to
migraine in a recent large-scale study[40]. Collectively, these results identify loci not previously
reported in the GWAS Catalog and characterize how genetic effects at both novel and
previously reported loci vary across early childhood.

Figure captions

Figure 1: Temperament trajectories and diagnostic profiles in the MoBa subsample
a. Temperament measurements and diagnostic counts. Temperament (*emotionality*, *activity*, *shyness*, and *sociability*) data were available for 45,028 children at 1.5 years, 36,435 at 3 years and 25,969 at 5 years (left *y*-axis). The number of recorded diagnoses increased with age (right *y*-axis). Counts are cut off at age 18 years. Approximately 71% of the cohort had reached this age at the time of data extraction. **b.** Most common diagnoses. The ten most frequent mental and behavioral disorder diagnoses recorded until the end of 2024. Full ICD-10 codes are listed in Supplementary Data 1. ADHD corresponds to F90.0 (Disorders of activity and attention). PTSD refers to Post-traumatic stress disorder (F43.1), and OCD (mixed, F42.2) refers to Obsessive–compulsive disorder with mixed obsessions and compulsions. **c.** Temperament dynamics. Each temperament trait reflects the average of several 5-level questionnaire items. Points and error bars depict mean values and interquartile ranges at each timepoint. **d.** Temporal stability. Pearson correlations between all pairs of timepoints are shown for each temperament trait. Colors denote the three timepoint comparisons. **e.** Cross-trait similarity. Heatmap displays pairwise correlations between temperament traits, averaged across timepoints. **f.** Cross-trait similarity across time. Lines trace correlations between each pair of temperament traits. Line colors correspond to the mean correlation strength of each pair. Together, the four temperament traits show complementary dynamics that set the stage for examining their links to later diagnoses.

Figure 2: Deviations from predicted temperament trajectories
a. Spline basis functions modeling non-linear age effects. Age-dependent changes in temperament were modeled using natural splines with three basis functions. **b.** Model evaluation. Bar plots display correlations between predicted and observed temperament scores, averaged across timepoints. Error bars indicate the minimum and maximum correlations across timepoints. **c.** Modeled temperament trajectories. Mean *activity* scores are shown for each month with available data, colored by timepoint. The estimated population-level *activity* trajectory from the FEMA-Long model is overlaid. The grayscale bar indicates the number of observations available at each month. **d.** Deviations from trajectories. A schematic illustration of how temperament deviations were derived, based on the average difference between observed values and the covariate-adjusted predicted trajectory across the three timepoints. Collectively, the modeled trajectories capture normative temperament development while preserving substantial individual-level variation for quantification as deviations.

Figure 3: Multivariate links between temperament deviations and diagnoses
Canonical Partial Least Squares (PLS) analysis identified latent dimensions capturing shared
variation between temperament deviations and mental and behavioral disorders diagnoses.
Temperament loadings were defined as correlations between temperament PLS scores and
the four deviation measures, and diagnosis loadings as correlations between diagnosis PLS
scores and 266 ICD-10 subcategory diagnoses. **a**. First dimension spotlights ADHD.
Temperament loadings for the first latent temperament dimension are shown with 95%
confidence intervals obtained from 10,000 bootstrap resamplings (left). Diagnosis loadings
are displayed on the right, grouped and colored according to ten overarching diagnostic
categories. **b**. Second dimension highlights Asperger syndrome and social phobia.
Temperament and diagnosis loadings for the second latent temperament dimension are shown
using the same conventions as above. **c.** Hyperkinetic disorders dominate diagnosis
association. Diagnosis loadings are shown for models fitted to ICD-10 diagnostic categories,
for the first dimension (left) and second dimension (right). Together, two latent dimensions
link distinct patterns of temperament deviations with structured diagnostic profiles across
childhood and adolescence.
340
Figure 4: Decoding ADHD diagnosis using latent dimensions
a. ADHD risk across temperament traits. Hazard ratio of the latent temperament dimensions
is displayed along with hazard ratios of temperament deviations. Error bars depict 95%
confidence interval. **b.** ADHD risk and first latent temperament dimension. Kaplan–Meier
curves show cumulative ADHD risk for three groups stratified by the first latent dimension.
Curves correspond to low, intermediate, and high scores on the first dimension. Cumulative
risk is calculated as one minus the probability of remaining ADHD-free up to each age. **c.**
Sex-specific ADHD risk. Kaplan–Meier curves display the cumulative probability of
receiving an ADHD diagnosis separately for males and females, reflecting sex-specific
differences in ADHD risk across development. Taken together, latent dimensions capture
variation in ADHD risk and are associated with differences in the timing of diagnosis.

Figure 5: Dynamic genetic effects linking early temperament and ADHD

a. Temporal dynamics of genetic variance. Genetic variance estimates for each temperament trait at 1.5, 3, and 5 years, as derived from FEMA-Long. **b.** Enrichment of GWAS signal across main and age-dependent genetic effects. Each panel shows a deviation-based Q-Q plot, where the *x*-axis represents the difference between observed and expected $-\log_{10}(p)$. Curves show main genetic effects and age-dependent effects (see legend). Leftward deviation of the main-effect curve indicates enrichment of main genetic effects, whereas rightward deviation of the longitudinal-effect curve indicates enrichment of age-dependent effects. Deviations in the opposite direction reflect deflation. **c.** Shared SNPs between *activity* and ADHD. Miami plots show SNPs jointly associated with ADHD and *activity* for the main (bottom) and age-dependent (up) genetic effects. The *y*-axis displays $-\log_{10}$ conjFDR *p*-values. Black circles mark significant lead SNPs within linkage disequilibrium blocks. **d.** Comparison of association strength for main and age-dependent effects. Transformed conjFDR *p*-values for main and age-dependent genetic effects are plotted for all SNPs. **e.** Lead SNP overlap between temperament traits and ADHD. Bar plots show the number of significant lead SNPs jointly associated with ADHD and each temperament trait, separately for main and age-dependent effects. Dashed segments indicate SNPs previously reported in the GWAS Catalog. **f.** Trajectories of age-dependent effects. Estimated age-dependent genetic effects are shown across age for all significant lead SNPs identified in the conjFDR analyses. **g.** Gene-level overlap between temperament traits and ADHD. Bar plots show the number of genes corresponding to lead SNPs shared between ADHD and each temperament trait. Together, the results highlight dynamic genetic influences on temperament, with several effects emerging from genomic regions distinct from those implicated by age-independent (main) genetic effects.

Tables

Table 1: EAS questionnaire items and corresponding temperament traits

Each item is mapped to one of the EAS domains (*emotionality*, *activity*, *shyness*, *sociability*).
Items marked with an asterisk were reverse-scored so that higher values consistently indicate
higher levels of the corresponding temperament trait. All items were rated on a 1–5 scale.

Temperament trait	Question
Emotionality	Cries easily
Emotionality	Reacts intensely when upset
Emotionality	Gets upset or sad easily
Activity	Off and running as soon as wakes up
Activity	Always on the go
Activity	Prefers quiet, inactive games*
Shyness	Very friendly with strangers*
Shyness	Takes a long time to warm up to strangers
Shyness	Very sociable*
Sociability	Likes to be with people
Sociability	Prefers playing with others rather than alone

Table 2: Sample sizes after application of inclusion criteria

Sample sizes at each temperament assessment wave (1.5, 3, and 5 years) after applying each inclusion criterion sequentially. Criteria included: available genotype data from MoBaPsychGen pipeline v115[34], recorded biological sex, valid assessment age (child aged 1–6 years at questionnaire completion), and complete EAS questionnaire data (all items answered). The final row shows the analysis sample contributing valid temperament data at each timepoint. Of the final analysis sample, 18,651 children contributed valid temperament data at all three timepoints.

	1.5 years	3 years	5 years
Initial MoBa cohort	75,876	58,460	41,408
+ Recorded biological sex	66,103	53,737	39,677
+ Available genotype data	47,868	39,059	28,636
+ Valid assessment age (1–6 years)	47,868	39,059	28,636
+ Complete EAS data	45,028	36,435	25,969

Discussion

In this study, we leveraged a longitudinal birth cohort to characterize how early-childhood temperament deviations relate to psychiatric outcomes in early adulthood. We identified two complementary temperament deviation dimensions associated with risk profiles spanning multiple diagnoses, including ADHD, Asperger syndrome, and social phobia. Notably, ADHD was linked to both dimensions, which suggests that the disorder may arise from multiple developmental pathways. Additionally, we uncovered a shared genetic basis between temperament trajectories and ADHD. By modeling temperament trajectories, we identified age-dependent genetic effects at shared loci. Genetic liability for psychopathology thus appears to be developmentally regulated, with effects that may be missed by cross-sectional approaches.

Temperament is only moderately stable across early childhood[41] ($r = 0.27$–0.53 in our data), meaning that a single measurement cannot distinguish a child's stable behavioral tendency from age-specific fluctuation. By modeling temperament across three assessments, we evaluated each child against population-level developmental norms and averaged the resulting deviations across timepoints. These deviations quantify what clinicians observe informally: some children are consistently more active or emotional than expected for their age and sex. Whereas prior approaches linked individual traits to specific disorders at single timepoints[10], entering deviations into a multivariate analysis revealed that coordinated patterns across traits mapped onto distinct patterns of psychiatric risk. More pronounced deviations were associated with a higher hazard of diagnosis. The trajectory-deviation approach thus captures dimensions of transdiagnostic risk that are invisible to single-timepoint, single-trait designs.

The two identified temperament deviation dimensions captured distinct but complementary clinical patterns. In the first, elevated *activity* and *emotionality* alongside reduced *sociability* were associated with ADHD and related neurodevelopmental diagnoses including Tourette's syndrome, hyperkinetic conduct disorder, and specific reading disorder. These co-occurring diagnoses are consistent with the high comorbidity observed between ADHD and other neurodevelopmental conditions[42,43]. In the second, elevated *shyness* and *emotionality* alongside reduced *activity* and *sociability* were most strongly associated with Asperger syndrome, social phobia, and generalized anxiety, capturing a shared phenotype of social withdrawal that spans neurodevelopmental and internalizing boundaries[30,44,45]. What differentiated the two profiles was *activity* versus *shyness*: elevated *activity* characterized the externalizing-like profile, while elevated *shyness* characterized the social withdrawal profile. *Emotionality* was elevated in both, which suggests a transdiagnostic role that cuts across domains[46–48]. These complementary patterns align with hierarchical models that frame early developmental variation as broad vulnerability rather than disorder-specific risk[49].

ADHD offers a clear illustration of this broad vulnerability pattern. It was linked to both latent dimensions, but with elevated *emotionality* in the first and reduced *emotionality* in the second. This means that children arriving at the same ADHD diagnosis depart from expected temperament development in opposite directions on *emotionality*: some through

heightened emotional reactivity, others through reduced reactivity. This divergence may help
explain the disorder's well-documented clinical heterogeneity[42,43]. If distinct temperament
profiles converge on the same diagnosis, then apparent heterogeneity within ADHD may
partly reflect different developmental origins. These origins may later diverge into different
comorbidity patterns, which may help explain why ADHD so frequently precedes more
differentiated psychiatric outcomes[50]. The strength of the phenotypic link between
temperament deviations and ADHD raises the question of whether they also share genetic
architecture.
442
Using the FEMA-Long framework to estimate age-dependent genetic effects, we
identified genetic loci jointly associated with temperament trajectories and ADHD. These
variants exhibited both main and age-dependent effects on temperament across early
childhood. The presence of both types of effects is consistent with classical quantitative
genetic evidence that genetic factors contribute to both stability and change in temperament
during development[17,51]. It also aligns with longstanding distinctions between genetic
influences that support continuity and those associated with developmental change[3,52,53]. Our
findings extend this work at the molecular level. Shared genetic influences between
temperament and ADHD are not uniform. Some loci showed predominantly main effects,
others predominantly age-dependent effects, and some showed evidence of both. Moreover,
the direction of age-dependent effects varied, with some increasing and others decreasing.
Together, these results indicate that the shared genetic architecture between temperament and
ADHD is partly dynamic, with effects that would be missed by analyses assuming static
genetic effects.
The implicated genes have been previously associated with cognitive ability,
educational attainment, and multiple psychiatric conditions beyond ADHD. This suggests
that the genetic overlap between temperament and ADHD is embedded within a broader
architecture spanning cognition, self-regulation, and neurodevelopmental risk[54,55]. The
genetic breadth parallels the phenotypic picture, where different temperament profiles
converge on the same diagnostic outcomes. Furthermore, the wide phenotypic reach of the
identified loci suggests that age-dependent genetic effects extend beyond the temperament-
ADHD link alone. More broadly, genetic liability for ADHD, and likely other disorders,
appears to be developmentally regulated.
A key question is whether current approaches to genetic risk prediction adequately
capture such developmental dynamics. Polygenic risk scores represent the primary tool for
translating genetic findings into individual risk prediction. They aggregate the effects of
many variants into a single measure of liability[56]. Current implementations, however,
typically assume that these genetic effects are constant across development[57]. As such, they
may not capture the developmentally varying influences identified here. Incorporating age-
dependent genetic effects into risk prediction frameworks could distinguish children whose
genetic risk is expressed early versus later. This would in turn enable more precisely timed
monitoring or intervention.
475
For clinical practice, monitoring deviations from expected developmental trajectories
may offer an additional signal for identifying children at elevated risk before formal

diagnosis. Much like pediatricians track height and weight against population norms and flag
children who deviate, repeated temperament assessment could allow clinicians to plot a
child's development against age-expected trajectories, with persistent departures signaling the
need for follow-up. In our analyses, such deviations were associated with an increased hazard
of psychiatric diagnosis across childhood and adolescence. Risk patterns differed by sex, with
males showing higher cumulative ADHD risk and females a higher risk for social phobia.
These patterns mirror well-established sex differences in the prevalence of both disorders[58,59]
and underscore the importance of incorporating biological and environmental context into
monitoring frameworks[60]. Temperament questionnaires such as the EAS are brief and
inexpensive. Their integration into routine developmental monitoring is therefore feasible in
principle. Prospective studies are now needed to test whether deviation-based monitoring,
combined with age-dependent polygenic risk scores, can translate into earlier identification
and improved outcomes.
491

Some limitations should be considered when interpreting our findings. Registry-based
psychiatric diagnoses capture clinically recognized disorders but lack symptom-level
granularity and reflect administrative timing rather than biological onset, which may
introduce lag into survival estimates. The birth cohort design favors capture of early-onset
diagnoses like ADHD over later-onset conditions like bipolar disorder or schizophrenia, and
cross-diagnostic comparisons are therefore limited. Temperament was assessed through
parent reports, which may be influenced by reporting biases. Our longitudinal, deviation-
based approach partially mitigates this concern by emphasizing within-individual change
rather than absolute levels, although systematic reporting biases could still influence
trajectory estimates. Loss to follow-up in this cohort is non-random and can lead to both
attenuation and inflation of associations[61]. Additionally, temperament measures may overlap
with early symptoms of the disorders they predict. *Activity* items, for instance, resemble early
ADHD symptoms. However, this reflects genuine continuity between normal temperament
variation and clinical disorder[62] rather than a methodological artifact. Finally, the cohort is
drawn from a European population-based sample and genetic analyses were restricted to
participants of European ancestry. Findings may therefore have reduced generalizability to
populations with different ancestral backgrounds, healthcare systems, or sociocultural
contexts.
510

In conclusion, we show that deviations from expected temperament trajectories in
early childhood are associated with psychiatric diagnoses across neurodevelopmental and
internalizing domains. ADHD emerged as a particularly salient outcome, associated with
distinct patterns of early temperament deviation. These associations are underpinned by
shared genetic architecture between temperament and ADHD that is itself partly dynamic,
with effects that vary across early childhood. Together, these findings suggest that early
psychopathology risk is best understood not as the presence of specific traits, but as
deviations from expected trajectories shaped by dynamically expressed genetic influences.
Linking trajectory deviations to long-term psychiatric risk supports a shift from reactive
diagnosis toward proactive, developmentally informed monitoring of mental health risk.

Methods

MoBa: A nationwide population cohort

The Norwegian Mother, Father and Child Cohort Study (MoBa) is a population-based pregnancy cohort study conducted by the Norwegian Institute of Public Health[33]. Participants were recruited from all over Norway from 1999–2008. The women consented to participation in 41% of the pregnancies. The cohort includes approximately 114,500 children, 95,200 mothers and 75,200 fathers. All participants provided written informed consent; children are included after consent from the mother, once they reach 18 years old. MoBa requires informed consent from the participant for further use and storage of the data. Umbilical cord blood collected at birth was used for genotyping[63]. MoBa was linked to the Medical Birth Registry of Norway (MBRN), a compulsory national registry containing clinician-reported information on all Norwegian births. The present study used version 12 of the quality-assured maternally reported questionnaire data, released in January 2019. Linkage to the Norwegian Patient Registry (NPR) provided ICD-10 diagnoses from 2008 through 2024, enabling longitudinal follow-up of health outcomes. Analyses were restricted to participants who had not withdrawn consent from MoBa as of April 25, 2025. Quality-controlled genotype data for 75,807 children were obtained from the MoBaPsychGen pipeline v115 (described previously[34]). Sex assigned at birth and birth year were extracted from the 6-month questionnaire. Family membership was defined by maternal identifier, such that children sharing the same mother were treated as a single family unit.

The establishment and data collection of MoBa were approved by the Norwegian Data Protection Agency and the Regional Committees for Medical and Health Research Ethics (REK). MoBa operates under the Norwegian Health Registry Act. The current study was approved by the MoBa administrative board at the Norwegian Institute of Public Health (NIPH) and REK (reference 2016/1226/REK Sør-Øst C).

Temperament assessments across early childhood

Temperament was assessed using maternally reported Emotionality, Activity, Shyness, and Sociability Temperament questionnaire (EAS)[13,14] administered when children were approximately 1.5, 3 and 5 years old. The EAS measures four temperament traits: *emotionality* (irritability/anger), *activity* (activity level), *shyness* (fear), and *sociability* (positive affect/including approach). The original 20-item questionnaire assesses these four traits using five items per trait. In MoBa, only 11 items were administered consistently across all three timepoints. Analyses were therefore restricted to these items (listed in Table 1). Mothers rated each item on a five-point scale from "not at all typical" to "very typical" of their child. For each temperament trait, we computed summary scores by averaging available item responses. Analyses included children with available genotype data, recorded biological sex, and at least one valid EAS assessment. Valid assessments were defined as questionnaires completed when the child was aged 1–6 years with no missing items. The sequential application of these criteria and resulting sample sizes are detailed in Table 2. After all

560 exclusions, 45,028 children contributed temperament data at 1.5 years, 36,435 at 3 years, and
561 25,969 at 5 years, with 18,651 children measured at all three timepoints.

Diagnostic data from the Norwegian Patient Registry

Diagnoses were obtained from the NPR[64], which records ICD-10 diagnostic codes[65]
from specialist healthcare encounters in Norway, including inpatient and outpatient hospital
services and privately contracted specialist care. In this study, we focused on mental and
behavioral disorders (ICD-10 codes, chapter V: F00–F99), as access to somatic diagnoses
was limited. Diagnostic information was available from January 2008 through December
2024. The NPR linkage procedure used in the present MoBa analyses is described in detail in
[66]. A complete list of diagnoses and their frequencies is provided in Supplementary Data 1.
We also extracted diagnoses from the KUHR database (Norwegian Control and
Payment of Health Reimbursements Database), which contains reimbursement records for
healthcare encounters funded through the Norwegian National Insurance Scheme, including
primary care consultations and some specialist services not captured by NPR. Notably, these
nationwide records are available from 2006, compared with 2008 for NPR. Yet, KUHR
contributed only a few additional diagnoses beyond those already recorded in NPR (Fig. S3).
Given this observation, and because NPR diagnoses reflect specialist healthcare assessments
with more comprehensive clinical evaluation, all main analyses were restricted to NPR
diagnoses.
ICD-10 codes follow a hierarchical structure ranging from broad categories to
increasingly specific subcodes. We used both detailed ICD-10 codes (four- and five-character
subcategory codes, e.g., F90.1) and the corresponding higher-level code categories (e.g.,
F90), depending on the analytic context. For each condition, the date of first recorded
diagnosis was used in all time-to-event analyses.

Longitudinal modeling of temperament trajectories

Temperament trajectories were modeled using the FEMA-Long framework
v3.0.0[35,36]. FEMA-Long is a scalable mixed-effects modeling framework specifically
designed for longitudinal population cohorts with correlated observations. The method
extends classical mixed-effects modeling by allowing random effects to vary over time,
which enables flexible estimation of the covariance structure without imposing restrictive
parametric assumptions on temporal dependence. In the present study, random effects were
specified to capture individual, family, and genetic similarity, enabling decomposition of
variance into individual, familial, and genetic components.
Each temperament trait was modeled separately. The covariates included age
(modeled nonlinearly as described below), biological sex, the first twenty genetic principal
components[34], and genetic batch indicators. All continuous covariates were mean-centered
and scaled to unit variance prior to modeling. Covariate effects, except for age, were modeled
as time-invariant fixed effects.
Rather than specifying age as a strictly linear effect, we used a natural cubic spline
basis expansion of age with an interior knot at 3 years and boundary knots at the minimum
and maximum observed ages, ensuring adequate flexibility across the early-childhood

window. To avoid collinearity with the intercept, the three basis functions were
reparameterized via singular value decomposition (SVD), yielding two orthogonal
components that, together with the intercept, span the same function space and ensure a full-
rank design matrix.
Genome-wide genetic relatedness was captured via a genetic relationship matrix
computed across 500,243 SNPs that were either directly genotyped in at least one imputation
batch or had an imputation INFO score exceeding 0.99785 across all imputation batches[34].
This matrix was included as a random-effect covariance component within the FEMA-Long
framework.
610
Genome-wide association analyses were performed using FEMA-Long framework
applied to 6,981,748 SNPs[34]. For each temperament trait, FEMA-Long estimated the main
effect of the SNP, as well as the age-dependent SNP effect (modeled as an interaction of the
SNP with the smooth basis functions). To test whether a SNP exhibited a significant age-
varying effect, we performed an omnibus Wald test jointly testing all spline coefficients
associated with that SNP (contrast matrix equal to the identity over the spline terms).
Manhattan plots display *p*-values for the main effect and Wald-test *p*-values for the age-
dependent effect, allowing visual comparison of static and dynamic genetic influences on
temperament. Genome-wide significance was defined using the conventional threshold of
$5 \times 10^{\square\square}$.

Multivariate analysis of temperament-diagnosis associations

Constructing the inputs for multivariate analysis

We quantified temperament deviations for the 18,651 children with complete
measurements across all three timepoints (Fig. S12). For each child, we generated expected
temperament values using the fixed-effect component of the FEMA-Long model, which
estimates fixed-effects coefficients while accounting for the covariance structure induced by
the random effects. Deviations were then computed as the signed difference between
observed and model-estimated temperament values and averaged across the three timepoints.
We retained the signed differences to capture persistent upward or downward deviations from
the expected developmental trajectory. This yielded a single deviation score capturing the
extent to which a child consistently differed from the model-estimated developmental
trajectory. Resulting deviation measures served as the basis for linking temperament
dynamics to healthcare diagnoses.
To obtain a stable, low-dimensional representation of the diagnostic data, we applied
principal component analysis (PCA) to the binary ICD-10 matrix. We retained components
accounting for 95% of the variance, which corresponded to the first 100 components (Fig.
S1). This resulted in two aligned input matrices for subsequent multivariate modeling: an
$18,651 \times 4$ matrix of temperament deviations and an $18,651 \times 100$ matrix of diagnostic
components.

Latent dimensions linking temperament and diagnosis

We leveraged canonical Partial Least Squares (PLS) to identify multivariate associations between temperament deviations and diagnostic profiles[67]. PLS extracts latent dimensions that maximize the covariance between the two high-dimensional datasets. These dimensions are orthogonal and ordered by the amount of cross-covariance they explain, with the first dimension capturing the dominant shared structure.

To determine how many PLS dimensions reflected signal rather than noise, we compared the observed covariance of each dimension with a permutation-derived null distribution. We generated the null distribution by repeating the full PLS procedure 10,000 times. In each iteration, we permuted the temperament matrix at the family level by shuffling family blocks (keeping siblings together), while leaving the diagnostic data unchanged, thereby preserving within-family dependence. For each iteration, we recorded the covariance explained by the first PLS dimension, yielding a null distribution of covariances expected under no association. For each observed PLS dimension, the p-value was computed as the proportion of permutations in which the permuted covariance exceeded the observed covariance. Dimensions with $p < 0.05$ were retained (Fig. S1).

Phenotypic contributions to latent dimensions

To quantify how individual variables contributed to each latent dimension, we computed PLS loadings as the Pearson correlation between the individual-level PLS scores and each original variable. These loadings capture both the strength and direction of association between a variable and a given latent dimension. Higher absolute loadings indicate variables that contribute more strongly to the underlying pattern of covariation, thereby identifying the traits and diagnoses that most prominently shape each PLS dimension.

Finally, we employed a cluster bootstrap resampling procedure to evaluate the stability of the PLS loadings. In each of 10,000 iterations, families were resampled with replacement to generate bootstrap samples of the same size as the original dataset, with all siblings from selected families retained together to preserve within-family dependence. The ordering and orientation of PLS dimensions were aligned across iterations to ensure comparability of loadings. This yielded 10,000 bootstrap estimates for each loading, forming an empirical sampling distribution. A loading was deemed robust if its two-sided 95% confidence interval (2.5th to 97.5th percentile) did not include zero, indicating a consistent contribution to the latent dimension across resamples.

Time-to-event analysis of ADHD diagnosis

Time-to-event analyses were performed using the *lifelines* Python package (v0.30.0). To examine associations between latent temperament dimensions and the timing of psychiatric diagnosis, we fitted Cox proportional hazards models[68] using age (in months) as the time scale. For each child, time-to-event was defined as either the age at first recorded diagnosis or the age at last registry entry for those without a diagnosis. Predictors included the PLS scores along with the covariates used in the FEMA-Long models (first twenty genetic principal components). Year of birth and sex were included as stratification variables, allowing the baseline hazard to vary across strata rather than assuming proportionality for

these predictors. Models used age-based delayed entry, such that participants entered the risk
set at age 5 and were followed until diagnosis or censoring at age 21. Left truncation at age 5
ensured that exposure measures were fully defined before participants entered the risk set.
We used cluster-robust variance estimates with family ID as the clustering variable
(cluster_col='fid', robust=True). Full model outputs are provided in Table S1.

We assessed the proportional hazards assumption using tests and visual inspection of
scaled Schoenfeld residuals (implemented in *lifelines*). Sex showed clear evidence of non-
proportional hazards for ADHD diagnosis (ICD-10: F90.0) and was therefore treated as a
stratification variable, allowing sex-specific baseline hazards. For the PLS dimensions,
diagnostics indicated only minor deviations from proportionality. Cox model estimates are
thus interpreted as average effects over the follow-up period. Full diagnostics are provided in
Fig. S13.

To visualize cumulative diagnosis risk across levels of the temperament-derived
dimensions, participants were stratified into quantile-based groups according to their latent
dimension scores. Kaplan–Meier curves were then constructed for each group to estimate the
survival function, that is, the probability of remaining diagnosis-free up to a given age.
Cumulative risk was expressed as *1-S(t)*, corresponding to the proportion of participants
diagnosed by age *t*. Furthermore, we examined sex differences by generating separate
Kaplan–Meier curves for males and females, without further stratification by PLS score.
Together, these plots provide complementary perspectives on how variation in the latent
temperament dimensions and sex relate to cumulative diagnosis risk over development.

As a sensitivity analysis, we generated Kaplan–Meier curves using a reference
covariate profile. Continuous predictors were fixed at their sample means, categorical
predictors at their reference categories and year of birth at the median stratum. Only the latent
dimension scores were varied. This allowed us to visualize diagnosis risk for a typical
participant and verify that the observed survival differences arose from variation in the latent
temperament dimensions rather than covariate heterogeneity (Fig. S14).

A limitation of NPR is that nationwide coverage began in 2008. For children born
before 2004, the registry gap potentially overlaps with the early risk period following
temperament assessment. For the example of ADHD, twelve children with ADHD diagnosis
in the analysis sample were born in 2002–2003. Therefore, their first recorded diagnosis
occurred at ages 6–7, but could have occurred earlier without being captured, leading to
overestimated time-to-event times. Excluding these twelve children from the time-to-event
analysis yielded estimates identical to the main analysis, confirming that this potential bias
does not materially affect the reported associations.

Identification of shared genetic loci

Given the strong phenotypic association between temperament and ADHD, we
examined whether the two phenotypes share genetic architecture. To identify jointly
associated loci, we applied conjFDR analysis[37,38]. ConjFDR extends the conditional FDR
(condFDR) framework by re-ranking association statistics for a primary phenotype

conditional on their association with a secondary phenotype. The re-ranking exploits cross-
trait pleiotropy to enhance detection of shared genetic signals. For each pair of traits,
condFDR is estimated in both directions, and conjFDR is defined as the maximum of the two
corresponding condFDR values, providing a conservative estimate of the probability that a
variant is a false positive jointly for both traits. Genomic inflation was controlled using the
standard empirical null correction implemented in the conjFDR framework[37,38]. Consistent
with prior studies, loci with conjFDR $p < 0.05$ were considered jointly associated. We
excluded variants from the major histocompatibility complex (MHC) region, the 8p23.1
locus, and the 17q21.31 inversion region to limit inflation driven by the extended linkage
disequilibrium structure characteristic of these regions.

As input to conjFDR analyses, we used summary statistics from genome-wide
association studies. ADHD summary statistics were obtained from the Psychiatric Genomics
Consortium[69] (PGC; 38,691 cases and 186,843 controls). For temperament, we used GWAS
summary statistics derived from FEMA-Long, including both main genetic effects and age-
dependent genetic effects.

To identify lead SNPs in each conjFDR analysis, we performed LD-based clumping
using a threshold of $r^2 < 0.6$ to obtain a set of approximately independent index variants. For
each index SNP, all variants in high linkage disequilibrium ($r^2 \geq 0.6$) were identified using
the 1000 Genomes Project European reference panel. Loci located within 250 kb of each
other were subsequently merged, and within each merged locus, the SNP with the lowest
conjFDR value was selected as the lead SNP.

Gene mapping and phenotypic annotation

To functionally characterize genetic loci identified in the conjFDR analyses, we
annotated candidate SNPs using the SNP2GENE pipeline implemented in FUMA[70].
Candidate SNPs were defined as all variants reaching conjFDR $p < 0.05$, including
independent lead SNPs as well as all SNPs in linkage disequilibrium with these lead variants
($r^2 \geq 0.6$). SNP-to-gene mapping was performed using the default FUMA settings based on
positional mapping. To contextualize the resulting genes, we queried their previously
reported trait associations in the GWAS Catalog[39]. For each mapped gene, we extracted the
number of reported associations per gene and the frequency of each trait across genes.

Acknowledgements

The Norwegian Mother, Father and Child Cohort Study is supported by the Norwegian Ministry of Health and Care Services and the Ministry of Education and Research. We are grateful to all the participating families in Norway who take part in this on-going cohort study. For generating high-quality genomic data, we thank the Norwegian Institute of Public Health (NIPH), the HARVEST collaboration, the NORMENT Centre at the University of Oslo, the Center for Diabetes Research at the University of Bergen, deCODE Genetics, the Research Council of Norway, the South-Eastern and Western Norway Regional Health Authorities, the ERC AdG, Stiftelsen KG Jebsen, the Trond Mohn Foundation, and the Novo Nordisk Foundation.

This work was partly performed on the TSD (Tjeneste for Sensitive Data) facilities, owned by the University of Oslo, operated and developed by the TSD service group at the University of Oslo, IT-Department (USIT). Computations were also performed on resources provided by UNINETT Sigma2 - the National Infrastructure for High-Performance Computing and Data Storage in Norway (NS9703S).

The authors were funded by the Research Council of Norway (296030, 324252, 324499, 326813, 334920, 351751); the South-Eastern Norway Regional Health Authority (2020060 (IES), 2025037 (IES)); European Union's Horizon 2020 Research and Innovation Programme (Grant No. 847776; CoMorMent and Grant No. 964874; RealMent); EU's Horizon Psych-STRATA project (#101057454); Stiftelsen KG Jebsen (SKGJ-MED-021), National Institutes of Health (NIH; U24DA041123; R01AG076838; U24DA055330; and OT2HL161847). JK was supported by a Marie Skłodowska-Curie Postdoctoral Fellowship under the European Union's Horizon Europe research and innovation programme (Grant Agreement No. 101150746). EPT was supported by a Rubicon Postdoctoral Fellowship of ZonMW (Grant No. 04520242420017). PP is supported by the National Institutes of Health grants U24DA041123 and U24DA055330, and by Wellcome Leap, CARE Program ("FEMA-AD").

Data availability

Data from the Norwegian Mother, Father and Child Cohort Study (MoBa) and the Medical Birth Registry of Norway are administered by the Norwegian Institute of Public Health and are not publicly available due to ethical and legal restrictions. Access may be granted to qualified researchers following approval from the Regional Committees for Medical and Health Research Ethics (REC), compliance with the EU General Data Protection Regulation (GDPR), and permission from the data owners. Participant consent does not allow for deposition of individual-level data in public repositories. Researchers seeking access for replication or further analyses must apply through the Norwegian health data access portal (helsedata.no) and enter a data access agreement with MoBa.

Code availability

FEMA-Long is available on GitHub at: https://github.com/cmig-research-group/cmig_tools.

conjFDR is available on GitHub at: https://github.com/precimed/pleiofdr.

Custom analysis software used in this work is available in a publicly accessible GitHub repository: https://github.com/jakubkopal/temperamet_deviations

Conflicts of interest

OAA has received speaker's honorarium from Lundbeck, Janssen, Otsuka, and Sunovion and is a consultant to Cortechs.ai. and Precision Health. AMD is Founding Director, holds equity in CorTechs Labs, Inc. (DBA Cortechs.ai), and serves on its Board of Directors. AMD is the President of J. Craig Venter Institute (JCVI) and is a member of the Board of Trustees of JCVI. He is an unpaid consultant for Oslo University Hospital. OF is a consultant to Precision Health. All other authors report no potential conflicts of interest.

References

1. Thomas, A. & Chess, S. *Temperament and Development*. xv, 270 (Brunner/Mazel, Oxford, England, 1977).

2. Rothbart, M. K. Temperament and the Pursuit of an Integrated Developmental Psychology. *Merrill-Palmer Q.* **50**, 492–505 (2004).

3. Saudino, K. J. Behavioral Genetics and Child Temperament. *J. Dev. Behav. Pediatr. JDBP* **26**, 214–223 (2005).

4. Goldsmith, H. H., Buss, K. A. & Lemery, K. S. Toddler and childhood temperament: expanded content, stronger genetic evidence, new evidence for the importance of environment. *Dev. Psychol.* **33**, 891–905 (1997).

5. Goldsmith, H. H. *et al.* Roundtable: what is temperament? Four approaches. *Child Dev.* **58**, 505–529 (1987).

6. Rothbart, M. K. *Becoming Who We Are: Temperament and Personality in Development*. xi, 324 (The Guilford Press, New York, NY, US, 2011).

7. Shiner, R. L. *et al.* What Is Temperament Now? Assessing Progress in Temperament

Research on the Twenty-Fifth Anniversary of Goldsmith et al. (). *Child Dev. Perspect.* **6**,

436–444 (2012).

8. Saudino, K. J. & Micalizzi, L. Emerging Trends in Behavioral Genetic Studies of Child

Temperament. *Child Dev. Perspect.* **9**, 144–148 (2015).

9. Posner, M. I., Rothbart, M. K. & Sheese, B. E. Attention genes. *Dev. Sci.* **10**, 24–29

(2007).

10. Rothbart, M. K. & Bates, J. E. Temperament. in *Handbook of child psychology: Social,*

emotional, and personality development, Vol. 3, 6th ed 99–166 (John Wiley & Sons, Inc.,

Hoboken, NJ, US, 2006).

11. Cicchetti, D. Development and psychopathology. in *Developmental psychopathology:*

Theory and method, Vol. 1, 2nd ed 1–23 (John Wiley & Sons, Inc., Hoboken, NJ, US,

2006).

12. Fox, N. A., Henderson, H. A., Rubin, K. H., Calkins, S. D. & Schmidt, L. A. Continuity

and discontinuity of behavioral inhibition and exuberance: psychophysiological and

behavioral influences across the first four years of life. *Child Dev.* **72**, 1–21 (2001).

13. Buss, A. H. & Plomin, R. *Temperament: Early Developing Personality Traits*. (Lawrence

Erlbaum, Hillsdale, NJ, 1984).

14. Mathiesen, K. S. & Tambs, K. The EAS temperament questionnaire--factor structure, age

trends, reliability, and stability in a Norwegian sample. *J. Child Psychol. Psychiatry* **40**,

431–439 (1999).

15. Ohashi, Y. & Kitamura, T. The EASI: Factor Structure and Measurement and Structural

Invariance between the Parent's Gender, the Child's Age, and Two Measurement Time

Points. *Psychology* **10**, 2177–2189 (2019).

16. Bould, H., Joinson, C., Sterne, J. & Araya, R. The Emotionality Activity Sociability

Temperament Survey: Factor analysis and temporal stability in a longitudinal cohort.

Personal. Individ. Differ. **54**, 628–633 (2013).

17. Saudino, K. J., Plomin, R. & DeFries, J. C. Tester-rated temperament at 14, 20 and 24

months: Environmental change and genetic continuity. *Br. J. Dev. Psychol.* **14**, 129–144

(1996).

18. Caspi, A., Elder Jr., G. H. & Bem, D. J. Moving away from the world: Life-course

patterns of shy children. *Dev. Psychol.* **24**, 824–831 (1988).

19. Wright, A. J. & Jackson, J. J. Childhood temperament and adulthood personality

differentially predict life outcomes. *Sci. Rep.* **12**, 10286 (2022).

20. Sayal, K., Heron, J., Maughan, B., Rowe, R. & Ramchandani, P. Infant temperament and

childhood psychiatric disorder: longitudinal study. *Child Care Health Dev.* **40**, 292–297

(2014).

21. Caspi, A., Moffitt, T. E., Newman, D. L. & Silva, P. A. Behavioral observations at age 3

years predict adult psychiatric disorders. Longitudinal evidence from a birth cohort. *Arch.*

Gen. Psychiatry **53**, 1033–1039 (1996).

22. Tang, A. *et al.* Shyness Trajectories across the First Four Decades Predict Mental Health

Outcomes. *J. Abnorm. Child Psychol.* **45**, 1621–1633 (2017).

23. Shephard, E. *et al.* Systematic Review and Meta-analysis: The Science of Early-Life

Precursors and Interventions for Attention-Deficit/Hyperactivity Disorder. *J. Am. Acad.*

Child Adolesc. Psychiatry **61**, 187–226 (2022).

24. Goodwin, A. *et al.* Behavioural Measures of Infant Activity but Not Attention Associate

with Later Preschool ADHD Traits. *Brain Sci.* **11**, (2021).

25. Joseph, H. M. *et al.* Research Review: A systematic review and meta-analysis of infant

and toddler temperament as predictors of childhood attention-deficit/hyperactivity

disorder. *J. Child Psychol. Psychiatry* **64**, 715–735 (2023).

26. Forbes, M. K., Rapee, R. M., Camberis, A.-L. & McMahon, C. A. Unique Associations

between Childhood Temperament Characteristics and Subsequent Psychopathology

Symptom Trajectories from Childhood to Early Adolescence. *J. Abnorm. Child Psychol.*

45, 1221–1233 (2017).

27. Chronis-Tuscano, A. *et al.* Stable early maternal report of behavioral inhibition predicts

lifetime social anxiety disorder in adolescence. *J. Am. Acad. Child Adolesc. Psychiatry*

48, 928–935 (2009).

28. Bakken, N. R. *et al.* Childhood temperamental, emotional, and behavioral characteristics

associated with mood and anxiety disorders in adolescence: A prospective study. *Acta*

Psychiatr. Scand. **147**, 217–228 (2023).

29. Kostyrka-Allchorne, K., Wass, S. V. & Sonuga-Barke, E. J. S. Research Review: Do

parent ratings of infant negative emotionality and self-regulation predict psychopathology

in childhood and adolescence? A systematic review and meta-analysis of prospective

longitudinal studies. *J. Child Psychol. Psychiatry* **61**, 401–416 (2020).

30. Kagan, J., Reznick, J. S. & Snidman, N. The physiology and psychology of behavioral

inhibition in children. *Child Dev.* **58**, 1459–1473 (1987).

31. De Pauw, S. S. W. & Mervielde, I. Temperament, personality and developmental

psychopathology: a review based on the conceptual dimensions underlying childhood

traits. *Child Psychiatry Hum. Dev.* **41**, 313–329 (2010).

32. Nigg, J. T. Temperament and developmental psychopathology. *J. Child Psychol.*

Psychiatry **47**, 395–422 (2006).

33. Brandlistuen, R. E. *et al.* Cohort Profile Update: The Norwegian Mother, Father and

Child Cohort (MoBa). *Int. J. Epidemiol.* **54**, dyaf139 (2025).

34. Corfield, E. C. *et al.* The Norwegian Mother, Father, and Child cohort study (MoBa)

genotyping data resource: MoBaPsychGen pipeline v.1. 2022.06.23.496289 Preprint at

https://doi.org/10.1101/2022.06.23.496289 (2024).

35. Parekh, P. *et al.* FEMA: Fast and efficient mixed-effects algorithm for large sample

whole-brain imaging data. *Hum. Brain Mapp.* **45**, e26579 (2024).

36. Parekh, P. *et al.* FEMA-Long: Modeling unstructured covariances for discovery of time-

dependent effects in large-scale longitudinal datasets. *BioRxiv Prepr. Serv. Biol.*

2025.05.09.653146 (2025) doi:10.1101/2025.05.09.653146.

37. Andreassen, O. A. *et al.* Improved Detection of Common Variants Associated with

Schizophrenia and Bipolar Disorder Using Pleiotropy-Informed Conditional False

Discovery Rate. *PLOS Genet.* **9**, e1003455 (2013).

38. Smeland, O. B. *et al.* Discovery of shared genomic loci using the conditional false

discovery rate approach. *Hum. Genet.* **139**, 85–94 (2020).

39. Sollis, E. *et al.* The NHGRI-EBI GWAS Catalog: knowledgebase and deposition

resource. *Nucleic Acids Res.* **51**, D977–D985 (2023).

40. Choquet, H. *et al.* New and sex-specific migraine susceptibility loci identified from a

multiethnic genome-wide meta-analysis. *Commun. Biol.* **4**, 864 (2021).

41. Bornstein, M. H. *et al.* Infant Temperament: Stability by Age, Gender, Birth Order, Term

Status, and SES. *Child Dev.* **86**, 844–863 (2015).

42. Willcutt, E. G. The prevalence of DSM-IV attention-deficit/hyperactivity disorder: a

meta-analytic review. *Neurother. J. Am. Soc. Exp. Neurother.* **9**, 490–499 (2012).

43. Faraone, S. V. *et al.* The World Federation of ADHD International Consensus Statement:

208 Evidence-based conclusions about the disorder. *Neurosci. Biobehav. Rev.* **128**, 789–

818 (2021).

44. Rubin, K. H., Coplan, R. J. & Bowker, J. C. Social withdrawal in childhood. *Annu. Rev.*

Psychol. **60**, 141–171 (2009).

45. Lai, M.-C., Lombardo, M. V. & Baron-Cohen, S. Autism. *The Lancet* **383**, 896–910

(2014).

46. Kotov, R. *et al.* The Hierarchical Taxonomy of Psychopathology (HiTOP): A

dimensional alternative to traditional nosologies. *J. Abnorm. Psychol.* **126**, 454–477

(2017).

47. Krueger, R. F. & Markon, K. E. Reinterpreting comorbidity: A model-based approach to

understanding and classifying psychopathology. *Annu. Rev. Clin. Psychol.* **2**, 111–133

(2006).

48. Barlow, D. H., Ellard, K. K., Sauer-Zavala, S., Bullis, J. R. & Carl, J. R. The Origins of

Neuroticism. *Perspect. Psychol. Sci. J. Assoc. Psychol. Sci.* **9**, 481–496 (2014).

49. Shiner, R. L. & Caspi, A. Temperament and the development of personality traits,

adaptations, and narratives. in *Handbook of temperament* 497–516 (The Guilford Press,

New York, NY, US, 2012).

50. Beauchaine, T. P. & McNulty, T. Comorbidities and continuities as ontogenic processes:

toward a developmental spectrum model of externalizing psychopathology. *Dev.*

Psychopathol. **25**, 1505–1528 (2013).

51. Saudino, K. J. & Cherny, S. S. Sources of continuity and change in observed

temperament. in *Infancy to early childhood: Genetic and environmental influences on*

developmental change 89–110 (Oxford University Press, New York, NY, US, 2001).

52. Plomin, R., Loehlin, J. C. & DeFries, J. C. Genetic and environmental components of

'environmental' influences. *Dev. Psychol.* **21**, 391–402 (1985).

53. Tucker-Drob, E. M., Briley, D. A. & Harden, K. P. Genetic and environmental influences

on cognition across development and context. *Curr. Dir. Psychol. Sci.* **22**, 349–355

(2013).

54. Allegrini, A. G. *et al.* Genomic prediction of cognitive traits in childhood and

adolescence. *Mol. Psychiatry* **24**, 819–827 (2019).

55. Plomin, R. & von Stumm, S. The new genetics of intelligence. *Nat. Rev. Genet.* **19**, 148–

159 (2018).

56. Choi, S. W., Mak, T. S.-H. & O'Reilly, P. F. Tutorial: a guide to performing polygenic

risk score analyses. *Nat. Protoc.* **15**, 2759–2772 (2020).

57. Lewis, C. M. & Vassos, E. Polygenic risk scores: from research tools to clinical

instruments. *Genome Med.* **12**, 44 (2020).

58. Polanczyk, G., de Lima, M. S., Horta, B. L., Biederman, J. & Rohde, L. A. The

worldwide prevalence of ADHD: a systematic review and metaregression analysis. *Am. J.*

Psychiatry **164**, 942–948 (2007).

59. McLean, C. P., Asnaani, A., Litz, B. T. & Hofmann, S. G. Gender differences in anxiety

disorders: prevalence, course of illness, comorbidity and burden of illness. *J. Psychiatr.*

Res. **45**, 1027–1035 (2011).

60. Kopal, J., Uddin, L. Q. & Bzdok, D. The end game: respecting major sources of

population diversity. *Nat. Methods* **20**, 1122–1128 (2023).

61. Vejrup, K., Magnus, P. & Magnus, M. Lost to follow-up in the Norwegian mother, father

and child cohort study. *Paediatr. Perinat. Epidemiol.* **36**, 300–309 (2022).

62. Lahey, B. B., Krueger, R. F., Rathouz, P. J., Waldman, I. D. & Zald, D. H. A hierarchical

causal taxonomy of psychopathology across the life span. *Psychol. Bull.* **143**, 142–186

(2017).

63. Magnus, P. *et al.* Cohort Profile Update: The Norwegian Mother and Child Cohort Study

(MoBa). *Int. J. Epidemiol.* **45**, 382–388 (2016).

64. Bakken, I. J., Ariansen, A. M. S., Knudsen, G. P., Johansen, K. I. & Vollset, S. E. The

Norwegian Patient Registry and the Norwegian Registry for Primary Health Care:

Research potential of two nationwide health-care registries. *Scand. J. Public Health* **48**,

49–55 (2020).

65. International statistical classification of diseases and related health problems.

https://iris.who.int/items/5858156c-f66b-43cc-be2c-37710e5cd2e6.

66. Birkenæs, V. *et al.* Youth psychotic experiences: psychometric evaluation and diagnostic

associations of the CAPE-16 in adolescents from the Norwegian Mother, Father and

Child Cohort. *J. Child Psychol. Psychiatry* **67**, 27–40 (2026).

67. Krishnan, A., Williams, L. J., McIntosh, A. R. & Abdi, H. Partial Least Squares (PLS)

methods for neuroimaging: a tutorial and review. *NeuroImage* **56**, 455–475 (2011).

68. Cox, D. R. Regression Models and Life-Tables. *J. R. Stat. Soc. Ser. B Methodol.* **34**, 187–

220 (1972).

69. Demontis, D. *et al.* Genome-wide analyses of ADHD identify 27 risk loci, refine the

genetic architecture and implicate several cognitive domains. *Nat. Genet.* **55**, 198–208

(2023).

70. Watanabe, K., Taskesen, E., van Bochoven, A. & Posthuma, D. Functional mapping and

annotation of genetic associations with FUMA. *Nat. Commun.* **8**, 1826 (2017).

a EAS temperament

b

c

d

e

f