Noisy periodicity in tropical respiratory disease dynamics

Fuhan Yang[1*], Ephraim Hanks[1], Jessica Conway[1], Ottar Bjornstad[1], Nguyen Thi Le Thanh[2],

Maciej F Boni[1,3], Joseph L Servadio[1,3]

1 Center for Infectious Disease Dynamics, Department of Biology, Pennsylvania State

University, University Park, PA, 16802

2 Oxford University Clinical Research Unit, Wellcome Trust Major Overseas Programme, Ho

Chi Minh City, Vietnam

3 Institute for Genomics and Evolutionary Medicine, Department of Biology, Temple University,

Philadelphia, PA, 19122

*Correspondence: fuhan_yang@outlook.com

Abstract

Infectious disease surveillance systems in tropical countries show that respiratory disease incidence generally manifests as year-round activity with weak fluctuations and irregular seasonality. Previously, using a ten-year time series of influenza-like illness (ILI) collected from outpatient clinics in Ho Chi Minh City (HCMC), Vietnam, we found a combination of nonannual and annual signals driving these dynamics, but with unknown mechanisms. In this study, we use seven stochastic dynamical models incorporating humidity, temperature, and school term to investigate plausible mechanisms behind these annual and nonannual incidence trends. We use iterated filtering to fit the models and evaluate the models by comparing how well they replicate the combination of annual and nonannual signals. We find that a model including specific humidity, temperature, and school term best fits our observed data from HCMC and partially reproduces the irregular seasonality. The estimated effects from specific humidity and temperature on transmission are nonlinearly negative but weak. School dismissal is associated with decreased transmission, but also with low magnitude. Under these weak external drivers, we hypothesize that stochasticity makes a strong sub-annual cycle more likely to be observed in ILI disease dynamics. Our study shows a possible mechanism for respiratory disease dynamics in the tropics. When the external drivers are weak, the seasonality of respiratory disease dynamics is prone to the influence of stochasticity.

Author Summary

Although the mechanisms driving seasonality of respiratory disease dynamics have been well-studied in temperate regions, they are unknown in the tropics. In this study, we used a 10-year influenza-like-illness (ILI) daily-reporting data set collected from outpatient clinics in Ho Chi Minh City (HCMC) in Vietnam to investigate the mechanisms associated with annual and nonannual (~215 days) periodic patterns in the data. By comparing seven mechanistic models against the data, we showed that the mechanism that best explains respiratory disease dynamics in HCMC is a stochastic susceptible-infected-recovered-susceptible (SIRS) model weakly driven by external drivers including specific humidity, temperature, and school term. The nonannual cycle's duration is consistent with the inferred duration of immunity of the model. By showing the nonannual cycle as strong as in the data is only observed in stochastic model, we showed that the observed respiratory disease dynamics in HCMC is under the influence of stochasticity when external drivers are weak.

1. Introduction

Seasonality of respiratory disease transmission manifests as regular winter epidemics in temperate parts of the world. The factors associated with this wintertime seasonality have been studied through laboratory experiments [1,2] and modeling studies [3–5]. The common understanding is that climate factors such as humidity [1] and human behaviors such as indoor crowding, movement [6] and school attendance [7–9] are the main drivers of seasonality in temperate regions [10,11]. Significant changes in climate and human activities forced by winter lead to epidemic onset through the increase of viral survival rate and human contact rates when gathering indoors during winter. Globally, the dynamics of respiratory virus transmission are less predictable outside of temperate regions. One obstacle is that there are fewer surveillance systems in subtropical and tropical regions; the lack of data makes it challenging to draw conclusions about seasonality. Another challenge is that seasonality patterns, which are very clear in temperate regions, differ across subtropical and tropical regions. Tropical regions tend to have more irregular seasonality than subtropical regions [12]. Studies have shown that respiratory disease dynamics in subtropical regions present as an annual pattern, a biannual pattern or both [13–15]. As latitude moves close to the equator, dynamics tend to show year-round activity and exhibit weaker fluctuation [14,16,17]. Most studies on respiratory disease dynamics in the tropics are descriptive, focusing on associations between seasonality and environmental drivers [2,13,18–22]. Few studies have examined the mechanisms behind these patterns.

Mechanistically, seasonality of infectious diseases is mainly influenced by four factors: the intrinsic cycle caused host-pathogen interaction [23–25] (like immune waning or pathogen

evolution), host behavior, environmental drivers, and stochasticity [26,27]. Seasonality could be

influenced by interactions of two or more of these factors, especially, the relative dominance of

the interacting factors [28,29]. In respiratory disease dynamics in temperate regions, the

regularly annual and substantial change in climate and host behavior may be the dominant

factors causing the dynamics to behave annually, overshadowing the influence from host-

pathogen interaction or stochasticity [30]. Our previous study found that respiratory disease

dynamics in the tropics showed weak fluctuation, with irregular seasonality and both annual and

nonannual signals [31]. With no winter in the tropics and mild changes in weather and human

behavior, the causes of these fluctuations are not obvious.

We are interested in identifying the mechanisms that cause the unique pattern of respiratory

disease dynamics seen in the tropics. In this study, we use seven stochastic dynamical models to

investigate the drivers of these dynamics using a 10-year daily respiratory disease reporting

dataset from Ho Chi Minh City (HCMC), Vietnam. We incorporate humidity, temperature, and a

term-time forcing factor representing school term; all models are based on stochastic versions of

a susceptible-infected-recovered-susceptible (SIRS) model. We compare their goodness-of-fit

using trajectory matching and their abilities to replicate the cycles observed in the data through

power spectrum comparisons. The results of this study show the contributions of environmental

factors, school term, and stochasticity to the dynamics, which allow us to identify which of the

considered eternal factors can best explain respiratory disease transmission patterns in the

tropics.

2. Results

We analyze a ten-year standardized daily influenza-like illness (ILI) time series from outpatient

community surveillance in HCMC, Vietnam, that has been validated in previous studies [16,31–

33] as the main data source in this study. Previous validations on these data included

examinations of trends to justify de-trending, evaluation of individual-clinic reporting

consistency [33], and evaluation of bias in clinical diagnosis of ILI [31]. Here we scale the time

series to represent observed case counts, whereas the previous study presented the time series as

a unitless trend in relative ILI incidence. The ILI case time series showed noisy and low-

amplitude fluctuations between 2010 and 2019 (Fig 1A). Spectral analysis reveals two periods of

comparable importance: signals with cycles of 215 days and 365 days (Fig 1B). This is

consistent with our previous results [31], showing that there is a combination of annual and

nonannual cycles in the weak fluctuations in ILI dynamics in HCMC.

2.1 Model comparisons through trajectory matching

We describe the respiratory disease dynamics as a model of susceptibles, infecteds and

recovereds allowing reinfection (SIRS) using multi-stage infection and recovery classes. We

select two I classes (i.e. the infectious period is gamma distributed with shape parameter 2) and

three R classes (i.e. duration of immune protection is gamma distributed with shape parameter 3)

from a range of possible number of I (1 to 2) and R (3 to 5) classes using partially-observed

Markov process framework (SI_2R_3S, Fig. S1, details in S2 Text) [34]. We compare the goodness-

of-fit of seven models to the data. Respectively, they are: (1) basic SI_2R_3S model without

external factors (SIRS); (2) SI_2R_3S model including absolute humidity covariate (SIRS with

AH); (3) SI_2R_3S model including smoothly-decreasing school term (SIRS with school term); (4)

SI_2R_3S model including absolute humidity and school term; (5) SI_2R_3S model including

quadratic ("U-shaped") effect of specific humidity on transmission, modulated by temperature

(SIRS with U-shaped SH and temperature); (6) SI_2R_3S model including linear relationship with

specific humidity, modulated by temperature (SIRS with linear SH and temperature); (7) SI_2R_3S

model including quadratic ("U-shaped") effect of specific humidity, modulated by temperature,

and effects of school term (SIRS with U-shaped SH temperature and school term).

The seven models are fitted to the entire ILI time series (January 1, 2010, to December 31, 2019)

using iterated filtering [35]. First, we evaluate goodness-of-fit using mean squared error (MSE)

between the median from 5000 simulations and the data. SIRS with U-shaped SH, temperature

and school term performs the best, indicating that this model best describes the observed data

from HCMC. The second-best model is SIRS with U-shaped SH and temperature. The fact that

both models share the same feature of the U-shaped effect of specific humidity indicates a level

of confidence for this relationship. The average across the 5000 MSE between the data and each

simulation is also evaluated to address the variability of individual simulations. With this

criterion, the best model is SIRS with U-shaped SH temperature, and the second best adds school

term. This means that both models are best able to reproduce the observed data, both across

individual simulations and when comparing an aggregate among simulations. In fact, when

comparing the median and simulation interval with the data, these two models are the only two

models that exhibit small and noisy fluctuations in the data while other models generally produce

a visually flat median line (Fig 2). These results suggest the U-shaped effect of specific humidity

as a potentially important mechanism of respiratory disease transmission in the tropics.

Based on the two best supported models, we find that the external drivers have a weak effect on

transmission. For SIRS with U-shaped SH and temperature, the parameters governing the U-

shaped relationship with SH are estimated to be small in magnitude ($a = 1.24 \times 10^{-6}$, 95% CI:

$[1.02 \times 10^{-6}, 1.29 \times 10^{-6}]$, $b = -1.16 \times 10^{-5}$, 95% CI: $[-1.21 \times 10^{-5}, -1.06 \times 10^{-5}]$, Table 1)

indicating changes in the transmission parameter β many orders of smaller than the estimated

value of β_0. Similarly, the modulation effect from temperature is small ($Texp = 0.088$, 95% CI:

[0.087,0.090]). These estimations overall show very narrow change in R_0 (ranging between 1.628

and 1.652) given values of SH and temperature in the entire range observed in HCMC (SH is

(9.43 g/kg – 20.65 g/kg) and temperature is (23.11 °C– 33.31°C)). Similarly, for SIRS with U-

shaped SH temperature and school term, the estimated parameters controlling the effect of SH (a

$= 9.36 \times 10^{-5}$, 95% CI: $[9.12 \times 10^{-5}, 9.48 \times 10^{-5}]$, $b = -4.50 \times 10^{-3}$, 95% CI: $[-4.58 \times 10^{-3}, -4.32 \times$

$10^{-3}]$), temperature ($Texp = 0.138$, 95% CI: [0.136,0.139]) and school term ($a\_school = 4.54 \times$

10^{-3}, 95% CI: $[4.47 \times 10^{-3}, 4.68 \times 10^{-3}]$) on R_0 are low in magnitude, showing estimated values of

R_0 ranging between 1.550 and 1.602. These results suggest a system near its endemic

equilibrium and weakly affected by external drivers.

Model/Parameters	SIRS	SIRS with AH	SIRS with school term	SIRS with AH and school term	SIRS with U-shaped SH and temperature	SIRS with U-shaped SH, temperature, and school term	SIRS with linear SH and temperature
β_0	0.227 (0.223,0.229)	0.213 (0.212,0.215)	0.204 (0.196,0.204)	0.255 (0.251, 0.255)	0.160 (0.160, 0.161)	0.168 (0.167,0.169)	0.160 (0.1600,0.1603)
a_AH		-0.55 (-0.57,0.53)		0.950 (0.941,0.951)			
w		-99.8 (-100, -99.4)		-23.78 (-24.07，-22.71)			
a_school			0.90×10^{-4} $(0.87 \times 10^{-4}, 0.93 \times 10^{-4})$	1.5×10^{-3} $(1.1 \times 10^{-3}, 1.8 \times 10^{-3})$		4.54×10^{-3} $(4.47 \times 10^{-3}, 4.68 \times 10^{-3})$	
σ			9.69 (9.42,9.89)	4.08 (3.37,4.12)		19.98 (19.47, 20.00)	
a					1.24×10^{-6} $(1.02 \times 10^{-6}, 1.29 \times 10^{-6})$	9.36×10^{-5} $(9.12 \times 10^{-5}, 9.48 \times 10^{-5})$	
b					-1.16×10^{-5} $(-1.21 \times 10^{-5}, -1.06 \times 10^{-5})$	-4.50×10^{-3} $(-4.58 \times 10^{-3}, -4.32 \times 10^{-3})$	-5.48×10^{-4} $(-6.00 \times 10^{-4}, -5.27 \times 10^{-4})$
c					0.752 (0.746, 0.760)	0.70 (0.68,0.71)	0.50 (0.50,0.503)
Tc					29.27 (29.19, 29.34)	29.59 (29.32, 29.74)	29.99 (29.81,30.00)
$Texp$					0.088 (0.087,0.090)	0.138 (0.136,0.139)	0.0028 (0.0026,0.003)
π	232.43 (231.89, 233.67)	230.42 (227.82, 231.54)	184.56 (183.12, 185.75)	325.94 (325.55,326.63)	216.89 (215.95, 217.79)	215.78 (215.28, 216.39)	149.77 (148.16, 151.40)
ρ	5.41×10^{-3} $(5.37 \times 10^{-3}, 5.80 \times 10^{-3})$	6.50×10^{-3} $(6.19 \times 10^{-3}, 6.70 \times 10^{-3})$	6.21×10^{-3} $(5.82 \times 10^{-3}, 6.22 \times 10^{-3})$	5.80×10^{-3} $(5.73 \times 10^{-3}, 5.82 \times 10^{-3})$	3.11×10^{-3} $(3.09 \times 10^{-5}, 3.15 \times 10^{-5})$	3.26×10^{-3} $(3.25 \times 10^{-3}, 3.27 \times 10^{-3})$	2.88×10^{-3} $(2.71 \times 10^{-3}, 3.12 \times 10^{-3})$
R_{0mean}	1.256 (1.232,1.266)	1.180 (1.170,1.190)	1.124 (1.079,1.125)	1.404 (1.382,1.406)	1.640 (1.632,1.647)	1.575 (1.560,1.590)	1.373 (1.371，1.378)

R_{0min}		1.180 (1.170,1.190)	1.123 (1.079,1.125)	1.396 (1.376,1.397)	1.628 (1.620,1.634)	1.550 (1.534,1.565)	1.371 (1.370, 1.377)
R_{0max}		1.180 (1.170,1.190)	1.124 (1.079,1.125)	1.404 (1.382,1.406)	1.652 (1.644,1.659)	1.602 (1.588,1.616)	1.378 (1.376, 1.382)
MSE from the median of simulations (rank)	63.49 (5)	63.38 (4)	68.00 (7)	63.11 (3)	61.81 (2)	60.95 (1)	64.20 (6)
Average MSE from each simulation (rank)	79.18 (6)	79.04 (5)	75.81 (4)	111.40 (7)	73.31 (1)	73.61 (2)	74.47 (3)

Table 1 Maximum-likelihood estimates (MLEs) of parameters from seven models, estimated R_0, MSE

between the median of 5000 simulations and the data, and the average of 5000 MSE between the data and

each simulation. The parentheses show 95% confidence intervals for each MLE, and model rank for two

MSE metrics.

2.2 Model comparison based on power spectrum

To examine a potential cause of the observed annual and nonannual cycle, we compare the power

spectrum from 5000 simulations of the fitted models with the power spectrum evaluated from the

data. The spectral densities for cycle lengths between 100 and 450 days are calculated for each

simulation, and the discrepancies between the data are measured as average weighted MSE (see

Methods 5.4).

First, we find that all models with external drivers outperformed the null model (SIRS) in

reproducing cycles similar to the data (Table 2). This indicates that the observed cycles are not

caused solely by the internal transmission clockwork, but also by external drivers contributing to

the dynamics. Second, two SIRS models with U-shaped SH and temperature models (with and

without school term) show improved ability in replicating the cyclic pattern of the data compared

to other models (Table 2). This suggests that the U-shaped relationship with specific humidity,

modulated by temperature, likely contributes to the periodicity of the respiratory disease

dynamics in HCMC.

Model	SIRS	SIRS with AH	SIRS with school term	SIRS with AH and school term	SIRS with U-shaped SH and temperature	SIRS with U-shaped SH, temperature and school term	SIRS with linear SH and temperature
Average weighted MSE across all the simulations (rank)	11.67 (7)	10.61 (4)	11.59 (6)	9.24 (3)	6.86 (2)	5.14 (1)	11.34 (5)

Table 2 Average weighted MSE in the power spectrum between 100 days and 450 days across 5000

simulations between each model and the data are shown, with the rank in the parentheses.

We also visualize the power spectrum of the data comparing to the simulations from the seven

fitted models (Fig 3). We find that the null model (SIRS) only captures the near-annual cycle

(around 330 days). The SIRS with AH, school term, or both, also only captures the near-annual

or annual cycles. Although SIRS with linear SH and temperature simulates a bimodal pattern in

the cycles same as the data, most of the simulations fail to reproduce the periodic components at

nonannual or annual cycles as strongly as observed in the data, indicating that the relationship

between specific humidity and respiratory disease dynamics in HCMC may not be linear, which

aligns with the results from trajectory matching. Only the models incorporating the U-shaped

relationship between SH and transmission modified by temperature (with or without school term)

are able to simulate the nonannual cycle at 215 days along with an annual cycle both with

comparable amplitudes. This is consistent with the distributions of cyclic summary statistics

simulated by the fitted models (S3 Text, Fig S2). Adding school term helps to better reproduce

the periodic pattern of the data (Table 2), especially at the annual cycle period (Fig 3). This

suggests that school term contributes to amplify the annual cycle, and the nonannual cycle may

be the consequences of interaction between climatic factors and respiratory disease dynamics.

We also validate the fitting between the best 10 simulations selected by MSE from trajectory

matching and the data. We find that these simulations capture the weak fluctuations well (Fig

S3). Combining the results from trajectory matching, we select SIRS with U-shaped SH,

temperature and school term as the best model capturing the fluctuations as well as reproducing

the cycles of respiratory disease dynamics in HCMC.

2.3 Characterization of respiratory disease dynamics in HCMC

We visualize the relationship between specific humidity and estimated R_0 from SIRS with U-

shaped SH, temperature and school term (Fig 4A). We find a weak negative relationship between

specific humidity and R_0. Note that the parameter governing the U-shape of specific humidity is

estimated to be very small ($a = 9.36 \times 10^{-5}$, 95% CI: [9.12×10^{-5}, 9.48×10^{-5}], Table 1), leading

to a relationship approximated to be negative and nearly linear ($b = -4.50 \times 10^{-3}$, 95% CI: [-4.58

$\times 10^{-3}$, -4.32×10^{-3}]) as shown in Fig 4A, which means R_0 will slightly decrease as the SH

increases. Similarly, we observe a nonlinear negative relationship between R_0 and temperature

(Fig 4B). As the threshold (Tc) determines whether the effect of temperature is amplifying or

inhibiting the effect of SH (see Methods 5.2.e), the estimated high Tc (29.59 °C, 95% CI:

[29.32 °C,29.74 °C]) suggests temperature will mostly inhibit the effect of SH on R_0. We can

observe that the higher temperature leads to even lower R_0 (Fig 4A). In addition, we observe R_0

during school recess is lower than the time of school in session (Fig 4B), while it overlaps with

the time when SH is high. However, the overall effect of SH, temperature and school term on R_0

is weak, leading to a small range of R_0 over time (1.550 - 1.602).

2.4 The effects of stochasticity

We compare the simulated power spectra between SIRS with U-shaped SH, temperature and

school term and its deterministic skeleton to evaluate the effect of stochasticity on the cycles of

the ILI data in HCMC. First, through the average weighted MSE of the power spectra, we find

that the power spectrum reproduced by the stochastic model is a slightly better fit than the power

spectrum produced by the deterministic version (Table 3), suggesting stochasticity may affect the

cycles of respiratory disease dynamics in HCMC. Comparing the power spectrum between data

and both models (Fig 5), the annual cycle is well-reproduced by both models, showing the

influence from these external annual drivers. However, only the stochastic version is possible to

reproduce the nonannual cycle at an amplitude equivalent to the data, or even higher, which the

deterministic skeleton failed to do. This indicates that the observed strong signal of the

nonannual cycle from the data may be produced under the influence of stochasticity. It is

possible that the ILI data in HCMC represents a real-world example that show stochasticity

contributes to the seasonality, along with the interaction between external annual drivers and the

intrinsic cycle of the disease dynamics.

Model	Deterministic SIRS with U-shaped SH temperature and school term	Stochastic SIRS with U-shaped SH temperature and school term
Average weighted MSE across all the simulations	5.73	5.14

242 Table 3 The average MSE in power spectrum from 5000 weighted MSE between the data and each

243 simulation, from stochastic SIRS U-shaped SH temperature and school term and its deterministic

244 skeleton.

3. Discussion

We use a 10-year daily ILI time series between 2010 and 2019 to explore the mechanistic forces

that drive the unique periodic pattern in respiratory disease dynamics in the tropics, using

HCMC, Vietnam, as an example. We compare the performance of seven models on their ability

to replicate trajectories and periodicities in the data. Overall, we find that an SIRS model that is

nonlinearly influenced by specific humidity, temperature and school term performs best among

these seven models. Through this model, we find that specific humidity, temperature and school

term drive the transmission negatively but weakly (Fig 4, estimated R_0 ranges from 1.550 to

1.602 given the entire range of SH, temperature, and school recess in the observed 10-year

period). We also find that the observed periodicities are the consequences of interactions between

annual external drivers, an intrinsic system cycle (~215 days), and stochasticity. Specifically, the

annual cycle is likely driven by the annual external factors (climate and school term), while the

nonannual cycle may be driven by stochasticity. Our study reveals a unique behavior of disease

dynamics when external drivers are weak.

This study is one of the first exploring the mechanistic causes of the irregular seasonality of

respiratory disease in the tropics [19–21, 39, 40]. There are challenges in studying respiratory

disease dynamics in the tropics for two main reasons. First, the absence of high-resolution data

over an extended time period makes it difficult to investigate the seasonality of respiratory

disease dynamics in the tropics. The ILI surveillance system analyzed here produced more than

66,000 data points from community clinicians over ten years, making it possible to conduct a

study of cyclic incidence. Second, it is challenging to provide mechanistic explanations of

disease dynamics when dynamical signals are weak. Many descriptive studies have conducted

and characterized ILI activity in the tropics as 'all-year-round' [14,20,41,42]. Our study is one of

the first studies that investigates the lack of regular seasonality of respiratory diseases dynamics

in the tropics from a mechanistic perspective.

Fitting noisy incidence data with mechanistic models is challenging. In this case, stochastic

models are useful because they can account for noise, which provides flexibility to capture short-

term fluctuations that deterministic models cannot catch. Our approach in using stochastic

models and stochastic optimization methods allow us to detect a weak dynamical signal behind

stochasticity, supporting the utility of stochastic models when fitting noisy data series.

3.1 The environmental drivers of respiratory viral infection in the tropics

Our analysis shows that a model with a U-shaped relationship between specific humidity

modulated by temperature and school term explains the data the best, consistent with the

observations in subtropical regions [37,39]. However, we have not observed the typical U-shape

as in previous studies [20,37,38,43], instead observing a monotonic relationship within the

observed SH values (the transmission-minimizing SH values were outside the observed range).

This may be caused by the estimated effect from specific humidity being weak ($a = 9.36 \times 10^{-5}$,

95% CI: [9.12×10^{-5}, 9.48×10^{-5}], $b = -4.50 \times 10^{-3}$, 95% CI: [-4.58×10^{-3}, -4.32×10^{-3}]). We

observe a nonlinear negative effect of temperature on transmission, in alignment with previous

observational studies [20] and laboratory findings [44], though this effect is small in magnitude

in our analysis ($Texp = 0.138$, 95% CI: [0.136,0.139]). School term-time forcing slightly

increases R_0 ($a\_school = 4.54 \times 10^{-3}$, 95% CI: [$4.47 \times 10^{-3}$, 4.68×10^{-3}]). Overall, our study

identifies effects of specific humidity, temperature, and school term on the transmission of

respiratory disease dynamics in the tropics, though the effect sizes are small.

3.2 Interplay between deterministic forcing and stochasticity

Distinct from respiratory disease dynamics in temperate regions that show a regular annual cycle,

the noisy and weak fluctuations with irregular cycles in tropical regions is likely caused by

different mechanisms. Our study found that even though the disease dynamics are weakly

influenced by the annual cycle of the external drivers, the intrinsic cycle can be observed (~ 215

days estimated), and stochasticity makes it more likely to see the strong signal of intrinsic cycle

(as in the data) (Fig 5). It is possible that the intrinsic cycle of the dynamics and stochasticity

dominates when external drivers are weak [28]. The estimated immune-waning duration (π) in

our analysis is estimated to be around 216 days for the two best-fit models, suggesting that

renewal of susceptibility on this time scale may have an influence on the sub-annual cycle seen

in the data. A brute force search for models producing this behavior shows that constructing this

effect in a model is non-trivial [45].This kind of the interplay has been widely observed in other

infectious diseases like whooping cough [27, 53], rubella [54], measles [53, 55], as well as in the

population dynamics in ecology such as flour beetles [56], coastal cod [57], and Dungeness crab

[58]. Our results also show that stochasticity could be a cause of the nonannual cycle in HCMC's

ILI activity, suggesting a complicated but remarkable tension in respiratory disease dynamics in

the tropics where irregular cyclic upswings would frequently be possible but not always

predictable. Perturbation analysis [26] is a natural future direction of this work to more precisely

quantify the effect of stochasticity on the cycles in ILI.

3.3 Limitations

A major limitation of our study is that our data set includes only symptomatic cases of ILI, which

omit asymptomatic infections and include an aggregation of multiple respiratory diseases.

However, we have observed consistently weak fluctuation in the ten years of the study period

without finding the pronounced regular peak (with a more than 5-fold difference) typically seen

in temperate regions [46]. Using our data from HCMC, we also found that ILI cases weakly

changed with specific humidity and temperature, suggesting that it is possible that the

transmission of the circulating pathogens has an overall weak relationship with environmental

drivers in the tropics. While it is also possible that virus interference may attenuate the

transmission of a single virus [47], we would require further virological data to understand the

transmission of respiratory pathogens in the tropics.

328   4. Conclusions

Our study proposed seven candidate models on the external determinants of the seasonality of

respiratory disease dynamics in the tropics. We explored and characterized the assumption that

were best able to explain the data through model-fitting as well as power spectrum comparison.

We found the dynamics to be weakly driven by specific humidity mediated by temperature as

well as by school term. We also found that while these external drivers are weak, the intrinsic

sub-annual cycle of respiratory disease dynamics in the tropics may be amplified by

stochasticity.

5. Methods

5.1 Data Acquisition

Between August, 2009 and December 31, 2019, the Oxford University Clinical Research Unit in Vietnam established a community-based mHealth surveillance system to monitor the transmission of respiratory viruses by recording the number of patients presenting with influenza-like-illness (ILI) symptoms in outpatient clinic settings across Ho Chi Minh City [31–33]. A total of 89 clinics participated in this study, and the percentage of ILI patients among the total patients was recorded daily during the period. We used data from 33 clinics for this study because more than 50% of their daily observations report non-zero ILI case counts. Data between January 1, 2010, and December 31, 2019, were used for analysis to remove potential effects of the 2009 H1N1 pandemic. We standardized and smoothed the daily percentage of patients presenting with ILI (details in S1 Text), then multiplied this time series by the 10-year mean of daily ILI cases (62.0 per day) to produce a daily count of ILI cases which we used in this study. The specific humidity and temperature data in HCMC from January 1, 2010 to December 31, 2019 were collected from NASA POWER project [48].

5.2 The Models

We use a discrete stochastic epidemic model to model the data, assuming the data is a partial

observation of a noisy epidemic process in HCMC. We assume that transmission follows

Susceptible – Infected – Recovered – Susceptible (SIRS) dynamics to allow the commonly

observed reinfection in respiratory diseases. We incorporate process stochasticity by defining the

outflow rate as a binomial draw from each compartment, assuming that the probability is an

Euler approximation of the outflow [49].

We include multiple stages in *I* and *R* to model a realistic memory-dependent process in infection

and recovery. To select the number of I and R stages, we use a particle filter to evaluate the

likelihood of the parameter combinations, including basic transmission rate β_0, reporting rate ρ,

and immune duration π, with each parameter sampled from an educated range (details in S2

Text, Table S1). Two I classes and three R classes are selected as they return the highest

likelihood (Fig.S1). The details of Euler-approximated transition rates along with the parameters

are listed in Table S2.

Our model can be described as a SI_2R_3S epidemic model, which is the model used throughout all

the analyses, and we simply abbreviate this as an SIRS model. Seven models are built based on

seven variations of the transmission rate β, describing seven assumptions about the transmission

of respiratory diseases in the tropics. These are listed below.

a.   Simple SIRS:

$$\beta = \beta_0$$ (Equation 1)

This model assumes constant transmission rate. β_0 will be estimated.

b.   SIRS with term-time forcing (SIRS with school term):

383
$$\beta = \beta_0 - a\_school \cdot e^{-\frac{(d-190)^2}{\sigma^2}}$$ (Equation 2)

This model assumes the transmission rate is only affected by school term. We assume the

transmission rate decreases during the summer vacation, which is from June 1 to August

15 every year. We choose an inverted bell curve (i.e, the shape of an inverted normal

distribution) to describe the smooth decrease of transmission during school recess, with

the lowest transmission at day 190 of the calendar year, which is July 10, the midpoint of

summer vacation. d represents the day of the calendar year. In this model, we estimate β_0

(baseline transmission, positive), a_school (peak magnitude of decrease, nonnegative),

and σ (steepness of bell curve, positive).

c.   SIRS with exponential relationship with absolute humidity (SIRS with AH):

394
$$\beta = \beta_0 + a\_AH \cdot e^{w*AH}$$ (Equation 3)

This model assumes the transmission rate is only affected by absolute humidity (AH).

The effect of AH is transformed through an exponential function and then added to the

baseline transmission rate β_0 [1,4]. In this model, a_AH is an offset term describing the

change in the transmission rate when absolute humidity is 0, and w is the rate of change

of viral viability due to absolute humidity. All of these parameters' values are estimated.

In temperate regions, it is observed that a_AH is negative, indicating transmission rate

decreases monotonically with increasing absolute humidity [3,50]. However, it has not

been clear if such relationship exists in the tropics, or if there exists a certain threshold of

absolute humidity when switching the relation from negative to positive. Given these

reasons, the signs of a_AH and w are not fixed in model fitting.

d. SIRS with exponential relationship with absolute humidity and term-time forcing (SIRS with

AH and school term):

408
$$\beta = \beta_0 - a_{school} \cdot e^{-\frac{(d-190)^2}{\sigma^2}} + a_{AH} \cdot e^{w*AH} \qquad \text{(Equation 4)}$$

409   This model assumes the transmission rate is under the composite additive effect of both

410   school term and absolute humidity. The assumptions about signs of parameters remain the

411   same as in the individual-effect models.

412   e. SIRS with U-shaped relationship with specific humidity modulated by temperature (SIRS

413   with U-shaped SH and temperature):

414
$$\beta = \beta_0 + (a \cdot SH^2 + b \cdot SH + c)\left(\frac{Tc}{temperature}\right)^{Texp} /D \qquad \text{(Equation 5)}$$

This model assumes the reproduction number R_0 has a non-monotonic relationship with

absolute humidity modulated by temperature. It has been observed that the transmission

of influenza in subtropical and tropical areas [37,39] possesses this relationship , thus it is

used here to test if the transmission in HCMC will follow the same pattern. Specific

humidity, a measure of absolute humidity, is applied here. This model assumes the effect

of SH on transmission rate is largest when SH is either very low or very high; a, b, c are

the coefficients controlling the effect and will be estimated. The effect is moderated by

temperature, depending on a threshold temperature Tc. When temperature is below Tc,

the effect of SH on transmission increases and the transmission rate further increases.

When temperature is above Tc, the effect is decreased, and the transmission is inhibited

at higher temperatures. The moderating strength of temperature is controlled by $Texp$.

Here we estimate the effect of specific humidity and temperature on R_0 on the basis of the

baseline transmission rate. We convert the effect to be on transmission rate by dividing

the infectious period D. The parameters to be estimated are a, b, c, Tc, and $Texp$. a, b, c

are restrained to be positive, negative, and positive, respectively to ensure a minimum

value of the parabolic curve and only positive-valued contributions to the transmission

parameter. Tc and $Texp$ are constrained to be positive to be valid in describing

temperature and its moderating strength.

f. SIRS with U-shaped relationship with specific humidity modulated by temperature and

school term (SIRS with U-shaped SH, temperature and school term):

$$\beta = \beta_0 + (a \cdot SH^2 + b \cdot SH + c)\left(\frac{Tc}{temperature}\right)^{Texp}/D - a\_school \cdot e^{-\frac{(d-190)^2}{\sigma^2}} \qquad \text{(Equation 6)}$$

The model assumes additive effect from U-shaped relationship of specific humidity

modulated by temperature and school term.

g. SIRS with linear relationship with specific humidity, modulated by temperature (SIRS with

linear SH and temperature):

$$\beta = \beta_0 + (b \cdot SH + c)\left(\frac{Tc}{temperature}\right)^{Texp}/D \qquad \text{(Equation 7)}$$

The model assumes a linear relationship between specific humidity and transmission,

modulated by temperature. This model is a variant of model e. The parameters to be

estimated are *b, c, Tc, Texp. b* and *c* are not constrained. Tc and $Texp$ are constrained to

be positive to be valid in describing temperature and the moderating strength.

Reporting rate ρ and immune duration π of each model are estimated altogether with the

parameters from each model. R_0 is calculated based on: $R_0 = \beta \times D$, where D represents the

duration of the infectious period, which is defined by:

$$D = \frac{\Delta t}{(1 - e^{-m}) + (1 - e^{-\gamma})}$$ (Equation 8)

accounting for the Euler-approximated recovery rate $(1 - e^{-\gamma})$ and natural death rate

$(1 - e^{-m})$, as well as time discretization, with $\Delta t = 1$ in the daily discrete time series.

The R_0 from each model will be calculated with the estimated β in this way.

The daily incidence derived from the fitted model is considered as the latent process. We assume

the observed data is a Poisson process of the latent process multiplied by the reporting rate ρ.

The initial ranges of all the parameters to estimate are in Table S1.

5.3 Inference framework

Our model-fitting process uses iterated filtering to maximize a Poisson likelihood function, and

we use a particle filter (Sequential Monte Carlo) to evaluate the likelihood. Because of the

stochastic nature of the particle filter, likelihood was evaluated 10 times to balance accuracy and

computational efficiency given each set of parameters; the standard error of the likelihood is also

calculated. We use iterated filtering (IF) algorithm [51] for parameter inference and Monte Carlo

likelihood profiling to estimate the 95% confidence interval of the MLEs [52].

Based on preliminary investigations of the likelihood space, we design an inference framework

of IF to efficiently search for MLE parameters. We fit the data with four rounds of iterated

filtering, conducted sequentially. We first sample 160 sets of parameters from a multivariate

uniform distribution for each parameter given educated ranges (Table S1). These sampled

parameter sets then undergo iterated filtering four times with different tuning parameters (Table

S3). Within the four IF runs, estimates with the 5% highest likelihood are selected as the initial

parameters for the next run. The first run quickly explores the entire parameter space and returns

the space with a higher likelihood. The subsequent runs conduct more thorough searches as the

parameter space is more refined. In this way, the estimates would converge to the global MLE

(or close) more efficiently. The likelihood broadly increases along four runs, validating the

framework (Fig S4).

The likelihood of the returned parameters from each run is evaluated using a particle filter. The

MLE was selected as the run with the highest likelihood and a standard error is smaller than 5.

All the analyses used the R package pomp [35] and were implemented in R 4.3.0.

5.4 The evaluation of discrepancies in power spectrum

The discrepancies in the power spectrum between the model and the data are measured through

average weighted MSE. Firstly, 5000 simulations are generated from each fitted model, and the

power spectrum between cycle lengths between 150 and 450 days is calculated for each

simulated incidence trajectory. Then, for one simulation, the error for each cycle between the

simulation and the data is calculated using MSE. To emphasize how well the model captures the

nonannual (215 days) and annual cycles (365 days) in the data, the MSE on these cycles is

multiplied by 10 to produce a weighted MSE. Next, the MSE across each cycle is summed to

represent the total error in the power spectrum between the data and the simulation. This

calculation is repeated for 5000 simulations for each model, and finally the MSE is averaged

across 5000 simulations to represent the average discrepancies in the cyclic pattern between

models and data, which we call average weighted MSE.

Data Availability

497   All the data and code are available at https://github.com/Fuhan-

498   Yang/hcmc_stochastic_ili_modeling.

References

1. Shaman J, Kohn M. Absolute humidity modulates influenza survival, transmission, and seasonality. Proc Natl Acad Sci USA. 2009;106: 3243–3248. doi:10.1073/pnas.0806852106

2. Irwin CK, Yoon KJ, Wang C, Hoff SJ, Zimmerman JJ, Denagamage T, et al. Using the systematic review methodology to evaluate factors that influence the persistence of influenza virus in environmental matrices. Appl Environ Microbiol. 2011;77: 1049–1060. doi:10.1128/aem.02733-09

3. Shaman J, Pitzer VE, Viboud C, Grenfell BT, Lipsitch M. Absolute Humidity and the Seasonal Onset of Influenza in the Continental United States. Ferguson NM, editor. PLoS Biol. 2010;8: e1000316. doi:10.1371/journal.pbio.1000316

4. Dalziel BD, Kissler S, Gog JR, Viboud C, Bjørnstad ON, Metcalf CJE, et al. Urbanization and humidity shape the intensity of influenza epidemics in U.S. cities. Science. 2018;362: 75–79. doi:10.1126/science.aat6030

5. Tamerius J, Nelson MI, Zhou SZ, Viboud C, Miller MA, Alonso WJ. Global Influenza Seasonality: Reconciling Patterns across Temperate and Tropical Regions. Environmental Health Perspectives. 2011;119: 439–445. doi:10.1289/ehp.1002383

6. Massaro E, Kondor D, Ratti C. Assessing the interplay between human mobility and mosquito borne diseases in urban environments. Sci Rep. 2019;9: 16911. doi:10.1038/s41598-019-53127-z

7. Kucharski AJ, Conlan AJK, Eames KTD. School's Out: Seasonal Variation in the Movement Patterns of School Children. PLOS ONE. 2015;10: e0128070. doi:10.1371/journal.pone.0128070

8. Lessler J, Grabowski MK, Grantz KH, Badillo-Goicoechea E, Metcalf CJE, Lupton-Smith C, et al. Household COVID-19 risk and in-person schooling. Science. 2021;372: 1092–1097. doi:10.1126/science.abh2939

523   9.   Vlachos J, Hertegård E, B. Svaleryd H. The effects of school closures on SARS-CoV-2 among

524   parents and teachers. Proceedings of the National Academy of Sciences. 2021;118: e2020834118.

525   doi:10.1073/pnas.2020834118

526   10.   Leung NHL. Transmissibility and transmission of respiratory viruses. Nat Rev Microbiol. 2021;19:

527   528–545. doi:10.1038/s41579-021-00535-6

528   11.   Moriyama M, Hugentobler WJ, Iwasaki A. Seasonality of Respiratory Viral Infections. Annu Rev

529   Virol. 2020;7: 83–101. doi:10.1146/annurev-virology-012420-022445

530   12.   Heraud J-M, Razanajatovo NH, Viboud C. Global circulation of respiratory viruses: from local

531   observations to global predictions. The Lancet Global Health. 2019;7: e982–e983.

532   doi:10.1016/S2214-109X(19)30277-3

533   13.   Berry I, Rahman M, Flora MS, Shirin T, Alamgir ASM, Khan MH, et al. Seasonality of influenza

534   and coseasonality with avian influenza in Bangladesh, 2010–19: a retrospective, time-series

535   analysis. The Lancet Global Health. 2022;10: e1150–e1158. doi:10.1016/S2214-109X(22)00212-1

536   14.   Saha S, Chadha M, Al Mamun A, Rahman M, Sturm-Ramirez K, Chittaganpitch M, et al. Influenza

537   seasonality and vaccination timing in tropical and subtropical areas of southern and south-eastern

538   Asia. Bull World Health Organ. 2014;92: 318–330. doi:10.2471/BLT.13.124412

539   15.   Yang L, Wong CM, Lau EHY, Chan KP, Ou CQ, Peiris JSM. Synchrony of Clinical and Laboratory

540   Surveillance for Influenza in Hong Kong. Myer L, editor. PLoS ONE. 2008;3: e1399.

541   doi:10.1371/journal.pone.0001399

542   16.   Thai PQ, Choisy M, Duong TN, Thiem VD, Yen NT, Hien NT, et al. Seasonality of absolute

543   humidity explains seasonality of influenza-like illness in Vietnam. Epidemics. 2015;13: 65–73.

544   doi:10.1016/j.epidem.2015.06.002

17. Chadha M, Hirve S, Bancej C, Barr I, Baumeister E, Caetano B, et al. Human respiratory syncytial

virus and influenza seasonality patterns—Early findings from the WHO global respiratory syncytial

virus surveillance. Influenza and Other Respiratory Viruses. 2020;14: 638–646.

doi:10.1111/irv.12726

18. Mahamat A, Dussart P, Bouix A, Carvalho L, Eltges F, Matheus S, et al. Climatic drivers of

seasonal influenza epidemics in French Guiana, 2006–2010. Journal of Infection. 2013;67: 141–

147. doi:10.1016/j.jinf.2013.03.018

19. Ali ST, Cowling BJ, Wong JY, Chen D, Shan S, Lau EHY, et al. Influenza seasonality and its

environmental driving factors in mainland China and Hong Kong. Science of The Total

Environment. 2022;818: 151724. doi:10.1016/j.scitotenv.2021.151724

20. Tamerius JD, Shaman J, Alonso WJ, Bloom-Feshbach K, Uejio CK, Comrie A, et al. Environmental

Predictors of Seasonal Influenza Epidemics across Temperate and Tropical Climates. Riley S,

editor. PLoS Pathog. 2013;9: e1003194. doi:10.1371/journal.ppat.1003194

21. Chong KC, Lee TC, Bialasiewicz S, Chen J, Smith DW, Choy WSC, et al. Association between

meteorological variations and activities of influenza A and B across different climate zones: a multi-

region modelling analysis across the globe. J Infect. 2020;80: 84–98. doi:10.1016/j.jinf.2019.09.013

22. Chan PKS, Mok H y., Lee T c., Chu IMT, Lam W-Y, Sung JJY. Seasonal influenza activity in Hong

Kong and its association with meteorological variations. Journal of Medical Virology. 2009;81:

1797–1806. doi:10.1002/jmv.21551

23. Domenech de Cellès M, Arduin H, Lévy-Bruhl D, Georges S, Souty C, Guillemot D, et al.

Unraveling the seasonal epidemiology of pneumococcus. Proceedings of the National Academy of

Sciences. 2019;116: 1802–1807. doi:10.1073/pnas.1812388116

24.   Rubin IN, Bushman M, Lipsitch M, Hanage WP. Seasonal forcing and waning immunity drive the

sub-annual periodicity of the COVID-19 epidemic. medRxiv. 2025; 2025.03.05.25323464.

doi:10.1101/2025.03.05.25323464

25.   Eclerová V, Sen D, Přibylová L. Seasonal synchronization and unpredictability in epidemic models

with waning immunity and healthcare thresholds. Sci Rep. 2025;15: 17190. doi:10.1038/s41598-

025-01467-4

26.   Nguyen HTH, Rohani P. Noise, nonlinearity and seasonality: the epidemics of whooping cough

revisited. J R Soc Interface. 2008;5: 403–413. doi:10.1098/rsif.2007.1168

27.   Bjørnstad O. Stochastics. In: Bjørnstad ON, editor. Epidemics: Models and Data Using R. Cham:

Springer International Publishing; 2023. pp. 165–184. doi:10.1007/978-3-031-12056-5_9

28.   Bjørnstad ON, Grenfell BT. Noisy Clockwork: Time Series Analysis of Population Fluctuations in

Animals. Science. 2001;293: 638–643. doi:10.1126/science.1062226

29.   Grenfell BT, Bjørnstad ON, Finkenstädt BF. Dynamics of Measles Epidemics: Scaling Noise,

Determinism, and Predictability with the Tsir Model. Ecological Monographs. 2002;72: 185–202.

doi:10.1890/0012-9615(2002)072%5B0185:DOMESN%5D2.0.CO;2

30.   Dushoff J, Plotkin JB, Levin SA, Earn DJD. Dynamical resonance can account for seasonality of

influenza epidemics. Proc Natl Acad Sci USA. 2004;101: 16915–16916.

doi:10.1073/pnas.0407293101

31.   Yang F, Servadio JL, Thanh NTL, Lam HM, Choisy M, Thai PQ, et al. A combination of annual

and nonannual forces drive respiratory disease in the tropics. BMJ Glob Health. 2023;8: e013054.

doi:10.1136/bmjgh-2023-013054

32.   Todd S, Huong NTC, Thanh NTL, Vy NHT, Hung NT, Thao TTN, et al. Primary care influenza-

589   like illness surveillance in Ho Chi Minh City, Vietnam 2013-2015. Influenza Other Respi Viruses.

590   2018;12: 623–631. doi:10.1111/irv.12574

591   33.   Lam HM, Wesolowski A, Hung NT, Nguyen TD, Nhat NTD, Todd S, et al. Nonannual seasonality

592   of influenza-like illness in a tropical urban setting. Influenza Resp Viruses. 2018;12: 742–754.

593   doi:10.1111/irv.12595

594   34.   Ionides EL, Bretó C, King AA. Inference for nonlinear dynamical systems. Proc Natl Acad Sci U S

595   A. 2006;103: 18438–18443. doi:10.1073/pnas.0603181103

596   35.   King AA, Nguyen D, Ionides EL. Statistical Inference for Partially Observed Markov Processes via

597   the *R* Package **pomp**. J Stat Soft. 2016;69. doi:10.18637/jss.v069.i12

598   36.   Servadio JL, Thai PQ, Choisy M, Boni MF. Repeatability and timing of tropical influenza

599   epidemics. Epidemiology; 2022 Nov. doi:10.1101/2022.11.04.22281944

600   37.   Yuan H, Kramer SC, Lau EHY, Cowling BJ, Yang W. Modeling influenza seasonality in the tropics

601   and subtropics. PLOS Computational Biology. 2021;17: e1009050.

602   doi:10.1371/journal.pcbi.1009050

603   38.   Deyle ER, Maher MC, Hernandez RD, Basu S, Sugihara G. Global environmental drivers of

604   influenza. Proc Natl Acad Sci USA. 2016;113: 13081–13086. doi:10.1073/pnas.1607747113

605   39.   Mahmud AS, Martinez PP, Baker RE. The impact of current and future climates on spatiotemporal

606   dynamics of influenza in a tropical setting. PNAS Nexus. 2023;2: pgad307.

607   doi:10.1093/pnasnexus/pgad307

40. Parino F, Gustani-Buss E, Bedford T, Suchard MA, Trovão NS, Rambaut A, et al. Integrating

dynamical modeling and phylogeographic inference to characterize global influenza circulation.

2024. doi:10.1101/2024.03.14.24303719

41. Hirve S, Newman LP, Paget J, Azziz-Baumgartner E, Fitzner J, Bhat N, et al. Influenza Seasonality

in the Tropics and Subtropics – When to Vaccinate? Goldstein E, editor. PLoS ONE. 2016;11:

e0153003. doi:10.1371/journal.pone.0153003

42. Kramer SC, Shaman J. Development and validation of influenza forecasting for 64 temperate and

tropical countries. Funk S, editor. PLoS Comput Biol. 2019;15: e1006742.

doi:10.1371/journal.pcbi.1006742

43. Zhang B, Huang W, Pei S, Zeng J, Shen W, Wang D, et al. Mechanisms for the circulation of

influenza A(H3N2) in China: A spatiotemporal modelling study. Viboud C, editor. PLoS Pathog.

2022;18: e1011046. doi:10.1371/journal.ppat.1011046

44. Lowen AC, Mubareka S, Steel J, Palese P. Influenza Virus Transmission Is Dependent on Relative

Humidity and Temperature. Baric RS, editor. PLoS Pathog. 2007;3: e151.

doi:10.1371/journal.ppat.0030151

45. Servadio J, Boni, M. Is there a canonical model for tropical influenza? Scientific Reports. 2026.

46. National, Regional, and State Level Outpatient Illness and Viral Surveillance. [cited 21 May 2024].

Available: https://gis.cdc.gov/grasp/fluview/fluportaldashboard.html

47. Nickbakhsh S, Mair C, Matthews L, Reeve R, Johnson PCD, Thorburn F, et al. Virus–virus

interactions impact the population dynamics of influenza and the common cold. Proc Natl Acad Sci

USA. 2019;116: 27142–27150. doi:10.1073/pnas.1911083116

48.   Sparks AH. nasapower: A NASA POWER Global Meteorology, Surface Solar Energy and

630   Climatology Data Client for R. Journal of Open Source Software. 2018;3: 1035.

631   doi:10.21105/joss.01035

632   49.   He D, Ionides EL, King AA. Plug-and-play inference for disease dynamics: measles in large and

633   small populations as a case study. J R Soc Interface. 2010;7: 271–283. doi:10.1098/rsif.2009.0151

634   50.   Yang W, Olson DR, Shaman J. Forecasting Influenza Outbreaks in Boroughs and Neighborhoods of

635   New York City. Lessler J, editor. PLoS Comput Biol. 2016;12: e1005201.

636   doi:10.1371/journal.pcbi.1005201

637   51.   Ionides EL, Nguyen D, Atchadé Y, Stoev S, King AA. Inference for dynamic and latent variable

638   models via iterated, perturbed Bayes maps. Proceedings of the National Academy of Sciences.

639   2015;112: 719–724. doi:10.1073/pnas.1410597112

640   52.   Ionides EL, Breto C, Park J, Smith RA, King AA. Monte Carlo profile confidence intervals for

641   dynamic systems. J R Soc Interface. 2017;14: 20170126. doi:10.1098/rsif.2017.0126

650 Figure 1. (A) The 7-day smoothed daily ILI incidence from 2010 to 2019 (A). (B) Discrete Fourier

651 transform of the cycles from 100 to 450 days in the ILI data. The two cycles (215 days and 365 days) with

652 the strongest spectral density are labeled with a circle.

Figure 2. Simulated trajectories from all seven models. The median (red) from 5000 simulated trajectories

is plotted with data (black) along with 95% simulation ranges (grey).

665　Figure 3. Power spectra from model simulations compared with data. The dark grey vertical bars show the

666　logarithm-transformed spectral density from the data. The mean and the 95% simulation interval of the

667　power spectrum from seven models are shown as black line and blue ribbon. Red dashed vertical lines

668　label cycles of 215 days and 365 days.

675 Figure 4. The effect of specific humidity, temperature, and school term on R_0 from the selected model.

676 (A) The relationship between specific humidity and R_0 under varying temperatures and school term

677 conditions. (B) The relationship between temperature and R_0 under varying specific humidity and school

678 term conditions. Triangles show values when school is in session, while circles show schools holidays.

Figure 5. Power spectra of the stochastic SIRS U-shaped SH temperature model with school term and its

deterministic skeleton. The power spectrum from the ILI data is shown as grey bars. The mean and the

95% simulation interval of the power spectrum is plotted as red and blue for the stochastic and

deterministic version of the model, respectively. The red vertical lines are the cycles at 215 days and 365

days.